



NATIONAL LIBRARY OF MEDICINE



NLM 00105585 0











# MEDICAL DIAGNOSIS

WITH

167  
SPECIAL REFERENCE TO PRACTICAL MEDICINE..

A


GUIDE TO THE KNOWLEDGE AND DISCRIMINATION OF DISEASES.

BY

J. M. DA COSTA, M.D., LL.D.,

PRESIDENT OF THE COLLEGE OF PHYSICIANS OF PHILADELPHIA; EMERITUS PROFESSOR OF PRACTICE OF MEDICINE AND OF CLINICAL MEDICINE AT THE JEFFERSON MEDICAL COLLEGE, PHILADELPHIA; PHYSICIAN TO THE PENNSYLVANIA HOSPITAL, ETC.

Illustrated with Engravings on Wood.



---

EIGHTH EDITION, REVISED.

---

PHILADELPHIA:

J. B. LIPPINCOTT COMPANY.

LONDON: 10 HENRIETTA STREET, COVENT GARDEN.

1895.

Annet

WBB

D118m

1895

Entered, according to Act of Congress, in the year 1890, by

J. M. DA COSTA, M.D.,

In the Office of the Librarian of Congress at Washington.

---

Copyright, 1895, by J. M. DA COSTA, M.D.



## EXTRACT FROM PREFACE TO THE FIRST EDITION.

---

MY chief aim in writing this work has been to furnish advanced students and young graduates of medicine with a guide that might be of service to them in their endeavors to discriminate disease. I have sought to offer to those members of the profession who are about to enter on its practical duties a book on Diagnosis of an essentially practical character,—one neither so meagre in detail as to be next to useless when they encounter the manifold and varying features of disease, nor so overladen with unnecessary detail as to be unwieldy and lacking in precise and readily-applicable knowledge.

In executing my undertaking, two plans offered themselves: either to describe morbid states in compliance with the usual pathological classification followed in treatises on the Practice of Medicine, or to group them according to their marked symptoms. The former plan would have been far the easier, but the latter seemed to me the more suitable for a volume of this kind; and, although it has involved much labor, and has rendered the task much more difficult of accomplishment, its advantages appeared to me so great that I have adopted it throughout. That this attempt at a purely clinical classification is not perfect, I am fully aware. But, with all its shortcomings, I venture to hope that it will not be devoid of value.

Some of the statements made may appear too absolute, and as not taking sufficient notice of the many exceptions which may arise. But it was impossible to avoid this without very lengthy discussion: and even in the lengthiest discussion all exceptions and all possible points of fallacy would not have been mentioned; for Nature does not limit herself in her irregularities any more than in her rules. The text must, therefore, be looked upon as treating only of general laws and of their most notable infractions; in fact, but as a series of etchings, with here and there a prominent figure shaded, but not as an attempt to reproduce the

colors of an original whose varied hues could not be closely copied, even by the hand of a master.

The main object of this work is, what its title implies, the consideration of Medical Diagnosis. In connection with this, I have endeavored to take cognizance of the prognosis of individual affections, and occasionally the record of cases has been introduced by way of elucidation. To have done this to a much greater extent, though in some respects desirable, would have swelled the work to an inordinate size.

The wood-cuts employed as illustrations are all original. Many are from sketches, or at least are based on sketches, taken directly from cases of interest.

PHILADELPHIA, April, 1864.

---

## PREFACE TO THE EIGHTH EDITION.

---

ANOTHER edition having been demanded, I have revised the work and altered some of its chapters. New matter has been inserted, old matter in parts condensed. I have endeavored to incorporate whatever of bacteriological interest appeared to be established as valuable for diagnostic purposes. The introduction of a number of additional wood-cuts and of temperature charts taken from cases actually observed will, I trust, conduce to greater clearness and accuracy. I gladly acknowledge the efficient aid of Dr. Eshner, Dr. Woodbury, and Dr. Leffmann in preparing this edition for the press. Since the work was last published, my attention has been called to a further Italian translation.

1700 WALNUT STREET,  
August, 1895.



33756

# CONTENTS.

## INTRODUCTION.

	PAGE
GENERAL CONSIDERATIONS . . . . .	17

## CHAPTER I.

### EXAMINATION OF PATIENTS, AND SOME SYMPTOMS OF GENERAL IMPORT.

General Considerations . . . . .	27
Position of the Body . . . . .	30
General Aspect—Expression of Countenance . . . . .	32
Skin . . . . .	34
Pulse . . . . .	35
Temperature of the Body . . . . .	43
Tongue . . . . .	52
Sensations of Patients . . . . .	55

## CHAPTER II.

### DISEASES OF THE BRAIN AND SPINAL CORD, AND OF THEIR NERVES.

General Considerations . . . . .	57
Cerebral Localization . . . . .	57
Sensory Centres, and Conducting Paths . . . . .	63
Spinal Localization . . . . .	65
Deranged Intellection . . . . .	67
Delirium . . . . .	67
Stupor . . . . .	70
Coma . . . . .	70
Insomnia . . . . .	72
Deranged Sensation . . . . .	72
Hyperæsthesia . . . . .	72
Anæsthesia . . . . .	74
Headache . . . . .	81
Vertigo . . . . .	83
Derangement of Special Senses . . . . .	87
Vision . . . . .	87
Hearing . . . . .	98
Deranged Reflexes . . . . .	99



	PAGE
Deranged Motion . . . . .	103
Paralysis . . . . .	103
Hemiplegia . . . . .	113
Monoplegia . . . . .	120
Paraplegia . . . . .	124
Sudden Paraplegia . . . . .	125
Spinal Hemorrhage . . . . .	125
Acute Ascending Paralysis . . . . .	126
Multiple Neuritis . . . . .	128
Gradual Paraplegia . . . . .	130
Spinal Congestion . . . . .	131
Spinal Anæmia . . . . .	131
Spinal Meningitis . . . . .	131
Myelitis . . . . .	132
Spinal Scleroses . . . . .	135
Tumors of the Cord . . . . .	136
Reflex Paraplegia . . . . .	137
Palsies usually Limited, though they may be General . . . . .	138
Hysterical Paralysis . . . . .	138
Rheumatic Paralysis . . . . .	140
Lead Palsy . . . . .	141
Diphtheritic Paralysis . . . . .	141
Syphilitic Paralysis . . . . .	142
Local Palsies . . . . .	144
Facial Palsy . . . . .	144
Paralysis of the Nerves of the Arm . . . . .	146
Bulbar Paralysis . . . . .	147
Palsies connected with Marked Muscular Wasting . . . . .	149
Progressive Muscular Atrophy . . . . .	149
Infantile Paralysis . . . . .	155
Ataxia . . . . .	159
Locomotor Ataxia . . . . .	159
Diseases of the Cerebellum . . . . .	165
Tremor . . . . .	166
Paralysis Agitans . . . . .	167
Multiple Cerebro-spinal Sclerosis . . . . .	167
Functional Tremors . . . . .	169
Spasms—Convulsions . . . . .	171
Deranged Nutrition and Secretion . . . . .	174
Acute Affections of which Delirium is a Prominent Symptom . . . . .	179
Acute Meningitis . . . . .	179
Tubercular Meningitis . . . . .	185
Cerebro-spinal Meningitis . . . . .	189
Delirium Tremens . . . . .	190
Acute Mania . . . . .	192
Diseases Marked by Sudden Loss of Consciousness and of Voluntary Motion . . . . .	194
Apoplexy . . . . .	194



# CONTENTS.

7

	PAGE
Sun-stroke . . . . .	210
Catalepsy . . . . .	212
Diseases marked by Convulsions or Spasms . . . . .	214
Epilepsy . . . . .	214
Chorea . . . . .	219
Hysteria . . . . .	224
Tetanus . . . . .	228
Functional Spasms . . . . .	233
Diseases characterized by Gradual Impairment of the Mental Faculties	
with Paralysis . . . . .	235
Chronic Softening . . . . .	235
Tumor . . . . .	240
General Paralysis . . . . .	246
Diseases characterized by Enlargement of the Head . . . . .	249
Chronic Hydrocephalus . . . . .	249
Hypertrophy of the Brain . . . . .	250
Diseases characterized by Enlargement of Various Parts . . . . .	251
Acromegaly . . . . .	251
Diseases characterized by Paroxysmal Pain . . . . .	252
Neuralgia in General . . . . .	252
Facial Neuralgia . . . . .	254
Hemicrania . . . . .	255
Sciatica . . . . .	256

## CHAPTER III.

### DISEASES OF THE UPPER AIR-PASSAGES.

#### SECTION I.

##### DISEASES OF THE NOSE AND ASSOCIATE ORGANS.

General Considerations . . . . .	260
----------------------------------	-----

#### SECTION II.

##### DISEASES OF THE LARYNX AND TRACHEA.

General Considerations . . . . .	266
Acute Laryngeal Affections . . . . .	274
Acute Laryngitis . . . . .	274
Edema of the Larynx . . . . .	277
Croup . . . . .	278
Chronic Laryngeal Affections . . . . .	286
Chronic Laryngitis . . . . .	286

## CHAPTER IV.

### DISEASES OF THE CHEST.

General Considerations . . . . .	297
----------------------------------	-----

## SECTION I.

## DISEASES OF THE LUNGS.

	PAGE
Different Methods of Physical Diagnosis, and the Physical Signs of	
Pulmonary Diseases . . . . .	299
Inspection . . . . .	299
Mensuration . . . . .	300
Palpation . . . . .	303
Percussion . . . . .	303
Percussion of the Healthy Chest . . . . .	309
Auscultation . . . . .	310
Sounds of Respiration in Health and in Disease . . . . .	313
Changes in the Vesicular Murmur . . . . .	314
Bronchial Respiration . . . . .	318
New or Adventitious Sounds . . . . .	320
Auscultation of the Voice . . . . .	325
Combination of the Physical Signs, and the Examination of Patients	
affected with Disease of the Lungs . . . . .	326
Principal Symptoms of Diseases of the Lungs . . . . .	330
Dyspnœa . . . . .	330
Cough . . . . .	335
The Sputum . . . . .	338
Hæmoptysis . . . . .	345
Diseases in which Clearness on Percussion is met with . . . . .	348
Acute Bronchitis . . . . .	348
Chronic Bronchitis . . . . .	352
Emphysema . . . . .	356
Diseases in which Dulness on Percussion occurs . . . . .	361
Phthisis . . . . .	361
Acute Affections of the Lungs accompanied by Dulness on Percussion . . . . .	384
Acute Phthisis . . . . .	384
Acute Pneumonia . . . . .	388
Acute Pleurisy . . . . .	404
Diseases presenting Dilatation of the Chest, Displacement of the Liver or	
Heart, and Dyspnœa . . . . .	410
Pneumothorax . . . . .	411
Chronic Pleurisy . . . . .	415
Diseases in which Retraction of the Chest occurs . . . . .	423
Chronic Pleurisy . . . . .	423

## SECTION II.

## DISEASES OF THE HEART.

General Considerations . . . . .	428
Examination of the Heart by the Different Methods of Physical Diagnosis	431
Inspection . . . . .	431
Palpation . . . . .	432
Percussion . . . . .	433
Auscultation . . . . .	435

# CONTENTS.

9

	PAGE
General and Local Symptoms of Diseases of the Heart . . . . .	445
Cardiac Dropsy . . . . .	446
Derangement of the Circulation . . . . .	446
Cardiac Pain . . . . .	448
Palpitation . . . . .	452
Functional Disorders of the Heart . . . . .	453
Disorders characterized by Palpitation, associated or not with	
Change of Rhythm . . . . .	454
Organic Diseases of the Heart . . . . .	460
Acute Diseases presenting Pain in the Cardiac Region; the Sym-	
ptoms of a Disturbed Circulation; and a Change in the	
Sounds of the Heart, or their Replacement by Murmurs .	460
Acute Endocarditis . . . . .	461
Acute Pericarditis . . . . .	468
Myocarditis . . . . .	476
Chronic Diseases attended with Increased Extent of Percussion	
Dulness, but with Normal or almost Normal Heart-Sounds	478
Hypertrophy . . . . .	478
Dilatation . . . . .	482
Fatty Degeneration . . . . .	484
Pericardial Effusion . . . . .	488
Diseases of the Heart exhibiting more or less of the Signs and	
Symptoms of Enlargement of the Organ, and accompanied	
by Endocardial Murmurs . . . . .	489
Valvular Affections . . . . .	489
Displacements of the Heart . . . . .	505

## SECTION III.

THORACIC ANEURISM . . . . .	506
-----------------------------	-----

## CHAPTER V.

### DISEASES OF THE MOUTH, PHARYNX, AND ŒSOPHAGUS.

Mouth . . . . .	519
Fauces . . . . .	522
Tonsillitis . . . . .	522
Diphtheria . . . . .	524
Mumps . . . . .	534
Chronic Sore Throat . . . . .	535
Pharynx and Œsophagus . . . . .	537

## CHAPTER VI.

### DISEASES OF THE ABDOMEN.

General Considerations . . . . .	541
Methods and General Results of Physical Examination of the Abdomen	542
Inspection . . . . .	542
Palpation . . . . .	543
Percussion . . . . .	544
Auscultation . . . . .	550



## SECTION I.

## DISEASES OF THE STOMACH.

	PAGE
General Considerations . . . . .	550
Loss of Appetite . . . . .	555
Excessive Acidity of the Stomach . . . . .	556
Flatulency . . . . .	557
Nausea and Vomiting . . . . .	558
Pain . . . . .	565
Diseases of the Stomach in which Pain and Soreness at the Epigastrium, and Vomiting, occur . . . . .	571
Acute Gastritis . . . . .	571
Chronic Diseases attended with Pain, Epigastric Tenderness, and Vomiting	574
Chronic Gastritis . . . . .	575
Gastric Ulcer . . . . .	576
Gastric Cancer . . . . .	582
Dilatation of the Stomach . . . . .	590

## SECTION II.

## DISEASES OF THE INTESTINES AND OF THE PERITONEUM.

General Considerations . . . . .	593
Alvine Discharges . . . . .	593
Diseases attended with Paroxysms of Pain referred chiefly to the Middle or Lower Part of the Abdomen, and not associated with Marked Tenderness or with Fever . . . . .	596
Colic . . . . .	596
Diseases attended with Pain and Marked Tenderness in the Umbilical Region or diffused over the Abdomen . . . . .	606
Acute Enteritis . . . . .	606
Acute Peritonitis . . . . .	609
Chronic Peritonitis . . . . .	621
Diseases attended with Pain and Tenderness in the Right Iliac Fossa . .	622
Appendicitis . . . . .	622
Disorders attended with Constipation, and of which it is a Prominent Symptom . . . . .	630
Intestinal Obstruction . . . . .	630
Habitual Constipation . . . . .	641
Disorders in which Morbid Discharges from the Bowels occur . . . . .	643
Diarrhoea . . . . .	643
Dysentery . . . . .	648
Intestinal Hemorrhage, or Melæna . . . . .	651
Fatty Diarrhoea . . . . .	653
Diseases attended with Vomiting and Purgings . . . . .	653
Cholera Infantum . . . . .	654
Cholera Morbus . . . . .	655
Cholera . . . . .	656

## SECTION III.

## DISEASES OF THE LIVER.

	PAGE
General Considerations . . . . .	661
Jaundice . . . . .	662
Acute Diseases of the Liver attended generally with Slight Enlargement of the Organ, and with more or less, though rarely very much	
Jaundice . . . . .	667
Acute Congestion . . . . .	667
Acute Hepatitis . . . . .	667
Inflammation of the Gall-Bladder and Gall-Ducts . . . . .	673
Acute Diseases characterized by a Decrease in the Size of the Liver and by Deep Jaundice . . . . .	676
Acute Yellow Atrophy . . . . .	676
Chronic Diseases attended with Enlargement of the Liver, and with slight or no Jaundice . . . . .	679
Chronic Congestion . . . . .	679
Chronic Hepatitis . . . . .	681
Abscess of the Liver . . . . .	682
Fatty Liver . . . . .	688
Waxy Liver . . . . .	688
Cancer of the Liver . . . . .	690
Hydatids of the Liver . . . . .	697
Chronic Diseases attended with Decreased Size of the Liver, and with	
Abdominal Dropsy . . . . .	702
Cirrhosis . . . . .	702
Chronic Atrophy of the Liver . . . . .	709

## SECTION IV.

## ABDOMINAL ENLARGEMENT.

General Abdominal Enlargement . . . . .	710
Ascites . . . . .	710
Chronic Tympanites . . . . .	717
Partial Abdominal Enlargement . . . . .	718
Abdominal Tumors . . . . .	718

## SECTION V.

## ABDOMINAL PULSATION.

Aortic Pulsation . . . . .	731
Abdominal Aneurism . . . . .	732

## CHAPTER VII.

## ON THE URINE, AND ON DISEASES OF THE URINARY ORGANS.

	PAGE
Urine . . . . .	736
Color . . . . .	740
Specific Gravity . . . . .	740
Reaction . . . . .	742
Changes in the Quantity of the more Important Constituents . . .	744
Presence of Abnormal Substances in the Urine . . . . .	757
Sediments . . . . .	782
Urinary Organs . . . . .	784
Diseases of the Kidney of which Pain is a Prominent Symptom .	784
Nephritis . . . . .	784
Nephralgia . . . . .	785
Diseases marked by an Albuminous Condition of the Urine, associated with more or less Dropsy . . . . .	791
Acute Bright's Disease . . . . .	791
Chronic Bright's Disease . . . . .	798
Diseases associated with Purulent Urine . . . . .	814
Acute Cystitis . . . . .	815
Chronic Cystitis . . . . .	816
Abscess of the Kidney . . . . .	816
Pyelitis . . . . .	819
Disorders in which a very large Amount of Urine is discharged . . . .	823
Diabetes . . . . .	823
Chronic Diuresis . . . . .	826
Disorders in which little or no Urine is discharged . . . . .	828
Suppression of Urine . . . . .	828
Retention of Urine . . . . .	829

## CHAPTER VIII.

## DROPSY.

Dropsy, according to its Seat and Extent . . . . .	830
Dropsy, according to its Causation . . . . .	832
Dropsy, according to the Rapidity of its Development . . . . .	835

## CHAPTER IX.

## DISEASES OF THE BLOOD-VESSELS.

Diseases of the Arteries . . . . .	836
Arteritis . . . . .	836
Atheromatous Changes . . . . .	838



	PAGE
Diseases of the Veins . . . . .	839
Phlebitis . . . . .	839
Diseases of the Capillaries . . . . .	840

## CHAPTER X.

## DISEASES OF THE BLOOD.

General Considerations . . . . .	842
Anæmia . . . . .	855
Pernicious Anæmia . . . . .	857
Leukæmia . . . . .	863
Addison's Disease . . . . .	869
Pyæmia . . . . .	872
Septicæmia . . . . .	876
Thrombosis and Embolism . . . . .	877
Scurvy . . . . .	884
Purpura . . . . .	886

## CHAPTER XI.

## RHEUMATISM AND GOUT.

Acute Rheumatism . . . . .	889
Chronic Rheumatism . . . . .	894
Gout . . . . .	898
Rheumatic Arthritis or Rheumatic Gout . . . . .	900
Rickets . . . . .	902

## CHAPTER XII.

## FEVERS.

General Considerations . . . . .	906
Continued Fevers . . . . .	908
Simple Continued Fever . . . . .	908
Catarrhal Fever . . . . .	909
Typhoid Fever . . . . .	913
Typhus Fever . . . . .	929
Cerebro-spinal Fever . . . . .	937
Relapsing Fever . . . . .	944
Periodical Fevers . . . . .	948
Intermittent Fever . . . . .	948
Remittent Fever . . . . .	954
Congestive Fever . . . . .	963
Yellow Fever . . . . .	969

	PAGE
Eruptive Fevers . . . . .	976
Scarlet Fever . . . . .	976
Measles . . . . .	981
Rubella . . . . .	985
Smallpox . . . . .	988
Dengue . . . . .	994
Erysipelas . . . . .	996

CHAPTER XIII.

DISEASES OF THE SKIN.

General Considerations . . . . .	1000
Erythematous Diseases . . . . .	1003
Papular Diseases . . . . .	1005
Vesicular Diseases . . . . .	1007
Bullous Diseases . . . . .	1010
Pustular Diseases . . . . .	1011
Squamous Diseases . . . . .	1013
Maculæ . . . . .	1015
New Growths . . . . .	1016
Hypertrophies . . . . .	1017
Parasitic Diseases . . . . .	1020
Altered Gland-secretions . . . . .	1023
Nervous Affections . . . . .	1025

CHAPTER XIV.

POISONS AND PARASITES.

Poisons . . . . .	1027
Acute Poisoning . . . . .	1027
Irritant Poisons . . . . .	1028
Narcotic Poisoning . . . . .	1032
Chronic Poisoning . . . . .	1038
Parasites . . . . .	1048
Vegetable Parasites . . . . .	1049
Animal Parasites . . . . .	1050
INDEX . . . . .	1067

# LIST OF ILLUSTRATIONS.

FIG.	PAGE	FIG.	PAGE
1. Sphygmograph of Marey . . .	39	34. Cavities in the Lung in Phthisis	367
2. Dudgeon's Sphygmograph . . .	40	35. Temperature Chart in Pneumonia	389
3. Sphygmogram enlarged . . .	41	36. Diagram illustrative of Perfect Pulmonary Consolidation, such as happens in the Second Stage of Pneumonia . . . . .	391
4. Self-registering Thermometer . .	44	37. Temperature Chart in Broncho- Pneumonia . . . . .	397
5. Seguin's Surface Thermometer . .	44	38. Diplococcus Pneumoniæ of Fraenkel . . . . .	399
6. Surface Thermometer, with coil at extremity . . . . .	44	39. Pneumococcus (Diplococcus) of Friedländer . . . . .	400
7. Temperature Chart in Simple Continued Fever . . . . .	47	40. Roughening of the Pleura from Inflammation . . . . .	404
8. Temperature Chart from a Case of Remittent Fever . . . . .	48	41. Large Effusion occupying the Left Pleural Cavity . . . . .	405
9. The Centres in the Human Brain	60	42. Physical Signs of Pneumothorax	412
10. Right Homonymous or Lateral Hemianopsia . . . . .	62	43. Topography of the Heart . . .	429
11. The Æsthesiometer . . . . .	78	44. Diagram showing the Points at which the Separate Valves may be listened to . . . . .	436
12. Mathieu's Dynamometer . . . .	108	45. Position of the Heart, and Dis- tention of the Pericardium with Fluid, in Pericarditis . .	469
13. Laryngoscopes . . . . .	268	46. Hypertrophied Heart, lying in its Position in the Chest . . . . .	480
14. Laryngoscopic Examination . .	269	47. Dilated Heart, the Right Ventri- cle opened . . . . .	484
15. Laryngeal Image, as seen in the Laryngoscope . . . . .	270	48. Narrowing of the Aortic Orifice by Vegetations . . . . .	494
16. The Stethometer . . . . .	301	49. Insufficient Mitral Valves per- mitting Regurgitation of the Blood . . . . .	496
17. The Stetho-Goniometer . . . .	302	50. Sphygmogram of Aortic Insuffi- ciency . . . . .	498
18. The Pleximeter . . . . .	304	51. Sphygmogram of Mitral Regur- gitation . . . . .	498
19. Percussion Hammer . . . . .	306	52. Klebs-Loeffler Bacilli . . . . .	525
20. The Ordinary Stethoscope . . .	311	53. Results of Abdominal Percussion	548
21. Hawksley's Stethoscope . . . .	311	54. Sarcinæ Ventriculi . . . . .	561
22. The Double Stethoscope . . . .	312	55. Comma-Bacillus of Koch, from Culture in Blood-Serum . . .	658
23. The Differential Stethoscope . .	312	56. Doremus's Ureometer . . . . .	746
24. Diagram illustrative of the Main Forms of Feeble Respiration .	315		
25. Diagram illustrative of Rales .	321		
26. Elastic Fibres of Pulmonary Tissue . . . . .	340		
27. A Spiral Magnified . . . . .	340		
28. Charcot-Leyden Crystals . . . .	341		
29. Tubercle-Bacilli in Sputum . .	343		
30. Tubercle-Bacilli (in colors) . .	344		
31. Casts from a Case of Plastic Bron- chitis . . . . .	355		
32. Appearance of the Chest in Em- physema . . . . .	357		
33. Beginning Infiltration in Phthi- sis . . . . .	365		



FIG.	PAGE	FIG.	PAGE
57. Greene's Ureometer . . . . .	746	75. Blood in Pernicious Anæmia . .	860
58. Crystals of Uric Acid . . . . .	749	76. Temperature in a Case of In-	
59. Mixed Urates . . . . .	751	fluenza . . . . .	911
60. Earthy Phosphates in the Urine.	753	77. Temperature in Typhoid Fever .	916
61. Calcium Oxalate Crystals . . . .	759	78. Eberth-Gaffky's Typhoid-Fever	
62. Albumin Test-Glass . . . . .	769	Bacillus, from a Potato Culture	917
63. Pus-Corpuscles . . . . .	779	79. Spirilla of Relapsing Fever . .	947
64. Epithelial Casts and Cells from		80. Temperature in Tertian Inter-	
the Kidneys in a Case of Acute		mittent Fever . . . . .	951
Bright's Disease . . . . .	792	81. Temperature Chart in Remittent	
65. Fatty Casts and Epithelial Cells		Fever . . . . .	955
filled with Fat, as seen in Dis-		82. Pigment in the Blood in Malarial	
charge from a Fatty Kidney .	807	Cachexia . . . . .	960
66. Hyaline or Waxy Casts from the		83. Hæmatozoa of Malaria . . . .	962
Urine . . . . .	809	84. Temperature in Yellow Fever .	972
67. Granular Casts, or Casts covered		85. Temperature in Variola . . . .	990
with Disintegrating Epithe-		86. Temperature in Varioloid . . .	994
lium and Granules . . . . .	810	87. Temperature Chart in Facial	
68. Artificial Capillary of Malassez.		Erysipelas . . . . .	997
Magnified 100 Diameters . .	844	88. Acarus Scabiei . . . . .	1021
69. Potain's Pipette . . . . .	845	89. Segments of Tænia Solium . .	1053
70. Graduated Moist-Chamber of		90. Heads of Tæniæ . . . . .	1053
Malassez . . . . .	846	91. Trichina in Recent Human	
71. Blood-Mixture, as seen with the		Muscle . . . . .	1057
Square Micrometer Ruling of		92. Trichina Capsule with Shell-like	
the Moist-Chamber of Malassez	847	Calcareous Deposits . . . .	1058
72. Daland's Hæmatokrit . . . . .	848	93. Encapsuled Chalky Concretions	
73. Hæmoglobinometer of Gowers .	852	in Muscle, due to Trichinæ . .	1059
74. Chart showing Blood Changes in		94. Trichina Spiralis, Magnified 300	
Chlorosis . . . . .	854	times . . . . .	1061

# MEDICAL DIAGNOSIS.

---

## INTRODUCTION.

### GENERAL CONSIDERATIONS.

THE study of any complicated subject leads of necessity to its arrangement into branches. Closely connected as these are, and forming always parts of a whole, they are not only capable of distinct treatment, but frequently become more intelligible as they are so treated. This is made very manifest in investigating disease. The extent of ground covered by the inquiry has rendered it imperative to map it out into various provinces, which, however intimately united, may be with convenience separately surveyed. One comprises the laws and facts common to individual affections; in another are gathered together all relating to their causes; another embraces the consideration of their detection and the full recognition of their nature. It is the purpose of these pages to examine this department somewhat minutely, and especially that portion of it coming within the range of the practitioner of medicine. In so doing it will become apparent how *diagnosis*, for such the distinction of disease is technically called, is partly a science, partly an art; a science, because it comprehensively takes account of general facts, and of principles based on those facts; an art, because it demands a cognizance of the means, and their application to arrive at the desired result.

To consider, then, medical diagnosis in all its bearings, it will be necessary not only to hold up to view the morbid states met with in the examination of the sick, but also to inquire in what manner they may be most readily recognized and explored, and how their differences may be made available in the discrimination

of one ailment from another. In a study of this kind, an investigation of symptoms plays unavoidably a prominent part. In truth, the detection of disease is the product of close observation of symptoms, and of correct deduction from these symptoms.

The first requirement therefore for an accurate diagnosis is to learn to recognize morbid signs. But the art of observation this implies is not easy, and cannot be thoroughly acquired except by practice. No one aspiring to become a skilful observer can trust exclusively to the light reflected from the writings of others: he must carry the torch in his own hands, and himself look into every recess. The knowledge obtained from reading is, however, serviceable in this way: it aids in overcoming one of the main difficulties at first experienced,—to know where to look and what to look for. There are in almost every affection some symptoms which can hardly escape the merest beginner; but also some which do not appear on the surface, and which to find taxes the skill of the experienced physician. And it is especially in this search after hidden signs that medical information as well as cultivated tact is demanded.

Now, to recognize the manifestations of disease, whether they are or are not readily perceptible, we have to employ our eyes and ears, our sense of touch and of smell. Formerly we could go no further than these senses unassisted would carry us. But science has lent its aid, and furnished means by the help of which we can detect clearly what before we could not detect at all, or of which at best we caught only a glimpse. We now possess instruments of precision by which we ascertain with accuracy the size of organs and their play. With thermometers we tell the degree of heat of various parts of the body. Specific-gravity bottles, and other measures devised for the purpose, inform us of the relative gravity of fluids. The microscope gives at a glance insight into matters which the naked eye fails even to perceive, shows us crystals in secretions, enables us to count the corpuscles in the blood, and to detect minute and disease-causing specific organisms. The laryngoscope demonstrates the appearance and the movements of the organ of speech. The ophthalmoscope informs us of the state of the vessels in the brain. And chemistry, with its marvellous teachings, is rendering our knowledge of many morbid states amazingly complete. Then the sagacity



of comparatively modern times has taught us to enlist the sense of hearing, and has demonstrated how a disciplined ear may detect the workings of disease in cavities into which the eye cannot penetrate. The effect of all these improved methods of study has been to give an immense impetus to clinical research, and to lead to the construction of a solid groundwork of experience in striking contrast with the looseness of former times. The advance in diagnosis thus attained forms, indeed, one of the most pleasing portions of medical history.

When, by means of the aided or unaided senses, the symptoms of the malady have been discovered, the next step toward a diagnosis is a proper appreciation of their significance and of their relation toward one another. Knowledge and, above all, the exercise of the reasoning faculties are now indispensable. The daily habit of investigating disease; a scrutinizing study of the anatomical lesions; chemistry, with its most searching analyses; the microscope, with the wonders it reveals,—are all of little use, unless we have been taught the necessity of placing in connection with one another the morbid signs they lay bare, and of considering in individual cases their respective value. Were it otherwise, the science of diagnosis would be simply a matter of memory. It is, however, this very analysis of symptoms and the lengthy process of induction attending it which make medical diagnosis so difficult and so unattractive to the beginner. He sees that by reflecting and reasoning on what are frequently but indirect manifestations he must find the seat and nature of disorders hidden from his view. Nor is it reasoning on the ascertained facts alone that is required; the premises may be but probabilities; for, in truth, diagnosis deals at times with the logic of probabilities as much as with the logic of patent facts.

Now, we are greatly aided in appreciating morbid signs, and in interpreting them correctly, by already existing knowledge. We look to landmarks which our predecessors have erected, and the gradually accumulated science of semeiology, rightly employed, furnishes the clue to the discovery of the disease. Thus the stores which medicine has laboriously collected during centuries can be used with advantage by all, and exist for the good of all.

But an acquaintance with semeiology is far from being the sole guide to diagnosis, nor does it at once help to a recognition of the

malady. There are few symptoms in themselves distinctive ; and often a symptom may be due to one of several causes. Semeiology informs us of these different causes ; but to find out the precise meaning of the abnormal manifestation in an individual case, we have to draw our inference from all the signs encountered ; to compare them with one another ; to seek out those that are in the background. We are thus arriving, step by step, at the explanation of the morbid appearances, the starting-point in deduction always being what is known of the affection the presence of which is suspected, and the symptoms of which we are contrasting with those before us. For the conclusion to be valid and exact, it is of course requisite that each part of the testimony have the proper position assigned to it. In reasoning correctly on symptoms, the same laws apply as in reasoning correctly on any other class of phenomena : the facts have to be sifted and weighed, not merely indiscriminately collected. And while this intellectual act is being performed, much collateral evidence is to be sought before a final judgment is given ; especially is it necessary to view the symptoms with constant reference to the age, sex, and habits of the patient, and to the circumstances amid which the disorder develops.

To accomplish all this effectually, the physician has need of much and varied knowledge. He must be master of something more than of the information supplied to him by semeiology. He must be an anatomist to pronounce with certainty on the seat of the malady ; a physiologist to appreciate the state of the great centres and the aberration of function. Above all, he must be a pathologist in the full sense of the term : he must understand the antagonism between diseases ; the frequency with which they coexist ; the influence of remedial agents on them ; and be cognizant of their natural history and of the general laws governing them,—for how else can he form an estimate of morbid action while in progress ? Then it is desirable that he should be aware of what are their current divisions and classifications. From what has already been represented, it is evident that he must also be a correct reasoner ; for even a good observer will, by bad reasoning, arrive at a faulty diagnosis ; just as sometimes a bad observer may, by the same process, blunder into the truth. There is, indeed, no end to the extent of knowledge which may be brought to bear in working out a conclusion regarding the char-

acter and seat of a malady. The habit of observation once acquired, information of the most varied kind will, by an accurate reasoner, be made tributary to the completeness of the diagnosis. Every fresh acquirement tends to enlarge our powers of insight. Just as in nature, the higher we ascend, the more fully lies the view before us.

Having thus indicated the elements of a thorough diagnosis, we may next inquire in what way this is most easily arrived at when at the bedside. The main facts of the case on which the deductions are to be based are of course first elicited. We lay hold of these main facts, and especially of those that are the most direct signs of the morbid action. They are coupled together, and the inquiry is started as to what organ they indicate as the seat of the malady. This often has been already determined by the very method of the examination; and we therefore proceed at once to investigate the precise nature of the disorder by analyzing the symptoms and the previous history. Sometimes, however, the site of the disease does not admit of being definitely fixed upon, or we can only in a general manner decide upon the function impaired. Again, as in idiopathic fevers, we may find no signs of local disease,—merely those of a general disturbance. In any of these instances clinical experience steps in to explain the phenomena as far as possible, and to inform us in what affections they occur. It may be only in one; then the desired goal is at once attained. But, as above stated, there are few signs in themselves pathognomonic. It is therefore to be ascertained which one of the disorders is before us that special pathology teaches may yield the symptoms encountered. One of these is taken up. Its symptoms are placed side by side with those present. They accord in some respects, but not in all. Moreover, in searching for some of the phenomena which the supposed malady gives rise to, these are not found. The view is abandoned, and another taken up. It agrees in all particulars. The diagnosis is made. Yet, when the diagnosis is thus arrived at, we have still to determine, before it can be considered as complete and can be acted upon, whether or not any other morbid state exists, and to take into account the patient's general condition and his individuality.

To cite a case in illustration. A person consults us for a cough



brought on by exposure. He has been ill for four or five days, having been previously in good health. We notice, on examining him, that his breathing is hurried, and that he has fever; the lower portion of one side of the chest is dull on percussion, and the respiration there is wanting; the action and sounds of the heart are normal. The facts point to the lung or its covering as the seat of the malady. We know, furthermore, from the history and the febrile symptoms, that we have to deal with an acute affection. What are the acute pulmonary affections? Acute bronchitis; acute phthisis; acute pleurisy; acute pneumonia. In all occur fever, cough, and disordered breathing. Is it acute pneumonia? No; for, notwithstanding there is in this complaint, in addition to the general symptoms mentioned, dulness on percussion, the dulness is associated with a blowing respiration; whereas in the case before us no respiration is heard. Let us look at the sputum, and see if it be tenacious and rusty-colored. It is not; it is thin and frothy. But acute pleurisy may explain all the signs. The patient, too, when questioned, states that he had at the onset a sharp pain in his side; and this, we are aware, takes place in pleurisy. The vocal vibrations, likewise, are noticed to be absent on the affected side of the chest, which, when measured, is found to be enlarged. This corresponds in all points with what happens in pleurisy in the stage of effusion. The disease *is*, therefore, acute pleurisy in the stage of effusion. We finish the diagnosis by ascertaining the existence or non-existence of other maladies, and by taking note of the severity of the complaint; that it has occurred in a young and robust person of good habits; and that the symptomatic fever is very active.

This process of arriving at an opinion is the simplest. It is one in which the investigation of the case is to some extent carried on while the deductions are being made. And it is astonishing how rapidly it may be performed by habit. The mind works unconsciously, and a decision is, to all appearance, formed intuitively, which surprises the inexperienced by its readiness and precision. This method aims, so far as the symptoms permit, at a direct diagnosis. But, in truth, it is often what is called *differential*: that is, it takes cognizance of and dwells on the essential signs by which one disease can be discriminated from another resembling it.



Sometimes, instead of attaining the desired result in the manner proposed, we are obliged to judge of the nature of the malady entirely by finding out what it is not. The various diseases capable of producing all, or even some, of the striking symptoms observed, are enumerated. They are one by one considered and set aside, until by this process of pure exclusion the mischief is brought to light. Thus, to use again the example just given, we should have to assign reasons why the disease is neither acute pneumonia, nor bronchitis, nor acute phthisis, and in this way determine it to be acute pleurisy. But to prove what a thing is by proving all that it is not, is a very tedious process, and we must be quite certain that really *all* morbid states which may give rise to the symptoms encountered are thought of and inquired into; otherwise our conclusion may be fallacious, though reasoned out in the most logical manner. Moreover, our knowledge of many pathological conditions is so imperfect that we are not fully cognizant of, or able at once to discern, the more characteristic signs; nor can the symptoms be taken hold of and arranged in such a way as shall permit us to make nice distinctions without a lengthy and laborious plan of procedure. Owing to these drawbacks, *diagnosis by exclusion* is not, on ordinary occasions, much employed, nor, indeed, is it to be recommended. Yet in difficult and obscure cases, where the accustomed pathway is blocked up, it may enable us to pass by obstacles otherwise insurmountable.

But can we by this or by any other road always reach a certain diagnosis? We cannot, and for several reasons. The patient may deceive us, wilfully or unintentionally. It may be necessary, for the confirmation of the opinion formed, to obtain an accurate history of the case, and circumstances may render this impossible. The disorder may be so rare that its symptoms are not understood. There may be several lesions present, the signs of one masking or neutralizing the signs of the other.

The first of the causes mentioned is a source of error difficult to guard against. To escape punishment, to avoid disagreeable duty, to excite compassion, to obtain a compliance with unreasonable wishes, or sometimes from the mere love of deception, symptoms may be stated to exist which do not exist, or may be imitated and artificially produced. Persons who thus feign disease are numerous. They are found in all occupations and in all classes of

society. They abound in the army and navy. Hysterical women and hypochondriacs help to swell the list. These, indeed, suffer mostly some inconvenience, but exaggerate it immensely, and, by deceiving themselves, end by deceiving, unless he be on his guard, their physician. On the other hand, disease actually in progress may be carefully concealed from motives of delicacy or from fear of the consequences.

An incorrect diagnosis from want of a proper history does not, on the whole, occur often. Patients are generally very willing to give a full account of themselves and of their distresses. Sometimes, however, the reverse happens. Pain or mental anxiety and sorrow may be wearing the body out while the sufferer obstinately persists in hiding the cause of his waning health. We meet also with individuals so stupid that the most elaborate cross-examination fails to elicit anything like a connected history. Again, we may be unable to do so from the patient having lost the power of speech. A man is brought into a hospital unconscious. It is of the utmost importance to know how long he has been in this state, and what were his prior symptoms: unless some friend can supply the information, the most valuable diagnostic data are wanting.

In the rarity of a disease we have a serious drawback to its recognition. This may occasion an error of diagnosis in a two-fold manner. The more distinctive symptoms may be so little understood, and the prominent features be so nearly identical with those of a malady with the manifestations of which we are well acquainted, that a conclusion of the presence of the latter forces itself almost immediately on the mind. Or, the disorder may give rise to phenomena wholly unknown, nothing but the autopsy revealing their true meaning. Every physician encounters such cases. It is true that the progress of science and the aggregation of clinical facts are from year to year bringing them into a narrower circle. Yet, are there not still diseases, nay, groups of diseases, that have eluded discovery to the manifold means of research of the present day, as they have to the accumulated experience of the past?

But the most serious obstacle to a precise diagnosis lies in the fact that frequently lesions coexist. Disease is a very complex state, and when one portion of the economy gets out of order, another is apt to follow. How close, for example, the connection

between affections of the heart and affections of the kidney ! Here it is comparatively easy to arrive at a conclusion, since we have the means of judging accurately of the condition of both organs. But there are instances in which it is very difficult, as when a part contiguous to one chronically affected is attacked with acute disease. A person applies for relief, presenting the symptoms of a severe local peritonitis. The inflammation spreads ; death results. The exciting cause of the inflammation is discovered to be a structural alteration of one of the abdominal viscera, the signs of which were completely merged in the more marked signs of the recent inflammation. And this disguisement is effected not only by the supervention of another and more acute complaint, but also sometimes by the prominence of those remote sympathetic derangements which an affection of any viscus may produce. Thus, the disturbed action of the heart in dyspeptic persons throws at times the symptoms of the gastric malady into the shade. Yet it must be admitted that errors of diagnosis from this source are not apt to occur to the careful physician. A thorough examination of the case is a safeguard against them.

These, then, are the various causes which render a diagnosis uncertain, or wholly unattainable. Let us add to them one that does so temporarily. There are disorders the early manifestations of which are so much alike that it is next to impossible to tell with which of several we have to deal. In fevers this often happens. Here, however, a few days, or even less time, will almost always solve the difficulty. But not so in other diseases. It is only after a much longer period, and by careful watching of the patient, that the appearance or disappearance of a striking symptom, or the greater prominence a hitherto indistinct sign assumes, inclines the scales toward one or the other of the affections between which judgment has been kept in suspense.

In some such instances, the treatment becomes the touchstone of the diagnosis. Now it may be asked, Does this demonstrate that the diagnosis of a case is not necessary for its treatment ? Not at all. It simply proves that we are sometimes obliged to aim at removing symptoms without understanding their source. But it does not prove that if we understood their source we should not be better able to remove the symptoms. The physician who undertakes to relieve disease simply by attempting to allay its

symptoms, regardless of their cause, and without understanding their true relation and significance, is groping in the dark. His treatment is vacillating ; drug replaces drug ; alleviation is taken for a cure ; and the experience obtained is utterly untrustworthy. One great advantage, indeed, of attending carefully to diagnosis is, that it enables us to use remedies knowingly and with decision ; to appreciate what they are effecting ; to abstain from such as must be injurious. There is less needless meddling, more calmness ; the treatment rises above the consideration of the moment, and takes into account what is for the patient's ultimate good. It is sometimes urged that the accurate detection of disease makes timid practitioners, and deprives them of confidence in medicines. More just is it to say that it shows how wide is the chasm between our acquaintance with morbid conditions and our acquaintance with remedies ; how far our skill to detect disease still outruns our power to cure it.

There is undoubtedly, however, a danger which may arise from paying very minute attention to diagnosis. The study of it is so interesting, and capable of being conducted so entirely without reference to other points, and especially to the treatment of the complaint, that some minds are carried away, and, lost in the pursuit of diagnostic knowledge, forget for what purposes chiefly this knowledge is profitable. Its main use is to enable us to foretell the course and probable issue of a malady, and to frame, with understanding, plans for its relief. Nor ought we ever to be unmindful how important it is, in basing the management of a disease on its diagnosis, to found that diagnosis on a general survey of all the circumstances ; how necessary not to assign prominence to minor points ; and how the extent of the affection, the circumstances under which it has occurred, the sympathetic disturbances produced, and the vital state of the patient, belong, rightly considered, quite as much to the diagnosis as the recognition of the precise seat and exact anatomical character of the malady, and are, in truth, frequently its more important part.



## CHAPTER I.

### THE EXAMINATION OF PATIENTS, SYMPTOMS OF GENERAL IMPORT, AND SOME OF THE INSTRUMENTS EMPLOYED IN THE DIAGNOSIS.

To elicit the facts of a case by a careful examination is, as has been stated, the first requisite for diagnosis. To conduct, however, a clinical inquiry with precision and facility, requires continual practice, and is rendered easier by following some well-digested plan. The advantage of adopting a method is clearly seen, if the attempts of a beginner be watched. He wanders in his search from one part of the body to another, attracted by different symptoms in turn; pointless question succeeds to pointless question; and a conclusion, almost certainly erroneous, is finally jumped at, or an acknowledgment made of inability to arrive at any.

Now, there are several ways which have been proposed to overcome this embarrassment. One of the principal consists in first questioning the patient with regard to his history. His age; his occupation; the diseases from his childhood up; his habits; his constitution; the affections hereditary in his family,—are all minutely inquired into. After this the origin and progress of the existing disorder are traced, and the remedies ascertained that have been used against it. The present condition is then explored; each organ or each system being in turn interrogated. The investigation is now regarded as complete; the facts are considered, and the diagnosis, prognosis, and treatment determined. This method of examining is termed the *synthetical* or historical. The *analytical* reverses the order. The present condition is first ascertained, and subsequently the patient's history or *anamnesis*. Both of these courses have something to recommend them, and to both there are objections. The synthetical method is the more purely scientific; but it is too full, and calls for too much labor, to meet the requirements of ordinary professional life. It is

much better adapted for recording cases in the pursuit simply of pathological knowledge, and decidedly the best where the history is obscure and the symptoms are ill defined. The plan which I habitually prefer is to take a general survey of the history and of the prominent symptoms, and, having thus obtained some clue to the part most likely to be affected, to explore this with care. For instance : we are brought to the bedside of a patient for the first time ; we inquire how long he has been ill ; how that illness began ; in what way he is now disturbed,—whether he has pain, or what is the main source of his annoyance. While questioning him, we are scanning his appearance, the position of the body, his movements, his manner of breathing. The hand is applied to the skin ; the pulse is felt ; the tongue is looked at ; the temperature is taken. Partly from this examination and partly from the history, some organ is fixed upon to be specially investigated : say pain in the epigastric region and vomiting are complained of,—our attention is directed to the stomach. We explore this organ, its physical state and its functions. Then we look to the parts that are anatomically or physiologically most nearly related to it, which are, in the case cited, the intestines and the liver. The examination is completed by taking heed of the condition of other portions of the body ; by reviewing the history of the case ; and by endeavoring to elicit fully such points as bear upon the diagnosis, which the mind, consciously or unconsciously, has already begun to frame. Then a balance between the symptoms is struck, the diagnosis is recast, modified, or extended, and the treatment is decided upon.

There is some repetition in this plan ; but it is the one which appears practically the most suitable. It has the advantage of bringing together the marked features of a case, and especially those most clearly indicative of the general or vital condition. But whatever scheme be chosen, it should, for us to become proficient in it, be as constantly and closely adhered to as the varying circumstances of disease will permit. Yet to acquire thoroughly the habit of examining with accuracy and care, and also to obtain the full fruits of experience, it is indispensable to keep written records. This, too, should, so far as possible, be done according to a uniform design, since it both prevents us from overlooking important symptoms and enables cases to be more readily com-

pared. I subjoin a schedule that is based on the plan of examination just mentioned.

Date of Examination ; Name ; Age ; Color ; Place of Birth ; Present Abode ; Occupation or social state ; In females, whether married or not, number of children, and date of last confinement ; how many miscarriages.

### HISTORY.

1. *History antecedent to present disease*: Constitution and General Health—Hereditary predisposition—Previous Diseases or Injuries or Taints—Habits and mode of life ; hygienic influences to which exposed, etc.
2. *History of present disease*: Its supposed exciting cause—Exposure to contagion—Date of seizure—Mode of invasion ; subsequent symptoms in order of succession—Previous treatment.

### PRESENT CONDITION OF PATIENT.

Height and Weight.

1. *General symptoms*:

Position { in bed—mode of lying,  
out of bed—movements ; gait and station.

Aspect { of body ;  
of countenance ;

Skin ;

Pulse ;

Temperature ;

Respiration ; frequency, character, etc. ;

Tongue ;

General state of Digestion { appetite ;  
thirst ;  
condition of bowels ;

General State of Urinary Secretion and Urinary Analysis ;

Sensations of patient : pain, etc.

2. *Examination of special regions, parts, and functions*, beginning with the one presumably the most affected, and embracing, whenever practicable, microscopical examination of the blood and bacteriological studies.

## DIAGNOSIS.

## TREATMENT.

## REMARKS.

The history is here placed first; then the symptoms of general import, such as those furnished by the pulse, the tongue, and the temperature, are made to precede the examination of special regions. These general symptoms are of great value in the recognition of disease, and of yet greater value in determining its treatment. They are more than the mere physical signs of textural affections; they indicate vital conditions, and partly from their importance, and partly from their not being linked to disease of any organ in particular, they demand a separate and detailed consideration.

**Position of the Body.**—By noting whether the patient is in bed or out of bed,—how he lies, or how he walks,—a general idea may be formed as to the acuteness of an attack, the impairment of strength it has produced, and sometimes even as to its nature. Let a person who has been actively attending to his usual occupation be suddenly confined to his bed, and the inference that the disease is an acute and a severe one will be commonly correct; certainly so, if no mishap to the organs of locomotion have necessitated a resort to the recumbent position. When the patient lies for a long time on his back, it is generally from exhaustion, or from paralysis, or it is owing to the pain which pressure or motion of any kind occasions. Such is the cause of the dorsal decubitus in peritonitis, and in rheumatism. Lying steadily on the back with a disposition to slip down in bed is a form of dorsal decubitus witnessed in low fevers. Lying fixedly upon one side may, as a rule, be looked upon as an indication that the action of the lung of this side is impeded, and that the respiration has to be carried on with the other. The patient may be confined to bed, yet unable to lie down in it, on account of the distress in breathing to which the recumbent posture gives rise: he leans forward, or sits erect. This necessity of breathing in the upright position, or “orthopnoea,” is a form of dyspnoea encountered especially in diseases of the heart, or where fluid is effused into the air-cells or into both pleural cavities.

If a person is able to be about, his posture and movements become important manifestations of his condition. The young and the strong walk erectly, quickly, and firmly; the aged and the



weak, stoopingly, slowly, and with difficulty. In diseases of the spine the body is bent; so, too, in affections of the larger joints of the lower extremities.

When, after a fever or any other prostrating malady, the patient leaves his bed, he totters, moves slowly, and is soon obliged to rest: returning strength brings with it a quicker and steadier gait. In some diseases of the brain the movements are very uncertain; in one-sided palsy the affected side lags, or its motions, if it can be moved at all, are laborious. Excessive and uncontrollable movements are observed in mania and in chorea; trembling motions in states of extreme debility, in shaking palsies, and in the delirium of drunkards; irregular motions and positions chiefly in hysteria.

The *gait* is always to be closely studied. We find it of special significance in affections of the nervous system and of the muscles. It is very erratic, from side to side, in locomotor ataxia, and there is almost total inability to walk in the dark. In paralysis agitans the tremors are associated with a festinating gait, each step becoming more rapid than the last, and a fall is only averted by seeking support. In spastic paraplegia the legs drag behind; in walking each leg is rigidly brought forward, the toes having a tendency to catch the ground. In pseudo hypertrophic paralysis occurs a peculiar oscillating or waddling gait, from weakness of the extensors of the knee and hips; there is also much difficulty in rising from the ground. In Thomsen's disease it may also be for some time impossible to rise from the floor, and the gait is at first impeded by tonic spasm of the muscles.

*Station*, or the power of preserving an erect position while standing, is often as characteristic as the gait. It should be noted while the eyes are open, while they are shut, and while the feet are placed alongside each other with the heels and toes touching. Under both the latter circumstances the station is always less certain and the swaying of the body more marked. Tested with an instrument invented by Weir Mitchell,\* Hinsdale† found in the normal man and woman the average sway, while the heels and toes were touching, to be about an inch in the forward and backward line, and three-quarters of an inch laterally.

---

\* Amer. Journ. Med. Sciences, 1887.

† Ibid., April, 1887.

Children sway to a greater extent than adults. Closing the eyes increases the sway about fifty per cent. In locomotor ataxia station is much disturbed and the sway greatly increased ; so it is in disease of the middle lobe of the cerebellum. In the attacks of aural vertigo all power of standing may be lost.

**General Aspect—Expression of Countenance.**—The eye notices rapidly whether the body is bulky or wasted, and whether the surface is discolored or otherwise changed. A bulky aspect of the whole body is the result of corpulency, or arises from universal anasarca. In some acute diseases, too, a general tumefaction may take place,—for example, in the exanthemata. A partial increase, or a swelling, arises from the local extravasation of fluid or air into the cellular tissues. If air, the tissues crepitate under the finger ; if fluid, the skin pits under pressure. A swelling may, further, proceed from an inflammatory thickening, or from a tumor or any morbid growth.

A diminution in bulk is a more frequent symptom than an augmentation. It may take place very rapidly, as witnessed in Asiatic cholera. More generally the wasting is gradual, and is an indication of defective nutrition. It occurs in the course of protracted fevers, and in most chronic diseases. In slowly fatal maladies, and in those attended with constant discharges,—for instance, in chronic diarrhœa,—the loss of flesh reaches its highest point.

Emaciation is most readily recognized in the face. But it is not the only striking alteration observable in the countenance when health has failed. There may be pallor, sallowness, a livid hue of the lips, a puffy appearance of the eyelids, a flush on the cheeks. Now, these changes in the features, added to the expression which pain or special trains of thought produce, make up that peculiar physiognomy of disease so pregnant with meaning. But I shall not attempt to describe in detail the cast or the play of features in the sick : the shades of expression are so numerous that they baffle description, and are to be learned only by continuous bedside experience. I shall merely set down a few broad facts which this experience teaches.

Among the countenances most frequently met with is that of apathy and stupor. The eye is dull and listless ; the face pale, or flushed with fever. This look is common in fevers of a low type,

and is often combined with blackish accumulations on the lips, gums, and teeth.

Unnatural fulness and congestion of the features are sometimes observed in enlargements of the heart, and oftener still in habitual drunkards. The same aspect is seen in apoplexy and in typhus fever. Local congestions on the cheeks and nose are met with in obstructive diseases of the liver, especially in cirrhosis, and in the endarteritis of old persons. A pinched expression is found when there is intense anxiety or pain, or a wasting malady attended with constant suffering. It is specially observed in acute peritoneal inflammation. When very marked, and accompanied by change of hue, it is the face which Hippocrates has so graphically described. In the great master's own words, "a sharp nose, hollow eyes, collapsed temples; the ears cold, contracted, and their lobes turned out; the skin about the forehead being rough, distended, and parched; the color of the whole face being green, black, livid, or lead-colored." This is the physiognomy of approaching death, and generally its speedy forerunner, except in those cases in which the expression proceeds from want of food, from protracted vigils, or from excessive discharge from the bowels.

The face of shock, with its great pallor, its anxious or frightened look, and its fixed or oscillating eye, often with a contracting pupil, is a face seen after severe injuries, and as such familiar to the surgeon. But in many of its main traits it may be also met with in diseases that make a sudden and overwhelming impression on the nervous system; for instance, it is at times encountered in cerebro-spinal fever and in cholera.

An aspect serious and dull on one side, while the other side is in full play, is witnessed in some instances of hemiplegia, and in paralysis of the facial branch of the seventh nerve. The difference in the cast of the features may escape observation when the face is in repose, but as soon as an attempt is made to laugh, it shows itself plainly.

Besides these lineaments, which may be said to be common to several diseases, we read frequently in the countenance the signs of special disorders. A dusky flush on the face, if associated with rapid breathing, is almost a certain indication of inflammation of the lung. Puffiness of the eyelids in a pallid person is



most apt to be expressive of Bright's disease. A bluish color of the lips shows plainly that the venous circulation is interfered with, or that the blood is but imperfectly aerated. The cyanosis is also recognized in the blueness of the nails and the duskiuess of the whole surface. Then there is the chronic pallor of the anæmias with the pearly eye and the yellowish tinge of the pallor in chlorosis; the straw-colored, anæmic hue of malignant disease; or we note the jaundiced, melancholy look of an hepatic affection; the downcast expression and mobility of the features in hysteria; the thickened upper lip, delicate skin, and fair complexion of scrofula; the sallow countenance and peculiar notched teeth that indicate inherited syphilis; and the various traits which tend to mark not only the special diathesis, but also the peculiar temperament, with the morbid tendencies that belong to it.

**Skin.**—By the state of the skin we can, to a great extent, judge of the activity of the circulation and of the character of the blood. Moreover, it is a fair index of the secretions, and of the condition of the system at large. In fever, along with the quickened circulation, the temperature of the skin is increased; the attending dryness is produced by defective perspiration. When, after pressure on the skin, the blood returns slowly to the surface, it denotes a sluggish capillary circulation; when rapidly, an active one. Coldness of the surface indicates a weakened capillary circulation, and is met with at the invasion of acute diseases, and when the nervous power is under the sway of some highly deleterious influence. If the heat of surface succeed a cold skin, we know that reaction has taken place, that the circulation has again become active. Protracted coldness, whether attended with dryness or with clamminess, is of evil augury: it implies seriously diminished vital force.

The cutaneous covering is pale whenever the blood is poor and watery. If this be seriously vitiated and largely deprived of its fibrin, as in putrid fevers, black spots are seen, due to extravasation. Ofttimes the surface is overspread with eruptions, some of which bear a close relation to disorders of internal organs, while others are connected with febrile or general maladies; and others, again, are owing to a disease of the texture itself.

Tension of the skin is met with in acute affections accompanied by active excitement. In wasting and prostrating ailments, on



the other hand, the skin feels very relaxed and soft ; and in those producing rapid emaciation, it is inelastic and lies in folds.

**Pulse.**—The study of the pulse has come down to us with the sanction of centuries ; and to feel the beat at the wrist is still, in the opinion of many, as indispensable to the understanding of a case as it was thought to be by the Arabs and in the Middle Ages. Yet the advance of science has shaken the belief in the paramount importance of the pulse. It has shown that, although a most valuable means of information, it is not exclusively to be relied upon, and has proved the many divisions and refinements of the physicians of by-gone days to be imaginary and useless. Indeed, were even all their distinctions founded in fact, we have now better ways of judging of many lesions than by feeling the radial artery.

The pulse enlightens us on the action of the heart, and on the state of the artery itself and of the blood. In a healthy adult a beat of some resistance is felt, recurring from sixty-five to seventy-five times in a minute. It becomes slower with advancing years, though it may rise in the very aged. The pulse of infancy is from one hundred and ten to one hundred and twenty ; that of a child three years old, from ninety to ninety-five. Warmth quickens the pulse ; so do rapid breathing, forced expiration, and the process of digestion. In the recumbent position and during sleep it falls. For purposes of comparison, the pulse should be, so far as possible, taken under similar conditions.

At the bedside we study in the pulse its frequency, its rhythm, its volume and strength, and its resistance.

Increased *frequency* of the pulse denotes increased frequency of the heart's action, and arises from any cause that excites the heart. Hence exercise, rapid breathing, mental emotion, or restlessness will occasion the number of beats to exceed the average of health as readily as fevers or acute inflammatory diseases. In great debility, too, the pulse rises ; and the more depressed the vital condition, the higher the pulse becomes. In exophthalmic goitre the pulse is generally very frequent, and rapid heart action may show itself without any other appreciable abnormal state, as in tachycardia, a disorder in which the pulse may considerably exceed two hundred beats in the minute. Under the influence of suggestion the cardiac action may be made very much more rapid

or slower.\* As a sequel of influenza there is often very rapid heart action. The heart may thus quicken from so many and such varied causes, acting temporarily or permanently, that increased frequency of pulse, taken by itself, has no significant diagnostic meaning.

A *slow* pulse, too, happens in many different states,—in cold, in exposure to wet, in icterus, in protracted convalescence from acute disease. It is also produced by an intense and prostrating shock, or is found coexisting with pressure on the brain, with melancholia, with atheroma, with fatty heart. A permanently slow pulse is also met with in irritative lesions of the cerebral centres, among them in spherical or pediculated thrombi, in altered state of the circulation in the bulb, and in injuries to the pneumo-gastric. It is not unusual in instances of very slow pulse, or bradycardia, to observe two or three abortive beats succeeding a strong beat. In some persons the pulse is naturally very slow.

The *rhythm* of the pulse is often perverted. Instead of the beats following one another in regular succession, they are unequal, or one or two intermit. An irregular pulse occurs from digestive disorder, from gout, from lithæmia, from the excessive use of tobacco, or from debility and nervous exhaustion; it is less frequently the indication of a cerebral or cardiac lesion. It is sometimes a difficult beat to count; and we must be careful not to regard at once a pulse as irregular because it appears to intermit. The seeming irregularity may be caused by the fingers slipping from the artery, which they are very apt to do after they have been on the vessel for some time.

Where every other beat is uneven in size, thus showing a beat of greater, followed regularly by one of lesser, altitude, though the rhythm may be regular, we have the *pulsus alternans*. Where a beat is dropped,—in other words, where the heart-beat is not transmitted to the artery with sufficient force to be felt,—it is designated as an abortive beat. Two imperfect or abortive beats occurring in rapid succession, and followed by a long pause and generally by a distinct beat, form a linked beat.

The *volume* and *strength* of the pulse are of much more importance than either its rhythm or its frequency. Volume and

---

\* Sgobbo-Nuovo Rivista, 1, 1892.

strength are often associated, and are much alike; but they are not identical. When the beat of the artery is large, we call it a *full* pulse. This is owing to the distention of the vessel with blood,—its complete expansion with every beat of the heart. A full pulse is, therefore, the pulse of plethora; the pulse of the young and robust in health, or in inflammatory diseases; the pulse in the early stages of fevers, or in obstruction of the capillaries. It is usually a pulse of power, just as its opposite, a *small* pulse, is usually the pulse of debility. Yet a full pulse may be produced by the distention of an artery which has lost its tone, and which the finger easily compresses. Such a pulse, the “gaseous pulse,” a pulse really of low tension, denotes exhaustion, and proves that a full pulse and a strong pulse are not always synonymous. Into the idea of *strength* something more than mere fulness enters. A strong pulse is a pulse heightened in all its natural characters. It has more fulness, but, in addition, more impulse, and less compressibility, than an ordinary pulse. A strong pulse, therefore, indicates activity of the contraction of the heart, and a normal, perhaps increased, tonicity of the arterial coats. It is found in active inflammations; also in hypertrophy of the heart. Its opposite, a weak pulse, betokens want of force, often want of healthy blood. It is generally small as well as weak. Yet as the full pulse is not always strong, neither is the small pulse always weak. The small, choked pulse of peritoneal inflammation may be fine and wiry, but it is not a weak pulse. We also find a small pulse of high tension in mitral stenosis and in contracted kidney.

The *resistance* or tension of the pulse is another valuable guide in the appreciation of morbid action. Is the pulse hard and resisting? is it soft and compressible? are questions on the solution of which the application of remedies may hang. A *hard*, tense pulse denotes increased contractility of the arteries, and high-wrought power. Be the beat full or small, slow or frequent, it tells us that the blood is being driven with force along the arterial system. But it also tells us that the irritation has implicated the coats of the arteries themselves, or that there is obstruction in the capillaries. A tense pulse is met with in active, violent inflammations, and sometimes, though not often, in states of extreme and continued excitement without inflammation. It

is almost needless to add that changes in the coats of the arteries may also be a cause of a hard and resistant beat, the common cause of the increased tension in elderly people. Where no local alterations are present, and where no acute symptoms explain the sympathetic disturbance of the heart and arterial system, the high arterial tension will be commonly found associated with hypertrophy of the left ventricle, with interstitial nephritis, with gout or lithæmia, or with blood-poisoning.

The opposite of the hard pulse is the *soft* or compressible pulse. This implies deficient impulsion, and loss of tone in the vessel; it is the pulse of low fevers, of debility, of cardiac weakness. But it is also, when following a tense state of the artery, the pulse which denotes returning health, and danger passed. In some persons the pulse has naturally a low tension, just as in others it has a high tension.

When the pulse is of low tension, and at the same time frequent, it may show double beats with each contraction of the heart. This dicrotic pulse is most often met with in fevers of a low form and preceding or during the continuance of hemorrhages. The rebound is chiefly due to the oscillation of the column of blood in the arteries, and is very much influenced by their elasticity. With lowered tension and increased elasticity of the tubes, dicrotism becomes obvious, especially with a rapid circulation. In old persons, in whom the coats of the arteries are inelastic, dicrotism is but feebly marked.

Such are the meanings attached to the various characters of the pulse. Yet they do not often present themselves thus isolated. The following are usually combined, and bear this explanation:

A hard, full, frequent pulse occurs in active inflammations, and in most of the acute diseases of robust persons.

A hard pulse, full or small, bounding or not, if unconnected with acute symptoms, leads to the suspicion of cardiac or of renal disease, or of an affection of the artery itself.

A tense, contracted, and frequent pulse is met with in a large group of inflammations below the diaphragm, as in enteritis, peritonitis, gastritis.

A frequent pulse, full or small, but not tense, is the pulse of most idiopathic fevers, and with marked low tension is also apt to be dicrotic.




A very frequent pulse, but very feeble and compressible, is the pulse of marked debility, of prostration, of collapse.

A pulse frequent, and changeable in its rhythm, is produced, for the most part, by perverted innervation in connection with gastric disorders, by tobacco, or by disease either of the heart or of the brain.

The appreciation of these different kinds of pulses requires considerable practice. But even this scarcely teaches us to estimate

FIG. 1.




Marey's sphygmograph attached to the wrist. Its tracings are shown by the white lines on the black background.

the exact degree of the alteration of the beat, certainly not with sufficient distinctness to convey to others an accurate idea, or even to be able ourselves to compare one observation with another. To attain these desirable results, instruments have been sought for by means of which the pulse can be examined with precision, its finer shades of difference recognized, and its movements recorded. The best of these instruments is the *sphygmograph* invented by Marey, which registers not only the frequency and regularity but the form of the pulse-wave, and which may be also used in the study of the cardiac impulse and of pulsatile tumors. Slight irregularities that wholly escape the finger are, through its aid, discerned with facility, and we tell at once in how far these irregularities belong to one beat or to a succession of beats. Double beats with each contraction of the heart, too, not appreciable to the hand, are easily detected. Indeed, the sphygmograph proves the phenomenon of dicrotism to exist in almost every person. The rebound may occur during the systole or the diastole of the vessel; and instead of one, there may be four or five of the secondary pulsations.

The mode of adjusting the instrument, and of proportioning the pressure of the spring, has something to do with the kind of

delineation obtained. To secure greater accuracy, Sanderson fixed the centre button at a definite pressure, and Mahomed added several serviceable contrivances, one of the chief of which is the causing of the amount of pressure employed to be accurately registered upon a dial. Still another modification, which, however, really makes use of a different principle, the displacing power of the artery rather than its lifting power, has been made by Holden. The movement thus obtained is from side to side. A sphygmograph marking extremely fine tracings is that of Pond. A rubber diaphragm takes the place of the spring of other sphygmographs, and is fixed to the artery by means of a holder. A delicate needle makes the tracing. The sphygmograph of Dud-

FIG. 2.



Dudgeon's sphygmograph.

geon (Fig. 2) is simple and very much employed. The system of levers is the same as in Marey's, but the slip of paper moves in a different direction.

To show the tracing distinctly, smoked glass or mica, or paper smoked over a lamp or by burning camphor, is much used; and the tracing may be preserved by dipping it in an alcoholic solution of shellac or of benzoin, or of a varnish of benzoin and methylated spirit, in the proportion of one to six. On every tracing the amount of pressure employed should be noted. Manifold have been the suggestions to obtain the steadiest application


of the instrument to the forearm and the greatest development of the tracing. Lorain \* has proved that raising the arm to a vertical position gives a much more ample trace ; and Richardson † shows that with the body in the horizontal line, the dicrotic wave becomes more prominent.

When we apply the sphygmograph for clinical purposes, we study in its tracing the line of ascent, the summit, and the line of descent. Each pulsation is composed of these three parts. The *line of ascent*, the upstroke, tells us the manner in which the blood enters the vessels. The more rapid the flow, and the more quickly the artery distends, the more strictly vertical the line. The force, too, is indicated by this line, or rather by its height : hence when the muscles of the heart contract powerfully, either from enlargement or from overaction, the line is both vertical and high.

Yet the strength of the ventricular contraction is far from being the only cause influencing the amplitude of the tracing. Indeed, as we may note in old persons, a large volume of the artery gives considerable height to the lines of ascent ; so does a long interval between the pulsations, or the obstruction of the vessel below the point where the observation is made. Low tension in the arteries or in the capillaries has the same effect ; whereas when the passage in the ultimate ramification of the vascular system is difficult, the lever descends slowly by a convex line, and is soon again raised by the next pulsation. When the contraction of the heart is feeble, the line of ascent is not vertical or high, but oblique and short. In aneurisms of the thoracic aorta—indeed, in an aneurism interposing anywhere between the heart and the radial artery—an oblique and short upstroke is also met with.

The line joining the summit of a series of pulsations, or the maxima of tension, is generally a straight line ; a similar imagi-

FIG. 3.



Sphygmogram enlarged.—*a, b*, upstroke, or line of ascent ; *a, b, c*, percussion-wave ; *c, d, e*, tidal or predicrotic wave ; *d, e, f*, aortic notch ; *e, f, g*, dicrotic wave ; *f, g*, diastolic period.

\* Le Pouls, Paris, 1870.

† The Asclepiad, 1886.

nary line connecting the bases, or the minima, is apt to run parallel to it; but irregularity of pulsation leads to irregular lines, and the lower line may be irregular while the upper is straight. Irregularity of the base line is seen in marked dyspnœa.

The *summit* of the pulsation informs us of the time during which the entrance of blood balances the onward flow. A pointed, distinct summit-wave belongs to vigorous contraction of the heart-muscle. The summit may be a horizontal line of some length. This broadening of the apex happens in high and prolonged arterial tension, such as from the slow contraction of a strong heart, fulness of the vessels, or obstruction in the capillaries; an extended plateau is also met with in induration or ossification of the arteries.

In some instances we find a little hooked point preceding the usually transverse mark of the summit. This occurs by the rapid movement of the lever, and is generally a sign of regurgitation coexisting with obstruction at the aortic valves. In aortic narrowing of marked degree the summit-wave is indistinct or absent; the line of ascent is oblique and gradual, and may show a break near the summit.

The *line of descent* is sometimes purely oblique, and the more rapidly the pressure is lessened in the arterial system, the more oblique is the line. It often shows a series of undulations. The first of these waves is called the tidal wave; it is still part of the systole and onward flow of the blood; the decided subsequent wave is specially called the dicrotic or great secondary wave. The closure of the aortic valves with the second sound of the heart happens just before the dicrotic wave; the exact time is marked by the aortic notch; the dicrotic wave represents the diastole of the heart. The tidal wave is large, but the dicrotism badly marked, in atheroma. In high arterial tension the dicrotic wave is also ill pronounced, and the line of descent is very gradual. In mitral narrowing, the line of descent is long, but is broken by small pulsations.

The sphygmograph requires much care and practice in its use; and with all the perfection of the instrument, its precise value for clinical research is still undetermined. I think it of much more avail in investigations on the exact action of medicines—where, indeed, it is of great value—than in aiding us materially in questions of diagnosis or in decisions on treatment. At all events, I do not



think that it supersedes the older and more usual means of research. Perhaps records of pulse-traces in which the amount of pressure has been carefully noted will enable us to judge more accurately than we can now of the state of the cardiac muscles in disease.

An instrument aiming at even greater accuracy than the ordinary sphygmograph is the *sphygmochronograph*.<sup>\*</sup> It is similar in its construction to the sphygmograph of Dudgeon, but it enables us to measure the curves of the tracings, and to ascertain the exact time of each part.

Normally *capillaries* do not pulsate. We judge of their dilatation by the flush, of their contraction by pallor. But in certain pathological conditions they beat, as may be observed in the capillary flush. We may note the *capillary pulsation* in instances of chlorosis and of aortic regurgitation. The capillary flush has generally to be brought about artificially by pressure on the skin, the nails, or the lips. We can then perceive the pink changing in color with each pulsation, or disappearing after it. The most marked changes are observable at the periphery of the pink patch. In those rare instances in which the capillary pulsation is regurgitant and of venous origin, as in tricuspid regurgitation, we find venous pulsation elsewhere, and the capillary pulsation precedes the radial beat.

**Temperature of the Body.**—The thermometry of disease has become indispensable. The thermometer used for clinical purposes should be very sensitive, and requires to be from time to time compared with a standard one, and verified. It may be straight, or curved. The scale, extending from about 85° to 115° Fahr., ought to be uniformly graduated; it should be divided so as to exhibit fifths of a degree. The most useful instrument is self-registering (Fig. 4); a straight thermometer, generally short, for convenience' sake, and having the mercury detached from the column. This detached part, or the index, is set by bringing it down below the lines of the scale by a rapid swing of the arm. After the thermometer has been in position for the required period, it is removed, and the end of the index farthest from the bulb records the temperature. A magnifying front allows the degrees to be easily read.

---

<sup>\*</sup> Jaquet, *Zeitschrift für Biologie*, 1891; and Mühl, *Deutsch. Arch. f. klin. Med.*, 1892, xlix.

FIG. 4.



Self-Registering Thermometer, showing the index marking 99° shortly after an observation.

FIG. 5.



Seguin's Surface Thermometer, modified to be self-registering.

FIG. 6.



Surface Thermometer, with coil at extremity. It may be, if necessary, kept in place by a thin elastic band.

Very delicate but fragile thermometers, registering in a minute or less, have of late come into use. Metallic thermometers are neither so cleanly nor so trustworthy as those made of glass.

As surface thermometers for localized thermometry various instruments have been suggested. I habitually employ one which has the mercury in a fine coil at the expanded extremity, and which is self-registering. The ordinary self-registering clinical thermometer may be made use of, with the bulb fitted into a piece of cork. Whatever instrument be resorted to, we should first obtain the heat of a corresponding or analogous well part, and then leave the bulb for five minutes on the suspected abnormal structure. Better still is it to apply two instruments at the same time; one on the sound, the other on the unsound side. In all observations the heat of the body, as ascertained in the axilla, should equally be noted. Thermo-electric apparatus have also been employed for surface thermometry, and certainly give very accurate results. But, with perhaps the exception of the instrument of Lombard,\* they are not sufficiently portable or easily enough managed for general use.

The surface temperature is, as a rule, lower by upwards of one or by several degrees than the general temperature. We find it so on the chest, on the abdomen, and on the head. The temperature, too, is not on corresponding sides entirely the same, at least not on the head. There is almost always a slight inequality in the temperature of the two sides of the head; Gray † demonstrates that when at rest the temperature of the left hemisphere is the higher, which accords with Broca's statement. And the observations of Amidon ‡ have shown that excessive use of a group of muscles may generate heat in the cortical region presiding over them, sufficient to manifest itself to surface thermometers placed on the scalp. The mean temperature of a healthy man's head is fixed by Maragliano and Seppili, as the result of many observations, at  $36.13^{\circ}$  Cent. ( $97.03^{\circ}$  Fahr.) for the left side of the head, and  $36.08^{\circ}$  Cent. ( $96.9^{\circ}$  Fahr.) for the right.§ These temperatures are much higher than those given by Broca and Gray, which

---

\* "On the Regional Temperature of the Head," London, 1879.

† Chicago Journal of Mental and Nervous Diseases, 1879.

‡ New York Archives of Medicine, April, 1880.

§ Translated in Alienist and Neurologist, St. Louis, Jan. 1880.

is accounted for by their having been taken in summer. Broca places the frontal region on the left side of the head at  $35.43^{\circ}$  Cent. ( $95.79^{\circ}$  Fahr.), on the right at  $35.22^{\circ}$  Cent. ( $95.39^{\circ}$  Fahr.); Gray's figures are somewhat lower. The parietal region on the right side is fixed by Broca at  $92.8^{\circ}$ ; by Gray at  $93.6^{\circ}$  on the right, and  $94.4^{\circ}$  on the left; the vertical by Gray at  $91.7^{\circ}$ , and the occipital at  $91.9^{\circ}$ ; the whole side of the head by Broca at about  $93^{\circ}$ ; the entire head at places remote from these points at  $93.5^{\circ}$  by Gray.\*

As regards the abdomen, Peter† places the normal mean of the parietes at  $35.5^{\circ}$  Cent. ( $95.9^{\circ}$  Fahr.), and the same observer records the normal temperature for the chest-walls at about  $36^{\circ}$  Cent. ( $96.8^{\circ}$  Fahr.). Certain diseases change the temperature locally. Thus, in neuralgia the heat near the painful points may be markedly raised. So, too, is it sometimes in some parts of the surface in hysterical women. In hemiplegia the paralyzed limb may show a higher temperature than the sound one. And over spots where there is inflammation or where decided tissue-change is going on there is a rise in local temperature. Weir Mitchell‡ has called attention to the manner in which posture affects surface temperature. It is, for instance, less by  $0.4^{\circ}$  C. to  $1^{\circ}$  C. on the dorsum and sole of the foot when standing than when lying down.

But to return to general thermometry. The clinical thermometer may be put under the tongue or in the rectum; but the most suitable site in adults is the axilla. The bulb is pressed into the armpit and kept in close contact with the skin for five minutes, except when the delicate minute thermometers are employed. The thermometer may be conveniently introduced just below the skin covering the edge of the pectoralis major muscle; and, to insure exactness, the axilla should be kept well covered. The best posture, as Ringer points out, is neither completely on the back nor on the side, but diagonally on the right or the left side.

In all cases of importance, not less than two observations should be made daily, and, so far as possible, every day at the same hour.

---

\* New York Archives of Medicine, 1879, vol. ii.


† Communication to the Académie de Médecine, quoted in Medical Times and Gazette, Dec. 1879.

‡ Medical News, Jan. 1894.



Between seven and nine o'clock in the morning, and about seven o'clock, or somewhat earlier, in the evening, are regarded as the most appropriate periods. If only a single observation be taken,

FIG. 7.




Temperature chart in simple continued fever. The *initial* stage, the stage in which the temperature rises to its height, is here attained on the second day; the *fastigium*, the stage at which it remains, with slight fluctuations, at its height, lasts until the sixth day; the *defervescence*, the stage when the fever-heat falls, is rapid, by crisis, and a subnormal point is, for a time, attained.

it is best done in the late afternoon or in the evening. In every record of the temperature the pulse and the respirations must also be noted. For the purposes of record it is best to use always clinical charts. To be graphic, these should not be too elaborate. Fig. 7 is a simple model. It can be arranged for thirty days or upward; by striking out the morning and evening marks, as many spaces as needed can be devoted to a case; or by using, as we do at the Pennsylvania Hospital, a marking in red ink in addition to the black line in which the morning and evening temperature is recorded, as near as may be always at the same hour, the supplementary temperature can easily be traced (Fig. 8). Where we

wish also to show the pulse and the respiration graphically, the chart of Crozer Griffith is excellent. In discussing pneumonia, further on, one of these charts is shown.

FIG. 8.



Temperature chart, from a case of remittent fever in a sailor at the Pennsylvania Hospital (No. 1570). The red lines show the intermediary temperatures.

In temperate climates the average heat of the body, as measured in the axilla, is estimated at 37° Centigrade; that of freshly-voided urine is about the same. Expressed in the scale used in this country and in England, the average heat of sheltered and internal parts of the body may be stated as 98.6° Fahr.\* This,

\* It may be useful, for the sake of comparison, to recall the fact that one degree of Fahrenheit is equal to five-ninths of a degree of the Centigrade thermometer, and four-ninths of a degree of Réaumur; and also that the freezing-point

at least, is the case in the axilla; in the rectum it is not quite  $1^{\circ}$  higher, and is very steady; in the mouth it is somewhat lower. In the groin, where, in children, it may be most convenient to take it, the temperature is apt to be lower than in the axilla.

The body temperature rises with the temperature of the air, and fluctuates slightly during the day, being in temperate climates, according to the most trustworthy observers, lowest between two and eight in the morning, and highest late in the afternoon. It is heightened by exercise and reduced by sustained mental exertion, and changes even when we are at rest.\* But, as a rule, with the exception of very active exercise, no cause save disease induces a variation of much more than  $1^{\circ}$ ; even in the extreme heat of tropical climates the animal heat does not surpass  $99.5^{\circ}$ . Thus a temperature above this, or more than a degree below the average stated, when persistent, indicates some morbid action in the economy. At all events, it does so in adults; in very aged persons a temperature of  $97^{\circ}$  may still be normal; while, on the other hand, the range may be as high as in infants.

In children, in whom the temperature, as a rule, is somewhat higher than in adults, the daily range is much greater. It falls rapidly in the evening, and is very much influenced by food and by crying. Immediately after birth the temperature is lowered; and in the new-born it is about  $99.8^{\circ}$  to  $100.4^{\circ}$  in the rectum. It falls from early infancy to puberty. The rectal temperature of young children ranges between  $99^{\circ}$  and  $99.7^{\circ}$ ; under six years of age the mean is  $99.4^{\circ}$ . But, as already stated, there are great variations in childhood. The maximum is attained in the afternoon; and in healthy children an evening fall is common. During the first three or four months of life the temperature, Henoch

---

of the first is placed at  $32^{\circ}$ ; that of the others at zero. To convert Centigrade into Fahrenheit, we multiply by 9 and divide by 5; to convert Réaumur, we multiply by 9 and divide by 4, and when above zero, in either case, add 32. To convert Fahrenheit above zero into Centigrade, we subtract 32, multiply by 5, and divide by 9.

\* See an instructive paper by Garrod, on the Minor Fluctuations of the Temperature of the Human Body, *Proc. Roy. Soc.*, May, 1869; an elaborate paper by Jaeger, *Deutsches Archiv für Klin. Med.*, Bd. xxix.; Goodhart, *Guy's Hospital Reports*, 3d series, vol. xv., particularly valuable as showing the variations during the prolonged application of the thermometer; and Boileau, *Clinical Thermometry in Hot Climates*, *Lancet*, Aug. 4, 1888.



asserts, has, from slight causes of faulty nutrition, a marked tendency to go below the normal. A further point, too, to be taken into account in those of all ages is, that the temperature is somewhat influenced by food and stimulants. And these are the elements which make deductions from single observations or comparatively slight changes untrustworthy. In high altitudes, as Keating \* has observed, there is a tendency to hyper-pyrexia.

In ordinary cases the pulse and temperature rise synchronously, and every degree above 98° Fahr. corresponds with an increase of ten beats of the pulse. The fever temperature ranges from 100° to 106°. When it exceeds this, the patient may be looked upon as in great danger, except the rise be due to malarial fever. Under these circumstances it is rapid, occurring in a person who but a few hours before was healthy. In typhoid fever a temperature of 105° is a proof of grave disease. In some severe cases of yellow fever the heat in the armpit has been noted as 108°.† In pneumonia a temperature above 104° is a symptom of a very serious seizure; so, too, is it in acute rheumatism a symptom either of danger or of some complication. Stability of temperature from morning to evening is a good sign; the temperature remaining the same from evening till morning is a sign that the patient is getting worse. In convalescence the temperature declines until it attains its norm, or even falls somewhat below this. If after the defervescence the thermometer again indicate a decided rise, it shows a return of the malady, or the supervention of some complication or new disorder; and the persistence of even a slight degree of abnormal heat after apparent convalescence is a sign of imperfect recovery, or of the existence of some lingering secondary complaint. Further, in cases of low fevers, the skin, particularly of the hands and feet, may feel cool at the same time that the instrument in the axilla marks 104°.

Specific forms of febrile diseases have their characteristic variations of temperature. In measles, for instance, the temperature rises toward the breaking out of the rash, reaches its height with the period of eruption, and in the twenty-four hours succeeding it falls rapidly. In scarlet fever the thermometer marks 105°, or

---

\* International Medical Magazine, Dec. 1892.

† Wragg, Charleston Medical Journal, vol. x.



upwards, at the beginning, and the heat only gradually subsides. Typhoid fever has its characteristic record; so have the malarial fevers theirs. The temperature of tetanus rises to great heights before death.

A temperature above  $107^{\circ}$  is almost certain to be the forerunner of a fatal issue. But recovery may take place. In a case of cerebral rheumatism under my charge\* the thermometer marked  $110^{\circ}$  in the axilla, yet the patient got well. In an instance of injury to the spine after a fall, reported by Teale,† the young lady lived though the temperature reached above  $122^{\circ}$  and ranged for days between  $112^{\circ}$  and  $114^{\circ}$ . A remarkable case has also been reported of hysteria and intercostal neuralgia, in which in one axilla the temperature registered  $117^{\circ}$  Fahr. and in the other  $110^{\circ}$ , but the patient recovered.‡ Galbraith§ has reported a case in which the thermometer registered  $151^{\circ}$ , and Jones|| that of a girl, fourteen years of age, in whom the temperature rose to over  $150^{\circ}$ . In neither instance was the extraordinary heat attended with evil results. Duckworth's report¶ of a case in which the thermometer marked  $228^{\circ}$  ( $108.9^{\circ}$  C.) is probably the highest on record. In all these extraordinary temperatures the possibility of deception practised by hysterical patients must be borne in mind. The temperature may be temporarily very high from emotion. I saw this once in a frightened child which had previously had but slight fever, and E. S. Tait has reported the same in the puerperal state.\*\*

On the other hand, the thermometer may show a depression in temperature below the normal. The body heat often falls at the beginning of acute peritonitis. It is low after severe loss of blood, or if exposure to cold happen in alcoholic intoxication, during convalescence from acute diseases, and in melancholia. It is depressed by various poisons, and has been observed down to  $93.9^{\circ}$  in carbolic acid poisoning.†† It is low in the insane. It may be

\* See Amer. Jour. of Med. Sci., Jan. 1875.

† Transact. Clinical Society of London, vol. viii.

‡ Philipson, London Lancet, April, 1880.

§ Journ. Amer. Med. Assoc., March, 1892.

|| Memphis Medical Monthly, Oct. 1891.

¶ Archives of Gynæcology, New York, Oct. 1891.

\*\* Obst. Soc. Transact., 1884.

†† Bäumler, in Quain's Dictionary of Medicine.

only a fraction above  $89^{\circ}$  in the axilla in cholera. From any other cause it rarely, even in extreme collapse, sinks below  $92^{\circ}$ .

Though having its widest range of applicability in fevers, in other than febrile states, too, the thermometer assists greatly in diagnosis and prognosis. It is invaluable, in many instances, in discriminating between functional and organic affections. It aids in the study of apoplexy, of palsies, and of hysterical affections, and tells the true story in cases of feigned disease. It also enables us to judge whether increased frequency of pulse be due to fever or to debility; and it indicates that sweating which is not preceded by a previous elevation of temperature is the result of exhaustion and not of fever. There is a continuous rise of the heat of the body in all cases in which a deposition of tubercle is taking place actively in any of its organs, and more especially in the lungs; while, on the other hand, I have noticed that in cancerous affections the heat of the body is but little influenced, and is sometimes even below the normal standard.

**Tongue.**—When a patient is told to put out his tongue, it is not to see whether this organ is the seat of disease, but because experience has taught that the tongue is a mirror, more or less perfect, of the condition of the digestive functions, and that it reflects the complexion of the nervous power and of the blood, and the state of the secretions. To judge of these varied circumstances, we have to examine the tongue in regard to its movements, its volume, its dryness or its humidity, its color, and its coating.

The *movements* of the tongue are impeded and tremulous in all conditions of the system attended with exhaustion. It is protruded slowly and with difficulty in fevers of a low type, and in nervous disorders which are accompanied by marked debility. The action of the muscles is seriously impaired in paralysis. In hemiplegia one side is crippled, and the tongue turns toward one of the corners of the mouth. When imperfect articulation is associated with difficulty in moving the organ, it commonly announces a serious cerebral lesion.

The *volume* of the tongue is changed by its own diseases; more rarely by the condition of the system at large, or by disturbances of the abdominal viscera. Yet a swollen or a broad and flabby tongue, on the sides of which the teeth leave their marks, is some-

times found in chronic ailments of the digestive organs, and as the result of the action of mercury, and of certain poisons. It is further observed in some affections of the brain, or as a consequence of a disturbed circulation attending diseases of the heart, and in distempers, like the plague, typhus, or scurvy, in which the blood is much altered. The tongue is sometimes observed to be swollen on one side only in consequence of catarrhal inflammation. This hemiglossitis affects the left side, and is supposed to be of neurotic origin.\* Loss of substance of the tongue, especially on its borders, is mostly due to syphilis. The ulceration is often associated with fissures.

*Dryness* of the tongue indicates deficient salivary secretion. In acute visceral inflammations, and still more frequently in febrile states, especially in the exanthemata and in typhoid fever, the tongue is dry; it may be so dry as to cause the papillæ to become prominent and the whole organ to appear roughened. This condition is one which, in acute diseases, is always to be dreaded, especially if the tongue be, in addition, of a dark color, glazy, or furred or fissured; for it is then a proof not only of generally arrested secretions, but also of depraved blood and of ebbing life-force. Yet a fissured tongue is not, by itself, indicative of great and imminent danger; it may occur in chronic affections of the liver, or in chronic inflammation of the intestines; and in some persons it is congenital. In estimating dryness of the tongue we must not overlook the fact that this may happen from persistent openness of the mouth, as during sleep, from obstruction of the nasal passages, or from coma. Among chronic diseases the tongue is most apt to be found dry in diabetes. A dry, incrusted, brown tongue is due to a continuous crust on and between the papillæ, which is filled with parasitic growths. It occurs in states of prostration with lowering of nutrition and tendency to sinking. Dickinson has calculated that a dry tongue is present in about fifty per cent. of fatal cases; more than any other it foretells death.† The opposite of dryness, *humidity*, is, unless excessive, a favorable sign. It is extremely so if it succeed dryness, because it is a proof that the secretions are being re-established.

---

\* Dyce Duckworth, *Liverpool Med.-Chir. Journ.*, July, 1883.

† *The Tongue as an Indication in Disease*, London, 1888.

There is a rare disease of the tongue known as xerostomia, occurring in women after middle life, in which the dryness of the tongue is so extreme that it may prevent speaking or swallowing. The tongue is cracked like alligator skin, and looks like raw beef.\*

The *color* of the tongue is subject to many variations. It is remarkably pale whenever the blood is watery and deficient in red globules. It is exceedingly red and shining in the exanthemata, especially in scarlet fever. The tongue is also very red if inflammation have attacked its substance, or the fauces, or the pharynx. It is bluish and livid when there is an obstruction to the flow of the venous blood or deficient aeration, as in some structural diseases of the heart and in dangerous cases of bronchitis or of pneumonia. A red, smooth tongue is a sign of failing nutrition. A tongue black in spots, the discoloration particularly marked about the middle of the dorsum, the papillæ enlarged, indicates a condition of parasitic origin.

As important as the color of the organ are the color and form of its *coating*. In health the tongue has hardly a discernible lining; disease quickly gives it one. In inflammation of the respiratory textures, at the beginning of fevers, in disorders of large portions of the abdominal mucous tract, the epithelium accumulates, and the tongue has a loaded, whitish appearance, due to excess of white epithelium on the papillæ with the intervals also more or less filled up. The coat is apt to be yellowish in disturbances of the liver, and of a brown or very dark hue when the blood is contaminated. But we must be sure, in drawing our inferences, that the abnormal aspect is not due to the food partaken of, or to medicine. Its color is also modified by the character of the occupation. Thus, as Chambers tells us, there is a smooth, orange-tinted coating on the tongues of tea-tasters. A local cause sometimes gives rise to a thick, opaque coat. For instance, decayed teeth may produce a yellow sheathing on one side. Affections of the fauces also occasion a deep-yellow hue. Again, there are many healthy persons who wake up every morning with their tongues covered, more especially at the back, with a heavy coating, which wears off after a meal.

---

\* See reports of cases in Sajous' Annual of the Univ. Med. Sci., vol. i. 1891, C-1.



In some diseases the epithelium, which is either formed in excessive quantities or not thrown off, collects between the papillæ, leaving them uncovered and prominent. This is especially noticed in scrofulous children. When the epithelium is sticky and adherent, it winds itself chiefly around the filiform papillæ, elongating them and giving to the surface of the organ a *furred* appearance. Although this kind of tongue, as almost every other variety, is met with now and then in persons who are not ill, yet it may generally be looked upon as denoting disease. It occurs sometimes in chronic diseases of the abdominal viscera, but much oftener in grave acute maladies. The tongue, on the other hand, may be bare of its epithelium or imperfectly covered with it. We meet with this in certain instances of scurvy, or in cases of chronic diarrhœa and dysentery with great prostration, in which the tongue is often found to be red, smooth, and dry, or in attendance on cachexias, as the malarial. Again, a denuded tongue is common in scarlet fever, and not infrequent in typhoid fever. In scarlet fever it has a strawberry look. This is sometimes also seen in pneumonia.

To sum up, before leaving the subject, the manifestations afforded by the tongue which are indicative of danger. They are, tremulous action; great dryness; a livid color; a very red, shining, or raw aspect; a heavy coating of a dark or black hue; a glazed, brown tongue. Any change from these to a more natural look bears a favorable interpretation.

The state of the *digestion* and the character of the discharges have so close a connection with the nutrition of the body that they become important general symptoms. But, for the sake of convenience, their value will be inquired into while discussing the diseases in the recognition of which they occupy the foremost place. A few words here, however, on the sensations of patients.

**Sensations of Patients.**—Sick persons are subject to many disagreeable feelings. They complain of chills, of heat, of languor, of restlessness, and of uneasiness; but their most constant complaint is of pain. Now, *pain* may be of various kinds; it may be dull or gnawing; it may be acute and lancinating. In its duration it may be permanent or remitting. A *dull* pain is generally persistent. It is most often present in congestions, in subacute and chronic inflammations, and where gradual changes

of tissues are taking place. It is the pain of chronic rheumatism, and shades off into the innumerable aches of this malady. The only acute affections in which it is apt to exist are inflammations of the parenchymatous viscera and of mucous membranes.

*Acute* pain is in every respect the reverse of dull pain. It is usually remittent, and not so fixed to one spot. It is met with in spasmodic affections, in neuralgia, and, with extremely sharp and lancing pangs, in malignant disease.

Pain varies much in intensity; it is sometimes so extreme as to cause death. We have to judge of its severity partly on the testimony of the sufferer, partly by the countenance, and partly by the attending functional disturbances. The latter are not to be overlooked, for they enable us, to some extent, to appreciate whether the torments are as great as they are represented to be.

The seat to which the pain is referred is far from being always the seat of the disease. A calculus in the bladder may produce dragging sensations extending down the thighs; inflammation of the hip-joint gives rise to pain in the knee; disorders of the liver occasion pain in the right shoulder. Pain felt at some part remote from that affected is either transmitted in the course of a nerve involved, or is sympathetic.

The same abnormal action does not always create the same kind of pain. Inflammation, for instance, causes different pain as it involves different structures: the pain from an inflamed pleura is not the same as that from an inflamed muscle. Speaking generally, the tissues themselves seem to determine the form of pain more certainly than does the precise character of the morbid process. Thus, pain in diseases of the periosteum and bones, no matter what may be the exact nature of the malady, is mostly boring and constant; in the serous membranes, sharp; in the mucous membranes, dull; and in the skin, burning or itching.

Pain produced by pressure is called *tenderness*. It indicates increased sensibility, and is most constantly associated with inflammation. Yet tenderness may be present without inflammation; the tenderness, for example, of the skin in hysteria. Commonly it is combined with pain occurring independently of pressure; but a part may be tender and not painful.

## CHAPTER II.

### DISEASES OF THE BRAIN AND SPINAL CORD, AND OF THEIR NERVES.

BEFORE entering upon a consideration of the affections of the nervous system it is proper to recall a few salient points connected with its structure and functions indispensable to a recognition of its derangements. We have constantly to bear in mind that there are in its composition nerve-cells composing ganglia, which are for the most part originators, and nerve-fibres, which are for the most part conductors, and, besides, a peripheral termination of these conductors, which forms a peripheral nervous system, chiefly concerned in receiving and distributing impressions. In the brain and spinal cord are the principal nervous centres which originate and control, and of the brain especially our knowledge of the subject of localization and special function of particular points has become so extended that it is made the basis of accurate diagnostic knowledge, which has of late years assumed the greatest practical importance.

#### *Cerebral Localization.*

Either for his own purposes or to co-operate with the surgeon, the physician must often satisfy himself of the exact seat of the lesion, and a knowledge of the centres in the brain is a necessity.

The methods by which, in animals, the locations of the cerebral centres have been determined have been principally those of electrical stimulation and the ablation of limited areas of the cortical gray matter. It has been thus ascertained that in the monkey the centre for the movements of the head as a whole is placed most anteriorly; extending from that part of the frontal lobe non-responsive to excitation—which is about opposite a point between the posterior and middle thirds of the superior frontal

convolution—to the arm-area, which, on the convex aspect of the hemisphere, abuts the face-area at its inferior border. The arm-area next occupies a somewhat irregularly shaped territory, comprising a space of the mesial surface directly posterior, bounded behind by the trunk-centre, and stretching upon the convex surface below the leg-area across the whole space of the ascending frontal and ascending parietal convolutions, to the interparietal sulcus. The trunk-centre upon the inner aspect occupies but a limited space, and is bounded posteriorly by the leg-centre, which is situated at the head and about the superior termination of the Rolandic fissure. The face-centres occupy the space about the lower termination of the Rolandic fissure, between the precentral and Sylvian fissures.

It has been found that these large centres are compound, that subdivisions control individual muscles or movements, and that the general order is such that upon the median aspect the centres arranged from before backward control successively the muscles of the head, shoulder, arm, trunk, leg, and feet. This is also true of the arrangement upon the convex or external surface of the hemisphere, the most anterior being those of the face. In man almost all peripheral movements have their cerebral reflex centres located with more or less exactness in certain cortical areas, and with more or less definite topographical relations to others. In correspondence with this law there seems to be another governing the position of the centres of the most specialized and differentiated muscles, whereby they are superposed upon those of the larger muscles and groups, of which they are in reality extensions, refinements, or specializations. Strong stimulus of a sub-centre will implicate a larger area and result in movement of larger related muscles.

The centres for movements of the trunk and head as a whole are not so clearly determined. But in a general way the arrangement of the cortical motor centres in man preserves the same plan and order as in the monkey. The facts in the case of man are chiefly the results of clinical and pathological investigation, and in consequence some indefiniteness still prevails.

The localization of human cortical centres is indicated in the annexed sketch. It should not be forgotten that in all such diagrammatic representations the picture represents the fact but



poorly. The two halves of the same brain are unlike. Moreover, there is never any hard and fast line dividing one centre from its neighbor. If they do not actually overlap, the centres certainly pass into one another by indefinable gradations. The strength of the stimulus, as has been stated, modifies the definiteness of limitation, and many facts go to show that the unaffected hemisphere has often a certain power of substitution, whereby it can take up the function of its injured fellow. Certain muscles, indeed, appear to be represented bilaterally in each hemisphere, whilst, on the other hand, there is, at least in the case of articulate speech, a location of the unique controlling centre singly upon one side or the other according as the person is right-handed or left-handed.

As in the monkey, the centres for voluntary motion of the opposite side of the body cluster about the fissure of Rolando. Upon the mesial surface the same order of arrangement from before backward is preserved, but there is less certainty as to the definite areas of the head and trunk. Externally, about the upper limit of the Rolandic fissure, and seated in the highest part of the motor region, is the leg-area, extending posteriorly to a somewhat indeterminate point of the superior parietal lobule, and inferiorly occupying the upper third of the ascending frontal and ascending parietal convolutions. It is almost certain that the centre for the foot is at the junction of the highest frontal and ascending frontal convolutions. The shoulder- and arm-area includes the middle third of the ascending frontal and ascending parietal convolutions, extending higher up the ascending frontal convolution.

The centre for the muscles of the trunk is supposed, in accordance with the experiments of Horsley and Schaefer, to be on the inner surface of the hemisphere. The centre for the neck region Munk and Wernicke located in the frontal lobe in front of the middle of the anterior central convolution.

In the lower third of the ascending frontal, perhaps extending to the ascending parietal, are located the centres of control of the facial movements; the centres of the lips and tongue lie at the lowest part of the ascending frontal. The centres for the movement of the head and eyes are not established. In the latter space, and extending into the posterior portion of the third left



frontal convolution, lies the centre for articulate speech, lesion of which causes motor aphasia. In connection with the subject of aphasia we may note that a separate centre is required to correspond to the clinical fact of psychic inability to write,—agraphia. Yet the location of this centre is not clearly established. There are, then, the two kinds of motor aphasia produced by lesions of the corresponding centres, which are called by some “aphemia,” or simply motor aphasia, and “agraphia.” But our complex power of thought-expression is made up of two other elements that are sensory; there must be psychical comprehension both of the heard and of the seen word. The centres intermediating these functions have been made out with some approach to definiteness. Lesions of the first temporal convolution produce word-deafness, or inability to comprehend the meaning of words though not deaf to other sounds. In the same way, word-blindness, or inability to understand the import of written or printed words, follows injury of the angular gyrus.

In reference to the cortical visual centre there can be little doubt that it is located in the occipital lobe, and especially in the cuneus. The production of hemianopsia from lesions of the occipital lobe, in accordance with the conclusions of Seguin,\* is shown in the accompanying diagram (Fig. 10.) Complete cortical blindness may be considered as a bilateral hemianopsia. The macula is also represented in the cortex. Dimness of sight in the opposite eye, with, as a rule, concentric diminution of the field, or crossed amblyopia, depends upon a lesion in the angular gyrus.

The centres for audition, smell, and taste are not absolutely determined. The auditory centre is most likely in the middle of the first temporo-sphenoidal convolution and related to the auditory nerve of the opposite side. The centre for smell is very probably on the medial surface of the temporal lobe at the anterior extremity of the uncinate convolution and in connection with the auditory nerve of the same side. The cortical centre for taste is most uncertain. The location of the centres of tactile or cutaneous sensation is also in dispute, but it appears probable that, if not identical with, they are at least contiguous to those of the motor functions of corresponding parts.† A geographical centre,


---

\* Journ. Nerv. and Ment. Dis., 1886, No. 1, and Nov. 1887.

† See Dana, Journ. Nerv. and Ment. Dis., Oct. 1883.

a centre for determining locality, is claimed to have its seat in the occipital lobe, near the visual centre. The psychical, or mental, processes have as their centres those parts of the cortex that have

FIG. 10.



**RIGHT HOMONYMOUS OR LATERAL HEMIANOPIA, FROM LESION OF THE LEFT VISUAL CENTRE OF THE CORTEX OR LEFT OPTIC TRACT.**—A, dark left nasal half-field from blind temporal half of retina; A', dark right temporal half-field from blind nasal half of retina; B, left eye; B', right eye; C, C', left and right optic nerves, composed of the crossed bundles of fibres; D, D', left and right crossed bundles; E, E', left and right occipital lobes; F, F', left and right posterior cornua; G, G', "optic radiation" of Gratiolet; H, H', optic chiasm; I, I', angular gyrus; K, region of optic thalamus, geniculate body, and quadrigeminal bodies, collectively termed the primary optic centres; M, M', cuneus of the occipital lobe, the cortical visual centre. The left cuneus and optic tract are shaded, to show lesion of these parts and the influence of the lesion upon the retinae.

not been found to possess any special motor or sensory function, and particularly the prefrontal lobes.

It is often a matter of much importance, especially with reference to brain surgery, to determine on the skull the seat of the underlying cerebral centres. Broca, Horsley, and Reid have specially investigated the subject, and from their and other researches we are sure of these facts:

Under the frontal bone lie almost the entire frontal, middle, and about three-quarters of the upper frontal convolutions. The temporal bone covers the temporal lobe, except its anterior extremity and its posterior fifth. The occipital bone covers the



greater part of the occipital lobe; the remainder of the cortex is beneath the parietal bone. The ascending frontal convolution starts somewhat lower than beneath the anterior inferior angle of the parietal bone in front of the prolonged line of the fissure of Rolando. In front of the precentral sulcus, the lower half of which is parallel to and behind the coronal suture, lies the root of the lower frontal; the root of the ascending parietal is behind the ascending frontal. The upper end of the fissure of Rolando corresponds to a point half an inch behind the middle of a line measured from the root of the nose upward to the occipital protuberance, and the fissure extends obliquely downward and forward, at an angle of  $67^\circ$ , to within a short distance of the fork of the Sylvian fissure. The fissure between the middle and lowest frontal convolutions is under the temporal ridge. The central convolutions are about an inch on each side of the fissure of Rolando; the centres for the leg, arm, and face lie on each side of the fissure. The angular gyrus is immediately behind the most prominent portion of the parietal eminence. The first temporal convolution is over the ear and mastoid process below the Sylvian line. This line is determined by drawing a line from the external angular process of the frontal bone to a point three-quarters of an inch below the most prominent part of the parietal bone.

### *Sensory Centres, and Conducting Paths.*

The sensory centres and the conducting paths by which the fibres unite the various parts of the brain, whether sensory or motor, and of the spinal cord, are not so definitely made out as the brain-centres are; particularly uncertain are we as to the course of the sensory paths in the medulla, pons, and peduncle.

The sensory centres for the muscular sense and the sense of touch are supposed by Horsley and others to be in layers of cells in the motor cortex. But the centres for sensory impression are also claimed to be the hippocampal convolution and the gyrus fornicatus, and, generally, the occipital and temporo-sphenoidal lobes.

Volitional impulses originate in the motor cortex, and pass by converging fibres through the white substance of the hemisphere to the internal capsule, thence beneath the optic thalamus, to enter the crus cerebri, and through the pons, reaching the medulla, where the larger number of fibres cross to the opposite

side of the cord to form the lateral or crossed pyramidal tract. The smaller fibres that continue onward form the anterior or direct pyramidal tract; these decussate in the cord at various levels. This constitutes the upper segment of the motor path of Gowers, which terminates in the ganglion-cells of the anterior horns of the cord. The lower segment consists in the fibres that originate in the efferent processes of the ganglion-cells and pass to their peripheral distribution in the muscles.

The fibres for the so-called cranial nerves leave the pyramidal columns as they approach the level of their nuclei on the opposite side of the medulla, to reach which they cross the median line somewhat in advance of the decussation of the remainder of the pyramidal tracts.

A lesion in any part of the upper segment of the motor path, between the cortical cells and the ganglion-cells of the anterior horns, is followed by descending degeneration in the pyramidal tracts. The resulting paralysis is attended with increased reflexes, unchanged electrical reactions, and little or no wasting of the muscles. A lesion in any part of the lower segment, between the gray matter in the cord and the terminations of the nerves in the muscles, gives rise to paralysis characterized by wasting, qualitative electrical changes, and impairment or abolition of the reflexes.

Sensory impressions reach the brain through the posterior roots of the cord, passing by the posterior and lateral columns in several tracts, most of which decussate in the cord.

There is reason to believe that the paths for common tactile impressions, for painful impressions, for the conveyance of thermal impressions, and of the muscular sense, are distinct; that for the first coursing through the posterior column, those for the second and third through the antero-lateral ascending tract, and those for the last through the postero-median column and the direct cerebellar tracts.

Lesions of the peripheral sensory segment are attended, in addition to the impairment of sensibility, with abolition of the related reflexes. Lesions of the cord involving the posterior and lateral columns are attended with ascending degeneration in the postero-median and postero-external columns, the direct cerebellar and the antero-lateral ascending tracts.

The sensory fibres for the muscular sense are supposed not to decussate in the cord.

*Spinal Localization.*

A centre for spasm is thought to be in the medulla at its junction with the pons, and is carried by the vagus ; the cardio-inhibitory centre is in the medulla ; the respiratory centre is in the medulla between the nuclei of the vagus and accessories ; the vasomotor centre is in the medulla ; so is the sweat centre in the medulla, with subordinate spinal centres.

The following facts will prove useful in localizing or determining the extent of a lesion of the spinal cord : Paralysis of the small rotators of the head and of the depressors of the hyoid bone points to involvement of the first and second cervical nerves ; paralysis of the levator anguli scapulæ to involvement of the third cervical ; paralysis of the sterno-mastoid, of the upper neck-muscles, and of the upper part of the trapezius to involvement of second, third, fourth, and fifth cervical ; paralysis of the diaphragm to involvement of the fourth and fifth cervical ; paralysis of the serratus, flexors of the elbow, and supinators of the forearm to involvement of the fifth and sixth cervical ; paralysis of the shoulder-muscles to involvement of the fourth, fifth, and sixth cervical ; paralysis of the extensors of the wrist and fingers to involvement of the sixth and seventh cervical ; paralysis of the extensors of the elbow, of the flexors of the wrist and fingers, and of the pronators of the forearm to involvement of the seventh and eighth cervical ; paralysis of the lower neck-muscles and of the middle part of the trapezius to involvement of the sixth, seventh, and eighth cervical and first dorsal ; paralysis of the muscles of the hand to involvement of the eighth cervical and first dorsal ; paralysis of the intercostals to involvement of the dorsal nerves from the first to the tenth ; paralysis of the lower part of the trapezius and of the dorsal muscles to involvement of the dorsal nerves from the second to the twelfth ; paralysis of the abdominal muscles to involvement of the dorsal nerves from the seventh to the twelfth, and also the first lumbar ; paralysis of the cremaster and flexors of the hip to involvement of the second and third lumbar ; paralysis of the extensors of the knee, of the adductors, extensors, and abductors of the hip to involvement of the fourth and fifth lumbar ; paralysis of the lumbar muscles to involvement of the second, third, fourth, and fifth lumbar nerves ; paralysis of



the peroneus longus, the flexors and extensors of the ankle to involvement of the fourth and fifth lumbar and first sacral nerves; paralysis of the flexors of the knee to involvement of the fifth lumbar and first sacral; paralysis of the intrinsic muscles of the foot to involvement of the first and second sacral; paralysis of the perineal and anal muscles to involvement of the third and fourth sacral nerves.

Loss of sensibility on the scalp points to involvement of the first, second, and third cervical nerves; on the neck and upper part of the chest to involvement of the second, third, fourth, and fifth; on the shoulder to involvement of the fourth and fifth; on the outer aspect of the arm to involvement of the fifth and sixth; on the radial aspect of the forearm and hand and on the thumb to involvement of the sixth and seventh; on the inner aspect of the arm, on the ulnar aspect of the forearm and hand, and on the tips of the fingers to involvement of the seventh and eighth cervical and first dorsal; on the front of the thorax to involvement of the dorsal nerves from the first to the tenth; over the ensiform cartilage to involvement of the sixth and seventh dorsal; on the abdomen to involvement of the dorsal nerves from the seventh to the twelfth, and also the first lumbar; at the umbilicus to involvement of the tenth dorsal; on the upper part of the buttock to involvement of the twelfth dorsal and first lumbar; in the groin and on the scrotum to involvement of the first and second lumbar; on the outer, anterior, and inner aspect of the thigh to involvement of the second, third, fourth, and fifth lumbar nerves; on the inner aspect of the leg to involvement of the fifth lumbar; on the lower part of the buttock, on the posterior aspect of the thigh, and on the anterior, posterior, outer aspect of the leg and foot to involvement of the fifth lumbar and the first, second, and third sacral; on the perineum and about the anus to involvement of the third, fourth, and fifth sacral; and on the skin between the coccyx and anus to involvement of the fifth sacral and the coccygeal.

Loss of the scapular reflex points to involvement of the fifth, sixth, seventh, and eighth cervical and first dorsal nerves; of the epigastric reflex to involvement of the fourth, fifth, sixth, and seventh dorsal; of the abdominal reflex to involvement of the dorsal nerves from the eighth to the twelfth, and also the first



lumbar nerve ; of the cremaster to involvement of the first, second, and third lumbar ; of the knee-jerk to involvement of the second, third, and fourth lumbar ; of the gluteal to involvement of the fourth and fifth lumbar and the first sacral ; of ankle-clonus to involvement of the fifth lumbar and first sacral ; and of the plantar to involvement of the first, second, and third sacral nerves.

Let us now look at the derangements of the nervous system. But first let us examine a few symptoms and morbid states having a general significance rather than a specific connection with any malady.

#### DERANGED INTELLECTION.

The great instrument of the intelligence, the brain, manifests its ailings, whether primary or merely sympathetic, by derangement of thought of every conceivable degree and kind,—from dulness and confusion of the intellect to its utter perversion and prostration. When one intellectual function is disturbed, generally all are, or soon become so ; yet we may find impairment of judgment and of imagination without deterioration of memory or of the powers of attention. One of the most marked signs of mental infirmity is a disordered memory. This is especially encountered in chronic cerebral diseases, or in such nervous affections of uncertain seat as epilepsy. Another signal of mental derangement is loss of judgment, or rather loss of power to appreciate the logical sequence of ideas ; still another is depression of mind, or its opposite, exaltation. All these abnormal conditions may happen in acute as well as in chronic maladies, but they are more striking in the latter, and become of more aid in the diagnosis ; and they may or may not be joined to appreciable textural changes. To the psychologist their significance is very great, as they are often the premonitory symptoms of that departure from mental health which terminates in confirmed insanity.

In acute disturbances of the brain, whether functional or organic, we meet with these striking phenomena connected with disordered intellection ; delirium, stupor, coma ; and with these we may consider insomnia.

**Delirium.**—This is a wandering of the mind, manifesting itself by the expression of ill-associated thoughts, of the incongruity of which the patient is not conscious. It most frequently occurs in those of susceptible nervous system, and is, in consequence, more

common in the young than in the old. It is almost invariably united with restlessness, and increases as night approaches.

The character of the delirium is various. There is first the *quiet* delirium, of a low or passive type. The patient mutters incoherent words, moans without any assignable reason, or lies silent, with his eyes open, his thoughts preoccupied with his vague illusions, and taking no notice of what goes on around him in the external world. If strongly aroused, he gives a rational answer, but not a long or a connected one, for he soon returns to his dreams and his ever-changing hallucinations. He picks at his bedclothes, moves in bed, and may even try to leave it, although he is easily prevented from so doing.

Then there is a delirium of somewhat more active type, still, on the whole, quiet; the patient wanders, yet not boisterously. He is irritable, and often does not show that his mind is disturbed, except in some one particular,—in irascibility about trifles, or in expressions and modes of thought foreign to his nature.

An active, *fierce* delirium presents different characteristics. The patient is wild, noisy; he sings, screams, gets out of bed; his face during the excitement becomes congested; the eye is bright, often fiery.

Now, all these forms of delirium occur in many different maladies, and are far from being of necessity linked to an organic cerebral affection. Nay, not even the most violent kind of mental wandering is positively indicative of a lesion of the brain; at least, not of such a lesion as can be determined by any of our present means of investigation. As a rule, we find the low, quiet delirium in conditions of vital exhaustion, particularly in those depressed states of the nervous system which are connected with quickened vascular action, and with a deterioration of the blood, as, for instance, in the low fevers. The fierce delirium may, however, be associated with prostration or depraved blood. Thus, the delirium of pneumonia is sometimes of a violent kind, owing to the maddening effect of the ill-oxygenated blood on the brain. In most of the ordinary fevers the delirium is of a moderate type; in inflammatory diseases of the brain and in acute mania it is fierce. The delirium of uræmia is apt to be active. If the delirium be due to cerebral disease, it is associated with headache; the headache of pyrexia generally disappears with the onset of delirium.

Delirium is not difficult of recognition ; yet we must be careful not to confound with it *night terrors*, those troubled dreams to which ailing children are so liable, and which occasion confusion of thought on first awaking, and until consciousness is fully aroused. Delirium is most likely to be mistaken for *insanity*. There is this palpable difference : an insane person is commonly in good health in all save his intellect ; a delirious person is ill, and exhibits evidences of his illness besides his delirium. It is true that, when the patient is first seen, doubt may arise ; but it is not generally of long duration. In the mania appearing occasionally after epileptic fits, or taking their place, there may be doubt until we obtain a clear history. Most perplexing are the cases in which insanity follows or attends inordinate drinking. But this is a subject which we shall discuss in reviewing mania a potu.

Another perplexing group of cases is furnished by the occurrence of that singular form of delirium which is met with at times in acute diseases, especially in fevers, and which, as it is apt to be associated with insufficient nutrition, has been called the *delirium of inanition*, or of collapse.\* Its outbreak is sudden, like an attack of mania, but it is found to be combined with a feeble pulse, with a skin bathed in perspiration, with cold hands and feet,—in a word, with the signs of great prostration or of collapse. The seizure happens usually early in the morning, and is unexpected, for it occurs commonly at the end of the febrile state, and when the condition of the skin and pulse bespeaks convalescence. The exhausted nervous centre betrays itself in the sudden mental wandering, which has generally this characteristic,—there is but one fixed delusion, and this one connected with the subjects which have most engrossed the mind before the illness. The seizure lasts from six to forty-eight hours, and at its termination the patient is apt to awake out of a sleep with a calm mind, remembering, perhaps, his hallucination as a vivid dream. There may be more than one attack, but this is not common ; and the duration is materially abridged by opium and by the employment of stimulants and nourishment. The form of delirium

---

\* See Weber, *Medico-Chirurg. Transact.*, 1865 ; Becquet, *Arch. Gén. de Médecine*, 1866 ; also the *Clinical Lectures of Chomel and of Trousseau ; Nothnagel, "Anæmia of the Brain," in Ziemssen's Cyclopædia.*



under consideration is not simply a sequel of febrile conditions. It may also succeed exhausting discharges and drains from the system, or inability to obtain or to digest the proper amount of food. Thus, it may happen in malignant diseases of the stomach; also in mere gastric irritability and persistent vomiting. The most marked instance of this kind of mental wandering I have encountered was associated with functional gastric disorder, which prevented enough food from being retained. In this patient the hallucination was on one subject,—a business matter which had been annoying him greatly just before his illness became decided.

Delirium is at times *simulated*. This differs from real delirium by the absence of all other signs of illness, and by the sameness of the mental wandering. In a case of feigned delirium I met with, the man whined when spoken to, and pretended to rave; but his ideas always ran on the same subject, and he was very solicitous about his food, and about other matters of which a delirious person takes no notice. Delirium is more or less continuous; once delirious, a patient remains so for some time, and until the exciting cause subsides. In this respect hysterical delirium is exceptional; it does not last long, or it intermits and then reappears.

**Stupor.**—A blunted state of mind, a partial, drowsy unconsciousness, constitutes the phenomenon called stupor. The patient lies in a deep slumber, from which he cannot be roused save with great difficulty, and when roused he answers reluctantly and briefly, and soon resumes his heavy sleep. The expression of his face is dull, yet now and then a ray of intelligence, excited by some object which attracts his attention or by some pleasant reverie, flits across his features.

Stupor is met with in several cerebral affections, and seems to be chiefly owing to a congestion of the brain. It is frequently seen in typhoid fever, immediately after an epileptic fit, or as the result of narcotic poisons, and is, in these states, also probably due to cerebral congestion. It may further be due to sedative drugs. But there is nothing pathognomonic about it in these various conditions, nothing by which we can judge positively of its origin.

**Coma.**—Coma is complete loss of consciousness: perception and volition are alike suspended, and there is an appearance



of the profoundest sleep. The face wears a confused look ; the pupils are sluggish, often dilated ; the mouth is open, the tongue dry. Sensation may be blunted, but is not destroyed ; nor is motion, for the patient moves when his skin is pinched or tickled.

Coma always betokens a serious disturbance of the functions of the brain. It is often witnessed in cerebral lesions, as from pressure from blood or fluid in brain-substance or in ventricles, more rarely from tumors, abscesses, or thrombosis. The most complete coma is seen in apoplexy ; it comes on quickly, and is attended with noisy respiration and a slow pulse. Another form of coma, scarcely less complete, is caused by narcotic poisoning ; it, however, does not appear suddenly, and when from opium is associated with contraction of the pupils. Profound intoxication with alcohol induces coma, but the attendant symptoms, as a rule, make the association clear. The coma of fevers and of acute diseases, whether cerebral or not, is also gradually produced, but, unlike that due to the toxical effect of opium, is ordinarily preceded for days by insomnia, by delirium, and by other signs of cerebral disturbance. The coma of epilepsy is recognized by its following epileptic seizures. In Bright's disease, among the nervous phenomena, of which coma as well as stupor and delirium may happen, the loss of consciousness is apt to occur subsequently to either of the two other morbid phenomena, and its cause is made manifest by finding albumen and tube-casts in the urine, and by the general evidences of uræmia. Uræmic coma may, however, come on suddenly and pass off suddenly. It is, as a general rule, associated with low temperature and dilated pupils. Coma also sometimes occurs in connection with diabetes. Under such circumstances examination of the urine will reveal the presence of sugar, perhaps also of acetone and of diacetic acid.

Sometimes a person appears to be comatose when his intellect is but little disordered. He may be paralyzed, and not have the power to communicate his ideas, from crippled articulation or in connection with aphasia. This state is distinguished from coma by noting that the patient's attention is always directed to the questions asked him, nay, that he strives to answer them, but cannot ; and that generally he has lost control over the muscular movements of one side or of both sides of the body. Coma must not be confused with *syncope*, which depends upon cerebral an-

æmia, and is usually of brief duration, and, except feeble heart action, is unattended with noteworthy symptoms.

**Insomnia.**—The deprivation of sleep is a concomitant of cerebral congestion and of the earlier stages of cerebral inflammation. But a person may be sleepless from excessive pain, from exhaustion, from grief, from mental excitement or fatigue, or from the too free use of coffee or of tea; sometimes insomnia is engendered by habitually working late at night. However, in several of these states congestion, of active or passive character, is, in all likelihood, the immediate cause of the wakefulness.

Insomnia often precedes or attends delirium, as appears in typhoid fever. Among purely nervous affections it is most marked in delirium tremens. It is a very troublesome symptom; but, occurring in so many abnormal conditions, it cannot be looked upon as having a distinct and specific diagnostic value.

#### DERANGED SENSATION.

The signs of perverted or impaired sensation are numerous. They may either be due to an alteration of the general sensibility or be the signals of a derangement of a nerve of special sense. Let us look at a few.

**Hyperæsthesia.**—An exalted sensibility of surface nerves,—of those of the skin, the mucous membranes, or even of those of deeper-seated structures,—in other words, a hyperæsthesia of these parts, is a symptom of importance; not so much, perhaps, on account of the light thrown by it on any particular disease, as because its presence makes it requisite to determine its origin and to separate its phenomena from those of inflammation. We may, as a rule, distinguish the peripheral sensitiveness from the tenderness of subjacent inflammation, by its extension over a larger surface; by deep pressure producing no more pain than a light touch; by the absence of signs of functional disturbance of the part involved apparently in inflammatory disease; by the uniformity of the painful sensation, no matter how long the duration of the disorder, though the sensitiveness exhibits distinct intermissions and exacerbations.

Hyperæsthesia is not closely connected with organic diseases of the brain or of the spinal cord. Indeed, it is in them not common, nor, as a rule, highly developed. By far the most usual causes

of hyperæsthesia are impoverished blood and hysteria; therefore conditions which bespeak lowered vital and nervous power. Sometimes hyperæsthesia is produced by rheumatism or by gout, by lithæmia, or by disturbance of the function of the kidney. It is further met with in epidemic influenza; in hydrophobia; in inflammations in internal cavities involving the ganglia of the great sympathetic; after the use of ergot and of opium; and in some diseases of the skin. It also attends paroxysms of neuralgia, as witnessed in the exquisite sensitiveness of the skin during an attack of *tic douloureux*; the painful spots, too, in the course of local neuralgias are, when not the result of neuritis, hyperæsthetical. Hyperæsthesia attends the irritative stage of inflammation of sensory nerves from whatever cause.

The seat of the heightened sensibility is ordinarily in the skin, in the distribution of the cutaneous nerves. Yet hyperæsthesia may affect the nerves of the special senses, manifesting itself, for instance, by intolerance of light or of sound. But this variety of hyperæsthesia need here be but alluded to, as we shall presently look more fully at the signs of disturbance of these nerves. Of the minute anatomical changes in hyperæsthesia we know nothing. The physiological basis for the increased sensation may be either in the peripheral nerves, or in the irritability of a cerebral centre or of the conducting fibres of the spinal cord, especially of those of the posterior columns. When a painful sensation is more acutely felt than normal, it is called "hyperalgesia." Sensibility to pain is most readily tested by a pinch or a prick, or by a wire brush with the faradaic current.

Let us now look at hyperæsthesia in connection with affections of the nervous system, especially with those of the brain and cords.

*Hyperæsthesia is general and combined with signs of organic disease.*—We find this in tumors pressing upon the pons Varolii and corpora quadrigemina, or in alterations or injuries of the posterior columns of the cord and those producing irritation in the course of the conducting fibres, in some cases of cerebral meningitis, and in spinal meningitis in which the posterior nerve-roots are implicated. We have in all these conditions a hyperæsthesia more or less extensive, and combined with other striking evidence of nervous disease, often with pain. But, in making up our minds as to the cause of the extended hyperæsthesia, the sen-



sitiveness in diffuse neuritis, in general neuralgias, and in reflected irritation to the posterior columns, especially in hysterical subjects, must always be remembered.

*Hyperæsthesia is limited to one side.*—An injury, or, at its beginning, a degeneration of only one posterior column, will give us increased sensibility on the same side as the lesion. Limited hyperæsthesia belongs much more closely to spinal than to cerebral disease. We also find it in connection with special neuralgias, and the sensitive skin shows augmented electrical sensibility. In some instances of limited as well as of more extended hyperæsthesia nothing abnormal can be detected, and the disorder must be, with our present knowledge, set down as a neurosis, one concerning which it remains uncertain whether it be of central or of peripheral origin.

**Anæsthesia.**—Loss of sensation, or anæsthesia, is of various degrees. It may be complete or partial,—a perfect absence of sensibility, or its mere benumbing. Not to speak of its meaning when displaying itself only in the organs of the special senses, we find it in diseases of the brain; in several of the neuroses; after large doses of Indian hemp, of lead, of arsenic; we see it ushering in attacks of neuralgia; accompanying or preceding cutaneous eruptions, such as elephantiasis or pemphigus; in hysteria, in syphilis, in rheumatism; and as the result of diphtheria, of pressure on nerve trunks, of peripheral neuritis or nerve irritation, and of disturbances of circulation and faulty blood conditions. In the mucous membranes, too, it may exist, in consequence either of the general causes just mentioned, or of some purely local irritation; and it may affect the muscles. But it does not attack these structures nearly as often as it does the skin: indeed, when we speak of anæsthesia without qualifying it, we mean that of the cutaneous nerves. In the parts affected with anæsthesia the nutrition is less active, and there is a feeling of numbness. The temperature is diminished, and, if the impaired sensibility be at all general, the patient is not susceptible to alternations of heat or cold. Frequently the circulation in the skin is retarded, occasioning a perceptible lividity and discoloration of the surface; or there are coexisting trophic changes, such as glazing of the skin, and grayness of hair. The electrical sensibility is diminished, as is made very manifest by the use of the wire brush with either



the faradaic or the galvanic current. In hysterical anæsthesia this is a particularly striking feature.

Loss of sensation has a much more constant connection with organic affections of the nervous centres than increased sensibility, which, however, may precede it. In the insane, especially in monomaniacs, anæsthesia is common, and ordinarily very extended: so, too, in general paralysis. Indeed, with few exceptions, an *extended* anæsthesia points to an affection of the nervous centres. It may in these organic cases be both general and very complete.\* *Localized* anæsthesia may usher in acute attacks of cerebral disease, and sometimes exists for years before any marked cerebral symptoms are perceived. Thus, a case of apoplexy was observed by Andral† in which deficient sensation was noticed at various portions of the thorax for a long time previous to the loss of consciousness; another in which the tips of the fingers were benumbed, and felt continually as if they had been subjected to intense cold. Forbes Winslow‡ mentions instances in which circumscribed conditions of impaired sensation were the premonitory symptoms of softening of the brain; the defective feeling being manifested in some cases in the skin, in others in the tongue and fauces.

If the defective sensibility be owing to a spinal malady, it is generally found in the lower extremities, and coexists with paralysis. Anæsthesia of spinal origin is usually indicative of the sensory conducting paths, principally in the posterior columns, having been disturbed or altered; when about the body, as in transverse dorsal myelitis, there is mostly present a zone of hyperæsthesia above the zone of anæsthesia. A limited area of anæsthesia, Allen Starr§ has demonstrated, is caused by a limited lesion in the spinal cord, and the situation and shape of the area of anæsthesia tell us the level of the lesion. In hysterical paraplegia, in paraplegia from hypnotic suggestion, or that following railroad or other injuries, the line of lost sensibility is, as Charcot|| has shown, very significant; it excludes the genital organs.

---

\* As in a case reported by Winter, quoted Schmidt's Jahrb., 1883, No. 1.

† Clinique Médicale, tome v.

‡ Obscure Diseases of the Brain, p. 549.

§ Am. Journ. Med. Sci., July, 1892.

|| Œuvres complètes, iii.

In accordance with the well-known law of the decussation of sensitive impressions in the cord, disease, if only of one posterior half, is followed by lost sensation on the opposite side of the body. One-sided anæsthesia, affecting even the face up to the middle line, is sometimes met with in hysterical subjects as the result of ovarian irritation, or after typhoid fever,\* and, though presumably cerebral, the pathology is unsettled. Hysterical hemianæsthesia is generally on the left side, and associated with tenderness over the ovary. But strictly-limited one-sided anæsthesia is more apt to be found in a distinct brain lesion, and the particular affection occasioning the "hemianæsthesia" is disease of the white substance just outside of the optic thalamus, of the posterior part of the internal capsule, on the side of the brain opposite to the side of the body which shows the anæsthesia, or damage to the fibres which conduct sensation through the pons or the crus. A lesion involving the upper part of the pons may give rise to "crossed anæsthesia,"—namely, loss of sensibility upon the same side of the face and upon the opposite side of the body. Hemianæsthesia is a not uncommon symptom between the attacks of hystero-epilepsy. Complete hemianæsthesia may be also associated with functional palsy.†

Whatever the clinical association, the insensibility is generally complete as to touch, pain, temperature, and electricity. Taste, smell, and hearing are also abolished on the one side, and the eye on the anæsthetic side loses its acuteness of vision and of perception of color. Color-blindness is complete or partial;‡ the degree of deafness corresponds with that of the cutaneous insensibility.§

A localized form of anæsthesia happens now and then in consequence of an affection of the fifth nerve. The extent of loss of sensation depends much upon the part of the nerve at which the cause of disturbance is seated. The skin of the nose and cheek may become devoid of sensation; the reflex movements of the muscles of the face may cease; the conjunctiva, or the whole surface of

---

\* Calmet, *Bulletin de la Société Médicale des Hôpitaux*, 1876.

† Wilks, *Guy's Hospital Reports*, 1883.

‡ Féré, *Archives de Neurologie*, Nos. 8 and 9, 1882.

§ Walton, *Brain*, January, 1883.

the eye, or one-half of the tongue, may be deprived of sensibility. Only one of these phenomena, or all conjointly, may be encountered, according as part of one, or one, or all of the branches of the fifth nerve are affected. Sometimes, as Romberg proves, *trigeminal anæsthesia* is of rheumatic origin. When it is complicated with disturbed functions of adjoining cerebral nerves, it may be assumed that the cause is seated at the base of the brain.

Anæsthesia is sometimes the result of *reflex* action. It may thus arise in disorders of any of the viscera, and from an irritation of any sensitive nerve. It has, for instance, been observed in both lower limbs in sciatica.\* But in many of these instances of supposed nerve irritation there is really a neuritis.

Diminished or lost sensibility to touch generally goes hand in hand with diminished or lost sensibility to pain, but the sensibility to pain may be augmented. This “*anæsthesia dolorosa*” is most commonly met with in multiple neuritis, and in spinal meningitis and myelitis from pressure.

Very often numbness and other altered sensations are complained of, and yet the whole is subjective; when tested, anæsthesia is not found. In endeavoring, indeed, to form an opinion of the existence or the completeness of anæsthesia, we do not trust to the patient's statements. We touch the part lightly with the finger or a feather while his eyes are shut, and the skin is pinched or a pin used to ascertain the extent of the impaired sensation. Or we resort to means by which we can make accurate comparisons; and one of the best is to pursue the method employed by Weber, which consists in determining how closely the points of a pair of compasses sheathed with cork may be approximated on the skin and yet be felt as two distinct points. An instrument for the same purpose, called the “*æsthesiometer*,” was invented by Sieveking (Fig. 11), and is very much the same as the lighter one of Brown-Séquard now in common use. An instrument combining the principle of the beam compass with that of the mathematical one has been contrived by Ogle,† and one with ivory points, by Manouvriez.‡ The points of the *æsthesiometer*,

---


\* Brown-Séquard, Central Nervous System, Tenth Lecture.

† Beale's Archives of Medicine, vol. i.

‡ Archives de Physiologie, 1876.

whether blunted or sharp, should be put down lightly and simultaneously, and parallel with the direction of the cutaneous

FIG. 11.



The aesthesiometer.

nerves ; at all events, the same relative direction should be preserved in making comparative estimates.

To understand any results obtained regarding the tactile sense, it is necessary that we should be aware how this differs in some parts of the body. Most works on physiology contain an account of the researches of Weber and of those who have prosecuted the inquiry he started ; yet a few of the conclusions may be here advantageously mentioned. At the tip of the tongue two points can be readily distinguished when separate from each other only  $\frac{1}{22}$  of an inch, or half a Paris line, 1.5 mm. ; at the palmar surface of the third phalanx the limit is one line ; on the palmar surface of the second phalanx, two lines, the same on the red surface of the lips ; on the palm of the hand, the cheek, and the extremity of the great toe, five lines ; on the back of the hand, at the knuckles, eight lines ; at the lower part of the forehead, ten lines ; on the skin over the patella and the dorsum of the foot, eighteen lines ; over the middle of the arm, the thigh, and over the spine, thirty lines ; on the back, 60 mm. is common. But these observations are found to vary somewhat even in healthy persons, some being able to distinguish at a shorter distance than others.

Besides the impairment or loss of tactile discrimination, the altered sensibility may show itself in the loss of the faculty of feeling pinching, pricking, and other acts which excite pain, "analgesia ;" or in insensibility to tickling ; or in the want of appreciation of heat or cold ; or in the loss of the sensation which attends muscular contraction, whether produced by the will or by an



electrical current. Now, it is of interest in individual cases to note which particular kind of sensibility is affected, though, as yet, we are not in possession of sufficient facts to draw, from the absence of one form of sensibility or the other, any positive conclusions as to the seat or character of the disease.

In affections of the base of the brain the patient feels three points instead of the two of the æsthesiometer.\* In sclerosis of the cord the sensation is retarded rather than lost.† A form of perverted sensibility, which may or may not be associated with anæsthesia, consists in the sensibility being more or less perfect, while there is doubt as to the side touched; indeed, the touch is commonly felt at a corresponding part of the other limb. This *allochiria*‡ is generally found in association with organic spinal disease; but it may also manifest itself in hysteria. A sufficient explanation of the erroneous reference of impressions is wanting. In a case recorded by Ferrier§ the reversal showed itself also in the reflex reactions. Tickling the sole of one foot caused retraction of the other; tickling the inside of one thigh produced flexion of the other. Occasionally a single sensory impression is perceived as two or more; this is known as “polyæsthesia.”

Sensibility to temperature has a close connection with sensibility to pain; but not always. There may be crossed paralysis of the thermal sense, while other senses are undisturbed.|| Sometimes the temperature sense is exaggerated or diminished, or much perverted, and cold objects feel hot, and the reverse. Then points may be found in the skin where only cold, others where only heat, is appreciated. To test heat, a heated spoon or a test-tube filled with hot water is the readiest means; to test cold, a sponge that has been dipped in cold water or a piece of ice is best.

*Muscular anæsthesia* has been mentioned. It is closely connected with the power we possess of estimating weight, the “muscular sense;” and the loss of ability of perceiving differences in small weights, or the impairment of the sense of muscular movement and effort, is its most common form. It is really distinct from the sensitiveness of the muscles to pressure or to electrical stimulation, which may be also wholly wanting. The loss of the

\* Brown-Séquard, Archives de Physiologie, t. i. No. 3. † Vulpian, *ibid.*

‡ Obersteiner, Brain, July, 1881.

§ Brain, October, 1882.

|| Case reported by Weir Mitchell, Trans. Assoc. Am. Phys., vol. vii. 1892.

power of appreciating muscular contraction, as well as the deficiency of sensation, is most readily tested by the use of the faradaic current; the contraction of the muscles produces no feeling.

Muscular anæsthesia is frequently combined with inability to determine the posture of a limb when the eyes are closed; it may or may not be associated with cutaneous anæsthesia. It is not uncommon in hysteria and in locomotor ataxia. Here the loss of the appreciation of the position of the limbs and of the sense of muscular effort are the usual varieties. When the muscles are completely paralyzed, the muscular sense cannot be tested. The muscular sense has been localized by Allen Starr and McCosh at the junction of the superior and inferior parietal convolutions, behind the posterior central convolution.\*

In testing for the muscular sense, the eyes of the person on whom the test is made should be kept closed, and objects used should be of uniform size. To detect the difference in weight, and thus the resistance to contraction, Gowers † recommends leather balls containing weights from two drachms to two pounds. The weights are placed in a bag, suspended by a string to the parts to be tested.

**Paræsthesia.**—This is a perversion of sensation, not an exaltation. It does not disclose itself by pain and tenderness, but in itching, in formication, in unnatural feelings of various kinds, such as the feeling of tingling, of pins and needles, of goose-flesh, of thrilling, of flushing, of the trickling of cold water, of shock-like sensations, of a sense of tightness, as in the girdle pain. It is generally purely subjective, though it may be influenced by touch. A form of paræsthesia is *acroparæsthesia*. This is chiefly characterized by numbness of the extremities. It is encountered in women at the menopause, and in those who do washing, scrubbing, sewing or other needlework. It may be also found in men, and by some is believed to be a neurosis, by others a neuritis.‡

The alterations of sensibility discussed manifest themselves chiefly in connection with external impressions. Let us now look at some abnormal sensations which are not objective, but subjective,—arising independently of external impressions. Headache and vertigo are of this character.

---

\* Amer. Journ. Med. Sci., Nov. 1894.

† Diseases of the Nervous System, vol. i., 2d ed., 1893.

‡ Sinkler, Medical News, Aug. 1894, p. 178.

**Headache.**—In every case of headache we must first ascertain that the pain really originates within the cranium, and that it is not owing to supra-orbital neuralgia ; to rheumatism of the scalp ; to disease of the bones ; to periostitis, syphilitic or otherwise ; or to affections of the ear. To accomplish this is generally not difficult. An inquiry into the history of the case, the locality of the pain, and its augmentation on pressure in most of the disorders named, furnish evidence which decides the source of the cephalalgia to be external to the cranium.

Another possible cause of headache, always to be kept in mind, has been made clear by the labors of eye-surgeons. It occurs in persons who have headache more or less intense, with abnormal sensations in the skin of the scalp, and at times vertigo and spasm of the eyelids and occipito-frontal muscle. The near use of their eyes increases their distress. When the eye is carefully examined, an optical defect is found, especially hyperopia or astigmatism. Again, we may have defective vision, with sleeplessness and severe headache, dependent on decayed teeth, and disappearing with their removal.\*

Having settled that none of these conditions are present, we have to determine the probable cause of the headache,—a question the solution of which depends frequently more upon the symptoms attending the pain than upon its character. But let us glance at some of the common causes and characteristics of intra-cranial headache.

Headache is a rarely absent symptom of *disease of the brain*. In acute inflammation it is generally agonizing, and, while subject to exacerbations, continuous ; it is associated with fever, with vomiting, although the tongue remains clear, and with delirium. In abscesses of the brain, in softening, and in similar affections which run a chronic course, the headache is less violent, and only occasionally paroxysmal ; it is usually accompanied by signs of disturbed intellection and of deranged motion. In tumor of the brain the headache is apt to be severe and paroxysmal, but intellection is not at first much affected. In congestion of the brain the pain is dull, increased by stooping or lying down, by long sleep, and by bodily or mental fatigue ; its concomitants are a flushed face, throb-

---

\* Case reported by Ogle, Medical Times and Gazette, Aug. 1872.



bing of the arteries of the neck, an eye-ground, as seen with the ophthalmoscope, in which the vessels, especially the veins, are turgid, and a heated head, with increased temperature, as shown by the surface thermometer. A form of congestive headache, apt to be relieved by bleeding at the nose, is often seen in young people at the age of puberty: the attacks are brought on by running or other violent exercise. In diseases of the meninges, especially those of a chronic character, the pain is constant and fixed, and sometimes very sharp. The latter kind of pain when persistent is significant either of disease of the membranes, or, at least, of parts of the superficial structure in contact with them, and is usually felt at the place on the head which corresponds to the seat of the lesion within the skull. Generally there is in meningeal affections coexisting heat of forehead, with signs of local vascular excitement.

*Nervous* or *neuralgic* headache is most common in women, especially in anæmic women. It is unremitting and very severe, yet of short duration; but after it is over there is great lassitude, and even some local soreness. It is not attended with rise of temperature, or with any signs of disturbance of the brain, except at times with a confusion of vision and an inability to carry on a connected train of thought. Anything that agitates the nervous system produces an attack; stimulants and food often relieve it. To the class of headache under consideration may be referred many cases of *megrin* or *migraine*.

But *migraine*, sick headache, or *hemicrania*, has ordinarily certain symptoms which set it apart. The pain is usually attended by nausea and vomiting, is generally at first one-sided, and is accompanied, or more often preceded, by visual disorder, such as a bright spot gradually enlarging. The disturbance of vision begins suddenly, lasting perhaps for half an hour before the headache begins, and is at times associated with tingling on one side, with difficulty in speech and confusion of ideas. The headache often begins in the temple, and is very severe; it spreads over the head, it may extend to the neck, or may leave the side originally affected to become agonizing on the other. There may be soreness of the head with the pain, and there is often pallor of the face, and a contraction of one pupil. Coldness of the extremities is not uncommon, and the patient vomits bile. This



bilious vomiting often terminates the attack, which comes on only in paroxysms. Migraine is frequently met with in gouty or lithæmic persons, and the urine is of high specific gravity, and contains an excess of uric acid and urates, though, during the attack itself, no such increase may be met with.

*Sympathetic* headache is found mainly in connection with disorders of the alimentary tube and of the uterus, and is often worse in the morning, before food has been taken.

Headache may be dependent upon various *poisons*, whether generated in the system or introduced from without; for instance, in diseases of the kidney, particularly Bright's disease, the retention of a large quantity of urea in the blood becomes the source of persistent pain in the head. In torpidity of the liver, in lead poisoning, in opium-eaters, in drunkards, after the use of strychnine or of large quantities of quinine, headache is common; and it is very likely that in persons with faulty assimilation certain ptomaines give rise to the headache.

In studying headache as a symptom, we must always note what influence position and movements of the head have on the pain: whether, for instance, stooping, swinging the head from side to side, or rising rapidly from the horizontal to the erect posture affect it, and cause it to be combined with vertiginous or other abnormal sensations. In headache connected with organic disease of the brain the pain is increased by whatever increases the blood-pressure,—by stooping, by coughing, by any effort. The site of pain bears no very definite relation to the site of lesion, except the lesion be near the surface. With severe paroxysms of pain vomiting often occurs. Headache increased by the erect posture and relieved by lying down bespeaks an anæmic condition of the brain.

**Vertigo.**—This is a transitory feeling of swimming of the head, a sense of falling, or illusory movements of external objects. The sensation is apt to occur whenever the circulation within the cranium is disturbed, and is often symptomatic of a disease of the heart, liver, kidneys, especially Bright's disease, or of an affection of the stomach, or of gout or lithæmia; or it accompanies anæmia, or follows long-continued and exhausting discharges. In the defective blood-supply to the brain produced by arterio-sclerosis the vertigo is attended by signs of the morbid process in other parts

of the body, and the tense pulse, increased blood-pressure, and accentuated second sound of the heart explain the cause of the giddiness. Extreme slowness of pulse, and a sensation of falling in a given direction, as in Ménière's disease, are not infrequent.\*

Vertigo may attend any disorder of the brain. The *cerebral* form is recognized in part by the absence of those affections of other organs which would induce the dizziness,—and among these we must not forget eye-strain and local palsies of the muscles of the eyeball,—in part by its being joined to an almost constantly present sense of uncertainty in movement, to headache, and to further signs of an encephalic malady. Moreover, it is usually objective in character: surrounding objects appear to the patient to move, not he himself; and, unlike the subjective vertigo so common in mere sympathetic disturbance of the brain, closing the eyes relieves it.

The most common form of vertigo, not arising from brain affection, is the so-called *stomachal vertigo*. It is apt to come on in paroxysms, sometimes in the middle of the night or in the early morning, and is associated with a dull, heavy ache in the head, and with more or less gastric disturbance, often following indiscretion in diet. Yet the tongue may be clean, and the digestive disorder so slight that it is only by the after-symptoms, by the relief afforded by attention to diet, and by remedies acting on the digestion, that we clearly make out the cause of the vertigo. Between the attacks the patient is free from the affection; though there are cases of more chronic kind, in which a certain amount of giddiness is present for long periods with only comparatively short intervals of freedom. Here food and stimulus are apt to relieve the giddiness, which exists often with symptoms not of violent indigestion, but of delayed and slow digestion, and may become aggravated into a severe attack if the stomach be for a long time empty. In the gastric vertigo there is no loss of consciousness. The pathology is obscure. Woakes† has endeavored to establish a direct nervous communication between the stomach and the

---

\* Grasset, *Vertige Cardio vasculaire*, Paris, 1890; Church, *Medical News*, June, 1892.

† Deafness, Giddiness, etc., 1879.

labyrinth to explain the vertigo. Others regard the irregularity in the cerebral circulation produced by the gastric disorder, anæmia or hyperæmia, as the cause.

Very similar to gastric vertigo is the vertigo of malassimilation in connection with *lithæmia*. The history of the case, the state of the urine with reference to uric acid, the striking change which follows diet and treatment that alter the formation and elimination of uric acid, distinguish lithæmic vertigo.

Another form of vertigo of eccentric origin is that associated with partial deafness or ringing in the ears. Again, there may be an affection of the internal ear, the semicircular canals of the labyrinth especially being the seat of an inflammation, and the vertigo set in suddenly. Its onset is apt to be associated with vomiting, with suddenly-developed tinnitus, with pain produced in the affected ear by the slightest noise, and with symptoms of apoplexy or a fainting condition. Such cases, to which Ménière particularly has called attention, at times very speedily terminate fatally. But the acute seizure, which is by far the most common beginning of the *aural vertigo*, may leave behind giddiness and a persistent unsteadiness in standing and walking, or a tendency to go forward or backward, or a reeling gait, which, with the intense vertigo, the vomiting, the persistent noises in the ears, the unimpaired consciousness, and the deafness, become very valuable signs of Ménière's disease. The deafness shows especially in defect of power of hearing vibration conducted through the skull. It is often one-sided, generally on the side of the marked tinnitus, and never absolute. Again, it may be noticed that there is deafness for certain groups of musical sounds, which Knapp accepts as proof that the disorder has extended to the cochlea.

In some instances the patient has a tendency to turn to one side or to walk round and round in a circle; and he is always miserable, although his general health suffers but little. The disturbance of the equilibrium is not always present; there may be disturbance of hearing without it. The vertigo is generally the most prominent symptom of the disease, and persistent vertigo not epileptic in character or obviously associated with an organic brain affection is nearly always aural. The dizziness is very apt to be severe, to come on in paroxysms, and to be excited by some effort or movement. It becomes associated with pallor,

with faintness, with vomiting, and in part it remains even between the paroxysms. During these the roaring in the ears may or may not be increased, but signs of eye-disturbance are very apt to show themselves. The disease may result from any process that involves the labyrinth and the nerve-endings. It is more common in men than in women, and is very rare in young persons. It may come on after cold and exposure, or originate in gout or in syphilis. It has also been observed in men working under ground and breathing compressed air.\* All cases of aural vertigo do not set in suddenly; some are slight, others are very severe and do not cease until the hearing is totally lost. Many cases progress slowly to recovery. Aural vertigo in its milder forms may be met with in affections of the ear, that have had their origin in catarrhal inflammation travelling along the Eustachian tube.

To return to vertigo connected with cerebral or cerebro-spinal disease. There is a kind which Trousseau especially has described. The abnormal sensation is very short in its duration, but severe; the patient momentarily loses all consciousness. The vertigo recurs at uncertain times: while actively engaged, sometimes while in bed and half asleep. The head feels heavy after an attack, and the mind is temporarily stupefied; otherwise the health is good. This type of vertigo is dangerous. It is often the *precursor of epilepsy*, and after a time becomes associated with convulsions.

Another kind of vertigo is that which arises from *overwork of the brain*. At times giddiness is the only symptom of disorder, *essential vertigo*, and is present for many years, the patient enjoying otherwise excellent health. I have known a number of such instances in which the tendency appeared to have been inherited. If it do not break out until late in life, it is a matter of more serious concern.

In *laryngeal vertigo* † there is a close connection with epileptic seizures. The chief symptoms are tickling or burning in the larynx, followed by vertigo, loss of consciousness, and spasmodic movements in the face and limbs. The larynx is healthy; but in

---

\* Curnow, *Lancet*, 1894, No. 3715, p. 1688.

† Gasquet, *Practitioner* for August, 1878; Charcot, *Progrès Médical*, No. 17, 1879.



a case observed by Sommerbrodt a polypus existed, the removal of which cured the affection.

Somewhat allied to vertigo is the condition known as *astasia-abasia*, the most marked characteristic of which is difficulty in standing and walking. Consciousness is not lost, but sometimes there is a sense of giddiness. The affection is most common in persons past middle life, especially if gouty and with degenerated vessels. It has also been observed in hysterical patients.

Besides headache and vertigo, there are various unnatural sensations, such as a feeling of momentary unconsciousness without giddiness; a feeling within the cranium of weight, of constriction; the feeling described as a rush of blood to the head; ocular spectra, and other false perceptions of many kinds and of every gradation. But I shall do no more than advert to this subject, and shall now consider some of the morbid phenomena of the special senses, particularly of the senses of sight and hearing.

#### DERANGEMENT OF SPECIAL SENSES.

**Vision.**—The sense of vision may be exalted, impaired, or perverted in disorders of the brain, whether organic or functional. It is exalted in inflammation; impaired, even totally lost, in softening, in tumors, in apoplexy, and during violent hysterical attacks simulating apoplexy. Perversions of the sense of vision are more frequent than its abolition, and probably more peculiar to cerebral affections. They are of all kinds,—some of great consequence, others of but little. *Musæ volitantes*, or the appearance of spots and various small objects floating before the eye, have the latter significance; for they may happen in almost any form of cerebral disturbance, also in anæmia, in cardiac maladies, in the neuroses, and in states of nervous exhaustion. They are simply the shadows of vitreous opacities or retinal vessels upon the retina, and have nothing to do with anything but the local condition, which is without significance. Of other manifestations of deranged sight, such as illusions, ocular spectra, and phantasms, I shall only state that they are more common in sick headache, and in derangement of the mind, temporary or permanent, than in recognizable organic disease of the brain. Yet they are found in affections of certain parts of the brain; for in

disease of the posterior lobes, as Hughlings Jackson has observed, colored vision and optical illusions are frequent.

The *appearance* of the eye is often of as much significance as the derangement of sight. In some cerebral maladies the eye has a fixed stare; in others the eyelids are constantly moving: but the latter is a sign more frequent in chorea, local spasm, and hysteria. Great brilliancy of the eye is often noticed in meningitis and in insanity.

*Derangements of the ocular mechanism* may be the result of remote causes, or, themselves primary, may become the starting-point of disorder elsewhere. In the first case their study is valuable to the general diagnostician as indicative of the seat, nature, or stage of many diseases in other parts of the system; in the second case the diagnosis as well as the therapeutics of the distant and related disease is dependent upon the appreciation of the ocular derangement. It thus becomes evident that the abnormalities of the visual mechanism are of the highest importance in many systemic affections, particularly in disease of the cerebro-spinal system, where it is always necessary to inquire as to derangements of the eyes and their significance.

Let us first briefly consider the *idiopathic derangements of the eye that induce derangements elsewhere*. Both in origin and in result these are essentially functional. So far as relates to the eye they consist chiefly either in abnormalities of refraction, classed under the general head of ametropia, and comprising hyperopia, astigmatism, myopia, and presbyopia, singly or combined; or in incoördination of the external ocular muscles, commonly called insufficiency. The results of ametropia and muscular insufficiency are conveniently called *eye-strain*; and the symptoms of these conditions must not be neglected by the physician, especially in view of the fact that eye-strain generally evinces itself not so much in ocular or visual symptoms as in functional nervous derangements often far removed and apparently disconnected. For example, it is a well-established fact that eye-strain is prone to produce headache, especially in young women after the age of puberty. These headaches are usually frontal, but may also be occipital, less frequently of the vertex or diffused. Eye-strain is at times the starting-point of choreic symptoms, and even of

genuine chorea. Cases have been reported\* by trustworthy observers showing that the same cause may produce functional gastric derangements, hysteria, melancholia, and even epilepsy. The lesson is obvious that when these or other functional affections do not yield to direct treatment, or when their origin is not otherwise explainable, we should at once proceed to exhaust the possibilities of a reflex neurosis due to ocular abnormality or to some other peripheral irritation.

Hyperopia and hyperopic astigmatism are much the most frequent sources of eye-strain, and by the aid of a mydriatic, followed by tests with the trial-lenses, the diagnosis of the existence and amount of the defect may be made. In the neurotic, or in those with intercurrent affections and weaknesses, the smallest amount may become the source of irritational strain. Muscular insufficiency is the next most frequent cause of ocular irritation, and its existence is at once and easily detected by a simple test. The correct diagnosis of its amount demands the offices of one specially skilled. Simple myopia produces no strain, but myopic astigmatism, and presbyopia, may sometimes cause it.

Turning now to the consideration of those *changes in the ocular mechanism* which indicate *effects and symptoms of disease elsewhere*, we find that disease in almost any part of the organism may give indications of its nature and location in the eyes. These symptoms, either singly or combined, are of a threefold nature :

Changes in the external appearances and visible to the naked eye.

Changes in the fundus oculi, or eye-ground, as revealed by the ophthalmoscope.

Defects of vision as shown by the subjective report of the patient.

The first and last set of symptoms require no very considerable special training to study, but the use of the ophthalmoscope does demand it, and often to such a degree that many are unfortunately compelled to forego invaluable knowledge.

I. Among the *external ocular abnormalities of the eyes* visible

---

\* For example, Clinical Illustrations of Reflex Ocular Neuroses, by Gould, Amer. Journ. Med. Sci., January, 1890.

to the examiner, exception must of course first be made of such local diseases as have no systemic relations, such as ecchymoses, congestions or inflammations of the lids and conjunctiva, trachoma, glaucoma (with an unusual hardness and anæsthesia of the eyeball, impaired vision, dilated pupil, etc.), cataract, congenital anomalies, etc. Herpes zoster ophthalmicus, a peripheral neuritis of the ophthalmic branch of the fifth nerve, is a dangerous and painful malady, often, if not always, owing to local causes. Exophthalmos is either due to local disease or is present as one of the main symptoms of the affection called exophthalmic goitre. A case recorded by Hale White \* seems to locate the focal lesion in the medulla, in the central part of the floor of the fourth ventricle, near the nucleus of the sixth nerve.

Next in importance is a class of diseases due to external infection that generally points to a source of contagion elsewhere in the organism. Cases of localized tuberculosis of the conjunctiva have been reported wherein the handkerchief has perhaps carried the bacillus to the eye. Gonorrhœal ophthalmia is a constantly recurring disease in ophthalmic practice; but the most frequent and frightful is the ophthalmia of the new-born,—ophthalmia neonatorum,—due to infection during labor with the vaginal discharges of the mother. It is said that the greater part of the blindness of the world is due to this wholly preventable disease.

Affections of the conjunctiva or lids may have their origin in diseases of the adjacent skin or mucous membrane, and extend to the eyes by simple contiguity of structure. There is reason to believe that a close connection may frequently exist between hay-fever, catarrhal and other diseases of the nasal mucous membrane, and similar conditions of the conjunctiva.

Arcus senilis, a ring of grayish tissue-change about the corneal limbus, betokens generalized atheromatous or fatty degeneration, chiefly arterial or cardiac. Interstitial or diffused keratitis is nearly always the result of inherited syphilis. In rubeola, scarlatina, smallpox, and erysipelas, the external ocular structures may be injured or destroyed by the same causes that produce the skin-lesions, or by extension of the disease to the eyes from the skin.

---

\* Brit. Med. Journ., March 30, 1889.



Of the remaining affections of the external parts of the eye indicative of general or internal disease, the most important are those pertaining to the muscles of the eye or movements of the globe. They easily fall into two groups,—those of the external and those of the internal muscles.

*Strabismus*, or squint, may be due to local causes, such as injuries, or cold, etc., but it usually arises from a lack of equal or balanced power among the twelve external muscles, and to ametropia and anisometropia. The distinctive subjective characteristic of squint is double vision; but so numerous are the possible combinations that it is often difficult, if not impossible, to tell just what nerves or muscles are implicated, and the exact seat of the lesion. In examining for strabismus we observe whether the eyeball is turned inward or outward. In paralysis of the external rectus we have ordinarily an internal or convergent squint, in paralysis of the internal rectus an external or divergent strabismus. In palsy of the superior rectus there is inability to raise the eyeball in a proper manner above the horizontal level; inability to lower it below indicates palsy of the inferior rectus. Strabismus due to local causes must be distinguished from true paralytic squint due to more centrally located lesions. It must also be distinguished from spastic action of the muscles caused by irritative intracranial injuries. In both the latter cases there is a conjugate or common movement of both eyes to one side or to the other, called *conjugate lateral deviation*; the head often shares in the lateral movement. In spastic irritative lesions of the cortex the eyes are turned from the side of the injury; in paralytic or destructive lesions they are turned toward it. The eyes, as it has been said, look at the lesion in paralysis, away from it in spasm. The symptom, however, owing to its frequently temporary existence, and also to the fact that it may arise as an indirect symptom, must not be relied upon except in conjunction with others and when continuing at least for several weeks.\* The seat of the lesion may be in the cortex, the internal capsule, or the pons; in the latter case the

---

\* The direct symptoms are those intimately dependent upon the lesion of a part; the indirect or distant symptoms are those due to disturbances of circulation; to pressure, to the reflex or inhibitory effects at other points than the seat of injury. The less marked the cerebral symptoms, the more probable is it that the paralyzes are direct.

symptoms are direct and the deviation of the eyes is the reverse of that given above: the eyes in paralysis look away from the lesion; in spasm, toward it. If in lesions of the pons the sixth nerve nucleus be included, there is, of course, paralysis of the external rectus, so that the corresponding eye cannot be rotated outward past the middle line, whilst the other eye cannot be rotated inward past the middle line. This associated movement of the other eye will not be impaired if the injury to the sixth nerve be between the nucleus and the globe.

Owing to the peculiar position of its nucleus and the long course of exit of the *sixth nerve*, its exclusive paralysis is the most frequent of single nerve palsies. It is peculiarly liable to paralysis from indirect or pressure causes, but if connected with paralysis of the opposite side of the body and with other symptoms of brain disease, it clearly points to a lesion of the pons. In consequence of the close anatomical relations of their nuclei, palsies of the sixth and facial nerves are frequently associated. Other nerves originating in the pons are, of course, liable to implication in varying degrees. Next to the sixth the *third nerve* is the one most often paralyzed, and, in proportion to the numbers of twigs involved and the completeness of their palsy, is there a probability of a lesion at the base of the brain, though the location can be rendered certain only by a study of associated palsies and symptoms. The various paralyses of the external ocular muscles are usually attended with double vision, *diplopia*.

*Ptosis* may exist either with or without involvement of other third-nerve branches, but in any case the value of the droop of the upper eyelid as a localizing symptom is somewhat indeterminate. If of one eye alone, ptosis usually indicates a cortical lesion, unless due to evidently local causes. In paralysis of the third nerve we have, besides the ptosis, dilatation of the pupil of moderate extent. Inability to close the eyelids is associated with paralysis of the facial nerve.

As regards the nature of the lesion, the ocular symptoms generally give little definite indication, and, at all events, must be considered in relation with others and with the history of the case.

*Abnormalities of the pupils* are understood by remembering that

the third nerve controls the contractile mechanism and the cervical sympathetic the dilating mechanism. Hence an unusual diminution or increase of either innervation, especially of the first, causes alterations of the pupils at once. Irritative cerebral lesions thus produce contraction, whilst lesions which destroy cerebral function produce morbid dilatation. The state of the pupil in tumors, hemorrhage, and inflammatory conditions of the brain may thus furnish us with most serviceable indications of the extent and destructiveness of the injury. When but one pupil is abnormal, the rule above given serves to indicate lesion of the corresponding half of the cerebrum, irritative or paralytic according to the degree of the injury. Yet one-sided contraction, like one-sided dilatation, may also be owing to tumors at the root of the neck. Hemorrhage or effusion into the pons or lateral ventricles, when small or irritative, produces contraction; but if large, permanent dilatation. Certain drugs, such as opium, contract the pupil; belladonna and chloral dilate it. We also find dilatation of both pupils in chlorosis. If the foot be pricked, the pupils dilate, provided the iris be uninjured and the sensory columns be intact. In epileptics this reflex excitability is greatly diminished.\*

The pupillary reaction to light may be useful in diagnosing the location of a lesion, whether beyond the corpora quadrigemina or not. If beyond, the pupillary reflex will be retained, despite the loss of sight. Lesions of the spinal cord and of the sympathetic nerve produce results the reverse of cerebral disease: irritative lesions dilate, paralytic lesions contract. In this connection the Argyll-Robertson pupil—the light-reflex lost, the accommodative reflex retained, of a myotic pupil—is of value as indicating, often early, sclerosis of the posterior columns of the cord. When hemianopsia is due to disease of the optic tract, the pupil fails to react to the stimulation of light reflected upon the blind half of the retina; contracting, however, if the lesion be situated in the cerebral hemisphere. *Paralysis of the accommodation* may exist independently of pupillary involvement, and its significance is that of paralysis of other branches of the third nerve. Paralysis of many or of all the

---

\* Lawson, West Riding Reports, vol. iv.



muscles of both eyes, *ophthalmoplegia*, is usually due to a lesion of the nuclei of the supplying nerves.

II. *Abnormal changes in the fundus of the eye* may be of great diagnostic value, and in almost every case of circulatory or nervous disease the ophthalmoscope gives valuable hints concerning the general disorder. This is rendered exceptionally true by the fact that these changes are most frequently symptomatic, and, with few exceptions, do not arise from local disease.

We should invariably examine with the ophthalmoscope the eyes of patients suspected of having disease of any part of the cerebro-spinal nervous system. Changes in the eye, indeed, often occur early enough to be the first certain sign of disease, and this, too, without any impairment of sight; on the other hand, lesions indicating cerebral or other organic affection have been found in cases in which failure of sight was alone complained of, the cause being unsuspected. But particularly is the ophthalmoscope valuable in enabling us to differentiate organic from functional affections. It tells us of extension of congestion or of inflammation of the brain to the internal structures of the eye, or of the amount of resistance offered to the circulation within the cranium. This resistance may either arise from a marked "coarse" lesion, or may make itself felt through the sympathetic nervous system.

The changes in connection with organic disease have been observed chiefly in the retina, the optic disk, and the choroid. In using the *ophthalmoscope for medical diagnosis* we pay particular attention to these structures; especially do we note the disk, its color and size, and the pigment around its edges, the region of the macula, the size and appearance of the arteries and veins, whether diminished, enlarged, or tortuous, whether there are exudations or hemorrhages in the course of the vessels, and in what part of the eye-ground the patches are most marked.

*Hyperæmia*, or increased redness, is due to local causes; and the fundus-changes in myopia, astigmatism, retinitis pigmentosa, and some forms of choroiditis are also to be excepted. In *diseases of the blood* and the blood-making organs, the indications are remarkably clear. *Retinal hemorrhages* are a common concomitant of such general diseases as albuminuria, diabetes, anæmias, cardiac valvular disease, arterial atheromatous and fatty degenerations,



chronic malaria, and other febrile conditions. *Embolism of the central artery of the retina*, causing unilateral blindness, points to cardiac valvular disease. There is a grayish discoloration about the macula, with a central cherry-red spot. Poverty of the blood, *simple anæmia*, is at once recognized by the transparency of the blood-columns, and *leukæmia* and *pernicious anæmia* produce characteristic changes in the eye-ground, especially the last, with retinal œdema and hemorrhages, disk-discoloration, arterial pallor, and venous distention. *Albuminuric retinitis* is not invariable in albuminuria, but, when present, renders the prognosis more serious. The typical fundus-changes consist in an early stage of haziness of the papilla and central part of the fundus, slight hemorrhages, and faint grayish discolorations. Later, white dots or splotches are grouped about the macula, or, flame-like, radiate from it. Striate hemorrhages are scattered over the fundus, the papilla is œdematous, and its limits are obscured. The ophthalmoscopic signs of *diabetic retinitis* are very similar to the last. Visual disturbances, however, do not, in either case, stand in any exact ratio to the defects of the eye-ground.

*Atrophy of the optic nerve*, recognizable by the whiteness or discoloration of the disk, failure of vision, even to blindness, etc., may sometimes seem to have no remote causes, but is commonly associated with, or a result of, diseases or lesions of the spinal cord or the brain, toxic substances in the blood, papillitis, etc.

*Papillitis*, optic neuritis, "choked disk," is a symptom of most decided diagnostic value. The picture is easily recognized, consisting in a swollen red disk, the edges and vessels of which are obscured by a "woolly," striate blurring extending to the adjacent retina. This condition is always symptomatic, and in the large majority of cases points to tumor of the brain, though other intracranial diseases may produce it. From papillitis, however, nothing can be argued as to the nature or location of the tumor or other affection. Its existence—and it is often not a late symptom—at once demands careful inquiry and energetic treatment. This is particularly true because unimpaired vision may coexist with even a severe papillitis. Optic neuritis has been observed after measles and scarlet fever, also after malaria and typhoid fever.\*

---

\* White, Journal of the Am. Med. Assoc., Oct. 1893.

*Choroidal inflammations* are chiefly distinguishable by the striking color and pigment changes of the fundus. Plastic choroiditis is commonly secondary to meningeal affections and prostrating fevers; purulent choroiditis, to local or general infection or septicæmia. Disseminated and central choroiditis, or choroidoretinitis, is frequently the result of syphilis. The choroid is peculiarly liable to become the seat of tuberculous growths.

III. Passing now to the consideration of purely *subjective visual derangements*, it becomes highly necessary to determine first whether such defects are due to refraction-errors, insufficiencies, and other local causes, or if they are secondary and symptomatic. Unless other indications are present, the complaint of headache, especially if frontal, weariness or pains of the eyes after near-work, affections of the lids and conjunctiva, conjoined with general irritability and functional gastric derangements, almost invariably indicate eye-strain as primary. Simple inability to see distant objects clearly, without other symptoms local or general, indicates myopia.

*Amblyopia*, due to the excessive indulgence in tobacco or alcohol, has but a single objective sign: an unusual pallor of the temporal portion of the papilla. There is deterioration of visual acuity, to which subnormal color-perception may be added. Amblyopia sometimes occurs also as a manifestation of hysteria and in association with migraine. It has further been observed as a symptom of intoxication with quinine and iodoform, or after sexual excesses; or the defective acuteness of vision shows itself as a day-blindness or as a night-blindness; or takes the form of contracted fields of vision, or of color-blindness. Marked visual deterioration of a single eye should lead to inquiry for extra-local causes. It may be due to disease of the corresponding optic nerve. When ametropia has been excluded and the above-described ophthalmoscopic signs are wanting, the cause must be sought in disease of other organs. Paresis, and even paralysis of the accommodation, and visual failure, are not infrequent as *reflex neuroses* from peripheral irritation of other parts. Cases of abnormalities of dentition and other dental troubles producing such visual defects have been frequently reported. Menstrual difficulties, masturbation, the influence of pregnancy and lactation, may sometimes account for obscure ocular troubles. Hemeralopia, night-blindness, due to

deficient nutrition of the general system, has been traced to insufficient food.\*

Modifications of the color-fields have been found chiefly in hysterical patients. The field for red and green, always the narrower, shows the restriction most markedly.

The most important ocular sign of cerebral disease, and one invariably pointing to intracranial affection, is *hemianopsia*, or loss of vision of the halves of the fields. The most common variety is that called homonymous lateral hemianopsia, in which the loss is either of the temporal half of one eye and of the nasal half of the other, or *vice versa*, a vertical line nearly through the centre being the dividing line. There are three other forms of hemianopsia, called temporal, nasal, and altitudinal, in which the half-fields are respectively the two temporal, the two nasal, with the dividing line, as previously, perpendicular, or the two dark half-fields are the upper or the lower halves, with the dividing line horizontal. These three varieties are seldom met with, and, from the peculiar anatomical relations of the optic chiasm or commissure, are readily recognized as the results of lesions of this part, either at one side or the other, above or below. Homonymous lateral hemianopsia always indicates lesion beyond the chiasm. If the hemianopsia be "relative,"—involve only a part of the perceptions of light, form, and color, the three constituent factors of ordinary vision, and believed to have special subcentres or strata in the general visual centre,—it must necessarily proceed from a partial lesion of the common visual centre situate in the cuneus of the occipital lobe.† But if the hemianopsia be absolute,—with complete loss of light, form, and color sense,—the lesion may be either one affecting the entire visual centre of one side, or one rendering wholly functionless the fibres of one radiation, internal capsule, or optic tract. If the latter were the case there would almost certainly be other intercurrent or general symptoms, such as paralysis of other cranial nerves, hemianæsthesia, some form of aphasia, or hemiplegic symptoms. A symptom of great value in locating the lesion of

---

\* See article by Kubli, *Archiv für Augenheilk.*, June, 1887, who describes three hundred and twenty cases occurring during the Russian church-fasts.

† Seguin limits the centre to the cuneus; Nothnagel makes it include also the posterior portion of the superior occipital convolution.

hemianopsia is the hemiopic pupil. Convergence of a narrow cone of light upon the insensitive half of the retina yields no pupillary reflex if the lesion be in the optic tract; if the pupil, under such stimulus, contract, the lesion must be beyond the tract. The intracranial affection giving rise to the hemianopsia may be of malarial origin, and it and the associate cerebral symptoms will disappear under active antimalarial treatment.\*

*Mind-blindness*, physical vision, but failure to realize the psychological import of the things seen, sometimes a symptom of general paralysis and obscure cerebral disease, indicates a cortical lesion in the occipital or occipito-temporal lobe, near by if not conterminous with the visual centre.

**Hearing.**—As regards the *sense of hearing*, the same may be said as of vision. It, too, is perverted and impaired in various cerebral affections. Yet, to be certain that the cause of the difficulty is cerebral, the ear must first be examined with reference to any physical imperfection; and in doing so we may by means of the otoscope get an idea of the vascularity of the drum, and be led from this to infer the condition of the vessels of the brain. We must also examine the throat and the condition of the Eustachian tube, for catarrhal inflammation extending to the middle ear may give rise to a form of aural vertigo.

Great acuteness of hearing and intolerance of sound are generally symptoms of extreme nervous irritability, or of beginning cerebral inflammation. Deafness may be owing to softening of portions of the brain; but Ferrier tells us that it is not met with in destructive lesions of the cortex. Deafness is also found as a temporary and by no means unfavorable symptom in the continued fevers. Imaginary sounds and ringing noises in the ear, or *tinnitus aurium*, are frequent accompaniments of cerebral disorders. But the latter is encountered in so many different conditions—in diseases of the cerebral vessels, in congestion of the brain, in Ménière's disease, in affections of the heart, in anæmia—that it is a sign of little moment; and, in truth, its most usual cause is local,—namely, an accumulation of wax in the meatus.

---

\* See my paper on Malarial Paralysis, with eye examinations by Harlan, in *International Clinics*, vol. iii., Ser. I., Oct. 1891.



## DERANGED REFLEXES.

Derangement of the reflex action plays a most important part in the study of diseases of the nervous system. Each action is brought about by a sensory nerve that conveys the impression to the centre, by a motor nerve that transmits the impulse from the centre to the periphery, and by a reflex centre between the two in the spinal cord connecting the roots of the sensory and motor nerves, which with them forms the "reflex arc." The reflex centre is to some extent under brain control.

There are two forms of reflexes to be especially studied,—the cutaneous or superficial, produced by stimulating the skin, and the deep reflexes, the muscle or tendon reflexes, evoked by tapping muscles or tendons.

The *superficial* may be almost everywhere excited by tickling or gently stimulating the skin. The most usual ones to be noted are the reflex of the sole of the foot, the *plantar reflex*; and that of the palm of the hand, the *palmar reflex*. The former, when normal, attests the integrity of the reflex arc at the lower end of the cord; the palmar reflex, contraction of digital flexors by tickling the palm, indicates a normal state of the reflex arc through a greater part of the cervical enlargement. Other superficial reflexes which may be mentioned are the *cremaster reflex*, the drawing up of the testicle excited by stimulating the front and inner side of the thigh, and originating in the cord at a point between the first and second lumbar pairs; the *gluteal reflex*, the contraction caused by irritating the skin over the buttock, and showing the integrity of the cord at the fourth and fifth lumbar nerves; the *abdominal reflex*, a contraction in the abdominal walls caused by scratching the skin on the side of the abdomen, and depending on the action of the cord from the eighth to the twelfth dorsal nerve; the *epigastric reflex*, an epigastric dimpling produced by stimulating the side of the chest in the fifth or sixth intercostal space, and indicating integrity of the cord from the fourth to the seventh pair of dorsal nerves; the *scapular reflex*, a contraction by stimulation of the scapular muscles, and bespeaking the integrity of the reflex arc at the level of the upper two or three dorsal and lower two or three cervical nerves; the *erector spinal*, showing itself by stimulating the skin along the border of the erector

spinal muscle, the contraction of these muscles showing the healthy state of the cord in the dorsal region. Other reflexes of indeterminate utility are the *erector spinæ reflex*, a local contraction of these muscles produced by stimulation of the skin along their border, proving that the reflex arc is intact in the dorsal region of the cord; the *platysma reflex*, dilatation of the pupil upon pinching the platysma myoides muscle; the *jaw-jerk* or clonus, rarely present in health, elicited by suddenly depressing or tapping with a hammer on the inferior maxilla, when clonic movements of the lower jaw are produced; the *peroneal reflex*, a stroke upon these muscles when in tension, or when the foot is bent inward, causing a reflex movement. To these may be added the *tendo Achillis reflex* or *front-tap contraction*, described by Gowers, a reflex contraction of the gastrocnemius when the muscles upon the anterior part of the leg are struck, the leg being extended and the foot flexed by the hand upon the sole. It is considered a delicate test of heightened spinal irritability. Among *cranial reflexes*, the more noteworthy are the *iris-contraction* upon exposure of the retina to light; the *eyelid-closure* from irritation of the conjunctiva; the *pharyngeal*, *laryngeal*, and *palatal reflexes* (cough, swallowing, etc.) from irritation of these parts; and *nasal reflexes*, as in sneezing. The *aural reflexes* are of some value in appreciating disease of the cervical part of the cord.\* In disease these superficial reflexes are often absent. Thus, disease of one cerebral hemisphere diminishes or destroys them on the other side, the paralyzed side of the body. In pregnancy all reflexes are increased. The superficial reflexes are much influenced, increased or diminished, by psychical causes.†

The reflex phenomena connected with the tendons give us the best illustration of the so-called *deep reflexes*. The tendon of the patella is the one most readily studied; and if, the body being bent forward, we strike abruptly the tendon of the patella just below the knee-cap, after rendering the ligamentum patellæ tense by flexing the knee at a right angle while one knee-joint rests upon the other, a sudden contraction takes place in the quadriceps femoris muscle, and the foot is jerked upwards. When very slight, the knee-jerk is most readily elicited by a tap with the percussion

---

\* Amer. Journ. Med. Sci., Dec. 1888.

† Jendrassik, Deutsch Archiv f. klin. Med., April, 18: 4.

hammer. This reflex is due to a muscle reflex action dependent upon the spinal cord.

The knee-jerk is found in health, and is markedly increased in disease of the pyramidal tracts, in heightened irritability of the gray substance of the spinal cord, in many tumors of the brain, in cerebro-spinal sclerosis, in lateral sclerosis, after epileptic seizures or unilateral convulsions, in spinal irritability.\* It is absent in locomotor ataxia, even at an extremely early stage of this affection. It is also abolished in affections of the anterior gray cornua, in infantile paralysis, in destructive lesions of the lower part of the cord, in advanced stages of pseudo-hypertrophic paralysis, in alcoholic paraplegia, and, temporarily at least, as pointed out by Hughlings Jackson, in meningitis and in instances of emphysema and other maladies in which the blood has become venous to an extreme degree,† and disappears in certain general constitutional affections, as in diabetes and in diphtheria.‡ I have also known it very exceptionally to be absent in healthy persons, in one instance in three brothers.

The *biceps reflex* is developed by tapping the tendon of the biceps. This leads to contraction of the biceps muscle. Its meaning is the same as that of the knee-jerk. Tapping on the front of the wrist gives rise to contraction in the flexors of the fingers; striking the tendon of the *triceps* above the olecranon causes contraction in the triceps. This is especially marked in the irritable muscle of the early and late rigidity of hemiplegia. Another deep reflex is the *periosteal*. It is produced by tapping the bones of the forearm or leg, which gives rise to active contraction of the muscles, and indicates a disease of the spinal cord, especially amyotrophic lateral sclerosis.

In some instances of disease the reflex phenomena are produced on the side opposite to the one acted on. These *crossed* reflexes are not unfrequently met with in posterior spinal sclerosis, and are not merely associated contractions. A secondary stimulation of a motor centre in the opposite side of the cord has been suggested as the cause in a case of transferred patellar tendon reflex.§ A tap on the tibia near its middle generally induces contractions of

---

\* Hughlings Jackson, *Med. Times and Gaz.*, Feb. 1881.

† *Brit. Med. Journ.*, 1892, No. 1614.

‡ Marie et Guinon, *Revue de Méd.*, July, 1886.

§ McLane Hamilton, *Archives of Medicine*, New York, Dec. 1883.

the quadriceps femoris; and it is often followed by contractions of the quadriceps of the opposite leg when both the pyramidal tracts are diseased.\*

The phenomenon called *reinforcement of a reflex* may have its use and significance in the diagnosis of doubtful or obscure cases. In testing the muscular power of the hand by the dynamometer, it is well known that when one hand is fatigued it has greater power if the other hand be forcibly and synchronously clenched than if acting alone. In the same way it has been shown that any reflex is heightened by coincident muscular exertion of other parts than those being tested. Thus, if a desired reflex be weak or difficult to elicit, it may be brought out by muscular tension of some other member or part of the body. Strong sensation of the skin acts in the same way to reinforce coincident reflexes. So slight an outlay of force as that of winking will increase the force of the knee-jerk, if correctly timed.† When the muscle is cut off from connection with the spinal centres, as in the late stages of locomotor ataxia, the reflex and any reinforcement are alike impossible.

Very similar to the knee phenomenon is the foot phenomenon, or *ankle clonus*, although its reflex character is even more doubtful. Gowers,‡ indeed, has made it probable that it is largely due to an exaggerated irritability of the muscles. It is produced if the foot be suddenly brought into complete flexion by the hand pressed against the sole, the leg being semiflexed, and still more readily if subsequently the tendo Achillis be quickly tapped. A kind of convulsive shaking of the foot results, dependent on alternate contraction and relaxation of the anterior tibial and calf muscles. Ankle clonus is at times, but not often, observed in healthy persons, although it is susceptible of being cultivated; in lateral sclerosis it is developed to an extraordinary degree. Indeed, it is marked in the class of affections in which the knee reflex is excessive. When produced by sudden passive tension alone of the muscle, it is indicative of structural change in the spinal cord.§

\* Ross, *op. cit.*, vol. i.

† Mitchell and Lewis, Tendon- and Muscle-Jerk, Amer. Journ. Med. Sci., vol. xcii., 1886.

‡ Medico-Chirurg. Transact., 1879.

§ Gowers, Diseases of the Nervous System



*Wrist clonus* may be induced in the late rigidity of hemiplegia by pressing the hand backwards so as to produce extreme extension at the wrist.

If a muscle be suddenly relaxed, a slow tonic contraction follows which may last for some minutes. The phenomenon is best witnessed in the *tibialis anticus*, but is rarely seen in the muscles of the arm. This *paradoxical muscular contraction* has no definitely ascertained value. It is sometimes met with in the early stages of locomotor ataxia.

#### DERANGED MOTION.

The chief manifestations of deranged motion resolve themselves into the phenomena called paralysis, ataxia, tremor, spasms, and convulsions.

##### Paralysis.

When we speak of paralysis, we mean a loss of the power of motion, although there is the impulse of the will to move the affected part. It is true, there is also a paralysis of sensation, a complete anaesthesia, which may be conjoined with the paralysis of motion; but the latter often happens alone, and is the morbid state meant when we use the word paralysis without qualifying it. A slight, incomplete paralysis is called "paresis," and this term is especially employed when the loss of power exists without demonstrable organic change.

Paralysis is nearly always of nervous origin. It may be general, or it may be partial. It may affect the majority of the muscles of the frame, or be limited to one muscle. It may be strictly confined to one side, or exist solely in the lower half of the body. It may come on rapidly, or appear slowly. But under any circumstances it is not a disease, but a symptom. The causes which give rise to paralysis may be thus summed up:

*Paralysis due to a lesion or any morbid condition of the nervous centres.*—Hemorrhage into or softening of the central nervous textures, or any other process which materially alters or compresses them, or interrupts the main conducting paths, occasions loss of power in the part over which their influence in health extends. The complete paralysis attending most of the diseases of the brain and of the spinal cord belongs, therefore, in this category.

But besides these palsies of organic origin there are *functional palsies*, dependent upon what, so far as we are aware, is simply a functional derangement of the great centres of innervation. Hysterical paralysis, and that occurring after overwork or excesses, and from nervous exhaustion, are examples.

*Paralysis due to a lesion in the course of a nerve.*—The nervous force may be properly generated, but the nerve-fibres may be incapable of conducting it. For instance, if a nerve be wounded or compressed, paralysis of the muscles which it supplies takes place. Palsy from this cause is local, and is apt to show marked nutritive changes in the affected part, such as glossy fingers and swollen joints, and to be associated with pain.

*Paralysis due to an affection of the nerves at their extremities.*—An illustration of such a disorder is the palsy resulting from exposure to cold. Peripheral palsies lead quickly to atrophy of the muscles. They are, from their very nature, local, and commonly remain so. But many peripheral nerves may become implicated, and extensive palsies result, as seen in multiple neuritis.

Motor paralysis due to cold may be met with as a family affection. It has been noticed as thus happening in twenty-two persons, and is clearly described by Rich.\* On exposure to cold and damp, especially after depressing conditions, the muscles become fixed and immovable in toxic spasm and enlarged. Respiration, cardiac action, cerebral phenomena, and sensibility are unchanged; the muscles of deglutition are affected if cold substances be swallowed. There is intense desire to urinate, but no derangement of micturition; the sphincters are undisturbed. Motor power returns gradually and progressively on exposure to heat. Recovery is followed by a sense of exhaustion. The disease is hereditary through many generations. It affects both males and females, and does not skip a generation.

*Paralysis due to reflex action.*—Here the paralysis is produced through the reflex centres, which reflect the irritation communicated to them to parts healthy in themselves. We see excitation of the dental nerves in teething children, disorders of the intestines both in adults and in children, or disease of the bladder, urethra,

---

\* Medical News, Aug. 25, 1894.

prepuce, uterus, lungs, or pleura, or irritation of the nerves of the skin, occasion paralysis, and a wound of a nerve on one side of the body lead to palsy on the other. The cause of the affection is peripheral irritation, or rather, as was insisted upon by Leyden, and is more and more believed, a true neuritis of ascending type.

Paralysis brought on by reflex action is rarely of long duration. It develops gradually, is increased or diminished as the causes which produce it increase or diminish, and, as a rule, soon disappears after the source of disturbance is removed. It may affect almost any part of the body, and assumes often the paralytic form.

*Paralysis due to serious interference with the circulation.*—This kind of palsy is observed if the principal artery of a part be obliterated. It is sometimes noticed as a transient phenomenon after the ligation of a large artery. If the vascular supply of the brain be interfered with by the occlusion of a vessel, whether by embolism or by thrombosis, the hemiplegia that results is more permanent and very marked. Among the circulatory disturbances that may lead to palsies we must not forget to look for the altered blood-tension produced by disease of the heart, and the degeneration of the vessels caused by Bright's disease.

*Paralysis due to a morbid state of the muscles.*—It is doubtful if it be correct to call that paralysis in which the nervous system is not primarily or particularly concerned. Yet certain forms of rheumatic palsy and of muscular atrophy in which the nervous implication is uncertain, but which entail loss of muscular power, may be classed here.

A loss of muscular power simulating paralysis is seen in *myasthenia gravis pseudo-paralytica*, the designation given by Jolly\* to a peculiar condition characterized by undue readiness of fatigue of voluntary muscles after ordinary functional activity. In fatal cases no distinctive or constant lesion has been found.

*Paralysis due to the presence of poisons in the system.*—The toxic effects of lead, of arsenic, of mercury, of alcohol, and of sulphuret of carbon, may exhibit themselves by producing palsy.

---

\* Berliner Klinische Wochenschrift, 1893, No. 1, p. 1.

Malarial poison, and poisons formed in the system, such as that of rheumatism or of gout, may act in the same way. The former occasions that singular "intermittent paralysis" which may come on either as one of the phenomena of a fit of ague, or as an apparently independent complaint, which may assume either the quotidian or tertian type, and in which both sensation and motion may be affected. How these poisons operate, whether by interfering with the nutrition of the nervous centres and weakening their generating force, or by enfeebling the conducting power of the nerves, or, as some of them undoubtedly do, by setting up a neuritis, is not fully determined. The palsies coming under this head, being for the most part functional, are not ordinarily intractable. Those due to malaria show the malarial corpuscles in the blood,\* and yield speedily to decided doses of quinia. Similar to the palsies of poisons and certain cachexias are the palsies after acute diseases. Yet structural changes have been found in these paralyses of supposed blood origin, and many of them are owing to neuritis.

All cases of periodic paralysis are not due to malaria; a number of instances have, indeed, been recorded which were probably not.† They are characterized by transitory and recurring muscular weakness of varying degree and distribution, but without other constant or distinctive symptoms. Sometimes there is diminished electric irritability of the affected muscles. There may be also enfeeblement of the reflexes, sensory phenomena, and increased thirst. In some cases, further, a family tendency is present.

In the parts affected with paralysis the nutrition and secretion are disturbed and the circulation is sluggish. They are frequently swollen and œdematous, the pulse is weaker than in the sound members, and the sensation is impaired. The nails grow slowly, so do the hairs; the perspiration is defective; the skin feels cold, is prone to break from the effect of pressure, or even indepen-

---

\* See a paper published by me in the *International Clinics*, 1891, vol. iii.

† See Westphal, *Berliner Klinische Wochenschrift*, 1885, Nos. 31, 32; Goldflam, *Wiener Medicinische Presse*, 1890, Nos. 36-39; Burr, *University Medical Magazine*, Aug. 1893, p. 836; Schacknowitsch, cited by Burr. *loc. cit.*; Cousot, *Vratch*, No. 32, p. 537; *London Medical Record*, 1884; and Hirsh, *Deutsche Medicinische Wochenschrift*, 1894, No. 32, p. 646.



dently of it, and the ulcers, if they heal at all, heal but tardily. The condition of the muscles is various. In some cases they are completely relaxed, in others rigid; at times they become agitated with convulsive movements. These phenomena are most evident in palsies of organic origin, especially in those dependent upon a brain-lesion, and in those due to disease of the spinal cord in which anæsthesia is present. Where hyperæsthesia occurs, the increased sensibility is attended with a larger supply of blood and a higher local temperature.

At times there are involuntary movements in the paralyzed parts. Thus, in cases of hemiplegia there may be automatic movements in the palsied arm when the patient sneezes, or some action in the muscles of the face to cause expressions in connection with those of the sound side. Again, muscles which are commonly associated in bilateral action no longer exhibit this, or an attempt only at action is perceptible; or certain muscles which are in the habit of acting together to cause a particular result do so no longer. Further, rotation of the head and neck to the same side as the one to which the eyes are directed—lateral or conjugate deviation of the eyes—is observed to be peculiar; the unopposed muscles turn the head and eyes towards the unparalyzed side. This symptom is mostly transitory, but is generally found in sudden marked hemiplegia.


Having examined some of the general traits and the causes of paralysis, let us investigate its chief varieties with reference to their significance and diagnosis. In so doing, it will be convenient to be guided by the marked coarse features rather than by the presumed origin.

But before inspecting these we shall briefly inquire into the *mode in which palsies are investigated at the bedside*. We ascertain, of course, the size, appearance, and feel of the stricken part; take notice of its growth, and of the nutritive changes, such as alterations in look and action of the skin, the presence on it of eruptions and of breaks, the state of the cutaneous circulation, of the nails, the hair, and the joints. Then we test the sensibility to contact, to tickling, to pinching, to heat and cold; measure the tactile sense by the æsthesiometer; ascertain the muscular sense; and carefully note any reflex movements that may be produced in the apparently lifeless limb, contrasting them with those of the sound

limb. We next, where minuteness of investigation is desirable, ascertain the surface temperature; and pass on to a thorough study of the condition of the muscles and of muscular motion.

Now, in examining the muscles we do not find them more wasted than their disuse will account for; certainly not in palsies of cerebral origin. Moreover, we generally observe them to be flaccid,—rigidity, especially early rigidity, being rare; but a stiffening associated with pain in attempts to straighten the contracted part is not so rare where the palsies have been of long standing, and has had, as we shall see presently, a special meaning attached to it. Then, irrespective of the condition of flaccidity or stiffening, we must look into the degree of abolition of muscular motion, carefully contrasting it, when one-sided, as indeed we must all the phenomena under investigation, with the movements of the other side. Is the motion completely abolished, or only impaired? what muscles particularly are affected? are concerted movements possible? and how is the gait, if we are testing the muscles of the legs, during these movements? Moreover, what amount of muscular effort is required to overcome special resistance? how is the balancing power? and how are delicate and combined movements executed when the eyesight is withdrawn? When the power in the arms is only impaired, not lost, we ascertain the degree roughly by the strength of the grasp. But we can do so accurately by a dynamometer. Of these, the best is that of Mathieu (Fig. 12), consisting of a steel ring, slightly elastic, which

FIG. 12.



is pressed firmly in the hand and records the pressure. The degree of swaying of the patient may also be accurately measured.

But the most valuable agent to judge of the state of the muscles is *electricity*, especially the forms of it known as the induced current, or “faradization,” and the constant current, or “galvan-

ization," and the action of each must be separately studied. The parts to be examined should occupy similar positions, and the muscles of the sides to be compared should be in equal condition of tension. We must begin with a weak current, and the wet electrodes are placed, one on the muscle itself, the other on some other part of the muscle or some indifferent point. This is the *direct excitation* of the muscle. Or the muscular action may be evoked by stimulating the motor nerve supplying the muscle to be tested. This is *indirect excitation*; and in healthy muscles the same strength of current will produce the same amount of contraction whether muscle or motor nerve be stimulated. It is also important to break the current by slow interruptions, and, especially in employing the galvanic current, to compare the positive (anodal) and the negative (cathodal) opening and closing contractions of the diseased with those of the sound side. In both currents, too, we should ascertain what the quantitative changes are,—whether the muscles react under a feebler current than is usual, or require one of great strength to move them. The response depends upon the integrity of both the muscle and the motor nerve. If this be preserved, contraction takes place with every change in a current of sufficient force. With every interruption and with each establishment of the current the muscle can be seen to contract in health, provided the intervals between breaking and making of the current be not too short. If, however, sufficient time do not elapse for the contraction to give way to relaxation, a tetanic state results. The readiness of response to the faradaic current is alike, whichever pole is applied to the muscle or nerve, and also when the current is made and broken; but, with the galvanic current, the readiness of reaction varies both with the electrode used and with the making and breaking of the current.

*Diminished or lost electro-muscular contractility* is a most valuable sign in destructive diseases of the cord. Indeed, speaking in general terms, we may say that it belongs to spinal palsies, while the electro-muscular contractility is intact in cerebral palsies. But this is only true of spinal palsies when the muscles are separated entirely from the influence of the cord: those supplied by nerves having their origin in healthy spinal texture preserve their normal irritability. In fact, if the uninjured part of the

cord have become irritated, or more vascular, the muscles having a nervous connection with it may show increased susceptibility to the electric current, and more energetic contraction. We also find diminished electro-muscular contractility when the nerve itself is injured, and it then comes on very quickly; when there is a mere local change in the muscular texture of the helpless part; and as the result of certain poisons, as of opium, lead, rheumatism, or other blood-poisons, which lower the power of nerve, of muscle, or of nerve-centre. We find it also when there has been long disuse of a limb, as in old cases of hysterical palsy, and even of cerebral palsy. But under these circumstances it is temporary, not permanent; for using the battery for a few days makes the greatest change in the electro-muscular contractility.

As already stated, the electro-muscular contractility is normal in the forms of palsy due to brain disease. The palsied limb may have, indeed, its muscles more powerfully convulsed by a current of the same intensity than those of the sound side are, and then we may infer, as Todd \* and Althaus † have shown, that the paralysis is due to brain disease of an irritative character. In recent hemiplegias, whatever their origin, increase of electric excitability is not uncommon. The response of muscle to faradaic stimulation is called *faradaic excitability*; and the remarks made are based on the effects obtained by faradization.

With reference to the galvanic or continuous current, or *galvanic excitability*, we find that in a healthy state of the muscles the galvanic current will give the same results as faradization, whether muscle itself or its motor nerve be acted on. Healthy muscle and nerve react most readily to galvanic stimulation when the negative pole is applied and the current is broken, and, successively, when the positive pole is applied and the current is either broken or made, and, finally, when the negative pole is applied and the current is made. This so-called "normal formula" may be represented graphically as follows:

K.Cl.C; { An.Cl.C.; K.O.C.; in which K (kathode) stands for  
 An.O.C.;

the negative pole, An (anode) for the positive pole, Cl (closure)

---

\* Clinical Lectures on the Nervous System.

† Medical Electricity.



for the making, and O (opening) for the breaking of the current, and C for the muscular contraction. In diseased conditions galvanism may show the same or it may show different reactions from faradism. The muscles of a palsied part may respond actively to galvanization and not at all to faradization. We observe this when the muscular tissue has begun to atrophy and to degenerate in consequence of extensive lesions of the cord, in degenerative affections of the motor roots, in traumatic nerve lesions, and in diseases of the peripheral nerves. While the faradaic excitability declines or is lost, the galvanic excitability not only remains, but may be even exaggerated; and in this "reaction of degeneration" (De. R.) there are also complete changes in the normal laws of electric muscular contraction: the anodal closing contraction equals or even exceeds the cathodal closing contraction, the cathodal opening contraction declines in the same manner. There is a deviation from the normal order of response, and thus we note qualitative and not merely quantitative modifications. Again, we may find dissimilarities by interrupting the galvanic current, and these may vary whether the current be rapidly or slowly broken. Thus, Russell Reynolds\* has shown us that in certain instances of facial palsy from exposure to cold, or in paralysis of the limbs from the same cause, or in lead palsy, the muscles act as little under the rapidly-interrupted galvanic current as under faradization; but if the galvanic current be slowly interrupted, they exhibit a greater amount of irritability than do the healthy muscles. In these cases it is found that the muscles are primarily affected, and the application of slowly-interrupted galvanism is rapidly of much service. It is, indeed, well in all cases of palsy, whatever be the form of battery employed, to note the differences in the contraction of the muscles produced by slow or by rapid interruptions. *Static* or Franklinic electricity may also be employed for purposes of diagnosis. We meet with instances where muscles contract under its use which do not respond to either the faradaic or the galvanic current.†

As already stated, a muscle may be indirectly acted on; one

---

\* Clinical Uses of Electricity, London, 1873.

† See an excellent summary of the diagnostic value of Franklinic electricity by Bernhardt, Samml. Klin. Vort., No. 41, Feb. 1892.

moistened electrode is placed over the motor nerve which controls the muscle, the other over its body. In inflammation of the nerve both galvanic and faradaic irritability of the muscle is increased; in destructive injuries it lessens and disappears. It is always well to note the indirect as well as the direct muscle excitation. But it has not, for purposes of diagnosis, proved itself as generally valuable. We should endeavor to place the one or other of the sponges exactly over the seat of chief nerve-supply in the muscle; and the ascertainment of the nerve point or points that correspond with the entrance of the motor nerves into the muscles has been made a matter of much study. Experience, indeed, proves that from these motor points, determined with infinite care and labor by Ziemssen,\* the readiest control of the muscles is obtained.

When the muscles react under electricity the contraction is felt, and the "electro-muscular sensibility" is more decided the stronger the contraction. Hence we almost always find increased electro-muscular contractility with increased electro-muscular sensibility. But the latter may exist alone, as we mostly observe in myalgias. On the other hand, the relationship between diminished contractility and sensibility may be changed, as we find, for instance, in the striking want of sensibility to the current in hysterical paralysis. The electric reactions of the skin, well tested by a metallic brush, as a rule go hand in hand with the reactions of the muscles, increase in sensitiveness with them, decrease with them.

Such are the chief facts with reference to the diagnostic applications of electricity in paralysis. There is yet another mode of investigation which we constantly bring into use, one also in which the action of the muscles particularly gives us valuable information concerning the state of the nervous system,—the testing of the *reflex excitability*. But we have already examined into the derangement of the reflex system, and shall only here add a few general clinical facts. We find the reflex excitability diminished in disease of the gray substance of the cord, in disease of the sensory root-fibres, which thus become incapable of conducting the impression, and in disease of the motor fibres, which fail to impart

---

\* Die Electricität in der Medicin; also Tibbit's Handbook of Electricity, and Bartholow's Medical Electricity.

the motor impulse. In the latter case there is coexisting paralysis of motion; in the second, anæsthesia. Increase of reflex excitability, producing twitching or even violent irregular movement on very slight stimulation, is found in all irritative lesions which have increased the excitability of the gray substance of the cord, as when this is disturbed by inflammation, or compressed by a tumor, or heightened by certain drugs, such as strychnine. Increase of reflex excitability is also found in parts below a lesion, when this gives rise to descending degeneration in the pyramidal tracts.

As regards the action of the brain, there are instances in which, if all power of appreciating impressions be lost, as in overwhelming cerebral apoplexies, reflex action may be everywhere suspended. On the other hand, we find signs of reflex action manifesting themselves by irritation transferred from diseased to healthy parts of the brain, producing spasms or palsy phenomena, alluded to in the sketch of the seat of cerebral lesions. Nor must we from a clinical point of view omit to mention the reflex actions excited in other parts of the body, as from diseases of bones and joints, or the muscular contractions in the legs during catheterization or in colics. Here the seat of the perverted reflex action is entirely in the reflex areas of the cord.

All these remarks tell us how to examine paralysis. Having now studied the modes in which this is investigated, I shall merely recall that to find out the cause of the difficulty we have to take into account the history of the case, and the attending symptoms, nervous and otherwise; and in eliciting these we should never forget to bring out prominently those shown us by the ophthalmoscope, and by examination of the urine and of the heart.

Let us return to the clinical study of palsies.

#### HEMIPLEGIA.

We shall first consider that form which almost always results from brain disease,—hemiplegia, or one-sided palsy. This state of things may affect all the voluntary muscles on one side of the body; but it generally exists only in those of the limbs and face; the eye, neck, and trunk muscles escape. Neither the legs nor the arms can move, and the muscles of the face on the side corresponding to the paralyzed limbs are motionless. The cheek hangs; the

mouth is drawn toward the healthy side, because the muscles on the other are powerless to resist; the tongue, when protruded, is ordinarily slowly pushed out toward the palsied side; the articulation is imperfect.

But the rule with respect to the face being paralyzed on the same side as the rest of the body has its exceptions. Indeed, when we reflect that the nerves which supply the facial muscles are given off above the point of decussation of the nervous fibres in the cord, it seems perplexing that it should be a rule at all. The solution of the question lies in the crossing of the facial nerves. Should, then, the lesion be seated in the brain above this crossing, both face and body are paralyzed on the side opposite to the diseased spot. Should, however, the lesion involve the facial nerve-fibres at a point below or after the decussation, there will be paralysis of the face on one side, and of the limbs on the other, the facial palsy being direct, and that of the body being crossed.

Now, according to Gubler,\* this *cross paralysis* is always indicative of a lesion of the pons Varolii, close to which the facial nerves originate, and through which the nerve-fibres for the limbs pass before they decussate lower down. But we must remember that there are rare cases of "alternating hemiplegia," due to a combination of lesions, one affecting a cerebral lobe on one side and the facial nerve on the other. Even when the lesion is unilateral, we may meet with exceptional cases; and, as Bastian† points out, the lesion may be situated in the pons, the palsy of face and limbs not being alternate, provided the disease occur in the upper or anterior part of one lateral half, implicating the fibres of the facial above their sites of decussation. With reference to the other cerebral nerves, should we find any of them paralyzed on one side and the body on the other, we shall generally be correct in assuming that the palsy is not due to disease on both sides of the brain, but is rather a disturbance of the affected nerve near its origin or in its course, and on the side on which the brain is injured, while the paralysis of the limbs is on the opposite side. Anatomical researches which have traced con-

---

\* De l'hémiplégie alterne envisagée comme signe de lésion de la protubérance annulaire, *Gaz. Hebdom.*, 1856, 1859.

† Paralysis from Brain Disease.



necting nuclei on the floor of the fourth ventricle and elsewhere explain these alternating palsies.

Hemiplegia, as already stated, results, in the vast majority of instances, from cerebral diseases. Hence we find it commonly associated with disordered mental powers, and other signs of a brain-lesion. The superficial reflexes are, as a rule, though not invariably, diminished; the deep reflexes are exaggerated. The rectum and the bladder perform their functions.

Hemiplegia caused by an affection of one-half of the spinal cord, near its beginning, is not combined with a decay of the mental faculties, but the muscles of the chest and abdomen are involved in the paralysis, which they are not in cerebral hemiplegia, unless the lesion be very extensive. Then in *spinal hemiplegia* there is a zone of anæsthesia on a level with the lesion, and coexisting anæsthesia, as Brown-Séquard has shown, on the side opposite to the lesion and the muscular palsy, and the temperature sense is impaired, as is the sensibility to pain; the palsied limb gives evidences of vaso-motor paralysis, has at first a higher temperature, and is hyperæsthetic; reflex action is increased on the side of the lesion, the muscular sense is impaired, and the umbilicus is with every act of inspiration drawn toward the sound side. We possess a further test in electricity: unlike what we find in cerebral paralysis, the electro-muscular contractility is greatly lessened or is lost. Spinal hemiplegia, or "hemiparaplegia," as it is more often called if the lesion be low down, occurs from injuries, tumors, syphilitic disease of the cord, and localized sclerosis.\* According to Romberg, spinal hemiplegia is more persistent in the leg than it is in the arm. In hemiplegia due to cerebral disease recovery is more rapid and more nearly perfect in the leg than in the arm.

But supposing that we have settled the hemiplegia to be cerebral, the points next to be investigated are, where is the lesion situated? and what is its nature? Now, the former question, concerning the *anatomical diagnosis*, may be answered in a general way by stating that the disease is on the side opposite to the palsy,

---

\* Cases by Charcot and Gombault, and by Troisier, *Archives de Physiologie*, 1873; by Riegel, *Berlin. Klin. Wochenschrift*, 1873, and by F. Remak, *ib.*, 1877. See also Gowers, *Diseases of the Nervous System*, 2d ed., 1893.

if the lesion, as it almost always is, be seated above the point of decussation of the pyramidal columns of the medulla; for a lesion below the decussation gives rise to palsy on the same side, and a lesion on a level with the decussation, to double-sided palsy.

Lesions of the posterior segment of the internal capsule give rise to typical hemiplegia, sometimes with hemianæsthesia and loss of the special senses.

Lesions of the corpus striatum only cause motor and sensory symptoms when they involve the internal capsule. The same is true of disease of the optic thalamus, except that mobile spasm and incoördination of movement have been observed to follow lesions of its middle third.

The nearer the lesion to the surface, the more marked are the mental phenomena, the greater is the tendency to spasms in the limbs, but the more limited is the palsy; and the farther the disease extends toward the internal capsule, the more extensive does the paralysis of motion become. We may further distinguish the palsy which ensues from that caused by an affection lower down, as of the *pons Varolii*, by observing that, besides the peculiar crossed paralysis of the face and limbs, we find giddiness and a tendency to vomit; either loss of the conjugate movement of the eyes toward the side of the lesion, or conjugate spasm with nystagmus; jerkings of the muscles of the face on the side opposite to the injury; sensations of tickling in the face; one-sided facial anæsthesia, with a loss of sense of taste on the corresponding side, though with unimpaired motion of the tongue; rigidity of the limbs, and spasm of the muscles supplied by the fifth nerve; disturbance of respiration and of the heart; albuminuria; glycosuria; high temperature. Should we encounter paralysis of sensibility and motion on one side of the body, and both sides of the face be palsied as to motion and sensation; should the recti muscles of the eye be paralyzed, and taste be lost over the anterior part of the tongue, we may infer that the injury is seated rather above the lower portions of the pons, and affects the spot where the facial nerve and part of the trigeminal cross.\* Hyperpyrexia is not uncommon after the onset of an acute lesion of the

---

\* Brown-Séquard, Dublin Quart. Journ., May, 1865.

pons, and in acute lesions convulsions\* also are common, as is marked contraction of the pupils. In lesions involving the central parts of the pons, paralysis, mostly unequal, of both sides of the body, with impaired sensation, irregular facial palsy, difficulty in deglutition and articulation, is the rule.

Lesions of the lower and inner part of the *crus cerebri* Hermann Weber has taught us to recognize by an alternate paralysis, in which the third nerve is palsied on the affected side of the brain, showing us want of action of the muscles of the eyeball, except the external rectus and superior oblique, with a dilated pupil, a tongue deviating to the paralyzed side, some difficulty in articulation, the body-palsy marked in the arm and leg, and co-existing with local temperature higher by several degrees, vasomotor disturbance, and very defective sensation.

Acute lesions of the medulla are likely to destroy life; in case they do not, the resulting symptoms are often bilateral and include derangement of the functions of the bulbar nerves.

Besides these well-attested facts, the brilliant researches of the day on the localization of cerebral functions have solved, and are still solving, many problems as important to the physician as to the physiologist. Let us look at some of the additions to pathological knowledge which appear the most certain, premising that in localization only symptoms that are permanent are of value, since any lesion, an acute lesion especially, may for the time being cause vascular or inhibitory disturbance in adjacent parts. We must also be mindful of Broadbent's law, that one-sided movements can be excited from either hemisphere, and that the loss may be soon compensated by the hemisphere with which they are not habitually associated. This becomes often manifest in damages of the cortex.

We shall first glance at lesions of the motor zone, including the ascending frontal and parietal convolutions, the anterior two-thirds of the superior parietal lobule, and the paracentral lobule, parts supplied by branches of the middle cerebral artery. A lesion of these cortical structures causes paralysis of motion without loss of sensation. The hemiplegia is more or less complete according to the extent of the motor area involved. It is

---

\* Gowers, Diseases of the Nervous System.

on the opposite side to that of the disease, and neither the nutrition nor the electric contractility of the palsied muscles is impaired.

The *cortical hemiplegia*, when sudden, is less frequently accompanied by loss of consciousness, is rarely complete from the first, affecting, perhaps, at the onset only the face, an arm, or a leg, and is soon followed by rigidity of the palsied parts. But, on the other hand, it is more apt to be transitory, to show slighter differences in temperature between the two sides, and to be accompanied by localized pain in the head, which may be elicited by percussion over the seat of lesion.\* Limited palsies, monoplegias, are much more common in disease of the cortex than in disease of deeper parts. The leg alone is affected in lesions of the medial cortex or those near to the longitudinal fissure. Irritative lesions of the cortex have as their most characteristic sign unilateral convulsions. In disease of the middle third of the central convolutions the convulsions generally begin in the hand. Disease of the ascending frontal convolution, behind the inferior frontal, gives rise to paralysis of the face, lips, and tongue.

Lesions confined to any one of the *gray central ganglia*, where the internal capsule is not involved, do not afford any special feature by which they may be distinguished from common cerebral hemiplegia. There is paralysis of motion only, which, Charcot † tells us, is generally transitory. If the anterior two-thirds of the posterior limb of the *internal capsule* be involved, the palsy is still exclusively of motion, though it is more or less persistent, and ultimately accompanied by muscular contractions; if the posterior third be also involved, we have in addition cerebral hemianæsthesia. Smell may also be lost on the anæsthetic side, and hemianopsia be met with. In disease of the angle and posterior segment of the internal capsule we have hemiplegia of the ordinary type. Indeed, in lesion of the corpus striatum the hemiplegia is permanent only if the internal capsule be involved in the damage.

A lesion of *one optic tract* or of the cortical visual centre in the occipital lobe will cause bilateral homonymous hemianopsia; a

---

\* Ferrier, *Localization of Cerebral Disease*, 1879.

† Lectures on *Localization in Diseases of the Brain*, New York, 1878.



similar effect is sometimes produced by a lesion of the corpora geniculata. There may be considerable hebetude, but no other marked symptom of an affection of the brain except hemianopsia. In lesions, also, of the *præfrontal lobes*, that part which, in its relation to the skull, is roughly bounded by the coronal suture, there is no disorder either of motility or of sensibility. The manifestations are simply those of restlessness and unsteadiness of mind, impairment of judgment and reason, and other psychical disturbances. There is no motor paralysis except of the foot. Late in the case, among pressure and invasion symptoms, we may find motor aphasia, nystagmus, and unilateral convulsions.\* In disease of the *temporo-sphenoidal* lobe we have deafness in the ear opposite to the lesion, and sometimes convulsions with preceding auditory aura. There is no hemiplegia.

The nature of the paralyzing lesion, the *pathological diagnosis*, can be arrived at only by a careful scrutiny of all the facts of the case. A sudden paralysis occurring simultaneously with coma almost always has its origin in an apoplectic effusion, more rarely in cerebral embolism or thrombosis. A sudden paralysis without coma is generally due to plugging of the vessels. A gradual development of palsy indicates some chronic cerebral disease, such as chronic encephalitis, or a tumor, or any affection compressing the nervous substance. We may also gain much knowledge by carefully exploring the organs of circulation and the kidneys. Thus, a paralysis found to be conjoined to a cardiac malady or to a diseased state of the arteries is, in all likelihood, owing to a clogging of one of the cerebral arteries, and to consequent tissue-change in the cerebral structures. When the kidneys are seriously disordered, the hemiplegia is likely to be caused by some chronic disease of the brain or its vessels, the result of an altered nutrition. The uræmic condition itself seems also capable of causing loss of power, sometimes of hemiplegic type.

A further clue to the character of the cerebral lesion is obtained by examining the palsied muscles. Todd † has taught us that when the paralyzed limbs exhibit a rigid state from the moment of, or soon after, the attack, we may from the *early rigidity* assume

---

\* Mills, *Cerebral Localization in its Practical Relations*, 1889.

† *Clinical Lectures on the Nervous System*.

the lesion to be of an irritative nature, such as an inflammation, or a compression of healthy brain-tissue by an apoplectic clot or by an accumulation of puriform fluid in the subarachnoid spaces. When the muscular contraction does not take place until late in the complaint, *late rigidity* is before us, and becomes associated with wasting of the muscles. The opinion of the day connects this late rigidity with a descending degeneration of the motor tracts. It depends on active muscular contracture, and lessens during sleep and if the limb be immersed in hot water. It is generally associated with excessive tendon reflexes. Under excitement the paralyzed arm and leg may be strongly flexed, and automatic movements may occur when the patient sneezes.\*

When hemiplegia has been of long standing, late rigidity may be combined with atrophy of the muscles and other nutritive changes that bespeak a secondary degeneration, spreading into the opposite lateral column of the spinal cord; also tremors, associated not unusually, as Charcot tells us, with diminution of sensibility on the palsied side; attacks of true spasms, happening particularly in the arms; joint disease and nodes; and choreic movements, a condition to which, under the name of "post-paralytic chorea," Weir Mitchell † has called attention. In some cases of hemiplegia there is much pain in the stricken limb. The pain may precede returning motion, and is thus of favorable augury. But in limited disease of the internal capsule affecting the sensory path the pain in the palsied limbs may persist through life.

Hemiplegia may be *feigned*.‡ But the results of electricity, especially where altered sensibility as well as defective motion is simulated, and the test proposed by Hughlings Jackson, that the arms do not, as in real hemiplegia, fall forward when the patient stoops, but are retained at the side, will usually detect the fraud.

#### MONOPLEGIA.

When we have limited lesions we have limited palsies, and researches on localization are teaching us more and more accurately to recognize the centres affected in these palsies of special

---

\* Ross, *Diseases of the Nervous System*, 1883, vol. i. p. 187.

† *Amer. Journ. Med. Sci.*, Oct. 1874; also *Med. News*, April, 1893.

‡ For an instructive case, see *London Lancet*, April, 1874.

parts, or of one limb, or of a group of movements. Of course, in making a diagnosis of the paralysis being due to disturbance of a special nerve-centre, we must be careful to exclude, as the cause of the local palsy, peripheral affections, and those in the course of the nerve supplying the stricken part, and also to make it clear that the lesion is not spinal of very circumscribed kind. In monoplegias the palsy is never complete. Furthermore, it is always important to endeavor in a given case to separate the symptoms which may be due to invasion of or to pressure on adjacent centres from the localizing symptoms of the main lesion. Let us now take up some of the limited palsies dependent on cerebral disease, especially in the motor areas of the cortex.

*One arm only is paralyzed.*—Here we find the lesion in the ascending parietal and the ascending frontal convolution on the side opposite to the palsy, and the disease is limited to the middle third of the convolutions. If the lesion be double, as in a case referred to by Bourdon,\* both arms are helpless. But, whether single or double, with the damaged motion there are unimpaired sensation and electro-motor contractility. Disease of the ascending frontal opposite the upper half of the inferior frontal convolution gives rise to palsy of the lower part of the face except the lips.

*One arm and the same side of the face are paralyzed.*—In this “brachio-facial monoplegia” the lesion is in the central region of the cortex, toward the middle or lower third of the ascending convolutions in the facial and arm centres. It is a pure motor palsy, associated, however, usually with aphasia when the disease is left-sided. The main movements of the muscles of the upper part of the arm are kept, while those of the hand are lost. Palsy, of cerebral origin, limited to *one side of the face*, without the arm being implicated, is rare; the cortical disease is in the centre for the facial region. The affection is usually left-sided, and is apt to become complicated with aphasia. The lower part of the face bears the brunt of the palsy; unlike Bell’s palsy, the orbicularis and the upper part of the face are but little, if at all, disturbed;† further, there is no disease of the temporal bone to

---

\* Bull. Soc. Anat., 1874.

† This was strikingly illustrated in a case reported by Guitéras, Phila. Med. Times, Nov. 1878.

explain the localized palsy by an injury to the facial nerve. The tongue is also very generally implicated.

*The leg only is paralyzed.*—This is a very rare form of paralysis, and presupposes a lesion limited to the motor centre for the leg. The centre for the leg and foot is fixed by the recent researches of Horsley and Schaefer as occupying the uppermost portion of the ascending frontal and parietal convolutions. In some of these cases of “crural monoplegia” on record the ascending parietal and superior-parietal convolutions have been found diseased. Sensation is not affected; the arm is apt to become gradually involved in the palsy: in Ferrier’s case \* the lesion was in the quadrilateral lobule on the internal aspect of the hemisphere and in the upper extremity of the ascending parietal and frontal convolutions.

There are many other kinds of limited palsies of cerebral origin, such as of the tongue, *glossoplegia*, of the face and tongue, *facio-lingual monoplegia*, of the eye muscles, *oculo-motor monoplegia*, and half blindness, *hemianopsia*, to all of which I can only refer, since our knowledge is not definite enough to lay down concise conclusions for diagnosis.† In part, too, they will be discussed further on. It must, however, be added that in all these limited palsies traceable to disease of the brain we are apt to have such symptoms as are common in brain affection,—head-

\* Brain, vol. iii., 1880.

† As bearing on these and other matters connected with localization, see especially the works of Ferrier and Charcot referred to; Hitzig, in *Klinische Vorträge*; many papers in the *Archives de Physiologie*, in the *West Riding Reports*, and in *Brain*; Hughlings Jackson, *Clinical and Physiological Researches on the Nervous System*; Pitres, *Lésions du Centre Ovale*, and papers with Charcot, *Revue de Méd.*, 1879 and 1883; Nothnagel, *Topische Diagnostik der Gehirnkrankheiten*, Berlin, 1879; Exner, *Untersuchungen über die Localisation der Functionen in der Grosshirnrinde*, Wien, 1881; Wernicke, *Fortsch. der Med.*, 1883; Gowers, *Diseases of the Nervous System*, 2d ed., 1893; Seguin, *Amer. Journ. Med. Sci.*, vol. xvi., 1888; Allen Starr, *Amer. Journ. Med. Sci.*, 1884, 1885, and since,—containing, besides other valuable matter, in July number, 1884, the collected American cases of Cortical Lesions,—also *Medical Record*, Feb. 1886; Mills, *Cerebral Localization in its Practical Relations*, *Transactions of the Congress of American Physicians and Surgeons*, vol. i., 1889; Horsley and Schaefer, *Philos. Transact. Roy. Soc.*, 1888; Dana, *Medical Record*, May, 1893; Wylie, *Edinb. Med. Journ.*, Feb. to Sept. 1893.



ache, giddiness, and the like. These aid us in understanding the nature of the disorder.

Perhaps, too, we shall receive help from a means of diagnosis inaugurated by Broca,—*cerebral thermometry*; and a higher local temperature will point to the region affected. But the observations are not as yet definite enough to warrant their adoption, and what makes them very difficult of application is, that the temperature of distant parts has been found to be influenced by excitation of the surface, and that the difference in the disease itself materially modifies the temperature of the head. Thus, in embolism\* we have a lower temperature over the part which ought to be supplied by the occluded vessel; in inflammation and tumor and abscess the temperature is higher. Again, as we know particularly by the elaborate researches of Lombard, emotional activity, as well as or even more than intellectual work, causes a rise of temperature, the rise sometimes exceeding  $0.18^{\circ}$  Fahr. ( $0.1^{\circ}$  Cent.). Active exercise, Amidon states, may do the same. Thus the patient should be examined when free from excitement and at rest. Various portions of the head must be selected as points for the application of the surface thermometer, and the corresponding regions compared. To appreciate the results observed in disease, we must remember that the frontal region in health on the left side, which always registers more, gives, Broca tells us,  $95.7^{\circ}$  Fahr. ( $35.43^{\circ}$  Cent.); the parietal,  $91.49^{\circ}$ ; Gray records, in accordance with Broca, the left occipital region as  $92.66^{\circ}$ . The fact has already been mentioned that Maragliano and Seppili, making their observations in summer, give the mean normal temperature as higher by nearly two degrees Fahr. It is so in the frontal regions, and in the occipital region the difference is much greater. These authors inform us that in the insane the temperature varies according to the form of insanity. The highest temperature is found on the left half of the head, and not materially different on the left frontal region, in furious mania,  $36.9^{\circ}$  Cent. ( $98.4^{\circ}$  Fahr.); in progressive paralysis,  $36.6^{\circ}$  ( $97.9^{\circ}$  Fahr.); in imbecility, idiocy, and simple mania,  $36.3^{\circ}$  ( $97.3^{\circ}$  Fahr.); in simple dementia,  $36^{\circ}$  ( $96.8^{\circ}$  Fahr.). In locating brain tumors several observers have made use of the thermometer.

---

\* Broca, Bulletin de l'Académie de Médecine, Dec. 1879.

Gray \* cites a case, and Mills † and Seguin ‡ have published several instances ; Eskridge § has recorded some elaborate studies of the head temperature in abscess and tubercular inflammation. On the whole, the head temperatures are more steady than those taken in the axilla.

#### PARAPLEGIA.

This differs from hemiplegia in the palsy occurring on both sides, yet being, in the vast majority of instances, limited to the lower extremities. Its almost invariable cause is a lesion of the spinal cord. In truth, if we call hemiplegia paralysis from brain disease, we may call paraplegia paralysis from spinal disease. Paraplegia is generally due to a marked organic lesion ; but there are cases in which it exists independently of any recognizable structural change, and in which it results from poisons, from fatigue, from excesses.

The disorder generally comes on slowly. At first the patient only loses the steadiness of his gait ; gradually he is deprived of all power of motion, but the intellect and the nerves of special sense remain unaffected. If the lesion be in the lumbar part of the cord, the palsy is confined to the lower extremities and to the pelvic muscles ; if the dorsal portion be attacked, we find, in addition, signs of paralysis of the abdominal walls and of the sphincters, tympanites, and somewhat impeded breathing. In diseases of the upper section of the cord there is coexisting palsy of the upper extremities, with dilated, sluggish pupils, and difficulty in deglutition and in respiration. In the muscles supplied by nerves which originate in healthy marrow, involuntary contractions or reflex phenomena can be induced,—are, indeed, generally exaggerated,—and the striking effects of strychnine, when given in doses sufficient to produce its peculiar muscular spasms, are manifested. The palsied muscles, in the majority of the

---

\* New York Medical Journal, Aug. 1878, and Chicago Journal of Mental and Nervous Diseases, Jan. 1879.

† Phila. Medical Times, Jan. 1879, and New York Medical Record, Aug. 1879.

‡ Amer. Journ. Med. Sci., vol. xevi., 1888.

§ Transactions of the College of Physicians of Philadelphia, Third Series, vol. vi., 1883.

affections occasioning the paraplegia, undergo wasting and do not respond to the electrical stimulus.

Paraplegia is generally more marked on one side than on the other, and the paralysis of motion is apt to be associated with complete anaesthesia. When, as sometimes happens, the mischief is limited to a lateral segment of any part of the cord, there is paralysis of motion on the same side of the body, and of sensation on the other. Preceding, or even attending, many cases of paraplegia, is a symptom which belongs exclusively to affections of the cord: a spasm of the flexor muscles of the lower limbs, so powerful that the anterior parts of the thighs come almost in contact with the abdomen, while the heels are drawn up so as to touch the back of the thighs.\*

Let us now take a cursory view of the different forms of spinal paraplegia.

#### SUDDEN PARAPLEGIA.

**Spinal Hemorrhage.**—Sometimes the paralysis occurs suddenly, and in consequence of an injury to the spine, of a displacement subsequent to a disease of the bones, of blood extravasated into the canal, of poisons, as the lathyrus sativus,† or of bulbar or spinal disorder from sudden displacement of the cerebro-spinal fluid following blows on the head.‡ When either of the first two causes has led to the sudden palsy, the diagnosis is materially aided by the history of the case, and by a close examination of the vertebral column. But if there be no signs of a disease of the bones or of the intervertebral cartilages, we may suspect a *spinal hemorrhage* to have produced the sudden and complete paraplegia, or the palsy which, though at first partial, rapidly becomes complete; and this suspicion becomes much strengthened if violent localized pain in the back exist or have preceded the rapidly developed palsy, if the patient be unable to retain his urine or faeces, and if the affected limbs be relaxed and largely deprived of sensation. These are the symptoms of apoplexy of the cord. The seat of pain corresponds to the seat of bleeding.

---

\* Brown-Séquard's Lectures on the Nervous Centres, p. 114.

† Irving, Indian Annals, No. 12, referred to in Brit. and For. Med.-Chir. Rev., Oct. 1860.

‡ Duret, Traumatismes cérébraux, Paris, 1878.

The pain occurs in distressing paroxysms and passes along the course of the nerves compressed by the extravasation. Where the hemorrhage is meningeal, there is more persistent pain, with rigidity of the spine, spasms of the legs, slighter disturbance of sensibility, and far less and less quickly increasing paralysis, and there is more apt to be spasmodic retention of urine. The absence of early fever distinguishes the spinal hemorrhage from spinal meningitis; subsequent fever bespeaks the occurrence of this as a complication. The muscular spasm is sometimes so severe that it has been mistaken for tetanus, which lacks the violent pain in the back. The most common causes of spinal hemorrhage are blows and falls on the back or falls on the feet. It is also met with in diseases with hemorrhagic tendencies, in convulsive affections, and in the course of myelitis. Hemorrhage into the membranes may result from the rupture of an aneurism of a vertebral artery.

But, besides these causes, others lead rapidly to paraplegia. Softening of the cord may have progressed latently until the degeneration destroys the continuity of the conducting tubules, when palsy at once takes place. Then there are cases following sexual excesses, cases for which neither during life nor after death can an organic cause be assigned,\* and which must therefore be viewed as due to enfeeblement of functional power. Similar cases of spinal paralysis, more or less complete, may occur after fatigue and violent exercise. In all instances of spinal palsy due to impaired nerve-power—or *spinal paresis*, as Handfield Jones † has termed this affection—the disorder is much more apt to come on quickly than gradually, and a tonic treatment is likely to be followed by decidedly good effects. But in regard to all these cases of functional palsy, the same as in regard to reflex palsies, science is clearly narrowing their number by finding some organic affection in the cord, often secondary to an ascending neuritis.

**Acute Ascending Paralysis.**—Yet another variety of paraplegia which may happen rapidly is that form which has been described as *acute ascending paralysis*, or Landry's paralysis. It may come on after fatigue and exposure in persons in perfect

---

\* For instance, Case XVIII. in Gull's series in Guy's Hosp. Rep., vol. iv., 3d Series.

† Functional Nervous Disorders.



health, generally in men between twenty and forty years of age. Usually there is little or no fever except at the onset. Numbness and tingling, and slight pain in the lower extremities, are soon followed by loss of muscular power, which, in turn, goes on rapidly, generally in a few days, to complete paraplegia. The legs are relaxed and immovable, the muscles of the trunk are next affected, then the upper extremities become implicated, and sensation, which at first was normal, is somewhat enfeebled, though never to a marked degree; occasionally the arms are involved before the legs. The patient is restless, sleepless, but his intelligence is, as a rule, unimpaired, and we find no bedsores and no palsy of the bladder or rectum. The respiration and circulation in the progress of the disease become embarrassed, there is acute enlargement of the spleen, and sudden death ensues within a month from the time of the seizure,\* or, indeed, the case may end fatally in less than a week. But all cases do not run so rapid a course; and, in truth, we meet with instances in which the disorder is rather chronic than acute, or is arrested. The muscles do not atrophy, and their electrical excitability is unimpaired, which is a very valuable diagnostic test. About the reflexes the statements are conflicting. It is most likely that at first both the superficial and the deep reflexes are absent, and that they do not return, certainly the knee-jerk does not, except when the paralysis passes away. Jaccoud† tells us that in the cases he observed the reflex movements were abolished.

In Mills's case‡ which recovered, both the superficial and deep reflexes were completely lost. The malady is generally looked upon as being an affection of the peripheral nerves, though the central nervous system is not infrequently involved. Toxic influences of invading micro-organisms are thought to give rise to it.§

The disease which most resembles acute ascending paralysis is acute progressive, or multiple neuritis. But here sensation is rapidly lost, and so is the electrical excitability.

---

\* As in the case reported by Hayem, *Travaux de la Société Médicale d'Observation*, tome ii., 1867; see also Leyden's *Klinik der Rückenmarks-krankheiten*.

† Clinique Médicale.

‡ Transactions of the Assoc. of Amer. Phys., vol. vii., 1892.

§ Cramer, *Centralblatt f. Path.*, Jan. 1892.

**Multiple Neuritis.**—When nerve after nerve rapidly inflames, or the inflammation occurs at one time, an extensive palsy is quickly developed, the nature of which we have only of late years recognized. The disease is an affection of the peripheral nerves, though it has the misleading symptoms of a spinal malady. It attacks both sexes, is most common between the ages of thirty and fifty, and, though it may follow altered blood-states or rheumatism, or be due to exposure, by far its most frequent cause is chronic alcoholism. It has been observed in the sequence of a number of infectious diseases and also as a result of the medicinal administration of arsenic.\* It has generally an acute or a subacute beginning, with decided increase in temperature. At first vague, then more decided pains are felt in the extremities, chiefly in the fingers and toes, and these pains soon become darting or burning. The pain is often preceded by tingling, is increased by motion, and is associated with tenderness of the affected nerve-trunks and with both skin and muscle tenderness of the parts to which they are distributed; finally this increased sensibility may give way to anæsthesia.

The palsy shows itself generally first in the arms, the earliest loss of power manifesting itself in the extensors of both sides. Soon the muscular weakness is seen also in the legs, and the trunk muscles and face muscles may become involved. The symmetrical character of the palsy is very noticeable, as are also the wrist-drop and the foot-drop. The parts affected waste, and lose their reflex excitability; the loss of the knee-jerk is especially pronounced. The muscles do not react to faradization, though they may to the galvanic current; often, indeed, they present the reaction of degeneration: the nerves are uninfluenced by the electric stimulus. Œdema of the arms and legs is frequent. Sometimes muscular incoördination is the most prominent symptom.

The disease may run on to complete palsy of the limbs in less than two weeks, and death result from paralysis of the respiratory muscles; or the affection may pass into a chronic condition, and a slow improvement, with return of power in the muscles, take place. In protracted cases contraction of unopposed muscles may

---

\* Ralston, *British Medical Journal*, No. 1714, p. 996.

occasion deformities, and there are arthritic adhesions, glossy skin, and thickening of the skin.

The diagnosis is generally easy. The tingling in the extremities, the cutaneous and muscular tenderness, the distribution of the symptoms, the early development of muscular weakness, and the palsy of the extensors distinguish the disease from *rheumatism*. In some instances, where it is difficult to elicit tenderness of nerve-trunks, or where this symptom is wanting, where the muscular tenderness is not marked, where, moreover, the palsy is slight and incoördination of movement is observed, the similarity to *locomotor ataxia* is great, and the eye-symptoms of this affection alone, if present, will help to a correct conclusion. Further, girdle-sense and lightning-pains are absent in peripheral neuritis, while wasting is not generally observed in posterior sclerosis. In ordinary cases the greatest resemblance is to those instances of *acute myelitis* which run a rapid course, and especially those in which muscular wasting is marked. To *acute ascending paralysis* intense cases of the disease also bear a strong likeness.

In the following table are contrasted the features of multiple neuritis, of acute myelitis, and of acute ascending paralysis.

MULTIPLE NEURITIS.	ACUTE MYELITIS.	ACUTE ASCENDING PARALYSIS.
Fever, with at first decided elevation of temperature.	Fever generally moderate.	Slight, if any, elevation of temperature.
Palsy begins in forearms, extends to legs and trunk; double-sided wrist-drop and foot-drop; palsies symmetrical.	Palsy generally affects only legs and lower part of trunk, though it may affect arms.	Paralysis rapidly extending from lower extremities; relaxed muscles.
Muscles atrophy. Trophic changes in skin and nails common; no bedsores.	Muscles atrophy rapidly. Trophic changes marked; bedsores.	No muscular atrophy. No trophic changes; no bedsores.
Marked pain and sensory disturbances, hyperæsthesia especially, later anæsthesia in the area of distribution of the inflamed nerves; muscular tenderness; tenderness of nerve-trunks.	No pain, or tenderness of nerve-trunks; complete anæsthesia below lesion; zones of hyperæsthesia corresponding to lesion.	No marked pain or more than dulling of sensation in affected parts; no tenderness of nerve-trunks.

MULTIPLE NEURITIS.	ACUTE MYELITIS.	ACUTE ASCENDING PARALYSIS.
Loss of electrical excitability. Generally reaction of degeneration.	Loss of electrical excitability. Generally reaction of degeneration.	No change in electrical excitability.
Reflex action lost, especially deep reflexes and muscle reflex.	Excessive reflex action, except in parts deriving nerve-supply from injured centres, there lost.	Absence of reflexes the rule.
Sphincters unaffected.	Sphincters affected early.	Sphincters nearly always escape.
No bulbar symptoms, though respiratory palsy may happen.	Bulbar symptoms rare; failure of respiratory power may happen.	Bulbar symptoms frequent.
Mind unaffected, except in the alcoholic and toxæmic cases.	Mind unaffected.	Mind remains clear.

The various causes of multiple neuritis give rise to some differences in the symptoms, by the close study of which we may infer the cause. Thus, in neuritis from metallic poisons the disorder is confined to the arms, as in lead-poisoning, or is found in the arms first and subsequently attacks the legs, as in the neuritis of arsenical poisoning. In malarial neuritis the legs are first attacked, and the neural malady may be confined to them. Neuralgic pain of irregular distribution, with sugar in the urine, is characteristic of the neuritis of diabetes. In alcoholic neuritis all the limbs are affected, the pains are very severe, and both sides of the face may become involved. Symptoms much the same with reference to the distribution of the motor and sensory disturbance happen in the multiple neuritis from cold. In the toxic cases the face is also apt to be involved, and the optic nerves seem to be in them only affected. In pyæmic and septicæmic cases there are the history and the pyrexia to guide us. In the multiple neuritis of influenza comparatively little pain occurs; but considerable palsy, some facial paralysis, and difficulty of swallowing are not uncommon.

#### GRADUAL PARAPLEGIA.

This occurs in congestion, in acute and chronic inflammation of the meninges, in myelitis, in softening, in atrophy, in sclerosis,



in compression of the cord, and from reflex irritation. These are some of the marks of discrimination :

**Spinal Congestion.**—In congestion of the cord there is dull pain, generally confined to the lumbar and sacral regions; the palsy progresses slowly from below upward, is preceded by aching in the legs, by tingling, by numbness, is incomplete, and is not combined with paralysis of the sphincters. Moreover, the difficulty in walking is much greater on arising after a night's rest, or indeed whenever the patient has been for any length of time in the recumbent posture. We may often, too, trace the congestion to some disturbance of the circulation, especially of the abdominal circulation; or to alterations in the composition of the blood, as in rheumatism, smallpox, or typhus; or we find it as a result of exposure to cold and wet, or of standing for a long time, or as a sequel of the malarial fevers.

**Spinal Anæmia.**—Similar in some of its symptoms to spinal congestion, though very dissimilar in its causation, is so-called spinal anæmia. A disease usually of young females, and forming part of a general anæmic condition, or following exhausting discharges, or associated with a uterine affection, it often shows the symptoms of hysterical or "irritable spine," or "spinal irritation." The traits distinguishing this malady from spinal congestion are, that in the former we have much more marked signs of head, chest, and abdominal distress, such as vertigo, palpitation, neuralgic chest and abdominal pains, nausea, and other dyspeptic symptoms. The inactive or slightly palsied limbs often ache,—though affections of motility are far from constant,—are not infrequently the seat of spasms, are sensitive to the touch, act better after having been in the recumbent posture; there is pain along the spine, and pressure on the spinous processes of the vertebræ shows marked tenderness. But that spinal anæmia is the determining cause of the symptoms is by no means proved.

**Spinal Meningitis.**—In inflammation of the meninges we encounter severe pain in the back, little influenced by pressure upon the spine, yet aggravated by movement, even by the acts of defecation and urination; sometimes a sensation as if a cord had been drawn around the belly; pains in the limbs similar to those of rheumatism; cutaneous hyperæsthesia or anæsthesia; muscular twitchings, rigidity, and contractions, more or less permanent

and painful; increased superficial and deep reflexes, when the disease is above the lumbar enlargement of the cord; very commonly distressing spasms in the muscles of the back, and spasms in the limbs occasioned by attempts to move them; rigidity of the spinal column; bedsores; dyspnœa; retention of urine; vasomotor derangement; yet only incomplete paralysis, or, indeed, none at all. In the acute form we have decided fever. When marked paraplegia follows the symptoms mentioned, we may suspect the development of myelitis, or that an effusion has taken place which compresses the spinal cord. Cases of spinal meningitis occur from falls and shocks, from exposure to cold, as a consequence of adjacent disease, and in the course of general infectious processes. They are not unusual among soldiers who have slept on damp ground.

As regards the special membranes involved, there is no certainty in diagnosis. The symptoms, save in the acute purulent forms of the disease, are slow in developing. In inflammation of the dura mater, *pachymeningitis*, the radiating pains are very severe, but there is less vertebral pain and stiffness in the back: these signs are seen in their fullest expression in inflammation of the pia mater and arachnoid. In inflammation of the inner surface of the dura mater, *pachymeningitis spinalis interna*, which particularly happens in the cervical region, the symptoms are chiefly referred thither; and stiffness of the neck, paralysis in the upper extremities, especially in the parts supplied by the median and ulnar nerves, claw-like hands, contractions, severe pains in the arms, spots of anæsthesia, and herpetic eruptions are common. At a later period, as the hypertrophic thickening of the dura mater extends and the cord is more and more compressed, the paralyzed muscles may undergo wasting. There is a hemorrhagic form of *pachymeningitis interna* having the same causes as hæmatoma of the dura mater of the brain, and often accompanying it.

**Myelitis.**—Myelitis presents many of the same symptoms as spinal meningitis, with which, in fact, it may be associated. Frequently the symptoms come on by slow degrees, and the paraplegia, a very distinctive symptom, gradually becomes complete. There is strong knee-jerk with ankle clonus. Contractions of the muscles are uncommon, and not permanent, unless late in the disease; the muscles are usually flaccid; there is comparatively

little pain, none on pressure at any part of the spine, or on motion, and anaesthesia sooner or later shows itself. Further, we generally, though not constantly, find the urine alkaline, and, as a rule, retention of urine and a want of control over the rectum exist, bed-sores form readily, and the temperature of the palsied is lower than that of the healthy parts.

In acute cases there are, as in acute spinal meningitis, raised temperature and a frequent pulse. The fever is moderate and irregular. There is pain in the back, not increased by movements, and pain in the limbs preceded by numbness or burning. In many instances we notice erection of the penis. Spasm in the extensor muscles is always of significance. Reflex movements in the relaxed palsied limbs, at first still easily excited, and excited, too, by irritation elsewhere applied, are gradually abolished as the process of inflammation and softening affects the gray matter of the cord. In dorsal myelitis the trunk reflexes are impaired, but the reflex excitability remains excessive in the parts supplied by nerves arising below the level of the greatly diseased centres. In disease of the lumbar enlargement the knee-jerks are wholly lost.

An altered sensibility to heat and cold, when, for instance, a sponge soaked in warm water or a piece of ice is applied to the spine over the inflamed spot, has been spoken of as a diagnostic test. In either case the sensation, when the diseased part is reached, changes to a burning sensation. This symptom is, however, far from constant, and cannot be accepted as conclusive. There is a zone of hyperaesthesia at the level of the lesion, and corresponding to this a zone of constriction or "girdle pain." Below the level of the lesion the loss of sensation is complete. The paraplegia, even in acute cases, is not suddenly developed. Yet we meet with marked exceptions. There are instances in which it comes on as rapidly as in spinal hemorrhage,\* and without attending loss of sensibility; or a paralysis of the bladder is the first symptom, and paralysis of motion and of sensation quickly follows.†

Myelitis may be the result of cold and exposure, of over-exertion, of syphilis, of peripheral irritation, of pressure, as from

---

\* Hayem, *Archives de Physiologie*, Sept. 1874.

† Erb, in *Ziemssen's Cyclopædia*, vol. xiii.

disease of the vertebræ, of tumors, connected with the bones or membranes, encroaching on the cord and setting up disease there, of injuries to the cord, or of concussion of the spine after railway accidents. It is sometimes met with in the course of measles and of smallpox, and of typhoid and typhus fevers and toxic blood-states, such as gout and syphilis. *Compression* as a cause has been noted in the cervical as well as in the other portions of the spine. Paralysis of the arms, with dilated or contracted pupil and very slow pulse, is among the chief symptoms of the "cervical paraplegia." Pain in the limbs, hyperæsthesia, muscular contraction, spasms, and great reflex irritability are among the earlier symptoms of this as of all the other forms of myelitis from pressure; but as the case progresses the reflex irritability is lost. Yet recovery, almost complete, is possible.\*

In looking at the symptoms which mark the extent and exact site of the inflammation, we find in the common form, where the disease affects a considerable portion of the thickness of the cord,—the *transverse myelitis*,—with the ordinary symptoms of complete paraplegia and anæsthesia, that the reflex excitability is lost in the parts supplied by the nerves coming from the affected part of the cord, and is preserved or increased in the parts supplied by nerves arising from the cord below the diseased area,† and that the muscles respond to the electric current. This is not the case in *central myelitis*, which, moreover, usually runs a rapid course, in which there is speedy loss of sensation and of reflex action, and in which muscular atrophy soon shows itself. In *disseminated myelitis*, a form where several foci of inflammation are present, there are lulls and exacerbations, the paralysis is not so constant nor so complete, although it may be in all four limbs, spastic symptoms are not uncommon, and the disease develops itself after acute maladies, as after smallpox. *Hemorrhagic myelitis* is usually central; the paraplegia comes on in less than an hour, and we can only distinguish it from pure hemorrhage into the cord if fever and other symptoms of an acute myelitis previously existed. In children the anterior cornua are apt to be affected, and the disease is known as *poliomyelitis*.

---

\* Buzzard, Brain, April, 1880.

† According to Bastian, total transverse lesion high up in the cord abolishes the knee-jerk.



*Softening of the cord* cannot be distinguished from myelitis; the inflammation is, in truth, the usual cause of the softening.

**Spinal Scleroses.**—Sclerosis of the spinal cord may be primary or secondary. The latter is represented by the descending or ascending degenerations that follow lesions of brain, cord, or posterior nerve-roots. In the former are included the so-called system-diseases,—posterior sclerosis, locomotor ataxia, and lateral sclerosis, spastic paralysis. In all of these it is believed that the morbid process takes place in the nerve-fibres, and that any excess of fibrous tissue is secondary. In the disseminated sclerosis of brain and cord, on the other hand, it is likely that increase of the fibrous tissue constitutes the first stage, and that cicatricial contraction causes destruction of the fibres. The sclerosis where brain and cord both suffer, we shall discuss with the forms of tremor; posterior sclerosis of the cord produces the symptoms of locomotor ataxia, not of palsy.

**LATERAL SCLEROSIS.**—Primary sclerosis of the lateral columns in which the anterior horns are not affected shows the group of symptoms described as *spasmodic dorsal tabes* by Charcot, or *spastic spinal paralysis* by Erb. It is characterized by a sensation of weakness in the back, a gradually increasing loss of muscular power in the lower extremities, proceeding slowly from below upwards, and associated with reflex spasms and persistent muscular contractions, with increased tendon reflex, but without impairment of sensibility, or trophic disturbances, or bedsores, or vesical disorder. The muscles are well nourished, or only very slightly wasted; the gait is very peculiar, the walk being on the toes, and as the foot touches the ground a trembling happens. There are no cerebral symptoms whatever; the electrical excitability is either normal or somewhat lessened. In rare instances the disease begins in the upper extremities; it is almost always of very slow development. Occasionally it terminates in recovery. It is most likely that the disease consists essentially in a primary sclerosis of the pyramidal tracts, or of the terminations of their fibres in the motor cells of the gray matter. But whether the group of symptoms may not be produced by various lesions of the cord is not settled. To an infantile form of degeneration of the lateral columns McLane Hamilton has called attention. Loss of power

in the lower extremities, muscular contractions without marked atrophy or greatly impaired electro-muscular contractility, such as happen in infantile paralysis, increased skin and tendon reflexes, and absence of sensory disturbances or brain-symptoms, are the chief signs of the affection.\*

When sclerosis affects the lateral columns, and is combined with degeneration of the great ganglion cells in the anterior horns of gray matter of the cord, the portion which has a controlling influence over nutrition, marked nutritive changes happen in the palsied part, such as we find in progressive muscular atrophy. But this *amyotrophic lateral sclerosis*, as Charcot has termed it, is from the onset an atrophy of a whole muscular group. It is a disease closely allied to progressive muscular atrophy, and affects as a rule the four limbs successively, beginning in the arms, produces strange deformities in the wasted and palsied limbs, which are often agitated by fibrillar movements, extends to the hypoglossal and to the pneumogastric nerves, and thus determines death.

**Tumors of the Cord.**—Tumors of the spinal cord, either growing from it or its membranes, or originating in the vertebræ and compressing the nerve-structure, occasion paraplegia. But the cause is beyond the reach of positive diagnosis. The symptoms vary with the situation and extent of the growth. They depend first upon the irritation, and later upon the compression, caused by the new formation. We suspect the affection if we have emaciation and signs of a grave constitutional malady attending the slowly progressing palsy, if this be more decided on one side than on the other, and if anæsthesia be found on the side opposite to that in which the palsy is marked and which is the seat of the tumor. Then severe pain over the locality of the disease occurs in cancerous new formations,—and most spinal tumors are cancerous,—and is aggravated in paroxysms. The pain is generally felt on one side first, and is associated with tenderness of the spine and muscular spasm. This is especially the case when the growth springs from the membranes. Yet, unless we have distinct evidence of tumors elsewhere, the diagnosis is never more than an uncertain one. If multiple tumors exist, it may

---

\* Transactions of the American Medical Association, 1879.

be made positive. Strong proofs of syphilitic infection point to the spinal symptoms being due to a syphilitic growth; and signs of scrofula, or tubercle in the lungs or in other internal organs, make it likely that similar morbid products are the cause of the palsy. Should a gradually progressing paralysis suddenly show symptoms of acute myelitis in a person with the constitutional cachexia just mentioned, we have an additional reason for supposing the affection to be tubercular and to be rapidly extending.\* Lymphadenomas elsewhere make it extremely likely that the spinal symptoms are due to one or several of them in the cord. Yet the spinal symptoms in the affection may really be due to myelitis. In all cases of suspected tumor we must be very careful that bone-disease is not the cause of the symptoms. The absence of sharp pain and the uniformity of the palsy on both sides are strong points of distinction as against tumors.

**Reflex Paraplegia.**—We cannot isolate this from the paraplegia of organic spinal origin with any certainty, unless we can discern the source of the irritation, obtain a clear history of the case, and satisfy ourselves of the absence of the special symptoms of an organic disease of the cord or of the peripheral nerves. Some features thought to be distinctive of this functional palsy are, that anæsthesia is exceptional; that the palsy is seldom complete; that some muscles are much more affected than others; that spasms in the paralyzed muscles are very uncommon; that there are no pains in the spine, produced either spontaneously, or by pressure, or by percussion, or by applying ice or a hot moist sponge; and that there is a correspondence between changes in the degree of the paralysis and changes in the visceral disease or the external irritation which is supposed to have produced the paraplegia. But it is certain that the condition of inhibition, in affections of the bladder and kidneys and other states, which is believed to cause the reflex paraplegia is very infrequent. Most of the cases are really due to an ascending neuritis. Some result from some morbid blood state, such as exists in gout and in diabetes.

Worms in the intestinal canal may give rise to reflex para-

---

\* See cases of Hayem, *Archives de Physiologie*, 1873; and Erb, in *Ziems-sen's Cyclopædia*.

plegia, which disappears with their expulsion. A form of reflex paralysis produced by intestinal disorder, and in which a motor and sensory paraplegia shows itself, has been carefully studied by Bartholow.\*

So much for paraplegia. We shall now examine some of the other clinical varieties of paralysis; beginning with a group in which the palsy is limited, though it may be general.

#### PALSIES USUALLY LIMITED, THOUGH THEY MAY BE GENERAL.

**Hysterical Paralysis.**—In hysterical paralysis there is no structural affection of the brain, yet all looks as if this were the case. This form of paralysis we distinguish from that of organic disease, by its occurrence in hysterical persons; its sudden appearance, and frequently its just as sudden disappearance; its coming on generally under the influence of some powerful emotion, often after an attack of hysterical convulsions; the absence of any signs of a serious lesion of the nervous centres, except the paralysis; the varying nature of the palsy, sometimes hemiplegia, sometimes paraplegia; its incomplete character, the patient being not infrequently able to move while under strong excitement; and the ease with which reflex movements are brought on in the seemingly helpless limb. Then there are nervous shiverings, noises in the ears, tingling sensation in the limbs, and vaso-motor disturbances showing themselves by slight swelling of the joints and elsewhere. The muscles, except in cases of long standing, contract perfectly both under the faradaic and the galvanic current. Duchenne pointed out that the electro-muscular sensibility is either diminished or abolished, while in cerebral paralysis it is intact. In some cases galvanic sensibility is lost.† We never find the reaction of degeneration. Hyperæsthesia, but much more generally anæsthesia, is observed, and this also may involve the special senses and affect the muscles. But muscular anæsthesia may be absent in hysteria. Rapid changes occur in the sensibility under strong electric currents, and there may be a transfer of the loss of sensation from the disordered side to

---

\* See Transact. of Phila. College of Physicians, Nov. 1883; also Barié, Arch. Gén. de Méd., Nov. 1881.

† Wood, Nervous Diseases and their Diagnosis, 887.



the healthy side, caused by stimulating the side of the hemianæsthesia,—by mustard or by the faradaic brush, or by certain metals, such as gold, or by wood,—or, indeed, by strong mental impressions.

The eye-symptoms, as Charcot has pointed out,\* are peculiar. There may be an amaurosis, but there is no alteration of the papilla; the constricted field of vision is concentric, not as in locomotor ataxia star-shaped, and red is the color that is seen longest, while in multiple sclerosis there is no loss of color or color-blindness. Then in hysteria the eye-brow on the affected side is lower than on the other side, while in true paralysis it is more raised on the side affected. Nystagmus is never observed in hysteria.

Persons affected with hysterical palsy are striking types of a nervous constitution, and, as Sir James Paget† mentions, show a singular readiness to be painfully fatigued by slight exertion. The palsy may seize only upon one limb, or upon part of one limb, or upon special muscles, as those of the pharynx and œsophagus, the larynx, the intestines, and the diaphragm; or it may, although it more rarely does, assume a hemiplegic or a paraplegic form. Hysterical hemiplegia presents a peculiarity in the gait, on which Todd‡ lays great stress. “In walking, when the palsy is pretty complete, the leg is drawn along as if lifeless, sweeping the ground.” It is not swung round, describing the arc of a circle, as it is in ordinary hemiplegia. The palsy is almost invariably left-sided. It is apt to be conjoined to left-sided ovarian tenderness, and to very decided anæsthesia, which passes beyond the paralyzed part to the nearest portion of skin and mucous membrane, though, as a rule, still limited to the same side. Thus we find the pituitary membrane of one nostril rendered insensible, if the loss of feeling affect the face.

In hysterical paraplegia we find the same incompleteness of the palsy and the same response to electric tests already mentioned, and we are also very apt to have the symptoms of spinal irrita-

---

\* International Clinics, vol. i., 2d Ser., 1892.

† Nervous Mimicry of Organic Diseases, in Clinical Lectures and Essays, London, 1875.

‡ Clinical Lectures on Paralysis and other Affections of the Nervous System, Lecture XIII.

tion. Hysterical contractions of the muscles especially affect the lower extremities, though they are not uncommon in the arm. These hysterical contractures, as they are now usually called, generally come on quickly, appear to be permanent, and to be associated with loss of power, but disappear as suddenly as they showed themselves. Yet they may really become permanent and combined with sclerosis of the cord, and we may find them associated with tremor, and with exaggerated knee-jerk. Ankle clonus has also been observed by Charcot as occurring in hysterical paralysis. Gowers, however, thinks that true uniformly persisting ankle clonus bespeaks secondary organic disease in the motor parts of the cord, while a spurious irregular clonus, now ceasing, now renewed by a fresh contraction of the muscle, is characteristic of hysteria.

Very similar to hysterical paraplegia is the *paraplegia produced by hypnotic suggestion*. Charcot and Longues\* have called attention to the fact that in this as well as in the functional paraplegia met with after railroad and other accidents, the so-called traumatic hysteria, there is a general anæsthesia in the palsied legs which follows the fold of the groin and excludes the genital organs. The same line also enables us to distinguish between hysterical paraplegias and those of organic origin.

One of the most difficult points of diagnosis with reference to hysteria is to distinguish the hysterical symptoms that arise in sclerosis and in myelitis, or that follow injuries to the nervous system, from the manifestations of pure hysteria. Nothing but a careful study of the individual case, of the history, of the reflexes, of the electric reactions, of the line of the anæsthesia, of the state of the muscles themselves, laying stress on the absence of muscular wasting in pure hysteria, will save from error.

**Rheumatic Paralysis.**—Rheumatic paralysis resembles hysterical paralysis in being ordinarily limited. It may affect any muscle or any group of muscles in the body; sometimes the rheumatic poison disorders the portio dura, and we observe, in consequence, facial palsy; or it may fasten on the radial nerve, and

---

\* Charcot, Œuvres Complètes, tom. iii. p. 448.

we have groups of muscles in the forearm palsied. Rheumatic paralysis is recognized by the history of the case; by the evidences of a rheumatic attack; by the rapid development of the palsy; by the pain which usually attends it; and by its being unaccompanied by symptoms strictly referable to a disease of the nerve-centres. It may or may not be attended by anæsthesia. The muscles themselves, certainly in those cases in which they, rather than a large nervous branch, are primarily and chiefly affected, are readily acted upon by electricity, unless their structure be altered; and the electro-muscular sensibility, though it may be lessened, is not abolished.

**Lead Palsy.**—Paralysis from lead poisoning occurs primarily, and sometimes only, in the extensor muscles of the arm, occasioning the well-known wrist-drop. It generally begins in the extensor communis, then affects the radial and ulnar extensors. Gradually other muscles become involved: there is loss of power in the ball of the thumb, in the deltoid, and in the triceps, but not in the supinator longus, or in the intercostal muscles, or in those of the lower extremities. The disturbed muscles on both sides of the body waste, entirely lose their irritability to electricity, and soon show the reaction of degeneration. The patient is weak; his movements are tremulous; he has the characteristic blue line on the gums, is obstinately constipated, is subject to colic, and lead can be found in the urine. Sometimes the poison seizes upon the brain, and epileptic convulsions and other signs of a serious cerebral affection appear, and we find marked optic neuritis. From the locality of the palsy, in addition to the accompanying symptoms and the knowledge of the man's employment, the diagnosis is usually arrived at with ease. Paralysis produced by an affection of the radial nerve shows the greatest similarity. Yet here the supinator muscles as well as the extensors, but upon one side only, are affected, which is not the case in lead paralysis, where both sides are affected and the patient can carry the hands supine.

Lead palsy is not often seen in children, but it may be met with.\*

**Diphtheritic Paralysis.**—Diphtheritic paralysis is a sequel

---

\* See cases of Sinkler, *Med. News*, July, 1894.

of diphtheria which follows an attack of the disease within a fortnight or two months, and, therefore, after the patient is to all appearance fully convalescent. It may be very localized, merely affecting the palate or the pharynx; or very general, fastening upon both of the lower extremities, and even upon the upper. When extensive, it is always ushered in by a change in the voice and a throat-palsy; there is difficulty in swallowing, fluids are regurgitated through the nose, and the saliva dribbles from the mouth. The eye-muscles are apt to be disturbed, and paralysis of accommodation and strabismus and double vision are not uncommon. The paralysis of the extremities ensues gradually; day by day the muscular power is more and more enfeebled. The loss of motion is often preceded by formication, and attended by a certain amount of anæsthesia. The faradaic electro-muscular contractility and sensibility are diminished, and the galvanic current shows mostly the same results. The palsy mends as slowly as it comes on; yet most cases fully recover. The brain itself is not directly affected; at least there were no symptoms of cerebral mischief in the cases which have come under my observation. In some cases rupture or plugging of a cerebral vessel takes place. More commonly the palsy is due to multiple neuritis. In children affected with diphtheritic paralysis regular bulbar crises may happen.\*

**Syphilitic Paralysis.**—Paralysis from syphilis we find in persons presenting signs of constitutional syphilis, and in whom any serious nervous disturbance points to a local manifestation of syphilis in the nervous centres. Not unusually the syphilitic exudation is localized in the course of one or of several nerves, and we have, for instance, paralysis of the sixth or paralysis of the fifth with or without paralysis of some other cerebral nerve. But as syphilis attacking the nervous system is chiefly characterized by a want of uniformity in the lesions it produces, so we observe dissimilar phenomena preceding or attending the palsies. Thus, we may or may not, though in point of fact we usually do, find the paralysis associated with pain in the head, with optic neuritis, with sleeplessness, vertigo, impaired memory, and sickness at the stomach. Decided vertigo is prone to take place

---

\* Guthrie, *Lancet*, April, 1891.



where the syphilitic affection, as it so often does, has led to disease of the vessels, and is apt to be the forerunner of local softenings and of hemiplegia. When disease of the membranes has happened, headache is generally severe, and local spasms or convulsions occur. The same symptoms are encountered when there is a growth in the hemisphere, which is very apt to be near the surface of the brain; though here again the form of mischief may be comparatively latent, the patient may have only occasionally convulsions, and the paralysis be slight or improving, yet a fatal coma may follow a few convulsions. Instances of this have come under my observation.

But, as a rule, syphilitic paralysis does not terminate fatally. In truth, the ease with which the palsy and its attending phenomena yield to treatment, if we except marked instances of hard nodules, forms one of the traits of the malady. Other common features, to speak in general terms and taking into account what has been said of the dissimilar character of the lesions, are—that it ordinarily affects persons younger than those in whom we find paralysis dependent upon disease of the nervous centres, and especially of the brain; and that its manifestations are shifting and capricious, and rarely symmetrical. These same traits characterize syphilitic affections of the nervous system in which paralysis is not among the symptoms. Paralysis of the third nerve is a frequent result of syphilis;\* but, as already stated, the poison may attack any part of the nervous system, and paraplegia dependent upon disease of the cord is not very uncommon. A progressive multiple palsy of cerebral origin, clearly affecting dissociated muscles, is usually syphilitic, and is mostly due to several patches of gummatous meningitis. At times a rapid, almost universal paralysis, as Buzzard notices, occurs in syphilitic subjects. This is very likely of peripheral origin. It is among the peculiar traits in syphilitic palsy that the lost electro-muscular contractility returns rapidly.†

Erb has called attention to an association of symptoms regarded as characteristic of syphilis of the cord. They come on gradually at first, but may rapidly increase. They are increased

---

\* Broadbent, *Lancet*, Jan. 1874.

† Engle, *Phila. Med. Times*, Dec. 1877.

reflexes, with but little muscular rigidity, slight spastic gait and muscular weakness, disorder of the bladder functions, and disturbances of sensation in the legs in the form of paræsthesia. The symptoms are confined to below the waist.

The mischief to the nervous system may not happen for years after the infection. It may be the result of an *inherited taint*. But such cases cannot be recognized unless there are other signs of syphilis than the suspected nervous symptoms; and chief among these signs are the evidences of periostitis in the long bones and of disseminated choroiditis in the fundus of the eye. Then there is that valuable test of congenital syphilis discovered by Mr. Hutchinson,—a malformation of the two upper central permanent incisors, which consists in their being narrower at their cutting edges than at their insertions, and often notched. The same observer has called attention to diffused opacity of the cornea and to diseased nails as being common among the manifestations of the inherited disease. Paralysis also may occur, as in the case reported by Bartlett;\* but it is very rare.

#### LOCAL PALSIES.

The forms of paralysis which have just been noticed are mainly such as are designated as partial. When the loss of power is very limited, the palsy is spoken of as *local*; most of these local palsies are peripheral, and the result of neuritis.

**Facial Palsy.**—Of the local paralyses, of particular importance—from its frequency—is facial, or Bell's palsy. The disease consists in an affection of the portio dura of the seventh nerve. In consequence of the derangement of this motor nerve, nearly all the muscles of one side of the face lose their faculty of motion, and, as it is their play which gives expression to the countenance, the appearance of the face is extraordinary. The eyelids are open and fixed; the features are rigidly composed on one side of the face, but reflect every change of feeling on the other; the mouth is distorted, being drawn to the unaffected side; the naso-labial fold is effaced; and in the old the furrows disappear from the forehead and the eye waters. In some cases the velum palati is involved in the paralysis. The impaired muscles

---

\* Clinical Society's Transactions, vol. iii.

waste; their electric irritability is diminished and degenerative reactions may be present. Sensation remains unaltered as long as the fifth nerve is not disturbed.

The causes of the palsy are such as influence the distressed nerve in its course or at its periphery: a wound; mumps; ear-disease; exposure to cold; rheumatism; syphilis. The most common cause is a neuritis from cold affecting the nerve within the Fallopian canal. The malady is easily discriminated from the facial palsy of disease of the brain by the inability to close the eyelids, owing to the paralysis of the orbicularis palpebrarum; by the absence of headache, of vertigo, of mental confusion, of loss of memory; by the much more complete though strictly local character of the paralysis, the affected muscles even failing to participate in bilateral or emotional movements; and, except in slight lesions of the nerve, by the lost electro-muscular contractility. In severe cases, indeed, the muscles soon cease to respond to faradization, while the galvanic irritability is preserved and even heightened, and the reaction of degeneration is very marked. Electric stimulation of the diseased nerve shows that it quickly loses its excitability both to the faradaic and the galvanic current.

The observations of Erb enable us to tell with considerable accuracy the exact part of the nerve affected. They take into account well-known anatomical and physiological facts, and lead to these conclusions. If there be complete palsy of all the facial branches with the exception of the posterior auricular nerve, the lesion is in the main trunk of the facial, exterior to the Fallopian canal. If the auricular nerve be also implicated, the lesion is within the Fallopian canal below the origin of the chorda tympani, the most common seat of the affection. If taste and salivary secretion be disturbed on the side of the tongue corresponding to the palsy of the face-muscles, the lesion is between the points where the chorda tympani and the tympanic branch are given off. If in addition the sense of hearing be abnormally increased, we may infer that the nerve is affected between the tympanic branch and the geniculate ganglion, and at the latter point palsy of the palate is superadded; and higher, up to the entrance into the brain, disorders of taste happen. Eventually implication of other cranial nerves, as of the auditory, also occurs.

Cases of facial-nerve palsy generally recover. Sometimes, how-

ever, the recovery is incomplete, and a rigidity with some contraction of the affected muscles takes place, which, when slight, may make the sound side appear relaxed, and the diseased side seem the normal one.

In rare instances the facial palsy is on both sides. Now, in this *double facial palsy* the lesion may be within the cranium, such as compression by a tumor, or may affect the nerves while passing through the medulla and pons in their farther course. When dependent simply on a local affection, and therefore limited to the manifestations of paralysis of the portio dura, we find the same causes at work which give rise to the one-sided disease. Exposure to cold and rheumatism are the most frequent; but syphilis is also among the causing elements. In an instance detailed by Todd in his clinical lectures, in which there was disease of the temporal bone, the portio mollis was also implicated. The face is immovable, or nearly so, and the palsy is generally more complete on the left side than on the right. The muscles do not respond to electricity, or respond imperfectly, and we notice, as in the one-sided malady, that a continuous current may excite their action, while faradization does not. Nay, the two sides may give different results in this respect,\* most likely caused by different conditions of exudation and of pressure on the affected nerves.

**Paralysis of the Nerves of the Arm.**—Paralysis of one or more nerves of the arm is very often encountered. It may happen from rheumatism, from cold developing a neuritis, from traumatism or fracture, or from the pressure of a growth; but its most common cause is accidental compression. A person falls asleep with his head on his arm, and a temporary palsy results; or it may follow the use of a crutch. In truth, the disorder may be taken as the type of the *palsies by compression*, and we find that the electro-muscular contractility depends on the severity of the nerve-lesion; as a rule, there is reaction of degeneration. Sensory symptoms are slight or wanting; often there are numbness and tingling.

---

\* Case of Baerwinkel, Schmidt's Jahrb., Bd. cxxxvi. No. 1. For other cases of double facial palsy, see Gairdner, Lancet, May 18, 1861; Pellet, Tra-vaux de la Société Médicale, 1867; Wright, British Medical Journal, Feb. 1869; Krüge, Neur. Cent., 1890.



The nerve most frequently paralyzed is the *musculo-spiral*, or its main branch the radial, and we observe palsy of the extensors of the wrist and the fingers and of the supinators. In the loss of power in these muscles, in the mode of onset, and in the unilateral involvement we find the differences between the palsy under consideration and the wrist-drop of lead palsy. When the *median* nerve suffers, the pronators, the radial flexor of the wrist, the flexors of the fingers,—except the ulnar half of the deep flexor,—the abductor and flexors of the thumb, and the first and second lumbricales are paralyzed; while sensibility is impaired or lost on the palmar aspect of the thumb, the index and middle fingers, and adjacent portions of the ring-finger, and often on the dorsal aspect of the last phalanx of the index and middle fingers.

Involvement of the *ulnar* nerve shows itself in palsy of the ulnar flexor of the wrist, the ulnar half of the deep flexor of the fingers, the muscles of the little finger, the interossei, the third and fourth lumbricales, the adductor and inner head of the short flexor of the thumb; and impairment of sensibility in the parts of the hand and fingers not supplied by the median and radial nerves. From those diseases of the spinal cord which begin with arm palsy, the local malady is distinguished by the tenderness in the course of the nerve, and the one-sided paralysis. The same signs separate this arm palsy from the loss of power in the wrists, arising from atrophy of the muscles in the overworked parts, occurring in undernourished persons, as in poorly-fed and hard-worked shoemakers.\*

About other local palsies, as of the pharynx and œsophagus, of the larynx, of one side of the palate, of the tongue, of the muscles of the eye, of the diaphragm, of isolated muscles of the trunk, and of the extremities, it is impossible here to enter into particulars. But there are some forms of local palsy which, from their striking interest, it is necessary to describe, the most important of which is the paralysis of the tongue and parts concerned in deglutition.

**Bulbar Paralysis.**—Bulbar palsy can scarcely be considered

---

\* Chambers on the Indigestions.

a local palsy. It has a close relation to progressive muscular atrophy; yet, from a clinical point of view, its main manifestations are those of disorder of special nerves. In this bulbar or *glosso-labio-laryngeal* paralysis, the first symptoms which are likely to attract attention are, that the tongue seems less supple and the utterance becomes nasal or thick, the food lodges between the teeth and cheek, and the saliva dribbles from the lips and corners of the mouth. As the paralysis progresses, articulate speech is almost lost, as is the reflex action in the throat; the shape of the tongue is altered, it generally dwindles, and at times shows twitching of its fibres, or lies motionless in the mouth, though, as a rule, it reacts to faradization, the posterior nares can no longer be closed by the velum and muscles of the posterior palatine arch; deglutition becomes very difficult, and the patient is tormented with hunger. Reflex irritability of the mucous membrane of the larynx is frequently lost; the respiratory movements are unusually weak, and fits of suffocation ensue. The general debility becomes extreme, and the patient is apt to perish by the sudden stoppage of the heart's action. The disease is unmistakable. Double facial palsy resembles it most; but here the tongue is not involved, and the eyelids remain open; on the other hand, in bulbar paralysis the lower part of the face only is motionless. This malady may have an acute beginning, and seemingly in cold; it is sometimes complicated with weakness of the muscles of one side of the body, or with muscular atrophy in the limbs and trunk, and sometimes with the symptoms of lateral sclerosis. As a rule, the mind remains clear. The affection is generally of rather slow development and slow but relentless progress; but it is not nearly so chronic a malady as progressive muscular atrophy, which may last from ten to twenty years, while the bulbar paralysis has, like lateral sclerosis, an average duration of from one to three years.\* Progressive bulbar paralysis has its seat of lesion in the medulla oblongata, in the motor nuclei, which undergo a degenerative atrophy; and we understand the main symptoms when we reflect on the nuclei which connect the hypoglossal, the spinal accessory, the vagus, and the facial.

---

\* Möbius, Schmidt's Jahrb., No. 2, 1882.

With reference to all these local palsies we are sometimes much perplexed to know if the palsy be the result of beginning disease of the brain or spinal cord, or if it be purely local. To speak first of the brain: the cerebral symptoms may not be marked, or they may be so contradictory as to afford no real help in diagnosis. When, however, we discover, as we generally can, that the palsy affects muscles which are supplied by different nerves and such as have no communication with one another, we may set down the complaint as having a central origin. As regards the distinction from spinal affections, the almost constantly single-sided character of the symptoms in local palsies, and their double-sided character in spinal affections, are very important.

#### PALSIES CONNECTED WITH MARKED MUSCULAR WASTING.

There is a group of palsies especially marked by wasting of the muscles. In some affections already discussed we have found wasting among the symptoms, as at times in myelitis, and in cervical pachymeningitis with considerable damage to the nerve-roots, where atrophy of the arms happens. Again, atrophy of the muscles of the trunk and limbs is often met with in the advanced stages of progressive bulbar paralysis. But in all these affections there are other and more distinctive symptoms. In some affections the wasting of the muscles is the pre-eminent feature. This is particularly the case in progressive muscular atrophy and in the essential paralysis of childhood.

Another important question which may arise—and with reference not only to limited but also to extended palsies—is, whether the loss of muscular power be not in reality dependent upon changes in the muscular tissue, and especially upon that change found in the disorder known as idiopathic or progressive muscular atrophy.

**Progressive Muscular Atrophy.**—This form of “wasting palsy” is due to chronic or subacute degenerative changes in the gray matter of the anterior horns of the spinal cord, particularly the large ganglion-cells, sometimes in association with similar changes in the peripheral motor nerves and the pyramidal tracts. The affected muscles undergo atrophy of varying degree and extent.

Progressive muscular atrophy is a disease of adults, and essentially of men. Its most striking sign is increasing inability to perform certain movements. When the muscle chiefly concerned in the attempted motion is examined, it is found to have dwindled. Soon other muscles follow; and their wasting, too, is accompanied by further muscular weakness. Portions of the disorganizing muscles twitch, much to the annoyance of the patient, and tapping them sharply causes a marked contraction of the fibres. The muscles of the face, as a rule, escape. In the affected part the circulation becomes languid; it is also very susceptible to cold, and indeed its temperature is lowered; there is a feeling of numbness in it, but, as the disease progresses, rarely pain; to pressure it is soft and yielding. The muscles most frequently attacked are those of the hand, the flexors and supinators of the forearm, the biceps, the deltoid, and the other muscles of the shoulder. Sometimes the disease begins in the trunk and the lower extremities; but it is most common to have it marked in the upper extremities and to find only weakness and spasm in the lower. Sometimes, also, bulbar symptoms, with weakness of the muscles of the lips and tongue and of the pharynx and larynx, appear, and changes are found in the medulla analogous to those present in the cord. The decrease of the muscular fibres gives rise to strange and palpable deformities, and, when the muscles of the trunk are involved, to extraordinary positions of the body, in consequence of all antagonism to the healthy muscles having been removed.

In the parts affected the reflex action is lost; even the deep reflexes disappear. We see this happening with the knee-jerk just so soon as the muscles of the legs become flaccid and begin to waste. To the electric currents, both faradaic and galvanic, the muscles respond feebly; still they respond, and in portions where there are many sound fibres they contract energetically. The degree of response depends, indeed, on the degree of disorganization and wasting. The excitability to the galvanic current is apt to remain much longer than that to faradization; the reaction of degeneration is likely to be present.

When we contrast progressive muscular atrophy with the forms of paralysis with which it may be confounded, we find several features at variance. From *cerebral hemiplegia* it differs by its



much more gradual invasion, by the rapidity but want of uniformity with which the muscular atrophy takes place, by the lost reflexes, by the diminished electric excitability, and by the absence of disordered intellect and of other signs of disease of the brain. Difficulty in articulation and in deglutition may occur in either. From *general spinal paralysis* it is diagnosticated by the spinal malady affecting primarily all the muscles of the lower extremities before those of the upper become involved. Then, too, if the spinal paralysis be due, as it so generally is when extensive, to myelitis, the alterations of sensibility, the totally lost electro-muscular contractility, and the affection of the sphincters are striking traits of difference.

The difficulty of distinguishing cases of *local paralysis* from progressive muscular atrophy is at times very great. Yet generally we may separate the latter, for instance from rheumatic palsy, by noticing that this affects a group of muscles rather than one muscle, or than one muscle here and another there. Further, the atrophied muscle in the rheumatic disorder is the seat of pain intensified by movement, and it contracts well under the electric stimulus. The same test by the electric current is of service in discriminating the muscular disease from hysterical paralysis, and from paralysis consequent upon injuries to nerve trunks and upon lead poisoning. In the first of these palsies the electrical contractility is, except temporarily in cases of old standing, intact, in the others it is abolished; while in progressive muscular atrophy, save when the wasting is extreme, it is simply enfeebled. Besides this test, the unimpaired sensibility, the capricious and unequal manner in which the atrophy seizes upon the muscles in this malady, the fibrillation, and the beginning of the wasting in the thenar muscles and the interossei are points to which we attach importance.

Hirst\* points out the occurrence of muscular atrophy as a phenomenon of *hysteria*. The peculiarity of this form of wasting is its unilateral or circumscribed character, though sometimes it is general. The recognition depends upon the psychic state of the patient and the occurrence of hysterical or hysterio-epileptic convulsions.

---

\* Deutsche Medicinische Wochenschrift, 1894, No. 21, p. 459.

The most difficult differential diagnosis we may be called upon to make is to distinguish certain cases of progressive muscular atrophy from *bulbar paralysis*. In truth, the two affections often coexist, and the features of each may be blurred to the last degree. The diagnosis depends upon the distribution of the symptoms, the morbid process being essentially the same in the two sets of cases. In the one the arms, and sometimes also the legs, suffer; in the other the tongue, the lips, the pharynx, and the larynx. Defective pronunciation points to the bulbar malady. Failure of the respiratory power is common to both.

Local atrophies may be mistaken for part of the general disease. There is, for instance, an affection, *unilateral progressive atrophy of the face*, in which gradual wasting of one side of the face occurs, of the soft parts first, and then of the deeper tissues. The facial hemiatrophy follows blows and contusions, abscess of the ear, influenza, typhoid fever, or, as in Cohen's case, an attack of erysipelas. It begins with a discoloration of circumscribed spots, a white or yellowish discoloration; the subcutaneous fat disappears, and the beard and eyelashes change. Sensation is, as a rule, not affected, nor are the electrical reactions changed.\* But in progressive muscular atrophy the face almost always escapes; if it be affected, it is so on both sides. *Acute or chronic joint-inflammations* are attended with weakness and wasting of the muscles moving the affected parts. The extensors usually suffer, occasionally also the flexors, and rarely distant muscles. Another limited atrophy is a wasting from *overuse of muscles*, seen especially in the small muscles of the hand. It shows no tendency to extend.

Paralyzed muscles atrophy, and may subsequently undergo degenerative change; but the distribution differs from that of progressive muscular atrophy, and we lay stress on the symptoms that usher in the paralytic state, and that attend it.

In the condition known as *syringomyelia*, in which the central gray matter of the spinal cord is replaced by gliomatous tissue that breaks down and gives rise to the formation of a cavity, we

---

\* See cases, *Journal of Nervous and Mental Diseases*, New York, March, 1880; *Schmidt's Jahrb.*, No. 7, 1881; *St. Louis Alienist*, April, 1881; and *Skyrme*, *Brit. Med. Journ.*, March, 1892.

have fibrillar contractions in the affected muscles and atrophy with resulting deformities. But symptoms of sensory derangement usually appear earlier and are more pronounced. Common sensibility may be unchanged, while there is inability to distinguish heat and cold, and often also to appreciate pain. The sphincters may or may not be disturbed; the knee-jerks are normal or exaggerated. The muscles waste, rapidly lose their faradaic excitability, and the reaction of degeneration is finally established. Yet the symptoms, on the whole, are of slow development, and show themselves chiefly in the arms and in the upper part of the trunk. There is unsteadiness of motion, with muscular weakness rather than paralysis, and trophic disturbances in the skin, such as thickenings, eruptions, ulcerations, are marked. In the legs there may be spastic paresis. Deviation of the spine is common; it was present in half the cases analyzed by Bruhl.\* A case presenting symptoms of syringomyelia has been recorded in which after death gummata were found on either side of the brachial enlargement of the cord.†

The disorder described by Morvan and called by his name, and also "painless whitlows," presents symptoms of syringomyelia in conjunction with those of peripheral neuritis. At first there may be neuralgic pains in the hands, followed by anæsthesia and muscular wasting, and the formation of whitlows that undergo ulceration, and are attended with necrosis of the phalanges.

There is another disease resembling progressive muscular atrophy which may be here mentioned, the singular affection endemic in parts of Japan, known there as *Kakke*, and probably identical with the disease called in India and Brazil *Beriberi*. The generally accepted view is that beriberi is an infectious disease, developing under conditions of high temperature and moisture, and presenting the symptoms of a multiple neuritis. Observations made in Japan render it likely that the cause of the neuritis is generally poisoning by damaged rice, and it is said that attention to the diet has almost banished the disease from the Japanese navy.‡ Four types of the malady are recognized,—an incom-

---

\* Etude de la syringomyélie, Paris, 1890.

† Beevor, Lancet, vol. ii. 1893, p. 1252.

‡ Takaki, Report of the Japanese Navy, 1886, quoted in Sajous' Annual, vol. i., 1888.



pletely developed or rudimentary form; an atrophic form; a dropsical form, with or without atrophy; a pernicious or cardiac form. The most conspicuous symptoms are impairment of motion, with wasting and diminution in mechanical and electrical irritability, sensory changes, circulatory disturbances, abolition of the knee-jerks, diminished secretion of urine, and albuminuria.

It is sometimes a matter of extreme difficulty to distinguish cases of what are called *idiopathic atrophy*, or primary muscular atrophy where there is no central nervous lesion, from the progressive muscular atrophy under consideration. When the former disease happens in children the distinction is not so difficult, for the age, and the circumstance that not unfrequently several members of a family are affected, in some of whom it may assume the pseudo-hypertrophic form, show what it is. But in adults there may be great uncertainty. The extremely slow progress of the disease, its not unusual beginning in childhood, the fact that the muscles of the forearm and hand escape, as a rule, while the face is often involved, as well as the latissimus and the lower half of the pectoralis, that it affects women as often as men, and that it is congenital, are some of the characteristic points. Fibrillary twitching of the muscles is wanting, and the electric reactions undergo only quantitative diminution proportionate to the degree of wasting.

The difference in age helps us to distinguish *pseudo-hypertrophic muscular paralysis* from progressive muscular atrophy. A disease almost exclusively of childhood,\* probably of developmental origin, and often occurring in several members of a single family, it is characterized by weakness in the lower limbs primarily, the muscles of which, and particularly the calves, increase greatly in size. The infra-spinatus, also, is frequently enlarged, while the muscles of the face and those of the hands usually escape, and the latissimus dorsi and the lower half of the pectoralis are often wasted or absent. Yet, notwithstanding the apparent hypertrophy, there is debility, with a waddling gait, the knee-jerk is lost, extraordinary attitudes are assumed in attempting to rise from the ground, and, as the disease progresses and becomes more general,

---

\* Destarac, *La Médecine Moderne*, 1894, No. 89, p. 1387, has reported a case of pseudo-hypertrophic paralysis in a man 68 years old, without hereditary predisposition.



complete paralysis may ensue, with rapid dwindling of the affected muscles. These show, in the stage of increase, large masses of interstitial fatty matter with an augmentation of the interstitial connective tissue, and the muscular fibres are in a state of granular degeneration. The wasted muscles undergo shortening and contraction, and various deformities result. The reflexes become enfeebled, and the electric reactions are diminished. The function of the sphincters is, as a rule, preserved; intelligence is not affected; and sensibility is unimpaired.

**Infantile Paralysis.**—In this disease, also known as essential paralysis of children, acute atrophic paralysis, and acute anterior poliomyelitis, rapid wasting of the muscles is the striking feature. It is pre-eminently an affection of early childhood, and, as shown by Wharton Sinkler, occurs much more commonly in summer than in winter. It happens most frequently during the first dentition, and is often ushered in by fever, and by convulsions and other cerebral symptoms. The palsy comes on quickly, generally before the fever-disturbance has passed away; or an entire limb, or even both legs and arms, may almost from the onset be affected. In any case the palsy becomes 'plainly discernible as the fever subsides. It is apt to begin in one limb and in a few days to become wide-spread. But it disappears, except from a particular region in which the muscles quickly waste.

Yet the palsy may at first shift; it passes away from some limbs, or fixes upon others or upon different groups on different sides of the body. It rarely, however, remains as palsy of more than one side, and is not associated with loss of sensibility. In the absence of sensory symptoms, as well as in the rapid development of the disease, we have the points of difference which separate infantile palsy of the upper extremities from the wasting and paralysis that may be due to pachymeningitis in the cervical region. There is often decided recovery within six months from the onset of infantile paralysis; although some loss of power may be permanent. The affected muscles are apt to begin to atrophy after the paralysis has lasted a month, and when their wasting is marked they no longer respond to the faradaic current, though they may still react strongly under the galvanic current; but gradually this excitability, too, is lost. Both the superficial and tendon reflexes are lowered or abolished. After six months or a

year some faradaic irritability is apt to return. The functions of the bladder and rectum are very seldom affected. In protracted cases, permanent shortening of muscles happens, contraction of the joints takes place, and atrophy of portions of the osseous system occurs, or rather a want of its development in the blighted parts, and various and striking deformities result.

Now, the onset of these cases, the febrile symptoms, the occasional retrocession from certain parts, and the subsequent course, separate infantile paralysis from *progressive muscular atrophy*. Then in forming a diagnosis we may take into account the extreme rarity with which children are attacked with progressive muscular atrophy,—a disease of adults, and pre-eminently of those of the male sex who use their muscles continuously and violently. But the affection may happen in children, and then, as Duchenne points out, is apt to show itself first in the muscles around the mouth. On the other hand, we must not forget that a disease identical with the essential palsy of children is met with in adults. Beginning acutely with febrile symptoms, headache, delirium, vomiting, and rheumatoid pain in the back, it leads within a few days or less to palsy with complete relaxation of the paralyzed muscles, yet without impaired sensibility; exhibits but passing vesical disorder; but shows soon disappearance of reflex irritability and wasting of the limbs, with or without paralytic contractions, and has the lesion which has been found in infantile palsy,—granular degeneration of the cells of the anterior horns. This acute atrophic spinal paralysis is not so uncommon as was formerly supposed, and under the title of *acute anterior poliomyelitis* we are now familiar with its clinical history, and have learned how often complete or nearly complete recovery from the threatening symptoms takes place. There is a prevailing belief that the disease is due to a systemic infection.

From the foregoing remarks it might be inferred that children are only subject to palsies that are spinal. But this is not the case. We find in them a whole group of *cerebral palsies*,—not nearly so frequent, it is true, as the spinal group, but palsies in which the lesion is cerebral, extending from any part of the cortex to the pyramidal tracts of the cord, and broadly distinguished from the spinal palsy by heightened reflexes, unchanged electrical reactions, loss of power with disordered movements or spasm, and

retarded growth of the affected parts. We may find either hemiplegia, bilateral hemiplegia, or paraplegia as the form of paralysis. In some instances the affection follows delivery with the forceps; like spinal infantile palsy, it has been observed after infectious diseases. Under the first condition it is probably due to meningeal hemorrhage; under the second, to either hemorrhage into brain or membranes, or to vascular occlusion. Sometimes the disease begins with fever accompanied by convulsions; these may be followed by marked coma. The hemiplegia is most persistent in the arm, and is apt to be associated with spastic contraction, producing a peculiar gait. Post-hemiplegic chorea and mobile spasm and athetosis were observed in a considerable number of cases analyzed in Osler's elaborate monograph.\* Convulsive seizures on the paralyzed side or general epilepsy are yet more common, and the intelligence is enfeebled.

In the bilateral form of hemiplegia the legs are more involved than the arms; spastic contractions of the muscles of the extremities are most marked; the mind is very much affected; sensation is not disordered. Destruction of the motor centres of the cortex is the essential lesion in bilateral spastic hemiplegia.† In the spastic cerebral paraplegia of children McNutt‡ found descending degeneration in the pyramidal tracts; the disease is limited to the lower extremities; there is no muscular wasting; the gait is stiff or cross-legged. The malady usually exists from birth, and follows a difficult labor. The intellect is impaired, though not always markedly so. Wood§ states the affection to be the result of sclerotic and atrophic changes in the brain.

Before proceeding, we will examine the main forms of paralysis which we have been studying, arranged in a tabular form, and chiefly with the view of ascertaining the seat of lesion, premising that the statements must be received rather as generally true than as absolutely so.

---

\* The Cerebral Palsies of Children, 1889. See, also, Sachs and Peterson, Study of Cerebral Palsies of Early Life, based upon one hundred and forty cases. Journ. of Nerv. and Ment. Dis., May, 1890; and Sachs, Samml. Klin. Vortr., No. 46, 1892.

† Osler, *op. cit.*

‡ Amer. Journ. of Med. Sci., vol. i., 1885.

§ Nervous Diseases and their Diagnosis.

## TABULAR VIEW OF PARALYSIS.

*Symptoms.**Seat of Lesion.*

Inability to move leg and arm of one side. Sensation unimpaired, unless posterior third of posterior limb of capsule involved. Paralysis of muscles of lower part of face; mouth drawn toward healthy side. Electro-muscular contractility preserved. Reflex excitability of the tendons exaggerated.	Corpus striatum, involving internal capsule, both on side opposite to the palsy.
Same symptoms, dependent on involvement of internal capsule. Mobile spasm and incoördination in paralyzed parts.	Optic thalamus.
Same symptoms, but paralysis of face, with anæsthesia, on opposite side to that of arm and leg, and usually marked; conjugate paralysis or spasm of eyes; difficulty in deglutition and articulation. Heightened temperature; convulsions; contracted pupil. Urine may contain sugar or albumen. Early rigidity of paralyzed muscles.	Pons Varolii, on side opposite to palsy of limbs. The part affected is below decussation of facial nerve.
Same symptoms, but face paralyzed on both sides.	Pons Varolii, and at level of decussation of facial nerve
Paralysis of arm and leg and lower part of face on one side; third nerve paralyzed on other side; defective sensation; vaso-motor disturbance.	Crus cerebri on side corresponding to paralysis of third nerve.
Paralysis of motion of face, arm, or leg, soon followed by rigidity; sensation may be impaired. Reflexes, superficial and deep, increased. Convulsions.	Cortical part of brain in motor zone on side opposite to palsy.
Motion more or less completely affected on both sides of body, except in face; paralysis of hypoglossal, glosso-pharyngeal, and spinal accessory nerves; often rapidly fatal.	Medulla oblongata.



- |   |  |
|---|--|
| <p>Both legs and lower part of trunk paralyzed as to motion; loss of sensation; some wasting of muscles; loss of power over bladder and rectum; reflex excitability in legs heightened, trunk reflexes impaired; electric contractility diminished or lost; trophic changes; paralysis of muscles of respiration in some instances.</p>   | <p>In the cord throughout its sections above the lumbar enlargement, as in transverse myelitis of the dorsal cord.</p> |
| <p>Both legs paralyzed, muscles of legs flaccid; feet extended; anæsthesia; incontinence of urine from the start. Superficial and deep reflexes lost. Rapid wasting of muscles. Reaction of degeneration. Trophic changes.</p>  | <p>In the cord in lumbar enlargement, as seen in myelitis of these parts.</p>  |
| <p>Arms as well as legs paralyzed; arms flaccid, legs spastic; otherwise symptoms much the same; affection of pupils.</p>   | <p>Cervical region of the cord, as in cervical myelitis.</p>   |
| <p>Paralysis irregular in degree and distribution, relaxation of muscles, sensation unimpaired, only transient loss of control over bladder and rectum; marked lowering or extinction of reflex excitability in the palsied muscles and tendons; lost electromuscular contractility to faradaic current; usually reaction of degeneration; rapid muscular atrophy; no bedsores; if disease become chronic, muscular contractions.</p> | <p>Anterior horns of the cord, as in degeneration of the cells in acute poliomyelitis.</p>                             |

### Ataxia.

Loss of co-ordination of muscular movement, which in the legs shows itself especially in the gait, and in the hands, in the difficulty of executing delicate movements, but which strangely contrasts with the muscular power that is present in the limbs, is found in some neurotic affections, as in general paralysis of the insane, multiple neuritis, and diphtheritic paralysis. But the ataxia is most constant and marked in locomotor ataxia.

**Locomotor Ataxia.**—In this disorder we have uncertainty of motion and seeming palsy; or, in the words of Duchenne, who

gave it the name of progressive locomotor ataxia, it consists in "a progressive abolition of the co-ordination of movement with apparent paralysis contrasting with the integrity of muscular force." The patient is not deprived of the power of motion, but of the power of controlling his motion: hence he staggers in his walk, or cannot walk at all without support; the muscles are obedient to the will, but the peripheral impressions by which motor impulses are guided are improperly or imperfectly conveyed.

Locomotor ataxia is identical with a form of palsy clearly recognized by Todd, and with the malady described by Romberg as *tabes dorsalis*; from the lesion it exhibits, it is often called *posterior sclerosis*, degeneration of the posterior columns of the cord and of the posterior nerve-roots being its main cause. A wasting of the nerve-fibres of the peripheral spinal sensory nerves has also been found.

The affection is a very chronic one, lasting many years. It may originate without assignable cause; or may follow alcoholic excess, or exposure to cold, or injury or inflammation of the cord; or is hereditary. It has been observed to follow pernicious anæmia.\* It is most frequently found to be associated with a history of syphilis. Among its early symptoms are piercing pains, lightning-like or similar to electric discharges, in the lower extremities; enfeeblement or loss of knee-jerk; disordered gait; diplopia or other disturbances of vision, which may be attended with the "Argyll-Robertson pupil,"—a small pupil that does not contract to light, but does contract during accommodation,—or with paralysis of the sixth or the third pair; and a zone in which sensation is greatly impaired on a level with the third, fourth, fifth, or sixth dorsal vertebra.†

Following these phenomena, or making its appearance with them, is a difficulty in co-ordinating movements and in maintaining the equilibrium of the body. It is manifest in attempting to walk with the eyes closed; and the patient is unable to take a step, or to stand erect with his feet in juxtaposition,

---

\* Putnam, Amer. Journ. of Med. Sci., March, 1895; also Burr, University Medical Magazine, April, 1895.

† Hitzig, in Ziemssen's Cyclopædia, article "Atrophy of Brain."

without swaying and losing his balance. This, the so-called *Romberg* symptom, is not pathognomonic, but it is very valuable in the diagnosis of the earlier stages, and so is the difficulty in placing the foot on small surfaces, in buttoning the clothes, or in walking backward.

Yet the stumbling gait is not connected with true paralysis. The muscles can act vigorously, are well nourished, contract readily when faradized, except in advanced stages of the disease, and show neither tremor nor spasm. The cutaneous reflexes are generally, yet not always, impaired; there is absence of the patellar tendon reflex in both knees. Sensibility is markedly diminished, pinching and pricking the foot may scarcely be felt, contact with the floor may not be appreciated, and the tactile sensibility may be almost gone; but all kinds of curious sensations are complained of. The power to appreciate differences of temperature may, though it does not always, remain, and there is a delay in the perception of pain. The muscles, too, lose their sensibility. It is not unusual to have pains in the region of the fifth nerve. The intellect is unimpaired, unless frequent attacks of vertigo and epileptic seizures should be among the symptoms. The eyesight fails more and more, there is loss of color-vision, and an atrophy of the optic nerve may produce irremediable loss of sight; the hearing, too, may become much affected; and signs of valvular disease of the heart, especially of the aortic valve, show themselves. The functions of the rectum and bladder are not markedly disordered, though retention of urine and sluggish action of the bladder are not infrequent. The sphincter ani is often weak, but constipation is common. There is loss of sexual power. Dropsy and local sweating are met with, and so is swelling of the joints, without redness and usually without pain.

But the joint affection may appear, as Charcot has taught us, before the loss of power of co-ordinating movement. In time, it may be rapidly, the articular extremities of the bones disappear, and the joints undergo a kind of dislocation. The shafts of the bones, too, show defects of nutrition, and spontaneous fractures happen. The teeth drop out of the atrophied alveolar processes, and so may parts of the bones themselves;\* the tendons

---

\* Newmark's case, *Med. News*, Jan. 26, 1895.

tear; the tongue may dwindle on one side; the spine becomes curved. Herpetic, bullous, and pemphigoid eruptions or ecchymoses may appear during or subsequent to exacerbations of the lightning pains. Perforating ulcer of the foot has also been observed among the trophic changes.

Among some of the less common symptoms is drooping of the eyelids, accompanied by weakness of all the muscles attached to the eyeball, so that the movements of the ball become much restricted or wholly lost.\* Another symptom, more frequent, however, is the occurrence of spasms and pain in the epigastric region, with attacks of vomiting. These gastric crises, as they have been termed, may be found to happen in those who complain much of fulness in the abdomen and of unsatisfied hunger. They have even been known to lead to vomiting of blood. Buzzard† shows the symptoms to be dependent upon sclerosis affecting the nucleus of the vagus. There is always in these gastric crises acid fermentation, but hydrochloric acid is also constantly found.‡ There are at times attacks of laryngeal spasm in ataxias. Arthropathies often happen in those who present laryngeal or gastric crises.

These two forms of crises are by far the most frequent. But in addition we have intestinal crises, urethral crises, rectal crises, genital crises, renal crises, cardiac crises, and others, in which, as the chief symptom, violent paroxysms of pain occur, that pass away and are found not to be connected with any organic change of the seemingly diseased part. The true meaning of these pain crises, as well as the distinction from the visceral affections they simulate, is detected in the absent knee-jerk and in the other symptoms of the ataxic malady.

In considering the diagnosis of locomotor ataxia, let us first examine how it differs from the *general paralysis* of the insane. Both maladies are very chronic in their course, and in both there is loss, or certainly impairment, of muscular co-ordination. In the one case, however, it exists with tremors, with thickness of speech, with dementia, with peculiar delusions, but without the

---

\* Hutchinson, Transact. Royal Medico-Chirurg. Soc., 1879.

† Diseases of the Nervous System, 1882.

‡ Cathelineau, Arch. Gén. de Méd., Avril, 1894.



eye phenomena, and without the sharp, peculiar pains of ataxia. Then, in this malady, the hands are rarely affected; indeed, should, in process of time, the upper extremities share in the disorder, there is in them often rather cutaneous anæsthesia, with some trembling, than an obvious failure of co-ordinating power. It must also be remembered that the two diseases sometimes exist in combination.

With reference to the distinction of progressive locomotor ataxia from most of the *diseases of the spinal cord*, it is only necessary to remark on the extreme rarity of muscular spasm in ataxia; from spinal paraplegia the result of myelitis it differs in the fact that the muscles act with strength, the patient can flex and extend his legs and kick vigorously, while in spinal myelitis the affected limbs cannot move, though the knee-jerk may be excessive. The lightning, or electric shock-like, pains are not entirely to be trusted to in diagnosis, for they may happen in acute myelitis as well as in spinal pachymeningitis and in disseminated sclerosis. The absence of the knee-jerk in locomotor ataxia is of great value. Its presence, in addition to the tremor, the nystagmus, and the scanning speech, distinguishes disseminated cerebro-spinal sclerosis. But we must not overlook the possibility of mixed symptoms existing from the different forms of sclerosis being combined. In the disease to which Gowers has given the name of *ataxic paraplegia* we have both disease of the posterior and lateral columns and a combination of the symptoms of spastic paraplegia and of locomotor ataxia. The knee-jerk is excessive, ankle-clonus is present, and there are extensor spasms in addition to weakness and the incoördination; but no lightning pains or loss of light reflex attend the ataxia, as in tabes.

There is a chronic degeneration of the spinal cord having its chief seat in the posterior columns and the lateral pyramidal tracts, which mostly develops in childhood. It is often hereditary, usually occurs in families, is probably congenital in origin, and has as its chief symptom ataxia. This disease is known, from the name of the observer who first accurately described it, as *Friedreich's ataxia*, and also as *hereditary ataxia*, and is of very long duration. The disorder of co-ordination shows first in the lower extremities, and advances upward, at last affecting the organs of speech. The patellar tendon reflex is generally

abolished; nystagmus and vertigo are frequent; while in the later stages spasms and contractions of muscles, curvature of the spine, want of control in keeping any part of the body quiet, and palsies, are not uncommon. Unlike what takes place in locomotor ataxia, we note no disorder of cutaneous sensibility, no lancinating pains, no atrophy of the optic nerves, no Argyll-Robertson pupil, no trophic lesions, no visceral disturbances.\*

From *diphtheritic paralysis* we distinguish tabes by the history of the malady, the absence of pain, and by the paralysis of accommodation and of the palate that precedes the muscular weakness. Loss of knee-jerk exists in both, and occasionally incoördination is met with in the former. In *multiple neuritis* this, too, may happen; but the marked muscular and nerve tenderness, the changed electric reactions, the normal pupils, the more decided loss of muscular power, and, usually, the evidence of alcoholism, tell the true meaning.

*A diminution or loss of the muscular sense*—that guiding sense by which we judge of the position of the limbs, by which we are conscious of their movements—occasions difficulty in diagnosis, since in locomotor ataxia the muscular sense may be also deficient. On the other hand, in the former morbid state the motion may be somewhat impaired, for, as in tabes, the feet may feel numb in standing and in walking, and the patient be unable to walk in the dark. But there is this difference: where merely the muscular sense is affected, he can walk and perform all movements, even those of a complex nature, without vacillation, so long as his eye is fixed on them and superintends and gives them direction; while in tabes the derangement of muscular co-ordination renders, even with the aid of sight, the movements uncertain and irregular. Then cutaneous anaesthesia is apt to coexist with this malady. The treatment, too, will throw light on a doubtful case: the local use of electricity will usually cure the loss of muscular sense in hysterical paralysis; it has no curative effect in ataxia.

Irrespective of the affection of muscular sense, the greatest similarity to locomotor ataxia I have seen has been in several

---

\* For an admirable analysis of cases, see Crozer Griffith's paper in the Transactions of the College of Physicians of Philadelphia, 1888. Sanger Brown, Chicago Medical Recorder, Feb. 1892, publishes a very striking family tree.

cases of *hysteria*; one in particular, in a very anæmic woman, resembled it closely; and it may be a question whether the nutrition of the parts affected in ataxia was not disordered, and the nervous structure functionally disturbed. I desire particularly to call attention to these cases, which can be distinguished by their history, the usual coexistence of anæmia, and the absence of severe darting pains. Yet pains may also happen in the hysterical complaint, as in a case I saw with Dr. Webb;\* but this is uncommon. Moreover, the apparent want of muscular co-ordination is more irregular in its manifestations, and the cases recover. So, I think, may cases of locomotor ataxia due to special causes. For I have seen cases in *syphilitic* patients, typical in everything except perhaps the severity of the neuralgic pain, essentially typical in the muscular phenomena and in the inability to walk with closed eyes, in which a gradual and nearly complete recovery took place. Here the lesion was probably removed or greatly influenced by the anti-syphilitic treatment, and a true sclerotic degeneration of the affected parts did not take place.

**Diseases of the Cerebellum.**—Diseases of the cerebellum produce many of the phenomena regarded as peculiar to locomotor ataxia. But the gait of the patient is that of a drunken man: when attempting to walk, he leans to one side, moves in arcs of a circle, or describes zigzags; and when standing erect, his body swings backward and forward, or from side to side, though his feet remain quietly fixed on the ground. In ataxia, on the other hand, the muscular contractions in the erect position or during attempts at walking are strong and sudden, more like spasms, yet not spasmodic, and have as their object to keep the body in the line of gravity; and the walk, though accomplished with difficulty, is straight, not reeling; the affected person, too, while he is walking, does not take his eyes off the ground or off his feet, from fear of falling; but he is not giddy. The peculiar gait is particularly found when the middle lobe is involved. Disease spreading from the cerebellum gives rise to hypoglossal, facial, and other local palsies. In diseases of the cerebellum we find vertiginous sensations, especially during attempts at loco-

---

\* American Journal of the Medical Sciences, Jan. 1876.



tion, which may be easier and straighter with the eyes shut than with them open ; vomiting, particularly at the onset of the complaint, aggravated or brought on by the erect posture ; nystagmus ; severe headache, occipital or frontal, when the head is bent ; defective vision, becoming very marked when an object is looked at for any length of time, or double vision, though the eye-disturbances may or may not be associated with choked disk or optic neuritis ; no diminution either of power of motion or of sensibility, unless from pressure on adjacent parts ; and in some instances rotary movements and hemiplegia. Rotary movements are regarded as a special proof of affection of the cerebellar peduncles. The knee-jerks are sometimes wanting, sometimes exaggerated. When the disease is localized in one hemisphere of the cerebellum, it may cause no symptoms and be beyond the reach of diagnosis. But with reference to the differential diagnosis of this, as of any other form of brain affection from locomotor ataxia, we may lay stress on the occurrence of the shooting pains in the latter, on other sensory phenomena, on the pupillary reaction, and on the absence of the knee-jerk.

### Tremor.

Any involuntary agitation of the body, or of part of it, without marked muscular contraction or impediment to voluntary movement, is called tremor. The trembling depends upon a weakening of the muscular and nervous systems. It is common in old age, in convalescence from debilitating diseases, in hysteria, in neurasthenia, and during chills. We also find it in workers in mercury or in lead or in arsenic ; in those who abuse alcoholic stimulants or coffee or tobacco, or who are addicted to the use of opium ; and in cases of exophthalmic goitre. In some instances, as we have seen, it is connected with an organic disease of the nervous centres, as in cerebro-spinal sclerosis ; and it constitutes the main symptom of the disorder known as shaking palsy or paralysis agitans.

Tremor is easily recognized. Yet it may be confounded with muscular twitchings. But it differs from these spasmodic movements by being more incessant, and unconnected with decided muscular contractions. In nervous, susceptible persons laboring



under an acute attack of disease, it is at times combined with great restlessness, and is apt to be mistaken for a convulsive state. Here again it may be distinguished by the absence of muscular contractions, and by the unintermitting irregular motions.

**Paralysis Agitans.**—Tremor is the chief symptom of paralysis agitans. The trembling consists of fine small movements, is combined with muscular weakness, or rather with slowness of muscular action, and, though increased by exertion and mental excitement, it persists during rest, though it ceases during sleep. It usually follows continuous mental strain, emotional shock, prolonged exposure to damp, or some depressing acute affection in elderly persons; it comes on slowly and progresses slowly; it ordinarily begins in the hand or foot and gradually becomes general. The disease lasts for years: as it advances, the patient loses his equilibrium in walking, leans forward or walks on the fore part of the foot, and is rapidly propelled forward, but he may be propelled to one or the other side or even backward.

The trembling takes place over the entire body, except in the head. It is in more or less continuous oscillations, at first, at least to a certain extent, controlled by the will. The muscles react to both the faradaic and the galvanic current. The expression of the countenance is vacant and fixed; the handwriting is tremulous, the voice piping, monotonous, the speech indistinct, at times hurried; the muscles of the extremities become rigid, especially the flexors, producing deformities like those of rheumatoid arthritis. Sensation is little, if at all, affected; there is great restlessness. Complaints are made of cramps, of muscular stiffness, especially in the extremities, and of a sense of excessive heat, associated, indeed, with increased temperature of the surface. There are no cerebral symptoms; yet hypochondriasis and loss of intellectual power occur as the disease progresses. The hands are apt to assume a position as in writing. The knee-jerk is, as a rule, normal; it may be increased. In exceptional instances tremors are absent. There is growing belief that paralysis agitans is dependent upon a diffuse sclerosis of brain and cord.

**Multiple Cerebro-spinal Sclerosis.**—Different is the palsy dependent upon multiple or disseminated *cerebro-spinal sclerosis*, or *Charcot's disease*. The symptoms of this vary somewhat, as the nodules of hardened tissue affect the brain or the cord first.

We have always tremor and paralysis, and sometimes sensory changes. The trembling may show itself from the start in the tongue or the eyeball, and with it we usually find headache, vertigo, failure of sight, nystagmus, amblyopia, impaired hearing, and at times gastralgia and vomiting. The want of power manifests itself in all the extremities, yet the lower exhibit the palsy most plainly, while the characteristic trembling is most evident in the arms; unlike paralysis agitans, the paresis or paralysis often precedes the tremor. Save in rare instances, the trembling is not witnessed except when the muscles are put into motion; stops, therefore, entirely or nearly so when they are at rest: it is usually tested by letting the patient pass a glass of water to his mouth. It occurs in decided jerks, and markedly affects the head, when this is moved at all. The gait is uncertain and tottering, and attempts at walking increase the tremor. The voice is weak, the speech slow and scanning; there is mental enfeeblement, with failure of memory. Sensation is at first not affected, nor are the sphincters; but we may have hyperæsthesia, or anæsthesia, or paræsthesia, and girdle pains. One or more cranial nerves may be involved in the sclerotic process, with resulting disturbance of function. The tendon reflexes are generally exaggerated, and foot clonus is not uncommon. A peculiarity pointed out by Charcot is that the pupils move under light, and that the papilla is yellowish. Toward the end, muscular cramps followed by contractions, and disorders of deglutition and of respiration, happen; or there may be attacks of an apoplectic character. It is in very advanced cases only that the electro-muscular contractility or the galvanic irritability of the nerves is decidedly diminished. Multiple sclerosis is most common between twenty-five and thirty-five, and lasts for years. One of its striking features is that long delusive periods of marked improvement occur.

The description given shows the dissimilarity between it and paralysis agitans. The most difficult diagnosis is as regards Friedreich's ataxia, when, by way of exception, disseminated sclerosis happens in the young. The disturbance of co-ordination in the former malady and the common loss of knee-jerk are the most obvious differences.

There is a form of disease in which the symptoms appear like those of disseminated sclerosis, and are yet due to an infectious

process, such as scarlet fever, measles, variola, typhoid fever, and influenza. The tremor aggravated by tension, the scanning speech, the drooping lip, the dull expression of face and general air of stupidity, the spastic gait with exaggerated deep reflexes, are common to both. Nystagmus has, however, not been observed in this *pseudo-disseminated sclerosis*, and the cases recover. They are in their general character, except in the special symptoms that approximate them to sclerosis, like the pseudo-ataxia which is also observed after various acute infectious diseases.\*

There are other, though far less common, forms of tremor, connected with organic disease, such as the *post-hemiplegic tremor* and the tremor in *spasmodic tubes*. In both, the history of the case and the attending muscular disorder, with the violent but rhythmical tremors on attempted motion in the latter affection, are of great significance. As an organic tremor, too, may be classed that of old age. In this *senile tremor* the trembling is most probably due to degenerative changes in the motor tract. At first it happens only on voluntary movement, stopping during repose and sleep, though ultimately it continues during rest as well as during motion. It begins in the hands, but extends markedly to the neck and head, and finally becomes very much like the tremor of paralysis agitans. Dana,† studying this and other forms of tremor with great accuracy by means of Dudgeon's sphygmograph, states senile tremor, indeed, to be the evidence of an abortive form of paralysis agitans.

**Functional Tremors.**—There is a group of tremors in which there is no organic cause, or at least the cause is so fine as to elude detection. Toxic tremors belong to this group, and we will look at their characteristics.

*Alcoholic tremor* occurs only on movement. It is irregular, and of considerable range. It is very pronounced in the arms, face, and tongue; in the legs it generally shows itself only when they are put in action, as in an attempt to stand. It is associated, in acute cases especially, with great restlessness, and muscular twitchings are not uncommon. The trembling is usually

---

\* See Westphal's paper in his *Archiv*, vol. xiv. p 87; also cases collected by Dawson Williams, *Med Chir Trans*, vol. lxxvii., 1894.

† *Medical News*, Dec. 1892.



worse in the morning. Then, too, in its diagnosis we lay stress on the habits of the patient.

*Tobacco tremor* is a fine tremor which more especially happens in the hands. It is sometimes seen in the tongue, which is smooth and shiny, and is apt to be combined with a relaxed skin, an irritable heart, and feebleness of sight.

*Lead tremor* is also a fine tremor. It is irregular in its distribution, usually seen in the hands, increased by motion, and not limited. It is often associated with beginning weakness of the extensor muscles of the forearm, with a blue line on the gums, and may involve the lips and tongue.

In *arsenical tremor* the trembling is wide-spread. There is also some difficulty in co-ordination, with beginning muscular paralysis, darting pains in the arms and legs, and diminution of tactile sensibility.

*Mercurial tremor*, another variety of tremor, usually appears first in the tongue and face, later extends to the arms and legs. It is increased by emotion and effort, and is recognized by observing that the trembling and the incessant movements stop when the shaking limb is supported. Then the gradual manner in which the disease appears, its occurrence among persons whose occupations predispose them to the absorption of mercury, the wakefulness, the disorder of the digestive organs, and the sponginess of the gums, form a group of phenomena very characteristic.

*Asthenic tremor*, such as follows debilitating disease, is fine, is induced by voluntary movement, and is most marked after exertion or fatigue.

*Hysterical tremor* may be fine and irregular, or coarse and rhythmical. It is usually induced by emotion and movement, although the second variety may occur independently.

The tremor that commonly attends *exophthalmic goitre* may be regular, but is often coarse and jerky, and occurs only on movement.

There is a form of functional tremor which is found to be unconnected with any obvious cause and may last through life. This *essential tremor*, to call it by that name, comes on often in young persons and lasts through life. It is generally fine, but sometimes irregular and unequal, and is apt to be associated with other hysterical manifestations; it is sometimes very severe, as



in the case recorded by Lloyd,\* in which marked hysterical anorexia coexisted. It shows itself most markedly in the hands, is made worse by excitement and by attempts at motion, and to a great extent, but not entirely, ceases during rest. It is not associated with any other motor disturbance, and I have known it in persons of high intellectual endowments. It may not come on until middle age, is not dangerous, but is not curable. In an instance that came under my observation the father and the son, a young man, both had it at the same time to an equal degree. Kindred to it is the *hereditary tremor* described by Dana, which also is a fine tremor, that does not interfere with co-ordination, and which affects especially the upper extremities. It begins in infancy or childhood and continues during a lifetime, without shortening life. It is often brought out by an infectious fever, ceases during sleep, and may become associated with slight contractures of the fingers.†

### Spasms—Convulsions.

Both these terms are applied to involuntary muscular contractions, with, perhaps, this difference: the word spasm is used when we wish to express the idea of less extensive muscular derangement, but especially when the muscles of organic life are believed to be involved; and convulsions, when the disorder affects the muscles of the whole body, or at least many muscles at once, and chiefly those of volition. Yet these are not distinctions that can be very strictly carried out, for the two phenomena often coexist, and, being produced by the same causes and obedient to the same laws, can hardly be separated.

Spasms may be clonic or tonic. In clonic spasms the muscles are agitated by successive contractions and relaxations of their fibres. In tonic spasms the muscles are rigidly set, and retain for a time their contraction, in spite of every effort on our part, or on the part of the patient, to relax them. The most marked type of this disorder is seen in tetanus; the most perfect illustration of clonic spasms is furnished by hysteria.

Convulsions may be accompanied by a loss of consciousness, and abolished sensibility, as in epilepsy; or they may coexist

---

\* Amer. Journ. of Med. Sci., Sept. 1893.

† Ibid., Oct. 1887.

with unclouded thought and unaltered sensibility, as in tetanus. What their immediate cause is, it is very difficult to determine; as yet we possess little positive knowledge; and concerning the portion of the nervous centres where they arise, or the structural changes that attend an attack, we are still ignorant. General evidence favors the cortex of the brain or the medulla as being the centres disturbed; but the irritation need not be direct, it may be reflected to them. Of their exciting cause we may say that, in those of susceptible nervous organizations, any extrinsic irritation, such as teething or disordered digestion, leads to a fit. Further causes are diseases of the brain; sudden interference with the circulation; profuse hemorrhages; anæmia; contaminated blood; the toxic influence of lead. Children often have convulsions as the precursors of febrile diseases. Convulsions have further been observed as a result of rupture of the stomach.\* In point of diagnosis it is of great importance to distinguish whether their inroad is or is not symptomatic of a cerebral lesion. If there have been a previous disorder of the intellectual functions, or any other manifestation of a brain affection, we may assume the convulsions to be the signal of cerebral mischief. But when no such phenomenon is met with, we are likely to find the source of irritation in some other portion of the body. Practically speaking, when convulsions are among the first signs of a malady, they are not apt to depend upon a disease of the brain; and even if recognized to form part of the symptoms of a cerebral lesion, we may conclude that the lesion has not reached its highest degree of development, but is still, as it were, irritative.

Besides separating convulsions or spasms in conformity with their centric or their eccentric origin, we must always attempt to ascertain the particular nature of the cause. If centric, is it congestion, inflammation, a tumor, sclerosis, or other lesion of brain or membranes? or is the convulsion due to influences the cognizance of which is not within our horizon? If eccentric, is it owing to an impure or impoverished blood, to retained poisons, to ptomaines, or is it peripheral from nerve lesion or intestinal or other visceral irritation? and what is the probable share the reflex system has in the visible disturbance of the muscles? To solve

---

\* O'Farrell, *Lancet*, vol. i., 1894, p. 1243.

these questions is often very difficult, and nothing but a careful analysis of all the phenomena of the case enables us even to approximate the truth.

Among the most extraordinary forms of spasm connected with increased reflex irritability of the cord is the so-called *saltatory spasm*, in which so violent a spasm of the legs takes place when the patient's feet touch the floor that he is thrown into the air. In some instances, as in one described by Bamberger, palpitation, dyspnœa, and inequality of the pupils coexisted. Other forms of tonic or clonic spasm happen in different parts of the body from reflex irritation of certain nerve-tracts, and these functional spasms produce for the time being the most singular contortions and deformities.

Rhythmic movements of the head, associated with nystagmus, are occasionally observed in infants and young children. The oscillation is sometimes horizontal, sometimes vertical, sometimes both. Many of these children are rachitic, some are epileptic. Occasionally there is an antecedent history of traumatism. In some cases the condition is connected with defective light in the crowded dwellings of the poor, necessitating almost constant artificial illumination.\*

Friedreich has described as "paramyoclonus multiplex" a condition of clonic spasm often recurring in paroxysms and involving the arms and legs and face and neck. The movements are increased by emotion, and may be controlled by voluntary effort.

Closely associated with spasms are other kinds of irregular muscular movements, such as cramps,—a contraction of short duration of one or of several muscles, occurring in paroxysms and attended with severe pain; rigidity,—a permanent tonic contraction of the muscles, often encountered in diseases of the brain; and the jerking movements of chorea. Now, some of these, especially localized spasm and even rigidity, have a strong connection with the seat and character of the lesion. Thus, broadly speaking, if we have spasm, perhaps alternating with chorea-like movements, confined to one arm, one leg, one group of muscles, we may infer an irritative lesion in the cortical motor area, affecting in this monospasm the centre presiding over the muscular motion of the

---

\* Lewi, Medical News, Nov. 10, 1894.



disordered parts. Early rigidity in the muscles, especially after hemorrhage, is apt to be associated with increased faradaic and reflex excitability, but the contracted muscles become relaxed during sleep; in late rigidity the contraction or "contracture" is due, barring the instances of hysterical contracture, to descending degeneration of the fibres of the pyramidal tract. In all forms of contracture from nervous causes the shortening of the muscles and the rigidity are increased by movements, whether voluntary or passive.

#### DERANGED NUTRITION AND SECRETION.

Among the subjects connected with the nervous system which have of late years received most attention, there is none of more interest than the association of its disorders with derangements of nutrition and secretion. Now, such are manifest in paralyzed limbs or after nerve-wounds. But these obvious alterations need here only be referred to; it is the intention to speak rather of the less palpable phenomena, the trophoneuroses in which, at first sight, the nervous system is not so distinctly concerned. For instance, there is to be noted the rapid development of blisters and bedsores in connection with marked cerebral and spinal lesions; the skin may become the seat of diverse eruptions, undergo modifications of color and structure, the secretions may be augmented or diminished, the muscles and joints show textural changes, swellings may happen affecting various portions of the body, either external or internal,—yet all be due to disturbed nervous influence, and the real disorder, therefore, be in parts very different from where it appears. Then we find the trophic symptoms of atrophy of the muscles in acute poliomyelitis and in Friedreich's ataxia, at times in the latter affection often associated with blueness and coldness of the feet from vaso-motor change.

To particularize with reference to a few of the derangements mentioned. There is the affection described as herpes zoster, in which the vesicles encircling half the circumference of the trunk are not a primary skin disorder, but the local expression of irritation of a nerve. They closely follow the distribution of some superficial sensory nerve, and this unilateral herpes is really but a sign of localized neuralgia,—most generally of a dorso-intercostal neuralgia. Then, again, we encounter instances of large vesicles or bullæ accompanying other neuralgias, as of the



sciatic; and attacks of erythema having their origin in facial neuralgia. Furthermore, various kinds of spots and blotches, and thickenings of the periosteum and of the skin, have been noticed after this and other forms of neuralgia; and we have eruptions of zoster in chronic myelitis; these in locomotor ataxia are limited to the limbs affected with the pain. Then, we may have eczema of nervous origin produced by reflex irritation in instances of disorders of the urinary organs;\* and ichthyosis of the lower extremities in chronic spinal diseases.

Oftentimes, too, these morbid appearances on the skin are combined with evidences of altered secretion. Thus, in a case related by Parrot,† in addition to the neuralgic paroxysms attended with sanguineous exudations at the painful parts, there occurred, at times, bloody sweating of the knees, thighs, hands, and face. Lachrymation was noticed in nearly half the cases of trigeminal neuralgia analyzed by Notta;‡ and one-sided furring of the tongue is a not uncommon phenomenon in this complaint. Associated with these evidences of altered secretion may be signs of altered nutrition, such as iritis, corneal clouding, and inflammation of the fascia or of the periosteum in contact with the aching nerve. Let us here add that these manifestations of perverted nutrition are not confined to neuralgic disorders. Trophic changes occur also in diseases of the central nervous system. Thus, inflammatory affections of the joints have been observed to follow cerebral hemorrhages, and various spinal maladies, particularly acute myelitis; local dryness of the skin occurs in unilateral atrophy of the face, and in some cases of syringo-myelia; a form of joint-mischief, of hydrarthrosis, has been specially described in locomotor ataxia by Charcot; affections of the joints have also been observed in syringo-myelia; and the perforating ulcer of the foot has been found by Ball§ and Fayard|| to be often connected with locomotor ataxia. Perforating ulcer of the foot has, however, also been noticed in Morvan's disease, an affection in which paronychias and recurring ulcerations and necrotic processes attend other trophic

---

\* Ord, St. Thomas's Hospital Reports, vol. vii., 1876.

† Gaz. Hebdom., 1859; Handfield Jones on Nervous Disorders.

‡ Arch. Gén. de Méd., 1854.

§ Trans. of Internat. Med. Congress, vol. ii., London, 1881.

|| Thèse de Paris, 1881.

changes\* of a peripheral neuritis. Then the elevation of temperature on the palsied side, the red spots or bluish-red skin or the intense flushing of the surface, the tendency to sloughing of the skin on pressure that the palsied limbs exhibit, the ready way in which it blisters, show the vaso-motor disorder.

Œdema happens also as a vaso-motor change. Weir Mitchell† points out swelling of the limbs in menstrual periods. Furthermore we find local œdematous swellings occurring in various parts of the body associated with intestinal disturbance, sometimes periodically and with an hereditary tendency, and this *angio-neurotic œdema* has been noticed by Osler‡ to affect members of a family for five generations.

Among the phenomena of altered secretion connected with nervous affections, one of the most striking is *excessive sweating*. In lesions of the cervical sympathetic on one side, we may have strictly unilateral sweating of the face and neck, the other side remaining perfectly dry;§ and greater vascularity and increased temperature are concomitants. In lesions of the abdominal ganglia, profuse sweating also happens, and is apt to be combined with impeded secretion from the mucous coats of the bowels, as we at times find in instances of abdominal aneurism. Not that excessive sweating, whether localized or general, is always linked to an affection of the great sympathetic ganglia. We find local sweatings limited to the hands and feet without any signs of other disorder. And general sweatings, irrespective of those of colliquative character attending phthisis, or of those of malarial diseases, happen after low fevers, in inactive states of the liver, and in some persons go on for years without obvious cause. It may be that in most if not in all of these cases the sympathetic system is really at fault, at least in so far that there is a reflex derangement of the vaso-motor nerves, and of course, then, of the subcutaneous blood-vessels and of the glands they supply.

But these are not questions which we can here consider. Indeed, the *why* and the *how* of all these changes of secretion and nutrition attending nervous affections are still very uncertain.

---

\* Monod and Reboul, Arch. Gén. de Méd., July, 1888.

† Amer. Journ. of Med. Sci., July, 1884.

‡ Ibid., April, 1888.

§ As in the case recorded by W. Ogle, Med.-Chir. Trans., vol. lli.

To return to the clinical phenomena. Besides the external manifestations of altered secretion and nutrition, there are certain changes in internal organs, the expression of nervous derangement. There is exophthalmic goitre; the pneumonia that results from injury to the vagus; the ophthalmia, which may even pass on to perforation of the cornea, that happens after paralysis of the trigeminus; the kidney disease which follows chronic spinal affections. And the Medicine of the Future will acquaint us with many more disorders of glands and viscera which originate in altered nerve-structure and in perverted power.

So much for the chief manifestations of nervous complaints. From the preceding pages it will have become apparent how many of them are functional, or are at least of necessity so regarded, though science is steadily narrowing their number, and how these functional disorders may be attended with the signs of as great disturbance as the organic maladies. In consequence of the uncertainty respecting the functional affections, doubt is thrown over any anatomical or pathological classification of nervous diseases. I subjoin a table of the main affections, arranged according to their supposed sites. It may not suit a strict critic, since, in several of the disorders regarded as functional, modern research has indicated the probable organic cause. But from the point of view of the physician it would be premature to recognize a fixed lesion, and I contend rather for the classification being useful clinically than unimpeachable pathologically. Nor will it be adhered to in the description of nervous affections, which will be traced according to divisions formed by groups of symptoms and not in obedience to a pathological classification.

TABLE OF THE AFFECTIONS OF THE BRAIN AND SPINAL CORD.

CEREBRAL.....	{	<i>Organic.....</i>	{	Hyperæmia.
			{	Anæmia.
			{	Meningitis in its various forms.
			{	Hydrocephalus.
			{	Abscess.
			{	Softening.
			{	Sclerosis.
			{	Hemorrhage (Apoplexy).
			{	Thrombosis.
			{	Embolism.
			{	Tumors, etc.

## TABLE OF THE AFFECTIONS OF THE BRAIN AND SPINAL CORD.

—Continued.

CEREBRAL .....	<i>Organic.....</i>	<ul style="list-style-type: none"> <li>Aneurism.</li> <li>Glosso-labio-laryngeal paralysis.</li> <li>Syphilitic affections.</li> </ul>
	<i>Functional.</i>	<ul style="list-style-type: none"> <li>Delirium.</li> <li>Insanity?</li> <li>Hypochondriasis.</li> <li>Headache.</li> <li>Trance.</li> </ul>
CEREBRO-SPINAL..	<i>Organic.....</i>	<ul style="list-style-type: none"> <li>Cerebro-spinal meningitis.</li> <li>Disseminated cerebro-spinal sclerosis.</li> <li>Paralysis agitans.</li> <li>Hydrophobia.</li> <li>Tetanus.</li> </ul>
	<i>Functional.</i>	<ul style="list-style-type: none"> <li>Occupation-neuroses.</li> <li>Epilepsy.</li> <li>Catalepsy.</li> <li>Ecstasy.</li> <li>Chorea.</li> <li>Hysteria?</li> <li>Neurasthenia.</li> </ul>
SPINAL .....	<i>Organic.....</i>	<ul style="list-style-type: none"> <li>Hyperæmia.</li> <li>Anæmia.</li> <li>Spinal meningitis.</li> <li>Myelitis in various forms.</li> <li>Softening.</li> <li>Atrophy.</li> <li>Sclerosis.</li> <li>Locomotor ataxia.</li> <li>Spastic paraplegia.</li> <li>Hereditary ataxia.</li> <li>Ataxic paraplegia.</li> <li>Spinal apoplexy.</li> <li>Tumors, etc.</li> <li>Syringo-myelia.</li> <li>Syphilitic affections.</li> <li>Progressive muscular atrophy.</li> </ul>
	<i>Functional.</i>	<ul style="list-style-type: none"> <li>Spinal irritation.</li> <li>Spinal exhaustion.</li> <li>Tremor.</li> <li>Reflex spasms due to irritation of the cord.</li> <li>Acute ascending paralysis.</li> <li>Myotonia.</li> </ul>



**Acute Affections of which Delirium is a Prominent Symptom.**

This clinical group embraces the different forms of meningeal inflammation, delirium tremens, and acute mania,—affections in all of which the brain is the seat of the disturbance.

**Acute Meningitis.**—By this term is understood an inflammation of the membranes of the brain, especially of the arachnoid and of the pia mater. The dura mater is far less frequently attacked; very rarely, unless the morbid action be of syphilitic origin, or have extended from the bones of the cranium, or have resulted from an injury.

The disease generally presents two well-marked stages. The first, or the stage of excitement, is characterized by intense headache, great restlessness, vomiting, a hard, frequent pulse, fever, injected eye, often with a contracted pupil, strabismus, an increased sensibility to light and sound, obstinate constipation, irregular respiration, and soon by active delirium, and by convulsions. The second stage is marked by an evident ebbing of the life-forces: the extremities are cold, the pupils dilated, the pulse is feeble and much slower, and intermitting, or becomes extremely rapid and thread-like; involuntary passages occur; there is utter loss of mind and of sensibility,—in one word, coma or collapse. In this stage the temperature may fall below the normal, or may exceed  $106^{\circ}$ .

Not every case, however, has all these symptoms, or goes at once from the stage of excitement to that of collapse. There may be a well-defined period of transition, during which the heat of skin, except of the head, diminishes, drowsiness appears, and the pulse sinks somewhat in frequency. Again, the disease may be arrested before the signs of prostration are very evident.

The attack may be preceded by sick stomach, buzzing in the ears, and vertigo, or it may set in with severe pain fixed to the forehead and increased by movement. In some cases it begins with delirium or convulsions. On the other hand, these signs may be absent.\* Among the symptoms of the affection, even in the earliest stages, a persistent pain attacking one or both knees, violent, intensified on motion, unrelieved by local means, and

---

\* In a paper by Church, in *St. Bartholomew's Hospital Reports*, vol. iv., several cases without delirium are narrated.

connected neither with swelling nor with any other change in the form or appearance of the joint, has been particularly noticed.\*

The malady may pass rapidly through its stages, so rapidly that their distinctive features become confused and blended. Generally it does not last less, or much more, than a week.

Acute meningitis is brought on by alcoholism, by exposure, by depressing cares, by intense application to study, by a blow or fall upon the head, by disease of adjacent structures, or by syphilis; or it may occur in the course of infectious processes, such as measles, scarlatina, smallpox, typhoid fever, acute rheumatism, pyemia, though it is rare under all these circumstances; finally, it may be due to pneumonia or to insolation. It sometimes affects mainly, or wholly, the coverings of the convex portion of the brain; at other times the inflammation is limited to the base. *Meningitis of the convexity* is apt to be purulent, and, if purulent, temperatures of  $104^{\circ}$  to  $105^{\circ}$  are usual. It generally comes on suddenly, and is found to be connected with disease of the bones of the skull, with ear disease, or to follow exposure to the rays of the sun. Severe headache, delirium, hyperæsthesia, rigidity of the neck, spasms in the facial muscles of one side and in one or both arms, and hemiplegic weakness are among the most marked symptoms.

According to Duchâtelet,† *meningitis of the base* may be discriminated by remissions in the delirium, and by the coexistence of spasmodic symptoms with profound and early coma. These signs, at all events, are said to be distinctive in children, who, more than adults, are disposed to this form of the complaint. In some cases acute muscular pains with defective motor power, a clear mind until late in the disorder, a temperature of  $105^{\circ}$ , have been specially noticed.‡ Moreover, the long duration of the malady,—for it lasts for weeks,—with the delirium of varying intensity, not occurring soon, the intervals of clearness, and the late and incomplete palsies, is regarded as very significant of this simple basilar meningitis.§ Then persistent vomiting, paralysis of cranial nerves, and early optic neuritis point to the base;

---

\* Lund, quoted in Amer. Journ. of Med. Sci., Oct. 1864.

† Inflammation de l'Arachnoïde, p. 230.

‡ Dowse, Medical Times and Gazette, Feb. 1874.

§ Huguenin, in Ziemssen's Cyclopædia.

optic neuritis is indeed rare in meningitis of the convexity. Yet there is no certainty in the diagnosis. Nor can we be sure of the membrane chiefly involved in the meningeal inflammation. Inflammation of the dura mater has the least severe and striking symptoms.

A form of inflammation of the cerebral meninges characterized by extravasation of blood between the dura mater and the pia-arachnoid is known as *hemorrhagic pachymeningitis*. It has been observed most commonly in the chronic insane, and in cases of chronic alcoholism. Among the symptoms to which this condition gives rise are apoplectiform seizures, headache, somnolence or coma, muscular weakness, nystagmus, smallness of the pupil, optic neuritis, headache, and vomiting.

Acute meningitis is not always easy of diagnosis. Leaving out for the present the other disorders belonging to the same group, such as acute mania and delirium tremens, it may be confounded with

CEREBRITIS ;

ACUTE SOFTENING ,

INTRACRANIAL TUMOR ;

HEAD SYMPTOMS OF CONTINUED FEVERS ;

HEAD SYMPTOMS OF ACUTE RHEUMATISM ;

HEAD SYMPTOMS OF ACUTE ULCERATIVE ENDOCARDITIS ;

HEAD SYMPTOMS OF PNEUMONIA ; OF PERICARDITIS.

*Cerebritis*.—There is little appreciable difference between acute inflammation of the brain-tissue and inflammation of the meninges. In truth, what we commonly call meningitis is not unfrequently also cerebritis ; since the diseased process extends readily from the tunics of the brain to the adjacent cerebral substance. We note acute cerebritis generally as the result of an injury to the head, of contiguous inflammation, or of septic influence. We may suspect the brain-structure to have become involved if the sense of vision or of hearing be suddenly perverted ; if the convulsions, the agitation of the limbs, and the tremors be very marked ; if they occur chiefly upon one side ; and if coma succeed rapidly to the period of excitement, and be accompanied or preceded by one-sided palsy.

*Acute Softening*.—The form of acute softening which simulates meningitis is that associated with delirium. Acute softening is

almost always the result of arterial occlusion from embolism or thrombosis, or of venous thrombosis; arterial thrombosis is by far the most common cause. The existence of disease of the heart or of the blood-vessels, or of contracted kidney, gives us for the most part the clue to the case. There is also apt to be aphasia and one-sided palsy, and the palsied side has often a decidedly higher temperature than the other side. The general temperature is that of fever, and may be high,  $104^{\circ}$  or more. In the cases of acute softening in very old persons, where an atheromatous state of the blood-vessels of the brain is the cause of a plug forming, the rapid softening that may follow is apt to be preceded by restlessness, some mental confusion, and signs of a general breaking up of nerve-force; it is soon associated with disturbances of the bladder and rectum, and leads to coma. In the cases which I have seen there was neither much headache nor febrile disorder.

*Intracranial Tumor.*—A rapidly growing tumor or one of latent course may give rise to symptoms resembling those of meningitis, but the early appearance of paralysis, the marked optic neuritis, the progressive character of the symptoms, and the absence of febrile phenomena should suggest the cause.

Occasionally disease of the ear, with or without extension to the membranes or sinuses of the brain, may be attended with deceptive symptoms the nature of which may become clear upon ocular examination.

*Head Symptoms of Continued Fevers.*—In all the varieties of continued fever, but especially in typhoid and typhus, cerebral symptoms at times arise which bear a strong resemblance to those of idiopathic meningitis; and such symptoms may appear without the examination of the dead body showing even traces of inflammation. How, then, are we to distinguish these fever cases from meningitis? or how ascertain if meningeal inflammation be really before us as a complication, as it sometimes is, of the fever? Unfortunately, there is no sign absolutely diagnostic. Neither an increase of phosphates in the urine nor cerebral auscultation affords us any help, for the blowing sound that is at times perceived is not constantly present in meningitis, and may be heard in health. As matters stand, a diagnosis can be established only by a careful consideration of all the symptoms, and of the history,



especially of the onset ; by searching for the eruption of typhus or typhoid fever ; and by taking note of the expression of the countenance. The character of the delirium will be of service ; it is ordinarily much more active when the brain or its membranes are inflamed, and is attended with throbbing of the arteries of the neck and face,—a symptom, however, not conclusive, for I have repeatedly noticed it in low fevers,—and not unfrequently with convulsions. The relation between headache and delirium may be of aid. In general diseases headache ceases when delirium sets in ; in meningitis the two coexist. Then, too, we may lay stress on optic neuritis ; on retraction of the head, if present ; on the more intense headache ; on the vomiting ; and we may attach some, but not too great, importance to the red line made by drawing the nail across the forehead,—the meningitic streak. The most valuable differential sign is the loss of the knee-jerk, a loss that is apt to happen, at least temporarily, in meningitis. But how difficult it may be ordinarily to arrive at a correct conclusion, unless we possess a full knowledge of all the circumstances, is shown by this case :

A man, about thirty-five years of age, was admitted into the Philadelphia Hospital some years ago, with a certificate that he was laboring under typhoid fever. No clue could be obtained to the history of the malady. He himself was not in a state to answer questions. His pulse was excessively feeble, and somewhat irregular ; the eye was not injected, but suffused and watery ; the pupils were sluggish, and the eyeballs in constant motion ; the tongue was dark, dry, and fissured ; the breath offensive. There appeared to be pain on pressure in the right iliac fossa, but the bowels were constipated, and no eruption could be detected. The most striking feature of the case was the delirium, which was noisy and violent and accompanied by great restlessness ; the man sang, screamed, and was constantly attempting to get out of bed. The malady did not seem to be typhoid fever ; the symptoms belonged more to inflammation of the brain ; but, knowing neither how nor when the delirium had begun, a positive conclusion was not reached. The patient died the day after his admission into the hospital. The autopsy showed the intestines to be sound. The membranes of the brain, after the dura mater was removed, were found to be opaque, and between the convolutions were shreds of lymph and a puriform liquid. There were only traces

of inflammation at the base, except in the neighborhood of the pons Varolii, where some lymphic effusion was discerned. The ventricles were filled with fluid, and the nervous structure in the neighborhood of the thalami and corpora striata was softened.

Subsequent to the man's death it was ascertained that he had been ill for only four days before he entered the ward; which fact, had it been previously known, would have materially assisted in arriving at the diagnosis. Irrespective of the difficulty of its recognition,—a difficulty which now, with our knowledge of the eye-ground and of the reflexes, would not be so great,—this case is of peculiar interest. It illustrates the possibility of the absence of convulsions and of paralysis notwithstanding the most evident cerebral disorganization.

*Head Symptoms of Acute Rheumatism.*—In rheumatic fever cerebral symptoms occasionally arise which may be referred to inflammation of the brain, or which, by their prominence, may mislead the physician, causing him to regard the signs of the rheumatism as of little importance, if indeed he do not wholly overlook them. The morbid manifestations are very much like those of acute meningitis: restlessness, headache, and violent delirium, succeeded by coma; besides, rheumatic involvement of the muscles at the back of the neck may cause retraction of the head. The delirium is commonly of gradual approach, but it may come on suddenly. Generally it does not appear until the patient has been suffering for at least a week with acute rheumatism; and the heavy sweats and swollen joints point out the malady with which it is combined.

Examinations of the head, in cases which have proved rapidly fatal, fail to detect, save in rare instances, any evidences of inflammatory action within the cranium. The abnormal signs are, as a rule, more properly attributable to the rheumatic poison seizing upon the brain, and to the altered condition of the blood. They are at times found to be connected with the setting in of inflammation of the membranes of the heart, or of pneumonia, or with albuminuria, or with plugs of fibrin in the capillaries of the brain, and are frequently associated with a very high temperature.\*

---

\* For a collection of cases, I may refer to a paper on Cerebral Rheumatism

*Head Symptoms of Acute Ulcerative Endocarditis.*—The severe headache, the delirium, the somnolence, which may attend ulcerative endocarditis may cause it to be confounded with meningitis. Generally, however, the fever is of a typhoid type; and the high temperature, the rigors, the marked swelling of the spleen, are very significant, and so of course are the cardiac murmurs.

*Head Symptoms of Pneumonia; of Pericarditis.*—In both these maladies delirium may be met with of a character so violent as to lead to the belief that the brain or its membranes are involved in an inflammatory disease. The diagnosis is cleared up by a careful examination of the chest. Then we may lay stress on the furious delirium being unattended with spasmodic movements or with paralysis. The form of pneumonia which is mostly associated with delirium is inflammation of the upper lobes. True meningitis sometimes attends pneumonia, and is with great difficulty distinguished from the mere disturbance of the cerebral circulation just mentioned, unless persistent vomiting, and pressure on a cranial nerve, or optic neuritis show us the real meaning of the brain affection.

**Tubercular Meningitis.**—This is a rare disease in adults; not a rare disease in children. Indeed, nearly all the cases of so-called acute hydrocephalus, and most of those of meningitis of the base, are instances of tubercular meningitis, or, to define the morbid state, of an inflammation of the meninges occurring in tubercular patients, and ordinarily accompanied by the deposition of tubercles at the base of the brain, and by effusion into the ventricles.

The premonitory signs of the malady are of great importance. The child has generally been ailing for some time; is restless, peevish, sleeps badly, complains of headache, and is troubled with a frequent, short cough, and with constipation. To these symptoms are soon added thirst, a slightly-coated tongue, vomiting, a dry, feverish skin, an accelerated pulse, and grinding of the teeth, constituting the prominent features of the first stage of the

---

which I published in the American Journal of the Medical Sciences, Jan. 1875. Dr. Posner, in the German translation of this book, points out that the use of salicylic acid, now so much employed, may give rise to confusing cerebral symptoms, such as headache, vertigo, hallucinations, even delirium.



affection. After four or five days the second stage is reached, and the brain symptoms become more clearly developed. The child shuns the light, puts the hand frequently to its head, and utters now and then a peculiar, sharp, distressing cry. At night the headache becomes worse, and is attended with fleeting delirium. A slight strabismus is observable, and the eyeballs oscillate. The pulse is very irregular in its rhythm, sometimes rapid and intermitting, then suddenly falling and becoming quite slow. The vomiting ceases, and there may be a remission in the symptoms, with restored intelligence; but the pulse remains irregular, the temperature is moderately elevated, the bowels are even more constipated than before, and the abdomen appears retracted. The third stage is one of complete stupor, accompanied or preceded by convulsions. The expression of the face is idiotic; the pupils are dilated; there is subsultus, and one side of the body is paralyzed. Deglutition is difficult; the surface is covered with cold sweats. This condition, so painful to behold, may last for days; repeated convulsions hasten its termination.

Can we distinguish this formidable complaint from *ordinary meningitis*? Seldom from meningitis of the base; generally from meningitis of the convexities. As regards the discrimination from the former malady, we are, it is true, sometimes enabled to pronounce the affection to be tubercular meningitis, if we are familiar with the patient's antecedents, and are cognizant, previous to the seizure, of the presence of scrofula of bones or joints, or of tubercle in any of the internal organs, or are able at the time to detect scrofulous glands or tubercular phthisis. But without knowledge of this kind a positive diagnosis is impossible: we have, notwithstanding symptoms of basilar meningitis, nothing to direct us except the probability that the case is tubercular, because most instances of meningitis of the base are of that nature. This uncertainty does not exist with reference to the usual form of simple meningeal inflammation. We may generally distinguish the tubercular malady by its occurrence in an unhealthy person; by its insidious approach; by the absence of violent delirium; by the appearance of convulsions, not early, but late in the disease; by the far less violent headache, and the less degree of febrile excitement; by the notable remissions in several of the cerebral signs; by the chest symptoms, and the long duration of



the affection. The ophthalmoscope gives no certain information ; tubercles are not commonly found in the eye-ground, only optic neuritis or choked disks.

Tubercular meningitis is ordinarily attended with an effusion of serum into the ventricles, and it is plain that many of the symptoms are attributable to pressure of the fluid on portions of the brain. Now, how can we separate the malady acute hydrocephalus, as it used to be called, from dropsy of the brain, or *chronic hydrocephalus*? Partly by the history of the case, and partly by the normal size of the head ; for the water on the brain is not sufficient in amount nor is it there long enough to produce an appreciable augmentation of the cranium. Then, in chronic hydrocephalus the symptoms manifest themselves for years, from childhood even to adult life. The signs of a profound cerebral lesion appear gradually, the special senses are by degrees enfeebled, but it is a long time before they are wholly abolished, or before complete loss of consciousness takes place.

As regards the diagnosis between tubercular meningitis and *acute hydrocephalus*, it need only be stated that the latter affection is in the vast majority of cases a synonyme for the former. Yet we occasionally meet with instances in which acute hydrocephalus occurs unconnected with tubercle. It then either runs a latent course, or appears as an acute malady with symptoms similar to those of acute meningitis, beginning either with fever or with convulsions, and often attended with intense restlessness, succeeded by drowsiness, and having periods of intermission of the symptoms and of apparent improvement. Toward the end convulsions are common. The complaint, unlike tubercular meningitis, happens in previously healthy children, begins suddenly, and is of short duration. But the effusion may remain, and the disorder lead to chronic hydrocephalus.

There is a functional disturbance of the brain of great importance to discriminate from tubercular meningitis,—the *hydrocephaloid* disease described by Marshall Hall. It has a stage of irritability, and a stage of torpor : a stage in which the little patient is restless, feverish, and a stage in which the countenance becomes pale, the breathing irregular, the voice husky, and the pupils are dilated and uninfluenced by light ; somnolence, coma, and even general convulsions occur. The symptoms indicate cerebral

anæmia and nervous exhaustion. They generally come on after an enfeebling attack of illness, especially subsequent to protracted diarrhœa or loss of blood; sometimes they follow premature weaning. In the history of the case; in the less tendency to vomiting; in the irregularity of the pulse; in the flaccid and hollow state of the fontanel, so dissimilar to its prominent and tense condition in inflammation; and in the arrest of the threatening signs by stimulants and by tonics,—we find the guides which enable us to decide against the existence of an organic disease of the brain or its membranes.

But other affections besides those of the brain may be confounded with tubercular meningitis, such as typhoid fever and pneumonia. From *typhoid fever* tubercular meningitis may be distinguished by the frequent vomiting; by the retracted abdomen, so unlike the swollen, tender belly of enteric fever; by the constipation instead of the diarrhœa; by the normal size of the spleen; by the irregularity of the pulse; by the occurrence of convulsions and anæsthesia and other signs of profound motor and sensorial disturbance, and by the lower heat, the thermometer seldom rising above 102°. I have never seen an eruption in tubercular meningitis; but Barthez and Rilliet speak of fugitive imperfectly-formed rose-spots being present in rare cases. The duration of the two complaints affords no help in diagnosis, since the one may last as long as the other.

Tubercular meningitis is often mistaken for the typhoid fever of childhood; indeed, there are many points of close resemblance between them. Yet, except in those rare cases of coexisting acute tuberculization of the intestines, we do not perceive in the cerebral disorder a tongue red at the edges, diarrhœa, and other manifestations of intestinal irritation; and vomiting and nausea are more prominent and protracted symptoms than in the febrile malady. But in this complaint the temperature is generally much higher; the pulse is quicker, yet not unequal and subject to such decided variations; delirium occurs much earlier, and is much more marked,—indeed, tubercular meningitis may run through all its stages without mental wandering.

In reviewing the maladies with which tubercular meningitis may be confounded, it is incumbent upon us to bear in mind the *inflammatory affections of the lungs*, which, in children especially,

are not uncommonly associated with delirium and other brain symptoms. But the cerebral phenomena take a different course; the febrile excitement is more intense; and an examination of the chest reveals the cause of the disturbance of the brain. Yet we must not overlook the fact that the signs of acute phthisis may be like those of acute bronchitis or of acute pneumonia; that hence it may become a very perplexing subject to determine the precise cause of the disordered respiration. In adults the difficulty is far less, because the demonstration of the existence or non-existence of pulmonary tubercle is much easier. As an important point in the diagnosis of the tubercular meningitis of children, with reference to the chest symptoms, Gee \* mentions that the chest heaves equally well on both sides, yet over a very large part, or even the whole, of one side, no respiratory sound is heard.

Tubercular meningitis is not so rare in adults as has been supposed, and presents, as Seitz in his admirable monograph has shown, marked features of pain in the head and temperature variations,† exhibiting a fever of moderate type, with irregular remissions. The deposit of tubercle both in adults and in children may not be confined to the head. Indeed, the observations of Liorilli ‡ teach that the spinal cord is frequently implicated.

The points of the differential diagnosis of the tubercular meningitis of adults are much the same as with reference to the disease in childhood. Yet one disorder is more apt to be confounded with it,—*hysteria*. Indeed, in young women the onset of the malady may develop very misleading hysterical symptoms. But on close examination we find the traits of the cerebral malady,—the temperature record of the attending fever, the unequal pupils, the divergent strabismus, the optic neuritis, the trophic changes in the skin, the incontinence of urine, the local beginning of the convulsions.

**Cerebro-spinal Meningitis.**—Now and then cases of meningitis are encountered in which the inflammation affects simultaneously the membranes of the brain and of the spine, and in

---

\* Reynolds's System of Medicine, vol. ii.

† Die Meningitis tuberculosa der Erwachsenen.

‡ Archives de Physiologie, 1870.



which the symptoms of the cerebral malady are found to be blended with severe pain along the vertebral column, with retraction of the head, with convulsions, with rigidity of the muscles, with perverted cutaneous sensibility,—in short, with the phenomena denoting spinal meningitis. But such sporadic cases are of rare occurrence. Generally cerebro-spinal meningitis is not met with save as an epidemic disease which presents itself in somewhat dissimilar forms, changing mainly as the cerebral or the spinal disturbance prevails, and belongs clearly to the group of fevers, with which it will be described. But here may be pointed out the extreme difficulty of recognition of the sporadic non-epidemic cases. The early retraction of the head, the eruptions, and the increasingly high temperature of cerebro-spinal fever are the most valuable diagnostic signs. Pneumonia, so common in this, may, as some cases mentioned by Gowers\* prove, also happen in the sporadic malady.

**Delirium Tremens.**—The prominent trait of this complaint is delirium, associated with trembling and with sleeplessness. It occurs in intemperate persons; yet such is not always the case, for we may find an affection identical with mania a potu in those who are not intemperate in the ordinary acceptation of the word, but whose nervous system has been racked by persistent mental anxiety, or by the use of other than alcoholic stimulants. I have seen such cases from the constant taking of chloral and of paraldehyde; and Levinstein notices the same in those who are addicted to the use of morphine.†

Generally, however, delirium tremens is brought on by the abuse of intoxicating liquors. It is a current belief, and one which has found much favor among habitual drinkers, that a diminution or a sudden discontinuance of the accustomed beverage is followed by an onset of delirium. This may perhaps happen; but, if I am to take as a standard the large number of cases of the disorder which have come under my care at the Philadelphia and Pennsylvania Hospitals, I should say that its appearance is most commonly preceded by a long-continued and unusually severe debauch, which finds its winding up in an attack

---

\* Diseases of the Nervous System.

† Die Morphiumsucht, Berlin, 1877.



of mania; hence that this occurs in consequence of an excess, rather than of a diminution, of the habitual stimulus.

Let us look a little more closely at the mental wandering. It is very rarely fierce; nor is the patient taken up wholly with his delusions. He pays a certain amount of attention to surrounding objects, answers, perhaps in a rambling manner, the questions put to him, but fancies that animals are running around on his bed or are crawling on the walls, and is thereby, or by some equally distressing illusion, kept in horror and in dread. Or he imagines himself to be engaged in his ordinary occupations, and gives minute directions as to what he wishes done; tries to get out of bed, yet is quite tractable when thwarted in his efforts. He is very restless, his hands are constantly moving, and his delirium, to use the graphic epithet of Watson, is a busy one. With it are associated great sleeplessness, a frequent, soft pulse, utter loss of appetite, a moist, coated tongue, and a clammy skin.

The tremor is irregular, wide in range, affects particularly the arms, face, and tongue, and is only induced on attempted movement. There is often, besides, spontaneous muscular twitching. The temperature is usually elevated, though rarely to a considerable degree; it seldom reaches  $103^{\circ}$ .

How are we to distinguish the malady from one to which it bears a certain resemblance,—*acute meningitis*? Taking clearly-expressed examples of each, we find the following marks of distinction: the pulse is different; tense and hard in meningeal inflammation, it is yielding and soft in delirium tremens. The skin and tongue are dry and feverish in the former affection, moist in the latter. Then the characteristics of the delirium are dissimilar: and in the one disease the mental wandering is combined with severe headache, but not with tremors; in the other, with tremors, but not with headache.

Yet in actual practice the diagnosis is not always so easy as it might appear to be at first sight, and here and there we meet with cases presenting symptoms the exact meaning of which it is puzzling to determine. The difficulty is mainly occasioned by extreme cerebral congestion, or by inflammatory action, having been produced by the same exciting cause that has brought on delirium tremens. In this blending of two morbid states, the pulse is, or soon becomes, tenser than in pure mania a potu; the temperature

is apt to be higher, and the irritability of the stomach more marked and more persistent. In some instances, convulsions, strabismus, and deep stupor—carefully to be distinguished from the sleep which often announces the termination of mania a potu—set all doubt at rest. But when these signs are not present, we have to judge of the mischief that is going on within the cranium chiefly by the vascular excitement, by the activity of the fever, and by the appearances of the eye-ground. Yet caution is necessary in accepting as evidence phenomena which may be of diverse origin: the marked fever may be the result of, what is very frequent in delirium tremens, an intercurrent or coexisting pneumonia, of a gastritis, or of a pulmonary apoplexy, as in a case I saw at the Philadelphia Hospital years since. Then, again, we must not overlook the fact that in instances of pneumonia of the apex a delirium very similar to that of mania a potu may happen.

There is another point connected with the diagnosis of the malady which it is necessary to mention, and chiefly for the purpose of calling attention to a common error. The fact that a person known to be of bad habits is affected with delirium is received as a sure indication that the mental delusions have been produced by the abuse of ardent spirits. But they may be owing to other causes: to fever; to a visceral inflammation; to acute mania. To avoid being deceived, we must lay stress rather on the special character of the delirium, and on the symptoms with which it is combined, than on its mere presence. In other words, delirium in inebriates is not of necessity the fruit of intemperance. In discussing acute mania we shall return to this subject.

When delirium tremens ends fatally, death takes place from exhaustion. The fatal issue is occasionally brought on by an intercurrent inflammation, especially of the lung, or by disorder of the kidneys and uræmia. Sometimes, after the subsidence of the urgent cerebral symptoms, the patient dies very unexpectedly, and there are no morbid appearances in the brain or its membranes to account for the abrupt extinction of life. In many instances, however, of these sudden deaths, a large amount of serum is found in the ventricles, or in the subarachnoid spaces.

**Acute Mania.**—It would be obviously out of place to attempt to give, in a work of this kind, a detailed account of any of the forms of insanity; but, in its acute variety especially, it resembles

other affections of the nervous system so closely that it cannot be wholly passed over.

There are mainly two disorders with which acute mania is liable to be confounded,—acute meningitis and delirium tremens; and we shall for our purposes best learn the manifestations of acute mania by contrasting it with these maladies.

From *acute meningitis* mania differs in these essential particulars: the premonitory symptoms of the former are headache, drowsiness, and often a sense of tingling and of numbness in the extremities; these signs are, however, soon succeeded by the severer headache, tense pulse, decided fever, and optical illusions of the developed disease. The premonitory symptoms of acute mania, on the other hand, have generally existed for a longer time before the marked outbreak; some singular change of manner or of mode of thought commonly precedes the first violent attack of insanity, except in those cases in which the overthrow of reason results from a sudden, great grief, or from a violent shock to the nervous system. Further, when the delusions have taken full possession of the mind, the patient attempts to act up to them, and his bodily strength enables him to do so. He has little if any fever; no spasms; his pupils are not contracted; his stomach is not irritable; he does not suffer from headache, or at least does not in any way complain of his head. It is needless to point out how all this differs from acute inflammation of the brain.

There is but little difficulty in discriminating between typical cases of *delirium tremens* and of acute mania. The anxious countenance, the alarm, the good-natured loquacity and restlessness of the patient, his moist skin, compressible pulse, and creamy tongue, are very different from the ravings and excitement, or the stubborn silence alternating with the wild hallucinations, of insanity. Yet there are cases in which it is not easy to tell if the delusions are really due to intemperance: cases of insanity excited by drink in persons predisposed to mania. It may, indeed, at first be impossible to decide upon their nature, and upon the share the drinking has in their production. A few days, however, ordinarily remove all uncertainty: the person who was thought to be merely delirious is seen to become frantic after an intermission of quiet, or, unlike what happens in mania a potu, to be still out of his mind after he has had several sound sleeps. In one instance, in



which much doubt existed as to the diagnosis, the patient solved the doubt by jumping out of bed after having been quietly sleeping for hours, and, in a state of wild excitement, knocking down the nurse who tried to prevent her from leaving the room. Furthermore, in *acute alcoholic mania* there is a strong tendency to homicide, while in *acute melancholia* induced by drink the tendency is to suicide.

### Diseases marked by Sudden Loss of Consciousness and of Voluntary Motion.

The chief diseases of this class are apoplexy, sun-stroke, and catalepsy. Epilepsy, too, might be here regarded ; but it will be more convenient to consider it with the convulsive affections.

**Apoplexy.**—This is coma coming on rapidly, and occurring nearly always as a result of intra-cerebral hemorrhage. It may be, however, also caused by sudden arrest of the blood-supply to the brain, by concussion, by congestion, and by laceration of the brain. Disease of the blood-vessels and miliary aneurisms are very commonly present, or we observe the malady in affections in which hemorrhages are prone to happen, such as in purpura, in scurvy, in pernicious anæmia, in leucocythæmia.

The malady has sometimes no prodromata ; but not unfrequently it is preceded by great depression of spirits, by attacks of loss of memory, by illusions, by vitiated perceptions, by vertigo, by odd sensations in the head, or by one-sided weakness or numbness.

The seizure is generally sudden, and the coma quickly developed. The patient falls to the ground, bereft of all consciousness. In other instances, before he sinks into the comatose sleep, there will be more or less pain in the head, sickness at the stomach, heaviness and confusion of thought, or even slight convulsions. Such gradual cases, Abercrombie tells us, are more dangerous than those of abrupt origin. Again, we may have convulsions a prominent feature almost from the onset.

When, whatever the beginning, the attack has reached its height, it presents these well-known features : the patient lies as if in a deep sleep, breathing laboriously and noisily, and each snoring inspiration is followed by a flapping of the cheeks in ex-



piration. The pulse is slow, full, at times irregular; the carotids throb violently, and the increased pulsation is particularly noticed in large extravasations; there is difficulty of deglutition; the pupils are immovable, and either contracted or dilated; the eye is half open; there is conjugate deviation. All thought, all sensation, all volition, is suspended; the limbs are motionless, flaccid, and, when lifted, fall passively and to all appearance lifeless to the ground. Occasionally their muscles are rigid; but, save when the apoplexy is very extensive, reflex contractions can be excited in them, or, to speak more accurately, on the paralyzed side the patellar tendon reflex is exaggerated, while the superficial reflexes are absent. In severe cases the insensibility becomes greater, the reflex action is abolished, the breathing becomes very irregular, of the Cheyne-Stokes variety, and involuntary discharges take place from bladder and rectum.

If the patient recover from the comatose state, he does so generally in a short time: in a few hours, unless the lesion be very great, the intellectual faculties begin to resume their sway, and all the functions of the body are slowly restored to their natural condition. Yet there is a palpable exception to this in the muscular system. Paralysis of one side is apt to remain.

The temperature variations in apoplexy may be turned to useful diagnostic account. The temperature of the body is at first somewhat lowered, but this is followed by a stationary normal period, and not unfrequently by a rapid rise, which again, as the patient recovers, is succeeded by a return to the natural body heat. In severe cases where large hemorrhages take place, the temperature seldom rises, or only rises to fall with the recurrence of the fatal bleeding, yet some apoplectic lesions of the pons and the medulla are throughout attended with elevation of temperature. If the stationary period be short or absent, and the body heat rise therefore almost continuously after the primary depression, the prospects of recovery are gloomy. From Dana's elaborate study,\* we know, indeed, that in fatal cases of apoplexy a rise of rectal temperature to from 100° to 102° F. occurs, as a rule, on the first day after the seizure. On the second day, if the cases be not immediately fatal, the temperature falls a little, averaging 101° or 101.5°

---

\* American Journal of the Medical Sciences, June, 1894.

F. In cases of acute softening, on the other hand, there is rarely any disturbance of temperature, even in fatal cases, on the first day. On the second day there is a slight rise to from  $99.5^{\circ}$  to  $100^{\circ}$ . As regards the surface temperature, it is noted that in instances of extensive intracranial hemorrhage the temperature upon the paralyzed side is somewhat higher than upon the sound side, while in cases of thrombosis or embolism there is no such difference.

Apoplexy is very apt to happen after dinner and during sleep, and is most common when sudden variations of temperature are most frequent. Liddell has shown that attacks are more usual in the spring. In New York he found the mortality greatest at that time of year.\* One attack of apoplexy is likely, sooner or later, to be followed by another; and the reason of this is, that the predisposing cause is generally of a persistent character,—an organic cardiac malady, especially hypertrophy of the left ventricle or tricuspid regurgitation; Bright's disease; degeneration of the cerebral arteries; disseminated sclerosis, or softening of the brain. It is likely that the extravasation of blood is generally due to the same immediate cause,—to rupture of miliary aneurisms on the minute diseased arteries.

Now, is there anything at the time of the apoplexy, or after its most urgent symptoms have passed away, by which we can recognize whether the pressure on the brain results from a clot, from vascular obstruction, or from turgescence of the cerebral vessels? And, again, do the morbid manifestations furnish any clue to the seat of the hemorrhage? With reference to the former question, all clinical experience forces us to admit that, in any of the states mentioned, the actual signs may be the same, and that we never can be quite certain of the non-existence of a clot. It is true that when the apoplectic symptoms abate rapidly; when thought, however confused, soon returns; when the limbs are not paralyzed, or are so but imperfectly and for a short time, we have strong reason for believing that congestion, simply, lies at the root of the disturbance; that, in other words, the case is one of those called simple apoplexy. But it is never possible to give a positive opinion, since a clot near the periphery of the brain may occasion the same phenomena as those specified. Attacks of cerebral con-

---

\* Treatise on Apoplexy, New York, 1873.

gestion with apoplectic symptoms happen in the general paralysis of the insane. The features of this point out their nature.

With regard to a rapid effusion of serum, the difficulty of distinction from hemorrhage is very great. In fact the only differential signs which were formerly claimed for *serous apoplexy*, namely, pallor of face and feebleness of pulse, are common in large sanguineous effusions; and we find absolutely no sign that can be looked upon as conclusive. In a case which came under my observation some years since,\* the respiration was not noisy, nor was there flapping of the cheeks, or the least discernible movement of any portion of the body; yet none of these points can be regarded as diagnostic. Most of the cases of so-called serous apoplexy are instances of Bright's disease with serous effusion into the brain, and the very existence of the disease is now, for the most part, denied.

The *seat* of the hemorrhage can be detected with more certainty than the cause of the cerebral pressure; it could be detected with greater certainty were it not that the extravasation so often takes place into an already diseased brain. The order of frequency in which hemorrhage occurs into various parts of the brain is as follows: central ganglia, cerebrum ovale, cortex, pons, cerebellum, medulla, crus cerebri. In the majority of instances the blood is effused into one of the corpora striata and the *internal capsule*, or at the same time into the optic thalami, and we find only one-sided paralysis. If the lesion be in both hemispheres, the palsy is on both sides of the body, although more complete on one side than on the other. Yet a double-sided palsy does not justify an absolute opinion that the extravasation of blood into the brain-substance is double-sided; it betokens also an extravasation into the ventricles. But *ventricular hemorrhage* is distinguished by profound coma and by tonic contraction of the muscles, or by tonic alternating with clonic spasms, and rigidity of the muscles either on one or on both sides occurs; the respiration is much embarrassed, and the breath-sounds are obscured by râles. It is common in the very young and in the old, and paralysis is frequently absent, though it may be general.† Ven-

---

\* Charleston Medical Journal and Review, March, 1859.

† Sanders, Amer. Journ. Med. Sci., July, 1881.

tricular hemorrhage is more often secondary than primary, the blood having torn its way into the cavity. Hemorrhage limited to the *thalamus* may give rise to no symptoms unless the internal capsule be damaged, when slight hemiplegia, hemianæsthesia, and hemianopsia, with mobile spasm and motor incoördination, are apt to show themselves.

Hemorrhage into the *corpora quadrigemina* presents most frequently this combination of symptoms: muscular incoördination, impairment of sight and alteration of the pupils. *Cerebellar hemorrhage* gives rise to very temporary loss of consciousness, to unsteadiness of movement, and to frequent vomiting; vision is not affected. In instances in which there is hemiplegia it may or may not be on the same side as the lesion. In hemorrhage into one-half of the *pons*, there is palsy of the extremities on one side, and of the face on the other.\* The pupils are often contracted, though they may be dilated and inactive. Disturbance of respiration is common. In lesions of the *pons*, too, as in those of the medulla, we have high and rapidly rising temperature almost from the onset, and we find an exception to the rule that the lateral deviation of the eyes and head, a sign so commonly present in apoplexy, is toward the side of the brain affection.† Anæsthesia and double-sided palsy are often met with, and initial convulsions are very common, and are sometimes limited to the legs. There is vomiting, as well as hyperpyrexia. Hemorrhage into the medulla is almost always immediately fatal; should the patient survive, symptoms of bulbar paralysis will be present.

In *cortical* bleedings we are apt to have localized convulsions and but slight palsy. Extravasation into the *cerebrum ovale* gives rise to similar symptoms if it occur just beneath the cortex. Hemorrhage limited to one *crus cerebri* causes paralysis of the extremities on the opposite side and of the third nerve on the same side as the lesion. Hemorrhage limited to the *arachnoid*, with the blood poured into the subarachnoid spaces, occasions ordinarily pain in the head, somnolency, and profound coma without paralysis, and without anæsthesia or slow pulse, but with relaxation of the muscles, and sometimes with convulsions; now

---

\* Gubler, *Gaz. Hebdom.*, 1858, 1859.

† Bastian, *Paralysis from Brain Disease*.



and then the symptoms assume, to all appearance, a remittent course. It is a very fatal form of apoplexy, occurring chiefly in new-born children, and after injuries to the head, or from the giving way of a diseased and widened artery, or in consequence of a rupture of one of the sinuses of the dura mater.

When the effusion of blood takes place between the dura mater and the arachnoid, it is generally the ultimate result of an inflammation and of subsequent changes of the inner surface of the dura mater; and on close inquiry the precursory symptoms of a disease of the membrane may be traced, perhaps, by the constant and localized pain, and the nocturnal restlessness. But the symptoms of the *hemorrhagic pachymeningitis* or *hæmatoma* are obscure. It happens generally after fifty years of age, in the decrepit, or in those suffering from pernicious anæmia, scurvy, emphysema, hooping-cough, alcoholism, or after head injuries. When the cyst ruptures in the thickened membrane, which it may not do for years, the signs are those of an apoplectic condition, lasting for eight or ten days.

What has been said of the symptoms pointing to the seat of lesion is exclusively based on well-attested clinical experience. The recent researches on the localization of the cerebral functions make our knowledge of the seat of the apoplexy more and more definite.

Let us now examine how the diagnosis of apoplexy can be determined, and how this condition may be distinguished from other states which produce rapid loss of consciousness, or sudden paralysis. Not to mention epilepsy,—the phenomena of which we shall farther on contrast with those of apoplexy, and shall observe to differ chiefly in the prominence of the convulsive seizures; or meningitis,—in which fever, headache, and other signs of an acute cerebral disease precede insensibility; or a tumor,—which, save in the rarest instances, leads only very gradually to a comatose condition; or sun-stroke,—exhibiting insensibility, yet also presenting points of contrast that will shortly engage our attention,—we find, excluding concussion and compression as belonging more strictly to surgical diagnosis, these morbid states liable to be mistaken for cerebral hemorrhage:

OBSTRUCTIONS OF THE CEREBRAL ARTERIES;

INSENSIBILITY FROM DRINK, OR FROM NARCOTIC POISONS;

URÆMIA ;  
DIABETIC COMA ;  
SYNCOPE ;  
ASPHYXIA ;  
ACUTE SOFTENING ;  
SUDDEN EXTENSIVE PARALYSIS ;  
PROTRACTED SLEEP ;  
CEREBRAL HYSTERIA.

*Obstructions of the Cerebral Arteries.*—Cerebral embolism or cerebral thrombosis will produce symptoms so similar to hemorrhage that in every case of apoplexy we must ask ourselves the question whether the coma be due to obstruction of the vessels or to their rupture. We may suspect that an arterial obstruction is the cause of the symptoms if the patient be young or in middle life ; if there be signs of a similar condition elsewhere, or if he be laboring under an acute or a subacute endocardial inflammation, or a chronic valvular affection in which fragments of vegetations may be broken off and washed into the vessels of the brain. We further hold this view if there be evidence of atheromatous disease of the arteries, or of syphilitic inflammation of the coats of the vessels ; if the circulation be for any reason enfeebled and the plasticity of the blood increased ; or if within a brief period several incomplete attacks have occurred before a complete comatose condition sets in. The usual locality of the impaction is in the middle cerebral artery ; and the consequences of the interrupted circulation are at once perceived in the motor area of the cortex or in the internal capsule. The interference with the circulation through the obstructed vessels gives rise to necrotic softening in the area deprived of blood. The palsy which ensues in connection with the apparently apoplectic phenomena is, with few exceptions, one-sided ; not infrequently it is limited to one member ; and the facial paralysis is on the same side with the paralysis of the limbs. Unlike what happens in cerebral hemorrhage, little, if any, fall of temperature occurs, but there are subsequently decided fever and severe headache, with greater heat on the palsied side. If the obstruction be in the left middle cerebral or its branches, which is more common than on the right side, aphasia is among the symptoms.

The hemiplegia is not of necessity attended with loss of con-

sciousness, or this is slight and of short duration; sometimes giddiness and incoherence take the place of unconsciousness; convulsions are not infrequent. The palsy is often quickly followed by gangrene of the extremities, or it is associated with disturbance of the kidneys, or with enlargement of the spleen and tenderness in the splenic region, due to changes in the organs, produced by an impaction of fibrin, if plugs be washed also into other arteries of the body. Post-paralytic phenomena, athetoid movements and the like, are more common after softening due to vascular occlusion than after cerebral hemorrhage. Occlusion of the basilar artery may give rise to bilateral palsy, often with involvement of the fifth and facial nerves, with impairment of articulation and deglutition; sometimes with ptosis, with contraction or dilatation of the pupils, with paralysis of the third and fourth nerves; and occasionally with loss of conjugate movement of the eyes and with hemianopsia. The temperature may be low at first and high subsequently.

The clinical distinction between cerebral embolism and cerebral thrombosis is even more difficult than that between vascular occlusion and cerebral hemorrhage; it may be impossible during life. Thrombosis from vascular disease is often attended with premonitory symptoms, such as headache, vertigo, numbness, tingling, weakness, impairment of memory, and psychic changes. In any case the diagnosis depends upon the recognition of the conditions that favor the occurrence of one or the other, and upon the character and distribution of the symptoms present.

Coagulation of the blood may take place in the veins and sinuses of the brain and give rise to grave symptoms. The condition may result from depressing or exhausting disease leading to weakness of the action of the heart, with slowing of the circulation and increased coagulability of the blood; or from adjacent disease, such as traumatism, caries, meningitis, tumor. In addition to such cerebral symptoms as headache, delirium, vomiting, convulsions, coma, palsy, there are present cedema of the scalp and forehead, distention of the communicating veins on the surface, and epistaxis, with paralysis of individual cranial nerves. The diagnosis also depends upon a recognition of the causative factors and the exclusion of other conditions.

*Insensibility from Drink, or from Narcotic Poisons.*—Both these

conditions are sometimes very difficult to distinguish from the coma of apoplexy ; and, if we are not cognizant of the circumstances preceding their development, we have only these points to guide us : in intoxication there is a strong smell of whiskey, gin, or whatever liquor has produced it, emanating from the mouth, and alcohol may be detected in the urine, points which would be conclusive were it not that apoplexy may come on in the drunken state ; and the man, although unconscious, is not often entirely bereft of all power of motion,—he is certainly not paralyzed. Moreover, the pulse is not slow, it is frequent ; the pupils are generally dilated ; the eye is injected, shows no lateral deviation ; there is often violent struggling, and the symptoms become suddenly much ameliorated after the inhalation of ammonia, or after the stomach has been emptied of its contents. In narcotic poisoning, especially if from opium, the pupils are very much contracted, and we are likely to encounter repeated vomiting, and a gradual intensification of the coma. The patient, however, unless death be close at hand, can be momentarily roused from his deep sleep ; and his calm, slow breathing is unlike the stertor of apoplexy. But when the hemorrhage has taken place into the pons Varolii, the diagnosis is very difficult, especially if the bleeding be extensive, for then we are apt to have a contraction of both pupils, and the respiration may not be stertorous ; nor is there always at first paralysis. Yet this subsequently appears, and thus the detection of the cause of the insensibility is rendered easier.\* A symptom of great diagnostic significance, too, is the occurrence of convulsions. Still, as Russell Reynolds shows, this may happen in opium poisoning, and is not very rare in children.

Nitrobenzole, which operates as a narcotic poison in vapor as well as in a liquid state, may, in rapidly fatal cases, produce coma, which may be mistaken for the insensibility of apoplexy. But the poison is detected by its strong odor, resembling that of bitter almonds.† Poisoning by drinking chloroform gives rise to many of the symptoms of apoplexy ; it is discerned by the odor of the breath, by the quick and tumultuous heart action that

---

\* See an interesting case mentioned by Hughlings Jackson in *London Hospital Reports*, vol. i., 1864.

† Taylor, *Guy's Hospital Reports*, vol. x, 3d Series.



accompanies the stertorous breathing, by the relaxation of the limbs, by the deathlike aspect of the face, by the widely-dilated pupils, and by the complete general anæsthesia.\* Chloral insensibility is often preceded by vertigo and pains in the legs and arms, and is attended with flushing of the face, injected conjunctiva, a weak intermittent heart; the pulse may, however, be slow and full. Hydrocyanic acid poisoning produces profound insensibility, often attended by convulsions, and by peculiar breathing,—short inspiration with labored, prolonged expiration. The breath has the characteristic odor of the acid.

*Uræmia.*—The strong point in the diagnosis is that the coma is preceded by convulsions; exceptional instances are few indeed. An examination of the urine conduces, of course, to certainty; but, for obvious reasons, it cannot always aid us at once. Moreover, albumen—not, however, in large amounts—may occur in the urine after an apoplectic stroke, and after convulsions not uræmic. Puffy eyelids and swollen ankles, coma not profound, peculiar stertor seeming to emanate from the mouth, pupils normal or dilated but equal, low body heat, are symptoms that belong to uræmic coma. Yet there are many cases of uræmic coma with high temperature, especially when tested in the rectum.† Unilateral convulsions or loss of power are indicative of cerebral mischief and tell against uræmia.

*Diabetic Coma.*—We meet at times with comatose symptoms in diabetes which simulate those of apoplexy, and which are apt to be connected with the so called “acetonuria,” or the poisoning that takes place in the organism from the formation of acetone. Further on we shall inquire into the tests for this substance. But it may here be stated that the diabetic coma generally begins not abruptly, but with somnolency which passes into coma; that it is often preceded by great oppression, and is attended with a rapid weak pulse, but not with hemiplegia or other local palsies.

*Syncope—Asphyxia.*—The loss of consciousness in either of these states is as striking as in apoplexy. But there is this decided difference: the suspension of thought and of volition in a fainting-fit is due to failure of the circulation: hence the pulse is

---

\* As in the case reported in L'Union Médicale, October, 1864.

† McBride, American Journal of Neurology, etc., 1883.

hardly or not at all felt, instead of being full, as it is in apoplexy. Further, the pallor of the face, the quiet or sighing respiration, the well-preserved reflexes, and the short duration of the syncope mark plainly the one affection from the other. And with reference to asphyxia, the turgid and livid face, the bluish lip, the distressed and embarrassed breathing preceding the convulsions, and the loss of consciousness, show clearly that the disturbance affects primarily the lungs, and not the brain.

*Acute Softening.*—This state is so closely connected with cerebral embolism or thrombosis that an appreciation of the history of the case, and the causes that lead to occlusion of the vessels, tells us the meaning of the cerebral symptoms. Rapid softening, too, at times happens around a clot. In acute softening the mental phenomena are always obvious, the mind is much more obtuse or impaired than it is after the shock of cerebral hemorrhage is over. Durand Fardel \* regards as a significant sign of acute softening an increased secretion from the mouth and eye.

*Sudden Extensive Paralysis without Coma.*—This is not a trait of apoplexy, but rather of occlusion of the large vessels. Sudden extensive paralysis without coma is ordinarily owing to softening of the brain, most apt to have followed this occlusion; but it may be due to hemorrhage into the spinal column. Palsy from this source, unlike that caused by cerebral hemorrhage, is almost invariably double-sided, is accompanied by severe spinal pain, and, if the extravasation have taken place into the spinal meninges, by tonic spasms, like those of tetanus.

*Protracted Sleep.*—While recovering from acute diseases, the sick often sleep profoundly and for a long time. Yet there is little likelihood of confounding this with the sleep of apoplexy; for the antecedent circumstances reveal the meaning of this restoration of nature. Sometimes, however, persons sink into a deep and prolonged slumber without any previous ailment. Medical literature furnishes a number of such instances. In one recorded by Cousins,† the tendency to somnolency lasted for years. The patient frequently slept three, and sometimes five, days at a time.

---

\* *Maladies des Vieillards*.

† *Medical Times and Gazette*, April, 1863. See also a somewhat similar case, *New York Medical Journal*, Dec. 1867.

When he awoke he was well. In a case which I saw with Dr. Weir Mitchell,\* the slumberer was aroused out of her trance several times by the exciting influence of electricity; but this finally lost its effect, and she relapsed into a sleep from which she awoke no more. These cases may give the impression of apoplexy, yet they do not resemble it strictly. They are unlike it in the gentle, noiseless breathing; in the feeble pulse; in the occasional motion of the body; and in the protracted unconsciousness. Then generally the patient can be roused sufficiently to take food. Prolonged somnolence is also among the marked symptoms of cerebral syphilis.† In some instances the disorder shows itself in a constant tendency to fall asleep for brief periods at a time. One patient I had slept repeatedly during the day about five minutes at a time on her feet. She could be awakened by strong efforts. The comparatively short duration of the spells of sleep, and the absence of evidences of hysteria usual in trances, distinguish these cases of *narcolepsy* from trance. The *narcolepsy* may be, however, associated with catalepsy, and there are cases of undoubtedly hysterical origin. Such a case probably was the one of the lethargic Irish fasting girl, who is said to have existed for nearly six weeks without food; the disorder was ushered in by hysterical fits.‡ The recurrence of the sleepy fits, their innocuousness, and the absence of progressive emaciation and of enlargement of the cervical glands distinguish *narcolepsy* from the dangerous sleeping sickness affecting negroes on the west coast of Africa.

*Cerebral Hysteria*.—The actual similitude and the points of contrast between this curious state and apoplexy may be learned from the following sketch:

A married lady, of a remarkably impressionable and nervous disposition, had been for many months suffering from amenorrhœa and from sluggish action of the bowels. She had also a constant cough, dependent upon tubercles in one of the lungs. She had been in very bad health, but by the steady employment

---

\* Described by him, Transactions of College of Physicians of Philadelphia, 1856.

† See cases in Lecture XVI., Buzzard on Diseases of the Nervous System, 1882.

‡ Lancet, July 15, 1893. Another sleeping girl is mentioned, *ibid.*, July 29, 1893.

of tonics, and the beneficial effects of a sea-voyage, her symptoms were much amended. She began to gain flesh, and to take exercise without fatigue. She was, however, troubled with headache, and with pain at the lower part of the abdomen. On one occasion in the evening I ordered her some cathartic medicine; and in the morning she was better than usual, and in the liveliest spirits. A few hours afterward, I was sent for, and found her insensible. She had complained of a sudden, sharp cramp near the umbilicus, and had then ceased to speak. She remained unconscious for about twelve hours; yet not wholly so, for every now and then she opened her eyelids, muttered a word or two, a pleasant smile flitted over her countenance, but she soon relapsed into her deep slumber. Her thumbs were drawn inward; she had occasional convulsive movements; the breathing was rapid, but not noisy; the pulse feeble,—at first slow, then frequent; her eyes squinted in the most decided manner. Stimulants and antispasmodics were freely given, but without much benefit, for she recovered from her lethargy only with the setting in of the most violent paroxysmal pains in the abdomen, shooting down the thigh, and accompanied by contractions of the muscles and by exquisite local tenderness. The next day, without much abatement of the suffering, she was perfectly conscious; but still she had an internal squint,—nay, was totally blind, and remained so for two days. During this time a menstrual discharge began, which in part relieved the abdominal pain. It is needless to point out how this display of hysteria differed from apoplexy.

**APHASIA.**—The faculty of speech may be interfered with by various lesions of the brain and of the pons and the medulla. Of these, some cause only disturbance of articulation, while others derange the higher speech processes. From the first result difficult or defective articulation; from the second, the group of phenomena included in the designation *aphasia*. Though this is really a mere symptom, it is so prominent as seemingly to constitute the disorder. By aphasia is meant loss of the faculty of expression of thought, in consequence of loss either of the faculty of speech, or of that of communicating thought by writing or by gestures. The patient may be deprived of the ability of expressing himself in one of these ways, or in all. The loss of speech is the most common, and is apt to be associated with a



very decided impairment of memory and an enfeeblement of intelligence. The disorder may be temporary, lasting but a few hours or some days, or it may continue for months or years. During its course the affected person is incapable of recalling words to give utterance to his ideas ; or if he can recall the words to the mind, and thus think, he cannot express them.

Very often the patient has but a few words at his control ; he says "yes" or "no" for everything, and appears angry that he can say no more ; or he uses wrong words, knowing perhaps that they are wrong, and sometimes only those of a profane kind ; or he confuses merely some syllables in the words he employs ; or he may not be able to utter a word, using altogether unintelligible expressions. Yet, while in this condition, there is no defect in the tongue, or lips, or palate, to account for the inability to talk ; they are as healthy as usual ; the act of swallowing is easily performed ; and even where the aphasia is complicated with hemiplegia, it is not difficult to discern that the imperfect articulation and thick speech attending the palsy—which, moreover, are apt to pass off within a short period after the seizure—are not the cause of the singular disturbance of expression ; a disturbance which will mostly show itself not simply by the failure to utter words, but also by the inability to recollect them and write them down. Indeed, it is necessary to bear in mind that, while these states may coexist, they also may be present separately. Thus, there are persons who can think, but cannot speak or write ; there are those who can think and write, but cannot speak ; and there are those who can think and speak, but cannot write or read. For the second group the term "aphemia" has been proposed ; for the third, the term "agraphia ;" for the fourth, "alexia." Most patients understand perfectly well what is said to them ; some can read to themselves ; and, unless the general intelligence be perceptibly affected, they can express themselves by signs and gestures. In some cases there is rather loss of memory, and forgetfulness and confusion, and perhaps a consequent use of wrong words ; but when prompted the word is at once spoken. Where the power of expression only is lost, but the words are still suggested by the memory, the term "motor aphasia" is used. Where the memory of words is altogether lost, it is customary to speak of the affection as "amnesic" or "sensory aphasia." Again, there are cases

in which words and ideas remain, but in which the power of forming correct sentences is greatly impaired or is lost. This has been named "akataphasia." \*

Slips of the tongue are by no means always to be regarded as aphasia, for very often these have a local cause, such as a sore tongue or lip, or a sharp tooth fretting the tongue, producing unusual sensations in the mouth.†

Aphasia is dependent upon disease situated in the frontal convolutions, and by Broca the lesion was correctly located in the seat of articulate language, in the posterior part of the third frontal convolution of the left side of the cerebrum. This explains why the hemiplegia which may accompany aphasia is almost invariably right-sided. But it may be left-sided, if the corresponding parts of the right hemisphere have become the main centre of speech, as happens not infrequently in left-handed persons. It has been further shown that the disturbance will be in the cortical substance of the speech-centre or in the conducting fibres, according to the form of aphasia. Where the memory of words is gone, it is in the former.

The function of speech is subserved through sensory and motor processes. The former have to do with the reception of impressions from without, principally through hearing and sight, and the centres for these are in right-handed persons respectively in the first temporal convolution and in the angular gyrus on the left side; the motor speech processes originate in Broca's convolution. In disease destroying the latter and causing motor aphasia, the patient understands, but can speak but few words or syllables; in disease affecting the sensory centres, the auditory and visual appreciation of words is impaired or lost. When the first left temporal convolution is the seat of lesion, "auditory aphasia," or "word-deafness," with loss of hearing in the opposite ear, results, and words are wrongly used, and speech fails to convey any ideas. The words are heard merely as sounds. The patient fails to recognize his mistakes in speech, and is unable to correctly repeat words spoken to him. Disease of the angular gyrus on the left side gives rise to "visual aphasia," or "word-blindness,"

---

\* Steinthal; also Kussmaul, in Ziemssen's Cyclopædia.

† Ord, St. Thomas's Hospital Reports, vol. iv.

or even to "mind-blindness," the patient being unable to recognize words or objects through vision. There thus result alexia and perhaps agraphia, together with loss of vision in the right half of the visual fields.

As regards the exact lesion, it is very various. Aphasia may be due to functional as well as to organic disease. In cases of aphasia of short duration and without palsy, there is probably merely congestion; in protracted cases, and those in which we find persisting hemiplegia, a large clot, or softening, or abscess, is likely to be present; embolism of the middle cerebral artery on the left side is prone to be the cause in cases that are associated with valvular disease of the heart and that have come on suddenly. Thrombosis from enfeebled nutrition will explain the aphasia which may be noticed during convalescence from grave acute maladies. That consequent upon congestions ends in more or less rapid recovery; in the other forms, usually, either no improvement follows, or only a very partial gain of words takes place. Occasionally we meet with aphasia in hysteria or in epilepsy, in acute infectious diseases and in uræmia, or we encounter aphasia intimately connected with a syphilitic cachexia,\* and dependent most probably upon disease of the arteries.

Transitory aphasia has been observed in the course of pneumonia. The complication usually appears toward the second or third day of the disease, being ordinarily preceded by headache and vertigo, and sometimes by numbness and tingling on the right side of the body. It may set in abruptly, without loss of consciousness, or be preceded by an apoplectiform seizure. There may be, in addition, transient palsy of the right side of the body. The manifestation is thought to be due to the action of the toxic products of the disease process.† Aphasia may become manifest subsequent to attacks of vertigo, or to a paralytic stroke preceded or not by the ordinary signs of an apoplectic fit. Under these circumstances the diagnosis cannot be definitely made until consciousness has returned, and we have an opportunity of examining the state of the mind, and of the tongue, and of the muscles concerned in articulation, remembering that if there be merely difficulty in articulation the case is not one of aphasia.

---

\* See Clin. Soc. Trans., vol. iii., and Arch. Gén. de Méd., Feb. 1871.

† Chantemesse, *Semaine Médicale*, 1893, No. 73, p. 582.

**Sun-stroke.**—Persons exposed to the scorching rays of the sun in midsummer often become dizzy, and fall to the ground insensible: they have had a sun-stroke. The attack either takes place while the patient is still exposed to the sun, or, in rarer instances, he reaches his home with a staggering gait and a suffused face, giddy, faint, suffering from a dull, oppressive pain in the head, having a constant desire to micturate, and after some hours becomes unconscious. However the onset, the insensibility which occurs is generally complete, although it may be so but for a few minutes. Associated with it are a frequent pulse, a skin harsh and warm and sometimes very hot on the forehead, shallow, noisy breathing, difficulty in swallowing, contracted or, more generally, dilated pupils, and relaxation of the limbs. Scanty urine, delirium, and convulsions, which may or may not depend on uræmia, are not uncommon.

When we contrast these symptoms with those of apoplexy, we find the following marks of distinction: the pulse is not slow and full, but frequent and often feeble; there is more difficulty in deglutition, but a less snoring respiration; the coma does not ordinarily remain as complete for so great a length of time, for soon the patient may, temporarily at least, be partially roused from his deep sleep; and no hemiplegia, no paralysis, either of the limbs or of the cheek, occurs. The temperature of the body is usually very high,  $104^{\circ}$  to  $109^{\circ}$ , and not below the normal, as it is at first in apoplexy. The after-symptoms, too, are different: in cerebral hemorrhage, paralysis; in sun-stroke, feebleness of movement, but no paralysis. In the former, no marked, persistent headache; in the latter, headache, more or less chronic, always aggravated by walking in the sun, and often for months accompanied by signs of an exhausted nervous system, and in some instances by epileptic convulsions.

The question with regard to the discrimination of these morbid states is one of great practical value, as on the conclusion arrived at depends our therapeutic action; and generally it is readily determined by paying attention to the variance in the symptoms mentioned. But it must be confessed that we sometimes meet with ambiguous cases,—cases in which the signs of nervous exhaustion produced by exposure to heat are blended with those of cerebral congestion or hemorrhage excited by the same cause, and



in which, when they terminate fatally, the autopsy shows not simply a changed blood, or pulmonary congestion, but turgescence of the cerebral vessels, or an extravasation. It may also be difficult to distinguish between sun-stroke and acute alcoholism, particularly because those who drink freely are very prone to the disease. The chief distinguishing trait is in the high temperature of sun-stroke, and the normal or lowered temperature of alcoholism.

The remarks just made refer to the most common form of sun-stroke,—that attended with more or less sudden loss of consciousness. But there are cases in which the abnormal manifestations come on gradually, and in which the patient at no time becomes insensible. I have seen a number of the kind: they were not unusual among officers sent home from the wearing summer campaigns of our civil war. The chief symptoms are intense headache, nausea, prostration, and inability to perform any work requiring sustained attention. All these signs appear after protracted exposure to the sun; and they mend but tardily. In truth, in the slowly-developed disorder the subsequent nervous exhaustion and the paroxysms of headache are often much more persistent than are the same phenomena when they follow what seems to be the more violent form of the malady. Among the sequelæ of these apparently incomplete attacks are irritability of the bladder, incontinence of urine, and irregular action of the heart. But nothing is as striking as the loss of mental and bodily energy.

The symptoms of “insolatio,” or sun-stroke, may be induced by prolonged atmospheric heat while the patient is in doors and not exposed to the rays of the sun. Such cases of heat-stroke are known to occur in India even at midnight. They may be preceded by a sense of extreme weariness, by inability to sleep, by loss of appetite, by constipation and frequent micturition, and by deficient perspiration; or the signs of exhaustion, followed by more or less complete insensibility, appear without distinct prodromes. Cases of the kind under consideration may or may not show an increased or high temperature; generally they do.

Then, again, we find cases of *heat exhaustion*, often seen in our hot summers, in which there is from the first great tendency to syncope; the skin is pale, cool, and moist, the temperature not increased, the pulse very feeble, the pupils dilated, and stimulants freely given rapidly relieve the urgent symptoms.

The nature of heat exhaustion, as of sun-stroke, is obscure. It is held by H. C. Wood\* to be a fever which is dependent upon heat. Certain it is that the heat centres are very much disturbed in the affection. In occasional instances meningitis rather than sun-stroke follows exposure to the sun, and we find the ordinary symptoms of meningeal inflammation.

**Catalepsy.**—This, as a rule, is a sudden suspension of thought, of sensibility, and of voluntary motion, during the continuance of which the muscles become rigid, although they retain the exact position they happen to be placed in. The uncommon complaint occurs in paroxysms, which may last but a few minutes or for several hours, and during which the most complete anæsthesia, not only of the skin, but also of the deeper tissues, may occur.† Often consciousness is lost, but it may be only obscured. Respiration is disordered, the circulation is feeble, reflex action is abolished, and the temperature is lowered. The disorder is met with mainly in females, especially in hysterical females, and may alternate with outbreaks of hysteria. But it may also exist in the male sex, and be in either hereditary. It has even been noticed as an epidemic in localities where there are many families closely connected by intermarriage.‡ Nervous exhaustion or sudden alarm predisposes to the seizures, which at times recur periodically and last from a few minutes to a few hours.

Catalepsy may be mistaken for apoplexy, or even for death. It differs from apoplexy by its constant recurrence; and, further, during an attack the eyes are wide open, the pupils, although dilated, are very susceptible to light, and there is an absence of stertorous breathing as well as of the characteristic relaxation of the muscles or of the paralysis of apoplexy,—for the limbs are outstretched, or held in every conceivable annoying or painful position; yet as soon as consciousness is restored, their movement fully returns. The pulse is not retarded; on the contrary, although feeble, it becomes very frequent.

The perplexing affection varies from a kindred state, *cestasy*, in

\* Thermic Fever, or Sunstroke.

† As in the case reported by Lasègue, Archives Générales de Médecine, tome i., 1864.

‡ Vogt, Schmidt's Jahrbücher, Bd. cxx. p. 301.

this: in the latter the loss of consciousness is not complete; the patient is merely insensible to external objects, because he is intensely absorbed in some vision present to his imagination, or in the contemplation of some subject to him of all-engrossing interest. But he is not statue-like; on the contrary, his countenance is animated and earnest, and he talks, declaims, sings.

There is a curious form of the disorder, which Sir Thomas Watson describes. It is an imperfect kind of catalepsy, called *daymare*, the affected person being incapable of moving or speaking, yet cognizant of all that goes on. These seizures of temporary deprivation of muscular power, without unconsciousness, are thought to depend upon a diseased state of the blood-vessels of the brain.

*Feigned catalepsy* may be distinguished from the true disease by the muscles quickly showing signs of fatigue, which they do not in real catalepsy. A pressure-drum, Charcot\* found, fixed at the extremity of the outstretched limb in a person who feigns, will in a few minutes, in place of the straight, regular line, show crooked, very undulating traces, and the same irregularity is seen in the tracings of the pneumograph applied to the chest.

Catalepsy may be artificially induced, as we know from the interesting experiments on hypnotism which of late years have been made. Catalepsies of particular groups of muscles, or partial catalepsies, can also be artificially excited.

In the rare condition known as *trance*, or *lethargy*, there exists a state resembling sleep, from which the person can be roused with difficulty, if at all. It is principally associated with hysteria, although it has been observed as a result of excessive mental application and after exhausting disease. The patient is usually pallid. The extremities are, as a rule, relaxed, although they may be rigid for a time, and there may even be convulsions. The eyelids are closed and the eyes turned upward and to one side; the pupils vary in size, but react to light. Reflex action is usually lowered. Respiration and circulation are greatly enfeebled; the peripheral temperature is subnormal. An attack lasts from a few hours to weeks. Cases in which it occurred

---

\* Third volume of Clinical Lectures, 1889.

without any obvious previous ailment have been mentioned while discussing protracted sleep. In *narcolepsy* there occur sudden short periods of day sleep, from which, however, the individual can be roused. These may recur spontaneously, or be induced by peripheral impressions; at times we find the condition marked in diabetic or gouty patients.

The affection described as *African lethargy*, or *sleeping sickness*, attacks exclusively negroes, principally on the west coast of Africa, and is characterized by somnolence of progressive degree, usually leading to great emaciation and to a fatal termination. The disease is associated with enlargement of the cervical glands and of the spleen, and is thought to be due to the presence in the blood of the *Filaria sanguinis*.

### Diseases marked by Convulsions or Spasms.

**Epilepsy.**—Epilepsy is a disease the chief manifestation of which consists in recurring attacks of sudden loss of consciousness, attended with convulsive movements. The patient falls to the ground, without thought, without feeling, without the power of voluntary motion. He utters often a short piercing cry, then a fearful struggle begins. The legs are stiff, and turned inward; the head is tossed backward, or from side to side; the mouth is distorted, the lips are covered with foam; the arms are outstretched and rigid, or thrown about with great force; the eyelids are half closed; the teeth are ground together, and the tongue is thrust between them, and often severely bitten. Gradually the convulsive movements become less violent and cease altogether, and the patient passes into a deep sleep, from which he awakes fatigued and exhausted, and dull in intellect. But these symptoms disappear, and he returns to his usual state of health.

Yet every paroxysm does not present the same phenomena, or run the same definite course. In many the attack is preceded by strange sensations: by a peculiar train of thought; by retching; by the feeling of a puff of air ascending from the extremities to the head. This “aura epileptica,” on which so much stress has been laid, is, however, far from constant. Moreover, it may exist and hardly be perceived: it may be an unfelt irritation starting from some peripheral nerve in any part of the skin, or from some organ not deeply seated, as the testicle, and its point of departure



may be detected by observing, during the fit, in what neighborhood the first, or the most violent, or the most prolonged contractions occur. In very rare instances sudden spasms of the face and chest occur, with arrest of respiration, and with a subsequent clonic convulsion, yet with so little unconsciousness that it remains doubtful whether it has happened at all.

Some seizures are very light,—a transient suspension of consciousness, a slight twitching of some of the muscles, a fixed gaze, perhaps a decided impression of vertigo, and all is over. These abortive fits, the *petit mal*, or minor attacks, are very apt to precede by some days a severe attack, or several of them may take the place of the more turbulent form of the disorder. And they, like the graver epileptic convulsion, may present strange irregularities. They may manifest themselves, for instance, only in bursts of unmeaning laughter;\* or intellectual derangement replaces the ordinary convulsive attack;† or there is mental wandering, with disposition to commit acts of violence. The attacks of epilepsy which are chiefly characterized by vertigo are distinguished from all other forms of vertigo by the loss of consciousness they also mostly present, and the absence of any giddiness in the intervals. In nocturnal epilepsy ecchymoses on the face, conjunctival extravasations, a severe headache on awakening, and a sore tongue, may indicate what has happened in the night.

The epileptic paroxysm does not always pass off without leaving some trace of the profound disturbance it has occasioned. It may be followed by hemiplegia. Whether this be due, as Hughlings Jackson‡ asserts, to exhaustion of the nerve-centres following the excessive discharge of the nerve force bringing about the convulsion, it is certain that the palsy is very transient. Another sequel of the attack is aphasia; another, loss of voice; another, abdominal tenderness. As regards palsy, however, we must remember that epileptic fits may follow hemiplegia due to a vascular lesion, so-called *post-hemiplegic epilepsy*.

In the intervals between the seizures the patient is not in reality well. His temper is irritable, and his mental faculties slowly but

---

\* George Paget, British Medical Journal, Feb. 1859.

† Thorne on Masked Epilepsy, St. Bartholomew's Hosp. Rep., vol. vi.

‡ After-Effects of Epileptic Discharges, West Riding Reports, 1876.

certainly deteriorate. The loss of memory, particularly, is very marked; and dementia is not an unusual complication of long-continued epilepsy. In some epileptics there is much excitement or a curious mental state preceding the seizures, or a violent and dangerous mania may follow them. Again, as I have noted in common with several observers, a temporary albuminuria is not unfrequently met with at the termination of the paroxysm.

True epilepsy is probably owing to functional or nutritional changes in the cortex of the brain, giving rise to excessive activity of nerve-cells leading to periodic discharges of nerve-force. Its most potent cause certainly is hereditary predisposition. Convulsions due to reflex irritation, to organic brain-disease, and to toxic blood-states may result in true epilepsy. It is thus that the malady originates in injuries of nerves, in diseases of the skin, of the stomach and intestines, and of the uterus, in the irritation of worms, or in consequence of congenital phimosis,\* or of chronic nasal catarrh.† Now, with reference to both the prognosis and the treatment, it is very important to discriminate between true epilepsy and convulsions of eccentric origin; and to arrive at a conclusion is possible only by a thorough examination of all the constitutional symptoms, and by ascertaining the starting-point and tracing the course of the aura. The cases in which the aura is interrupted and the paroxysm arrested by a ligature are well known. Nothnagel cites an instance in which the aura began with peculiar sensations in the stomach, and the attack was stopped by swallowing table-salt. Convulsions may further be symptomatic of a cerebral disorder,—such as a tumor, cysticerci lodged in the organ, a syphilitic affection of the membranes, or a disturbance of the brain produced by disease of the skull-cap,—in fact, of any disease-process affecting the cortex of the brain; or it may be due to watery blood, or vitiated blood full of abnormal ingredients, as in diseases of the kidneys, acting injuriously on the nutrition of the cerebral texture. During the paroxysm it is impossible to determine the character of the convulsions; but in the interval we may often do so by close attention to the history of the case, and by noting whether the patient enjoys the usual

---

\* Althaus, *Lancet*, Feb. 1867.

† Cases collected by Salinger, *Polyclinic*, June, 1887.

health of epileptic subjects, or presents signs of a chronic cerebral disorder, especially steady headache, palsies of cranial nerves, optic neuritis, vomiting. Romberg tells us that where affections of the bones of the head lie at the root of the complaint, the fits are readily induced by pressure upon the skull; and, further, that if there be disease residing in one of the cerebral hemispheres, the aura affects the opposite side of the body, and is generally confined to the upper extremity. Convulsions are often found in connection with ear disease, and especially with purulent otitis.\* In those who inherit syphilis idiopathic epilepsy may happen.

Limited convulsive seizures are connected with disease of special convolutions; and, as Hughlings Jackson † has shown, if we have a convulsion which is limited, or at least begins always in the same limited manner, either a tonic or a clonic spasm of a group of muscles, we may from this monospasm diagnose an irritative lesion in the motor centre presiding over the disturbed part, though in the hemisphere opposite to the spasm. The irritative lesion is most frequently a meningo-encephalitis; the centre involved becomes highly charged, a discharge takes place with the convulsion, and a temporary paralysis in the affected group of muscles results. The spasm most frequently originates in the hand, but we may also find the spasm limited to a group of muscles in the face, or in the leg. At first there is no loss of consciousness during the seizures, but as the spasms spread and become unilateral, consciousness is lost. Convulsions due to syphilitic disease are, for the most part, of the kind just described, and are the chief form of the cortical, or so-called *Jacksonian epilepsy*. In *masked epilepsy* there is often an epileptic vertigo, with loss of consciousness and with twitching of some muscles, but the patient does not fall.

Much has been said of the distinction between epilepsy and *convulsions*. Now, as regards the seizure itself, there is no appreciable difference: the only diversity consists in the recurrence of the attack after intervals of comparative health, and in the non-existence of any disturbance from which convulsions are likely to arise, such as reflex irritation, organic brain-disease, or a toxic blood-state. In children, who, as is well known, are particularly

---

\* Ormerod, *Brain*, April, 1883.

† *Medical Times and Gazette*, 1875.

subject to convulsions, the diagnosis may be a difficult matter; but the fits of epilepsy are distinguishable by the dulness of intellect, and the slow mental and bodily development, observable in the intervals. And we are not often called upon to make this differential diagnosis, because of the extreme rarity with which epilepsy occurs in the young; although many insist that it is more frequent than is supposed, basing this assumption on the generally-received fact that the history of epileptics shows them to have suffered greatly from convulsions during childhood.

The diseases which are most apt to be confounded with epilepsy are *hysteria* and *apoplexy*. The former—like all the rest of the group now under discussion, like chorea, like tetanus, like hydrophobia—is discriminated by the absence of that perfect suspension of consciousness that takes place in epileptic seizures; and there are other marks of distinction, to which we shall presently refer. In apoplexy, as in epilepsy, we meet with loss of consciousness, sometimes with convulsions. But these are, on the whole, rare, and coma precedes and does not follow them, as happens in epilepsy. Then, stertorous breathing and a slow, full pulse are not observed in epilepsy; for the breathing, although irregular and gasping, is not coarse and noisy, and the pulse is feeble, irregular, and frequent. Epileptic patients bite their tongues; this does not occur in apoplexy. In epilepsy the paroxysm seldom lasts longer than from ten to fifteen minutes before consciousness returns and before the convulsions cease; in apoplexy the insensibility is of much longer duration. Epilepsy is not usually followed by paralysis; apoplexy is commonly.

There is sometimes a close resemblance between syncope and abortive epilepsy, *petit mal*. But they occur under widely different conditions; and the loss as well as the return of consciousness is less abrupt in the one than in the other.

Epilepsy is at times *feigned*; yet impostors cannot feign it completely. They may bite their tongue; they may imitate the stertor, the foam at the mouth, the convulsions, the thumb drawn inward toward the palm, the confused air on awakening; they may simulate, although they rarely do so, the indifference to pain; yet there is one feature of the real attack they cannot copy,—the insensibility of the iris. No matter how skilful the dissembler, his pupils must contract when exposed to a strong light, they must



dilate when the stimulus is withdrawn. Unfortunately, there are several difficulties in making this test an absolute one. In the first place, the pupils, during a fit, cannot always be readily observed. In the second place, not in every case of epilepsy are they perfectly immovable; in some, though sluggish, they react to light. Again, as proved by Keen, violent muscular motion instantly dilates the pupil, and so long as the movement continues, so long will the iris act sluggishly, even when exposed to a bright light. Thus, muscular spasms alone, even when simulated, may cause the pupils to be dilated and inactive. A test said to be more generally useful is the administration of ether. When given to an epileptic, its first effect is to increase the violence of the spasm, but eventually the patient passes into the deep sleep produced by ether, without any of the prior cerebral excitement; while in the malingerer this manifests itself by talking and laughing,—in fact, in the usual way.\*

**Chorea.**—This spasmodic affection is chiefly met with in young persons, especially in girls approaching the age of puberty. It is characterized by irregular clonic spasms of groups of muscles under the influence of the will, and mainly of those on one side of the body, together with muscular incoördination. But the patient is not deprived of consciousness and of all power of voluntary motion. He knows what he is about, and can in part execute the movements he undertakes; yet his limbs are not completely under his control. They obey only his general directions, but not entirely, or at once; for the muscles jerk and pull as seem to them best, taking no heed of the time or the manner in which the will wishes any movement executed. In some cases the muscles of deglutition, of respiration, and of articulation become implicated; and difficulty in swallowing, in breathing, and in speaking occurs. A dilated pupil, too, acting sluggishly in response to light, may be met with among the phenomena of the malady. Often there are mental irritability and dulness of intellect. The urine contains urea and phosphates in excess.

Chorea is essentially a functional disorder of the nervous centres, especially of the cells of the cerebral cortex. In a large number of persons the malady is called into existence by an

---

\* Keen, Mitchell, and Morehouse, Amer. Journ. Med. Sci., Oct. 1864.

irritation of peripheral portions of the nervous system. Thus, a blow, a wound of a nerve, disorders of the uterus, painful menstruation, pregnancy, eye-strain, or gastric or intestinal affections may act as the exciting cause of the perverted muscular movements. In cases due to organic causes, endocarditis or plugging of the vessels leading to the corpus striatum is a common lesion, one-sided embolism giving rise to one-sided chorea.\* The association with vegetations on the valves is in fatal cases certainly very frequent.† It has, indeed, been suggested that the wild, maniacal delirium, with subsequent rapid emaciation, which we meet with in some instances of chorea, has its origin in embolism.‡ But all cases of chorea cannot be explained by minute cerebral embolism, as has been attempted.

Chorea may be produced by strong mental emotion, especially by fright. It may follow scarlet fever, but it is more often the sequence of rheumatic fever or arises from the same diathesis that attends or occasions rheumatism. Yet this is not, as some have alleged, its only cause; for in a number of persons affected with chorea we fail to detect any proof of a rheumatic diathesis. Still, the greatest prevalence of chorea in the spring of the year coincides, as Morris J. Lewis§ has shown, with the greatest prevalence of rheumatism. The action of special toxic substances on the brain is held by several to be the ultimate cause. As regards the cardiac complication, the presence of which is chiefly deduced from the existence of a murmur, the inference drawn from this sign, of its being due to organic disease, is not always accurate; for the murmur is often owing to anæmia, or dependent upon spasmodic action of the papillary muscles,—the same spasmodic action that is seen in the striated muscles of the face and of the extremities.

The disease is rarely fatal: but its duration is very variable; for, although it may be acute, lasting for six weeks or more, it may last for months, even for years, and relapses are frequent. There are in chronic cases no attending cerebral symptoms, yet the mental faculties are not in a perfectly healthy state. The

---

\* Hughlings Jackson, London Hospital Reports, vol. ii., and Edinburgh Medical Journal, Oct. 1868.

† Ogle, British and Foreign Medico-Chirurgical Review, 1868.

‡ Tuckwell, *ibid.*, Oct. 1867.

§ American Journal of the Medical Sciences, Sept. 1892, p. 251.

intellect of a choreic child develops slowly, and is evidently enfeebled while the disorder lasts. In some cases paralysis supervenes; but it is not permanent, nor, indeed, of long duration. But those who have been choreic remain subject to nervous disorders; and I have known several instances in which the complaint has been, in after-years, followed by epilepsy.

A chronic progressive form of chorea sometimes develops late in life, with, in many instances, a history of hereditary transmission. The movements in this *Huntington's chorea* usually appear first in the face and upper extremities, and gradually extend, and there is disturbance of speech. In the hereditary cases mental changes are common and sometimes pronounced, and gradual dementia is observed.

In some parts of Italy there has been noticed an affection characterized by sudden, shock-like muscular contractions, with progressive palsy and wasting, and known as *electrical chorea*. Of its nature and cause we have no definite knowledge.

So-called *hysterical chorea* consists in general spasmodic movements occurring in hysterical subjects, but the movements are far more regular and rhythmical than those of true chorea, and are usually deliberate and of wide range.

The diagnosis of chorea is generally easy. The peculiar habit some children or even older persons get into of winking, or jerking the head, or of making other strange movements, the "habit-chorea" or "habit-spasm," as it has been called, is distinguished by its gradual development, and its limitation to a single muscle, or group of muscles or of associated muscles. Chorea with loss of power on one side, "paralytic chorea," when confined to the arm is in children recognized by the occasional choreic movements, and by the loss of power which happens gradually.

Chorea from eye-strain is, as a ready test, discriminated by using atropine. Dr. Hansell employed this in many cases at my clinic with quick results. Atropine paralyzes the ciliary muscle; no effort of accommodation can then be made; therefore muscular twitching, as well as headache or other functional disturbances from disordered accommodation, must cease after an interval of time long enough to break up the habit; chorea from constitutional causes will, of course, be unaffected by atropine or other paralysis of the ciliary muscle.

Chorea differs from the *spasms of acute cerebral disease* by the absence of fever, of delirium, and of coma, though we must bear in mind that we sometimes have elevation of temperature and mania in the chorea of pregnancy ; from *epilepsy*, by its being continuous, by the non-existence of unconsciousness, and by the rarity with which the muscles jerk at a time when epileptic convulsions are frequent,—at night ; from *tetanus* it is chiefly distinguished by not exhibiting tonic spasm. *Paralysis agitans* is, like chorea, attended with disturbed muscular movements. But we find weakness of the muscles and persistent tremor rather than spasmodic contraction and want of control over muscular motion. Then the history of the case, and the signs of general decay associated with the trembling, clearly distinguish paralysis agitans. In *cerebro-spinal sclerosis*, the scanning speech, the increased patellar tendon-reflex, the nystagmus, the occurrence of the jerks only when the muscles are put into motion, unlike the abrupt and erratic movements of chorea, and a persistence in the direction given to the motion notwithstanding the oscillations, are most significant. Both affections, too, are encountered in persons older than are generally subject to chorea ; especially is paralysis agitans. Multiple sclerosis happens, however, also in children, and we meet with cases of paralysis agitans closely affiliated to chorea ; like it, too, originating in fright. But they differ in the motions repeating themselves rhythmically and symmetrically on the two sides of the body,\* and in presenting nothing of the irregular and rapidly changing character of the true choreic movements.

*Convulsive tremor*, a paroxysmal affection in which severe muscular tremor arises several times in a day, differs from chorea in not being continuous, as it occurs in attacks lasting from fifteen to twenty minutes, passing off gradually. The unrestrainable tremor affects the face, the arms, and the trunk, but not the lower extremities, and is associated with increased sensibility of the skin of the disturbed parts. Clonic spasms occurring also as sudden contractions or shocks, and affecting pre-eminently the upper part of the limbs, have been delineated by Friedreich as *myoclonus multiplex* ; the muscular irritability is much increased.

---

\* As in the case recorded by Sanders, Edin. Med. Journ., May, 1865.



In *athetosis*, the condition described by Hammond, there is continual motion of the fingers and toes, with inability to retain them in any position in which they may have been placed. Great tendency to distortion exists in the mobile spasm, and we find, on the whole, much resemblance to localized chorea. But the malady generally comes on with epileptic paroxysms; and headache, vertigo, slowness of speech and of thought, tremulousness of the tongue, numbness of the affected side, and pains in the limbs which are the seat of the spasms, give us a very different clinical picture from chorea. Athetosis is supposed to be due to disease of a cortical centre. It has been observed to be bilateral in idiotic children. Similar, too, is the mobile spasm that may be noticed in palsied limbs, the *post-hemiplegic chorea*. Gowers regards the spasm as identical with mobile spasm, in which slow, irregular movements conspicuous in the hand, and slow irregular incoördination occur.

*Facial spasm* differs from the spasmodic contractions of chorea in being always of equal intensity, and in the grimaces being strictly confined to the face, manifesting themselves in the same group of muscles, and generally existing only on one side of the face. In *convulsive tic*, as described by the French, the facial spasm is combined with signs of hysteria and with mental changes.

*Writers' cramp*, an affection in which every attempt at writing at once produces spasmodic action of the muscles of those fingers that are brought into play, is separated from chorea by its occurrence in individuals who have strained their muscles in using a pen continuously and rapidly; by the almost instant cessation of the spasm when the afflicted person ceases to write; and by the ease with which the fingers perform other motions and are capable of being used for every purpose except the one which has brought on the disorder. Pain, limited to the affected part, or more extensive, often attends this affection, at times induced only by writing, at other times spontaneous. A very analogous complaint, an "occupation neurosis," too, is encountered in seamstresses; also in telegraph-operators, particularly those who use the Morse instrument. These cramps, and all those of a similar kind caused by the occupation, such as in pianó-players, in violinists, and in type-writers, car-drivers, stone-masons, cigarette-

makers, shoe-fitters,\* have the same diagnostic sign that has just been mentioned as characteristic of writers' cramp—namely, that the spasm befalls only those muscles the overstrain of which has led to the affection, and that it ceases when the fatigued muscles are kept at rest or are brought into action for a different purpose. A form of cramp like that of writers' cramp happens in those engaged in preparing photographic plates;† and I have seen it in turners, engaged in what is called "oval turning."

There is a disorder, closely allied to chorea, which consists in repeated violent bobbings of the head, lasting many minutes at a time. These *salaam convulsions*, as Sir Charles Clarke called them, are a very obstinate complaint. They are most commonly met with in children, but have been known to occur in adults‡ and to lead frequently to impairment of the intellect.§

**Hysteria.**—This description of hysteria will deal chiefly with the symptoms of an hysterical paroxysm. Most of the local hysterical affections have been, or will be, considered in connection with the diseases they ape; and to discuss any questions relating to the nature of this perplexing malady, or to attempt to scrutinize or to interpret all the false and contradictory signals it hangs out, is, in a work of this kind, manifestly impossible.

An hysterical fit may set in suddenly, under the influence of some violent mental emotion; but more generally it is preceded by altered spirits, by a sensation of pressure, and of constriction at the pit of the stomach, which feeling ascends to the throat, and is likened by the patient to the rising of a ball. She becomes much agitated, sobs, laughs, cries, her muscles contract violently, or she lies motionless, and apparently without the power of motion, until her seeming insensibility is disturbed by something she disapproves of, or fears. The heart palpitates; the breathing is irregular and heaving,—on account, perhaps, of an affection of the larynx, but not of its temporary closure, which, as Marshall Hall tells us, so commonly ensues in epilepsy.

These hysterical outbursts differ from the spasms of *chorea* by their remissions, the patient remaining at times for months free

\* Moyer, *Med. News*, Feb. 1893.

† Napias, *Gazette Médicale de Paris*, No. 40, 1883.

‡ Levick, *Amer. Journ. Med. Sci.*, Jan. 1862.

§ Henry Barnes, *Liverpool and Manchester Hospital Reports*, 1873.

from the convulsive movements. Moreover, there is not even partial or apparent unconsciousness in chorea. It is true that this malady and hysteria are sometimes combined, or rather that chorea happens in hysterical subjects, and is then brought about by imitation, and is apt to come on suddenly; yet it is remarkable how rarely fits of hysteria take place in those affected with chorea.

It is sometimes very difficult to distinguish between paroxysms of hysteria and of *epilepsy*; and it becomes the more difficult if the epileptic seizures occur in hysterical patients. Yet there are ordinarily many well-marked points of distinction between the two maladies, as will be seen from this table:

## EPILEPSY.

Usually occurs without exciting cause.

Sets in with a scream.

Sudden and complete loss of consciousness.

Livid face; escape of frothy saliva from the mouth; eyelids half open; eyeballs rolling; grinding of the teeth; biting of the tongue; more or less insensibility of the pupils to light.

Distortion of countenance.

Patient evinces no feeling.

Aura epileptica.

Convulsions often more marked on one side than on the other; and at first tonic rather than clonic.

Movements unlike voluntary acts.

May pass urine involuntarily.

Paroxysm generally of short duration.

Paroxysm followed by a heavy, half-comatose sleep, by headache, and by dulness of intellect.

Frequently occurs at night.

No particular connection with uterine or ovarian disturbance, although a paroxysm often takes place at the menstrual period.

## HYSTERIA.

Often induced by emotion.

Noisy during attack.

Gradual and only partial or apparent unconsciousness.

Face flushed, or complexion unaltered; no froth on lips; eyelids closed; eyeballs fixed; neither grinding of the teeth nor biting of the tongue; pupils react readily.

No distortion of countenance.

Patient sighs, or laughs, or sobs.

Globus hystericus.

No such difference; convulsions tonic, followed by clonic.

Movements resemble voluntary acts.

Copious diuresis afterward.

Paroxysm generally of longer duration.

Paroxysm not followed specially by sleep; patient often, after attack terminates, wakeful and depressed in spirits.

Rarely occurs at night.

Often connected with disorders of the uterus or ovaries, or of menstruation.

There are, however, spasms that occur in hysterical patients which, though a functional nervous affection, appear like a blend-

ing of hysteria and epilepsy. Charcot\* particularly has called attention to this *hystero-epilepsy*, and describes its distinctive traits as consisting in premonitory symptoms of rather long duration, and exhibiting an aura which, starting in most cases from the ovarian region, advances progressively to the head. The cry is prolonged and modulated, not short like the epileptic cry. The convulsions are identical; but, instead of entering subsequently upon a stage of snoring, the hystero-epileptic sobs, laughs, gesticulates violently, or is delirious and subject to hallucinations. In the ovarian form of hystero-epilepsy, pressure upon the ovary will invariably modify the symptoms, if not completely arrest the attack; whereas in epilepsy no such effect is produced. In the cases of hystero-epilepsy with repeated attacks, the temperature scarcely rises above the normal, as it rapidly does under similar circumstances in epilepsy. There is no epileptic vertigo; there are no abortive fits. The malady is occasionally observed in men.

Hysteria is not an affection merely of paroxysms. In the intervals between them we find peculiar and significant manifestations of the strange complaint, which should be understood, lest they be taken as the signs of other maladies. We observe an extreme susceptibility of the nervous system, with defective will-power and imperfect self-control, varied hyperæsthesia, such as tenderness in the epigastrium or in the course of the spinal column or over the ovary; that peculiar pain in the left side which distresses so many hysterical and anæmic women; and sometimes local anæsthesia. Besides these, we encounter manifold local hysterical ailments, such as hysterical paralysis, hysterical aphonia, hysterical contractures, hysterical tremor, hysterical anorexia, hysterical peritonitis, hysterical affections of joints, hysterical pain in the forehead, hysterical suppression as well as hysterical retention of urine. Hysterical laughter has been found to occur on a large scale as a form of epidemic convulsion.† Mitchell and De Schweinitz‡ consider disturbance of color-sense common in hysteria. They have found reversal in the normal sequence of colors

---

\* Lectures on Diseases of the Nervous System. See also Richer, *Études cliniques sur Hystéro-Épilepsie*, Paris, 1881.

† D. W. Yandell, *Brain*, Oct. 1881.

‡ *Journal of Mental and Nervous Diseases*, vol. xix., No. 1, p. 1.



to be usually present in cases attended with anæsthesia. Muscular atrophy has been observed as a manifestation of hysteria,\* and a case has been recorded closely simulating syringomyelia.† Hysteria is met with in the male, especially after railway accidents. Hysterical paralysis may also happen in either sex, in the shape of hemiplegia, of monoplegia, or of paraplegia, and may be of very long duration.‡ As regards hysterical hemiplegia, it is remarkable that it does not affect the face; yet there may be an hysterical facial paralysis.§ Hysterical headache, Charcot tells us, like syphilitic, increases at night. The reflexes are apt to be much deranged. As Goodell || well says: strange and misleading reflexes come from the loss of brain control over the insubordinate lower nerve centres. In toxic hysteria, such as we observe after chronic lead or mercurial poisoning, tremor, anæsthesia, palsies of various kinds, and anorexia or hysterical vomiting are often observed,—much oftener than hysterical paroxysms.

The distinction between these hysterical pseudo-maladies and the diseases they simulate is far from being an easy task. We have to take into account the patient's age and sex; whether or not she has suffered from paroxysms of hysteria; how the pain is influenced by pressure; and the signs of functional disorder of the apparently affected part. We may thus avoid mistaking a phantom for a true disease. Yet there is another and opposite source of error quite as strenuously to be guarded against. The complaint may be really an organic one, occurring in an hysterical patient, and concealed, or exaggerated and complicated, by the symptoms of hysteria. In all such doubtful cases we must accord great weight to the extent of functional and constitutional disturbance accompanying the local morbid state. Then, too, hysterical symptoms may be prominent in certain brain affections. I have repeatedly noticed them in cases of cerebral embolism; and Brown-Séquard and Seguin ¶ have shown their frequent oc-

---

\* Hist. Deutsche medicin. Wochenschr., 1894, No. 21, p. 459.

† Wichmann, Berliner klinische Wochenschrift, 1895, No. 12, p. 252.

‡ See cases reported by Morton Prince, of twenty-nine, twenty-eight, and twenty-nine years' duration, Amer. Journ. of Med. Sci., July, 1892.

§ Babinski, Soc. Méd. des Hôp. de Paris, 1892.

|| Med. News, Jan. 6, 1894.

¶ Archives of Electrology and Neurology, May, 1875.

currence in lesions of the right hemisphere. In hysterical attacks connected with a cerebral neoplasm, the urea and the phosphates in the urine, Gilles de la Tourette\* shows, are diminished, while in epileptic seizures connected with brain tumors they are increased.

Hysteria is sometimes feigned,—feigned to elicit sympathy, or to procure compliance with wishes or caprices. Nor is the simulation of the disorder an outgrowth of our civilization. The epigrams of Martial prove how common the feigning of hysteria was among the Roman women.

**Tetanus.**—A very fatal disease marked by persistent rigid contraction of the voluntary muscles, particularly of those of the jaw, with violent brief exacerbations.

This distressing malady, as we see it, is generally *traumatic*, following a wound or an injury; for *idiopathic* tetanus is very seldom met with in temperate climates. But in hot countries, or in those in which sudden alternations of temperature are common, it is not a rare disease, and is indeed frequent among new-born children. The cases of idiopathic tetanus we encounter almost always follow exposure to cold. The malady is also seen in the puerperal state.

The symptoms of tetanus depend upon the action of a poison generated by a special micro-organism, the bacillus tetani. This is a long, slender bacillus, found commonly in the superficial layers of earth, and growing best in the absence of air and light. It is usually found only in the local lesion. In the so-called idiopathic cases the channel of infection eludes detection.

The muscles ordinarily first affected are those of the jaw and neck; there is a stiffness about them which the patient is apt to attribute to having caught cold. Sometimes, however, the disorder exhibits itself primarily in the external respiratory muscles. When the malady is fully developed, most of the muscles are stiff and hard, the jaw cannot be opened,—whence the common name of lock-jaw,—and there is much difficulty in speaking and in swallowing. The face is distorted, presenting the “*risus sardonius*.” With these symptoms we usually find rigidity of the muscles of the abdomen and of the limbs, and a distressing pain

---

\* Quoted, *Lancet*, May, 1893, p. 1083.

at the pit of the stomach, dependent upon spasm of the diaphragm. Besides the permanent contraction of the voluntary fibres, exacerbations of spasm take place, during which the muscles become very hard. These paroxysms are accompanied by intense pain, and recur with increased severity and frequency as the disease advances to a fatal termination. When at their height, the body becomes curved, the patient merely resting upon his head and heels. This is *opisthotonos*; while the setting of the jaw, especially when its muscles alone are affected, is called *trismus*. The trunk may be bent forward,—*emprosthotonos*; or to one side,—*pleurothotonos*; or the trunk and neck are rigidly extended in a straight line,—*orthotonos*. The spasm relaxes during natural sleep or induced narcosis. At the height of the attack the body is covered with copious perspiration.

Notwithstanding the striking muscular disorder and the exhausting pain, there is little constitutional disturbance; the pulse may be quickened, but it preserves its volume until the last stage is reached; and there is often no fever, nor is the intellect affected. But the temperature shows extraordinary variations. The thermometer may mark an increase of several degrees in the evening,\* and toward the end show a heat of 110°, even continuing to rise after death.

When tetanus results from an injury to the head, and more particularly in the distribution of the fifth nerve, there is often, in addition to the initial trismus, paralysis of the face on the same side of the face as the injury, and spasm on the opposite side.

Tetanus runs an acute or a chronic course. Some cases last three weeks, and when of such long duration are apt to recover. But generally the malady terminates fatally before the eighth day.

Few complaints are likely to be confounded with tetanus; yet these few resemble it closely in many respects. For instance, one of the freaks of hysteria is to take the appearance of tetanus; and tonic spasms dependent upon an affection of the spinal cord or medulla oblongata, strychnine poisoning, or hydrophobia, may accurately simulate its symptoms.

*Hysterical* tetanus is distinguished from the real disease by being

---

\* Ogle, Clinical Society's Transactions, 1872.

preceded by, or attended with, fits of hysteria ; by the age and sex of the patient ; by the absence of pain ; by the occasional occurrence of clonic instead of tonic spasms ; and by the intermission every now and then of all muscular rigidity. Moreover, the influence of the mind upon the seeming tetanus is very striking. If within hearing of the patient the employment of cold to the spine, or of the cautery, be threatened, or, better still, if the latter instrument be actually made ready for use before her, an extraordinary subsidence of all stiffening and starting of the limbs takes place. Hysterical trismus is more common than more extended hysterical tetanoid spasm, but, besides the symptoms of hysteria just mentioned, the absence of rigidity in the neck is very significant.

Tetanic spasms *symptomatic* of an affection of the spinal cord are separated from tetanus by the different history ; by no violent exacerbations being brought on, as they are in tetanus, by slight movements, or by an attempt at speaking, or by any reflex irritation ; by the absence of marked remissions ; by the rigidity being almost always limited to the extremities—except in the case of meningeal apoplexy in the cervical region, in which the tonic contraction in the upper extremity is associated with stiffness of the neck ; and by the setting in of palsy before the malady terminates.

In the tetanic spasms which may occur in scarlet fever, in typhus, in smallpox, or in pyæmia, and which are the result of an irritation of the cord produced by the poisoned blood, the rigidity runs so uncertain a course, appears so quickly, disappears so suddenly, perhaps not to reappear, or only to reappear after a considerable interval, that there is little likelihood of confounding the muscular disorder with tetanus. In *cerebro-spinal fever* the resemblance is much closer ; yet the whole history of the disorder, the state of the mind, and the progress of the case, are such as to prevent error. With muscular rheumatism tetanus can only be confounded at its onset ; but the muscles of the jaw are not rigid in rheumatism.

Another form of symptomatic rigidity requires to be distinguished from tetanus,—a local rigidity, owing to irritation of the nerve supplying the stiffened muscles ; as, for instance, a spasm from irritation of the peripheral or the central tract of the motor portion of the fifth, the so-called “ masticatory spasm ” of the face. This ailment may be of reflex origin, the exciting cause



being a decayed tooth, a wound, or exposure to cold; or it may exist in connection with apoplexy, or with an inflammation of the brain. Its main marks of distinction from the trismus of tetanus are, that it is purely local, is often of long continuance, is not painful, has no paroxysms of aggravation, is not combined with impaired deglutition, and is not dangerous.\* Similar spasm has also been observed as a result of irritative lesions of the pons, such as a new growth, or vascular disease, affecting the motor nucleus of the fifth nerve.

*Tetany* is characterized by tonic contractions, more especially of the legs and arms, and which may be intermittent, remittent, or continuous; the toes are apt to be flexed toward the soles; the hands become fixed; the spasm begins in the extremities. The jaws and the respiratory muscles are, unlike what we find in true tetanus, not affected,† or the jaws become so only toward the end.

The spasms are painful; they may occur several times in a day, or there may be weeks between them. They also can be produced, as Trousseau discovered, by pressure on the chief arteries and nerves of the affected limb. They are usually preceded by tingling or burning; in the intervals between them the muscles are readily excited to contraction and there is increased electrical excitability; the temperature, as a rule, remains normal throughout, although in severe paroxysms it may be elevated, and there may be copious perspiration. The contractions are bilateral, which distinguishes them from hysterical contractures. The malady happens chiefly in rachitic children, as a sequel of exhausting diarrhoea, after exposure to cold, and in nursing or in pregnant women; it has also been observed in connection with dilatation of the stomach. It has been described as occurring in an epidemic form, and the symptoms are like those of ergot poisoning.‡ The disease is not a common one in this country. Crozer Griffith§ has analyzed seventy-two cases reported in America.

\* Bright, in the second volume of his *Medical Reports*, gives the particulars of a case which illustrates many of the difficulties of diagnosis to which the affection may give rise.

† Wilks, *Guy's Hospital Reports*, 3d Series, vol. xvii.

‡ Stated in the German translation of this book.

§ *Trans. of the Assoc. of Amer. Phys.*, vol. ix., 1894.

The symptoms of *strychnine poisoning* are almost identical with those of tetanus ; yet there are some characteristic differences. The spasms from strychnine do not supervene upon exposure to cold, or upon a wound, but follow within about two hours or less the taking of some solid or liquid. They come on suddenly, with violence, and with epigastric pain and early reflex excitability. The tetanoid convulsions affect simultaneously nearly all the voluntary muscles of the body, but with greatest intensity those of the trunk and spine, producing very early—within a few minutes, commonly—a marked opisthotonus, which in tetanus does not appear, if it appear at all, for many hours or for days after the seizure. On the other hand, the stiffness of the jaws, which is among the very earliest signs of tetanus, is not at first perceived in strychnine poisoning, and, if it occur, occurs only imperfectly. Further, we do not see the frightful tetanic face, with its knit brow and horrid grin ; we do not observe intermissions in the convulsions, or difficulty in swallowing ; and in from ten minutes to two hours after the commencement of the attack the patient dies or recovers.

Finally, let us contrast tetanus with *hydrophobia*. Both showing the reflex functions of the spinal cord to be in an exalted condition ; both being spasmodic affections lasting ordinarily but a few days ; both taking place, the popular opinion to the contrary notwithstanding, at all periods of the year ; both presenting violent paroxysms of convulsions, which are often excited by the slightest touch or jar to the body ; both frequently occasioning torturing pain near the pit of the stomach ; both ensuing commonly upon an injury ; both usually augmenting in intensity from hour to hour, and scarcely within the reach of therapeutic measures,—these ghastly maladies are yet dissimilar. The one results from infection with a specific bacillus, often present in earth ; the other from infection with the virus of a rabid animal, most commonly the dog or the wolf. The one has a short, the other a long period of incubation. In the one, deglutition may be difficult ; in the other, it is next to impossible, all attempts at swallowing, especially of fluids, exciting the most distressing spasmodic dysphagia. In the one, early rigidity of the muscles of the jaw happens ; in the other, there is no such rigidity. In the one, the breathing may or may not be interfered with ; in the other, the spasms of respiration are almost as marked a feature as the spasms of deglutition.

Then the irritability of temper in hydrophobia; the increased reflexes; the fierce manner of the patient; his rabid, perhaps maniacal paroxysms; the constant thirst; the accumulation of stringy mucus about the angles of the mouth; the vomiting; the acute sensibility of the surface; the trembling of the muscles; the clonic instead of tonic spasms; the strangling sensation in the throat,—are phenomena too strikingly peculiar to render an error in diagnosis likely. The temperature is, as a rule, elevated, and in direct proportion to the intensity of the other symptoms. Toward the close it may reach a high degree, and it sometimes continues to rise after death. Some of the points here referred to serve also to distinguish hydrophobia from acute mania, and from hysteria. For, as in tetanus, so here, we find this erratic complaint simulating the terrible disease. In truth, it is the opinion of some, of Dulles\* especially, that the great majority of cases of supposed hydrophobia are of this character.

**Functional Spasms.**—There are spasms that take place in various parts of the body, sometimes clonic spasms, sometimes tonic spasms, which occur without apparent cause, and are more or less continuous or persistent. In time they may lead to contractures and deformity, or they may pass away. They may be of hysterical origin; but these are not now under discussion, rather the spasms that take place in one or both legs, sometimes in the arms, occasionally in the muscles of the face, which occur in those who are not hysterical subjects, and are not traceable to any lesions. Pressing on particular points may at once excite them; on the other hand, there are “pressure-points” which when acted on will cause the convulsive movements to be arrested. The trophic disturbance that attends them is usually very slight. Tonic contractions are very apt to alternate with clonic spasms, or there may be only complete tonic spasm during attempts at moving certain muscles. At times spasms of the internal muscles, as those of deglutition or respiration, may coexist; or the spasms may be limited to these muscles. The disorder is sometimes hereditary.

There is a curious form of spasm, a tonic contraction of the muscles, which impedes locomotion. It shows itself when the

---

\* Transactions of the College of Physicians of Philadelphia, 3d Ser., vol. xvi., 1894.

muscles are first put into action after a period of rest, or after an unexpected irritation, as striking the toes against a stone in walking, and is augmented by nervous dread about it. Happening, as it generally does, in the lower extremities, it leads there to muscular increase. This *Thomsen's disease*, or congenital or transient myotone, has been known to originate in sudden fright.\* It begins commonly at an early age, and is hereditary; it is persistent, although no organic cause for it has been detected. In one fatal case post-mortem examination disclosed the existence of hyperplasia of the muscular tissue, without appreciable lesion of the central or peripheral nervous apparatus.† The difficulty is most marked in the morning, on first rising, attempts at movement causing the muscles to become rigid and the joints fixed; yet if exertion be persevered in, the spasm becomes less and less, and continued walking is possible until after another period of rest. The spasm is aggravated by attention and emotion, and very rarely affects the muscles of the face. Electrical and mechanical irritability are heightened. Sensibility and reflex excitability are unaffected.

The chief difference between Thomsen's disease and *paramyotone*, which is also a family affection, is, that in the latter spasm is not started by voluntary movements and is more permanent; the marked spasms may last several hours. They are excited by cold and allayed by warmth.

---

\* Case of Schönfeld, Berl. Klin. Wochenschrift, July, 1883; and of Engel, Phila. Med. Times, Sept. 8, 1883. See also cases collected by Möbius, Schmidt's Jahrb., No. 6, 1883; and the cases by Deligny, Union Méd., 1885, xxxix. 50-52; Bernhardt, Beitrag zur Pathologie der sogenannten "Thomsen'schen Krankheit," Centralbl. f. Nervenhe., 1885, viii. 122-126; Fischer, Neurol. Centralbl., 1886, v. 73-78; Delmas, Maladie de Thomsen (Dysmyotonie congénitale), Journ. de Méd. de Bordeaux, 1886-87, xvi. 97-100; Buzard, Two Cases of Thomsen's Disease, Lancet, 1887, i. 972-974; Mibeleisen, Zur Casuistik der Myotonia congenita oder Thomsen'schen Krankheit, München. Med. Wochenschr., 1887, xxxiv. 433; Dana, Thomsen's Disease, Medical Record, April 21, 1888; Blumenau, Thomsen's Disease, Neurologisches Centralblatt, 1888, p. 679; Hale White, Guy's Hospital Reports, 1889; Delprat, Deutsch. Med. Wochenschr., Feb. 25, 1892; Dana, Journ. of Nervous and Mental Diseases, Feb. 1892; Huet, Nouvelle iconographie de la Salpêtrière, Paris, Feb. 1892.

† Déjerine and Sottas, Compt.-rend. hebdom. des séances de la Soc. de Biol., 1893, No. 23, p. 669.



A family type of congenital myotone is described by Eulenburg,\* and intermittent congenital myotone by Martin and Hausmann,† in both of which exposure to cold was followed by tonic spasm of various muscles. The irritability of the affected muscles was diminished. It is not impossible that future investigation may disclose some relationship between the spasmodic conditions here described and some of the forms of functional palsy previously detailed.

### Diseases characterized by Gradual Impairment of the Mental Faculties with Paralysis.

**Chronic Softening.**—There are two main forms of softening, —the red and the white. The former is inflammatory,—a circumscribed encephalitis,—and runs an acute course, with symptoms, as we have already mentioned, often closely simulating those of apoplexy, but sometimes with signs like those of the chronic malady, and differing in nothing but in their intensity and short duration. The second kind is chiefly dependent upon a change in the nutrition of the brain, and is nearly always linked to a diseased condition of the cerebral arteries and plugging of the vessels; it may be, however, caused, or at all events accompanied, by an inflammatory exudation infiltrated among the nervous pulp. These, briefly, are its early symptoms: gradual impairment of intelligence; weakening of memory; headache; vertigo; muscular debility; cutaneous hyperæsthesia or anæsthesia; formication and numbness; and slight and partial palsies, particularly of the muscles of one side of the mouth, or of one eyelid. Then there is not unfrequently defective articulation, with great irritability of temper, nausea and vomiting, extreme sensitiveness to sounds, and painful feelings in various parts of the body. As the local mischief advances, the paralysis becomes more universal, assuming generally the hemiplegic form; and spasms, either tonic or clonic, or epileptic convulsions, occur.

The mental decay proceeds steadily, and sometimes shows itself in a constant repetition of the same action or the same phrase. In

---

\* *Neurologisches Centralblatt*, 1886, No. 12; *Jahresber. über die Leist. u. Fortschr. i. d. ges. Med.*, xxi. 2, p. 164.

† *Arch. f. path. Anat. u. Physiol. u. f. Klin. Med.*, cxvii. 7, p. 587; *Jahresber. über die Leist. u. Fortschr. i. d. ges. Med.*, xxiv. 2, p. 76.

an old lady whom I attended, this was the most marked symptom : she was constantly complaining that her teeth needed attention, was perfectly satisfied when assured by the dentist that they did not, but soon reiterated her complaint. Beyond this, and a most painful sensitiveness to sound and to light, intense headache, nausea, and a progressive deterioration of memory and of the faculty of thought, she presented no signs of cerebral softening. She died without the occurrence of paralysis.

Softening of the brain may be caused by a diseased state of the cerebral vessels, or by their obstruction ; by long-continued grief ; by persistent mental labor ; by constitutional syphilis ; by frequently-repeated epileptic paroxysms ; and by an inflammatory disease spreading from the meninges to the brain, or taking place around new formations and old lesions. It may also be dependent upon cerebral hemorrhage. At all events, we frequently meet with it in connection with hemorrhage, and associated sometimes in such a manner as to make it a very perplexing matter to ascertain if the softening have followed the extravasation of blood, or if the extravasation have taken place into an already diseased brain. We may conclude the latter to have occurred, if signs of deranged intellection or sensation have preceded the attack and are more than can be explained by disease of the vessels ; if, after reaction from the shock, the patient, instead of mending in mind, exhibit unmistakable evidences of progressing mental decay ; and if convulsive movements or rigidity of the limbs appear. But let us, in passing, remark that a small clot breaking down the softened cerebral mass, yet not extending beyond the limits of the diseased texture, occasions no special signs,—occasions only the signs of a sudden giving way of nerve-pulp : paralysis without unconsciousness.

We shall next study how various other cerebral maladies, such as congestion, anæmia, abscess, and hardening, may be distinguished from softening.

*Congestion* is discriminated by its being very rarely a persistent state. It may be active or passive,—resulting on the one hand from an increased supply of blood, and on the other hand from interference with the venous return. An acute attack produces the symptoms of apoplexy ; a more lasting congestion is recognized by tracing the cause which has led to the fulness of the

vessels,—such as a disease of the heart or of the abdominal viscera,—and by noting that, although the patient suffers from dull headache, from disturbed sleep, from jerking of the muscles, from pulsation of the carotids, from vertigo, these signs are far from being constant, and come and go for a long time without any material disturbance of the functions of the brain being perceptible, in reference either to thought or to voluntary motion. The finding of optic neuritis or choked disk, or the presence of paralysis, would determine against congestion.

*Cerebral anæmia*, occurring suddenly, produces unconsciousness, or dizziness or stupor; or, if very general, and especially if associated with venous congestion, it may cause convulsions. When more gradually induced, it manifests itself by drowsiness, sighing respiration, distressing headache, often referred to the vertex; by the pale face and uninjected eye with large pupil; by derangement of the special senses; by the vertigo and the other symptoms of cerebral disorder being relieved in the recumbent position; and by the feeble pulse and cool forehead. Then, in tracing its history we are apt to find that it occurs in those who have been exhausted by debilitating diseases, or by repeated hemorrhages, or by albuminuria. The chief distinction from softening lies in the history of the case; the aspect of the patient, too, and the absence of palsies, or their passing nature, must be taken into account. But we must not forget that if the morbid condition be long continued, the ill-nourished brain softens. Anæmia is also the first stage of softening due to vascular occlusion.

Inflammation of the brain, *cerebritis*, unlike softening, is a disease of short duration. It usually follows traumatism, or adjacent or remote disease. If simple in character, it may give rise to softening; if attended with the presence of pyogenic micro-organisms, suppuration ensues and an abscess may be formed. If the inflammation be superficial, the membranes share in the process. In addition to the usual febrile phenomena, the symptom most commonly observed is headache; vomiting is occasionally present, while optic neuritis is rare. Sometimes convulsions and delirium occur. Other symptoms depend upon the situation of the inflammatory process. The diagnosis rests essentially upon the recognition of an adequate causative condition, such as traumatism, fracture or caries of the cranial bones, ear disease, or an

infectious process. Softening due to vascular occlusion is also attended with adjacent inflammation.

In rare cases there is a primary *acute hemorrhagic encephalitis* without obvious cause, though it is likely that the inflammation starts in the blood-vessels. In its main features it is similar to the acute encephalitis of children, where, however, the lesion is most apt to be cortical. In the primary acute encephalitis of adults punctiform hemorrhages are noticed.\* The chief symptoms are rapidly developing coma and hemiplegia. The knee-jerks are preserved, though there may be crossed tendon reflexes in the lower extremities. A significant feature is the extraordinarily high temperature, especially before death. In the latter respect, it is like what is sometimes observed in recent hemiplegia following embolism or hemorrhage. The disease progresses quickly to a fatal ending.

*Abscess* of the brain arises under the same conditions as cerebritis; but pyogenic micro-organisms play a very important part in the morbid process. Instead of simple, reddish, inflammatory softening, there results yellow, puriform softening. The most constant clinical association is with disease of the ear; suppurating processes in other parts of the body, such as abscesses in the lungs or fetid bronchitis, are also not infrequent causes. The symptoms are referable in part to the inflammatory process, in part to the presence of the purulent accumulation. The acute cases are likely to be progressively fatal; in cases which pursue a chronic course an initial inflammatory stage of brief duration is succeeded by a latent period, sometimes of considerable length, and this in turn by a terminal stage ending rapidly in death. Among the early symptoms are headache and vomiting, in association with febrile disturbance often attended with chills. Involvement of the cortex or subjacent white matter may cause local spasm; extensive disease, general convulsions. Paralysis and delirium may also be present. The acute period lasts from a few days to several weeks. In the latent stage, which may continue from a month to some years, decided manifestations may be wanting. Often there is headache; occasionally there are convulsions; at times slight mental disturbance exists. Elevation of temperature,

---

\* Strümpell, Deutsch. Arch. f. Klin. Med., Bd. xlvii, 1890, p. 53.



and recurrent rigors followed by sweats, also happen in abscess. Optic neuritis is as often absent as present. Constant headache and vomiting are among the most prominent symptoms. Though hemiplegia is met with not unusually, it is generally slight. The terminal stage which marks the rupture of the abscess may set in abruptly or gradually, with increase in the headache and mental symptoms, with vertigo, vomiting, derangement of consciousness, convulsions, and paralysis.

Few cases of abscess of the brain, as Lebert\* has shown, last longer than eight weeks. Abscess of the brain may be latent, and the sudden rupture of the abscess may give rise to symptoms undistinguishable from those of hemorrhage, undistinguishable unless we can infer an abscess from a disease of the bones of the head, or from some points in the history of the case.

*Atrophy* of the brain is especially observed in old age, and gives rise to the general decay of all cerebral functions noticed at this period of life. It is very generally connected with diffused sclerosis. As a rule, it occasions no distinctive symptoms. The brain is sometimes undersized from defective development. The diminution may be general, or unilateral, or even circumscribed. Partial atrophy is a common result of meningeal hemorrhage during birth. In some instances it follows meningitis early in life. Similar processes may also take place during intra-uterine existence. These varying conditions give rise to diverse symptoms, among the most common of which are mental defect, hemiplegia, convulsions, and mobile spasm.

There is yet, leaving tumors out of the question, another affection of the brain which may be confounded with softening: an *exhaustion of brain-power*, encountered among professional men, or those engaged in laborious literary undertakings. This sometimes comes on suddenly, with signs like those of a collapse; more generally it is slower in development. Its manifestations are a slight deterioration of memory, and an inability to read or write, save for a very short period, although the power of thought and of judgment is in no way perverted. Nor is the power of attention more than enfeebled: the sick man is fully capable of giving

---

\* Archiv für Path. Anat., Bd. x. See also Gull's paper in Guy's Hospital Reports, 3d Series, vol. iii.

heed to any subject, but he soon tires of it, and is obliged from very fatigue to desist. He passes sleepless nights, is subject to ringing in the ears, cannot bear much exercise, is troubled with irregular action of the heart, with a frequent desire to urinate, and with neuralgic pains in the face or a feeling of soreness in the head; but he does not lose flesh, and his digestion is unimpaired.

Many remain in this condition for months, and then slowly regain their health. What the precise disturbance of the brain consists in, is uncertain: it is possible that the nutrition of the organ has been interfered with from overuse and worry, and that the further continuance of mental toil and anxiety would have led to softening. The phenomena of this form of *neurasthenia*, as it is now customary to call the disorder, differ from those of softening by the absence of, or at least by the far less permanent and marked, headache, by the comparatively unimpaired intelligence, and by the non-occurrence of spasms or of paralysis, and of the causes that generally produce softening.

A condition comparable to the one just described occurs in those that have been subjected to persistent care and anxiety, or have suffered from long-continued or depressing illness, or have indulged in excesses, or have lost much sleep, or are the victims of digestive derangement or of eye-strain. A sense of fulness in the head, vertigo, languor, pain in the back, indisposition to effort, undue readiness of fatigue, tremulousness and general nervous and mental irritability, disturbed and unrefreshing sleep, and morbid fears, are common symptoms. Among women the menstrual function is often deranged.

Let us now consider the *diagnosis* of the chief varieties of softening. The more acute the symptoms, the greater is the likelihood of their being due to an inflammatory lesion; and in young subjects this probability becomes very great. A latency of the affection, its gradual manifestation, its existence in persons advanced in life, and in whom we have reason to suspect degeneration of the coats of the arteries, or, on the other hand, a history pointing to closure of the vessels by a plug, or to an embolus washed into them from a diseased heart, are facts which justify the conclusion that the softening is owing to depraved nutrition of the cerebral substance, and not to its inflammation.

**Tumor.**—Tumors of the brain give rise to a great diversity

of signs, according to their locality, their size, and their nature. Let us examine the group of symptoms by which we may infer their occurrence, and then see in how far an attempt to distinguish their seat and precise nature is likely to succeed.

The presence of a tumor in the brain is rendered probable if, in addition to vertigo, to vomiting or to a disposition to vomit, or to headache, violent, but paroxysmal and neuralgic in its character, we find impairment or loss of vision, or indeed anaesthesia of any special sense, and epileptiform convulsions not followed by any greater deterioration of health than previously existed; if with these signs of cerebral irritation the intellect is not at first markedly disordered, nor the articulation affected; and if paralyses do not show themselves until a very long time after the headache, and are even then limited to the muscles of the eyeball or of the face, or to the muscles of the extremities of one side of the body. As a further sign of cerebral tumor, we may class optic neuritis, or choked disk. It is a curious fact to be borne in mind that cerebral tumors occur in males more than twice as frequently as in females. It may also be noted that the larger number of cases are in the young or in those in the prime of life; the aged are remarkably exempt. The commonest forms of tumor are tuberculous, gliomatous, sarcomatous, and syphilitic. Less common are carcinomata and parasitic tumors. Before the evidence of a tumor is considered conclusive, we must exclude other chronic cerebral maladies, especially softening, abscesses, and chronic meningitis.

We separate chronic *softening* by noticing that the headache caused by a tumor is much more constant and violent, having paroxysmal exacerbations; that the intelligence remains for a long time intact, save, perhaps, in a weakening of the memory; that optic neuritis is a usual accompaniment; that motor and sensory disturbances are less frequent and prominent, but convulsions far more so. Further, cerebral tumor is more common in early life, chronic softening in late life. Disease of the heart or of the blood-vessels, or Bright's disease, points to the latter state. Remissions, or intervals of apparent improvement, occur in both morbid states; but they are more perfect and of longer duration in tumor than in softening.

The differential diagnosis between tumor and *abscess* is more

difficult. We may conclude the latter to exist, if the cephalalgia be sudden in its development, and uniform and general, instead of neuralgic and limited. Then, convulsions, drowsiness, paralysis, and coma succeed one another much more rapidly and, except convulsions, are present much more constantly in abscess than in tumor,—a malady running a very chronic course, and in which the patient does not remain drowsy or palsied after the epileptiform seizures.\* Further, optic neuritis and localizing symptoms are more common in tumor than in abscess, and this shows especially in the palsies of cranial nerves. If, moreover, we obtain the history of injury to the skull, or find a discharge from the ear, or pain upon pressure over the mastoid process, or a chronic disease about the head, or protracted suppuration in any part of the body, we may safely infer that an abscess, not a tumor, is the cause of the evident cerebral mischief. Abscess, like tumor, chiefly affects males.

*Chronic meningitis*, an affection sometimes complicating tumor, is discriminated by laying stress on its etiological relations,—such as blows upon the head, diseases of the bones, syphilis, rheumatism, or alcoholism,—and by observing its frequent though irregular accessions of fever, the great irritability of temper, the dulness of intellect, the loss of memory, and the nocturnal delirium. The pain, too, is, as a rule, somewhat duller and more diffused than in tumor, though, perhaps, more constant, and there is more vertigo; but the convulsions, on the other hand, are less distinctly epileptiform in type; yet convulsive movements of some muscles are very common, and may even be followed by incomplete paralysis. Meningitis may be excluded if optic neuritis or any marked alteration of the disks be found early in the case. Indeed, optic neuritis is mostly absent or is very slight in chronic meningitis.

*Thrombosis of the sinuses of the brain* may occasion partial palsies, with symptoms of cerebral pressure, like those of tumor, and cannot be distinguished except in the instances in which we find distention of the collateral circulation and injection

---

\* With reference to the epileptic fits, they may be absent. Thus, they occurred in only thirty-eight cases of abscess of the brain out of seventy-three collected by Gull and Sutton (see article "Abscess of Brain," in Reynolds's System of Medicine).



and œdema of the forehead and eyelids.\* Convulsions, further, are very rarely among the symptoms; and generally these are more similar to the manifestations of meningitis than of tumor. The condition is usually a result of enfeeblement of the circulation, in conjunction with increased coagulability of the blood; or is secondary to adjacent disease. In children with marasmus, or in adults with caries of the skull, marked cerebral phenomena may lead to the correct inference of thrombosis.

The precise *seat* of the tumor is difficult to determine. An affection of the special senses or of cranial nerves points to disease near to, or at, the base of the brain; and the probability of this view is much strengthened if there be paralysis of the face on the side opposite to that of the extremities, and if vigorous inspiration, during which the brain falls and presses the morbid mass against the walls of the base of the skull, cause or increase pain; whereas, so says Romberg, in tumors on the upper surface, forced expiration produces a like result. In cases of tumor of the *pons* or the *crus*, particularly when tubercular, incoördination of the arm similar to the jerky movement of disseminated sclerosis is met with; but it is unilateral, not bilateral as in sclerosis. In tumors of the *cerebellum* we have headache, severe, often bilious vomiting, nystagmus, staggering gait, also spasms, and rigidity; the knee-jerk may be absent or increased; there may be no marked alteration of the optic disks. Tumors in or near the *cortex* of the brain give rise to localized convulsions on the opposite side of the body. Then, in determining the exact position of brain tumors we must make use of the researches on the localization of the cerebral functions. The difficulty of applying this extending knowledge to the diagnosis of tumors at the bedside is, that they may give rise to circumscribed inflammation around them, or to irritation in even more remote parts, and that the special manifestations of the disorder of the part affected by the tumor are thus blurred or obscured.†

In endeavoring to determine the seat of the tumor it is necessary to distinguish as clearly as possible the difference between

---

\* Heubner, quoted in Schmidt's *Jahrbücher*, No. 1, 1869.

† Compare Nothnagel, *op. cit.*, and Bernhard, *Symptomatologie und Diagnostik der Gehirngeschwülste*, Berlin, 1881; and researches by Mills, in *Amer. Journ. of Med. Sci.*, in the last few years.

the results of generalized pressure or distant effects and those due to direct and localized influences. It is only the constant abnormal symptom that points out the location of the lesion. Paralyses, pareses, spasms, which change in intensity or affect now one, now another set of muscles or organs, show that the centres are disordered only indirectly and temporarily, and that the true position of the neoplasm is to be sought elsewhere. Another indication is derived from a consideration of the relative intensity of the different symptoms. The less complete a paralysis or the less energetic the spasm of a certain set of muscles, the less certain is the injury to be localized in their centres, and the reverse. Too much dependence must not be placed on the subjective location of the pain. Diffuse pressure may cause more pain at a point far removed from the growth than in its immediate neighborhood. But when spasm or paralysis of a limited set of muscles exists, as in cortical epilepsy, and the pain is located by the patient at a point corresponding to the topographical position of the corresponding centres, the deduction becomes quite certain that the lesion is at this point. When from other indications the inference is probable that the growth is in the cortical substance, the additional symptom of pain makes the diagnosis more sure. Yet another indication of location is gained when it is found that the symptoms are more narrowly limited either in extent or in kind; are, therefore, more strictly localizing.

It is manifest that in all tumors of the cortex, or of the white substance immediately beneath, the symptoms will be *unilateral* and include convulsions. When both sides of the body are about equally affected, the tumor must be placed at the base of the brain, unless the growths be multiple and situated in symmetrical parts of the two hemispheres. Where the symptoms are more intense upon one side of the body than upon the other, the weaker symptoms are to be attributed to the distant or indirect effects of pressure. Paralysis, of course, is a symptom of more profound disturbance than spasm or convulsive movement. The last is therefore probably due to an irritative or indirect effect, or to a slowly-growing neoplasm. The existence of *papillitis*, *optic neuritis*, or *choked disk*, is an almost certain sign of intracranial neoplasm. But, unfortunately, the symptom gives scarcely an indication either of the nature or of the seat of the new growth.

Yet since the papillitis may precede other symptoms, since also no deterioration of vision may have been noticed by the patient, an ophthalmoscopic examination should always be made when there is any suspicion of the existence of tumor.

Can we form an opinion of the *nature* of a tumor of the brain from any of the signs referable to the cerebral malady? We cannot: the character of the pain has been thought to be of great significance; but the testimony to prove that it is so, is in the highest degree unsatisfactory. We may sometimes, however, from the history of the case, or from the existence of some of the manifestations of special cachexia, draw a correct inference. In *gliomatous* brain tumors, Virchow has pointed out, there is often the history of a blow. They are usually single, and most common in the cerebral hemispheres, and occur next in frequency in the cerebellar hemisphere; then in the central ganglia, pons, medulla, crus, and corpora quadrigemina. Other sarcomata develop in the brain or in the membranes, or from the bones, particularly at the base. They differ from gliomata in being circumscribed and not infiltrating. Tuberculous growths are often multiple and most frequent in the cerebellum; they are also found in the pons, central ganglia, crus, medulla, and corpora quadrigemina.

If we find disease of the lungs, or any evidences of scrofula, and the patient be young, we shall probably be right in conjecturing the tumor of the brain to be a mass of *tubercle*; but if the sufferer be advanced in years, and exhibit tumors in various parts of the body, or other signs of a cancerous diathesis, we may with reasonable certainty presume the tumor within the skull to be *cancerous*. *Syphilitic* tumors are mostly cortical, very rarely cerebellar, grow rapidly, and are greatly influenced by antisyphilitic treatment. Other kinds of tumors and deposits can scarcely be said to be within the reach of diagnosis. *Cysts* seated in the superficial portions of the brain either occasion no symptoms, or give rise to headache, to attacks of vertigo, to vomiting, and to epileptic seizures, but very rarely to palsies. The symptoms mentioned are far more apt to be present when the cysts occupy the lateral ventricles; then epileptic convulsions especially are rarely absent.

The manifestations of an *aneurism* within the cranium are those of an ordinary tumor, and the affection is not distinguish-

able except when the symptoms are referable to the presence of a tumor in the course of a cerebral vessel, and we find present a cause of aneurism, or decided indications of disease of the vessels in other parts of the system.\* A small aneurism may occasion no symptoms; one large enough to exert pressure on adjacent structures may be attended with headache, often pulsating, usually continuous, sometimes paroxysmal; vertigo; mental dulness and irritability; occasionally convulsions; paralysis; rarely optic neuritis. Neither the presence nor the absence of a subjective feeling of pulsation and of a murmur has a positive significance; for, notwithstanding the cases of Jonathan Hutchinson† and Humble,‡ in which the diagnosis was made during life, the detection of a murmur, as I know from observation, is not a certain sign.

Aneurism of the internal carotid artery may cause blindness on the same side, paralysis of the third, and of the ophthalmic division of the fifth, nerve, impairment of the sense of smell, and hemiplegia. Aneurism of the anterior cerebral may occasion many of the same symptoms, although the muscles of the eyeball usually escape. Aneurism of the middle cerebral is usually attended with hemiplegia and convulsions. Aneurism of the basilar artery may cause extensive damage, with widespread paralysis, including the cranial nerves; convulsions are rare.

**General Paralysis.**—This fatal cerebral malady is the result of a diffuse interstitial meningo-encephalitis; the spinal cord may become secondarily affected. Clinically, the disorder is marked by impairment of the powers of locomotion; by an inability to articulate distinctly,—a symptom which precedes the deranged locomotion; by the meaningless countenance; and by failure of memory and complete perversion of the mental faculties, amounting, in fact, to insanity.

The palsy is peculiar: indeed, except toward the end there is, in the usual sense of the term, no palsy in the limbs at all; there is rather a want of control over their co-ordinate action, displaying

---

\* James H. Hutchinson, *Pennsylvania Hospital Reports*, vol. ii.

† *British Medical Journal*, April, 1875.

‡ *London Lancet*, Oct. 1875.



itself first in the hands by clumsiness of movements and irregular handwriting, and in the gait by uncertainty and a swaying from side to side when the patient attempts to walk. The impairment of the muscular movement gradually extends : tremulousness in the muscles of expression is noticed ; the speech becomes more inarticulate, until scarcely a word can be distinguished ; and the patient cannot rise without being assisted. In many cases reflex contraction of the tendo Achillis is wanting. As the disease advances, the cutaneous sensibility is greatly diminished or is lost. The pupils are unequal, and either markedly dilated or contracted. The mental derangement is generally manifested by an exaggerated sense of personal power or importance, and fancies of great wealth ; the moral feelings greatly deteriorate ; sometimes there are maniacal outbreaks and epileptic attacks, or alternating periods of excitement and depression. Deceptive remissions in the progress of the disease may take place, but the termination is invariably fatal. Death is often preceded by convulsive attacks and by coma, or by painful contractions of the muscles of the trunk or the extremities, or by obstinate diarrhoea, or by pulmonary affections. Pneumonia is especially common.\*

The early signs of general paralysis of the insane are difficult to recognize. A change in character, in power of mental attention, and in judgment, absent-mindedness, and weariness easily brought on by brain-work or by any physical exertion, are very significant in a middle-aged man, if joined to alteration in handwriting and some impairment in executing delicate muscular movements. With these symptoms there is commonly, as Folsom † mentions, loss of flesh.

In more advanced stages there is not much doubt about the malady. It differs from other forms of extensive general paralysis in being far less of a real palsy. It is certainly far less complete than the extensive paralyses which follow lesions of the upper portion of the spinal cord, or which are consequent upon the poison of lead, or of malaria, or of diphtheria. Its association with marked disturbance of the intellect furnishes, moreover, a differential test of great value, and not merely with reference to

---

\* Crichton Browne, *Brain*, Oct. 1883.

† *Transact. of Association of American Physicians*, 1889.

the general palsies just mentioned, but also as regards the trembling movements of old age, of progressive muscular atrophy, and of chronic alcoholism. In one of its forms, as Westphal mentions, there is a strong resemblance to locomotor ataxia in the signs of disturbed co-ordination, sensory impairment and absence of knee-jerks, with incontinence of urine and amaurosis; but the tremor in the muscles of the lips and face and the perverted mental state become of greatest significance. In some cases, however, changes in the posterior, as well as in the lateral, columns of the cord have been found after death, in addition to those present in the brain. On the other hand, the ataxia and the palsies distinguish the disease from mere senile dementia. Then, too, dementia paralytica is a disease of early manhood and of middle age, and often follows alcoholism or sexual excesses.

The defect in the articulation and the attending tremor of the lips, and in some instances the occurrence of apoplectiform seizures, accompanied by considerable elevation of temperature, may cause the disease to be mistaken for *cerebro-spinal sclerosis*. But in this affection, while the embarrassed, scanning speech coexists with great helplessness of manner, with oscillation of the eyeballs, with tremor manifesting itself only on motion, with paresis of the lower limbs, and finally with permanent contractions, we do not notice decided alienation of mind; there is nothing more than general enfeeblement and blunted emotional faculties.

*Paralysis agitans* may be confounded with general paralysis of the insane. But in paralysis agitans the voice is not really tremulous; there is rather a monotonous tone and uncertain utterance, which, with the fixed features, the sensation of excessive heat, the peculiar gait and attitude, the unaltered cutaneous sensibility, the tremor ever present except during sleep, the manner in which the patient when attempting to walk is propelled forward, and the very long duration of the symptoms, characterize the disease. The intellect becomes obscured toward the end of the malady, but not before. The cases most difficult to distinguish are those exceptional ones of general paralysis with altered character and enfeeblement of intelligence but without insanity, and in which the motor disorders are apt to be very pronounced.

### Diseases characterized by Enlargement of the Head.

**Chronic Hydrocephalus.**—The signs of dropsy of the brain are, progressive enlargement of the head, and a perversion or a gradual loss of one or several of the special senses, of the mental faculties, and of the power of voluntary motion. The child cannot bear the weight of the head; the gait is tottering and uncertain. The intellect, slowly but certainly, becomes deranged. As the malady advances, strabismus, partial palsies, epileptic convulsions, vomiting, cutaneous anæsthesia, and loss of sight, of smell, and of taste, are observable; the bowels become very constipated; and a copious secretion of tears and of saliva is not infrequent.

Before death takes place, which sometimes does not happen for years, the child ordinarily becomes idiotic. A few cases recover; fewer reach adult age with their brain compressed by the accumulated fluid; in still fewer the disease does not develop itself until after childhood. If the patient survive until adult age, the size of the skull is generally immense. I saw, some years since, a young man, twenty-two years of age, whose head measured fully two feet and a half in circumference. He could walk unaided, but often fell. He was half idiotic, and subject to epileptic fits; yet he had sufficient intelligence to understand what was said to him, and in his childish way to do as he was told.

Hydrocephalus may result from meningitis, from interference with the circulation through the veins of Galen, or from obstruction to the free movement of the cerebro-spinal fluid; occasionally no causative condition can be recognized.

The skull is sometimes very large without dropsy of the brain existing. The cranial bones may slowly thicken to an extraordinary degree from syphilis, or from unknown causes. The head may be overgrown, and its bones thickened and spongy, as in *rachitis*; or it may be large when there is no disease. These states differ from chronic hydrocephalus by the absence of cerebral symptoms; and in doubtful cases we may resort to the ophthalmoscope as a means of diagnosis. The vessels of the eye, even in the early stages of chronic hydrocephalus, enlarge, and in proportion as the serum compresses the brain we find an increase of

vascularity in the retina, with dilatation of its veins, and with an increase of the number of its vessels; complete or partial serous infiltration of the retina; and an atrophy, more or less perceptible, of the optic nerve. These lesions vary with the age of the disease and the amount of serous effusion; but none of them exist in rickets.\* Then in rickets the tendency is to spasm of the glottis, to diarrhoea,—not, as in hydrocephalus, to constipation, and the head is rather square-shaped than globular. In very rare instances the size of the head has been observed to be increased in the cerebral palsies of children, due to hemorrhage or embolism.

**Hypertrophy of the Brain.**—It is very questionable whether such a disease as a true hypertrophy of the brain exists. The enlargement, when not due to an unrecognized hydrocephalus, is mostly a congenital malformation, or is found in children in connection with changes in the brain of a sclerotic kind, or with those alterations caused by a defect of brain substance to which the name porencephalus has been given, but where, at the same time, in other portions of the brain extensive cell infiltrations and connective tissue changes may happen. The symptoms of hypertrophy of the brain are as obscure and doubtful as the nature of the malady. Undeveloped or failing intelligence, convulsions, varying palsies, are said to accompany the enlargement of the head, and when the fontanels are touched the sensation is that of a solid substance. In this respect the disease is unlike chronic hydrocephalus, as well as in the fact that the body does not waste as in dropsy of the brain. Mauthner† lays great stress on the different shapes of the head. In chronic hydrocephalus, he states, the forehead is the first to enlarge, and the posterior part of the skull does not expand until long afterward; in hypertrophy the reverse takes place. But this is of questionable value. West maintains that in hypertrophy there is no prominence, but an actual depression, of the anterior fontanel, and that a similar depression is observable at all the sutures. The disease has been observed in connection with rickets; it has been also noted to run in families.‡

---

\* Bouchut, *op. cit.*

† Krankheiten des Gehirns, etc, Vienna, 1844.

‡ See case of D'Espine, quoted in Schmidt's Jahrb., No. 3, 1882.



### Diseases characterized by Enlargement of Various Parts.

**Acromegaly.**—In this peculiar and uncommon affection, first described by Marie, enlargement of the hands and feet occurs, as well as of the head, and especially the face. Often bones and soft tissues both take part in the change, although the muscles may undergo wasting, with resulting weakness. The hands become broad and spade-like; the face assumes the shape of an elongated oval; the jaws, the malar bones, and the supra-orbital arches are prominent; the forehead is receding. Spinal curvature is common. The disease occurs in young adults and pursues a chronic course; in rare instances it has been met with in children, especially in imbeciles. In some cases a tumor of the pituitary body has been present, and in these headache, optic neuritis, and visual derangement have been observed. Sometimes changes in the thyroid gland have been noted, either enlargement or diminution in size. Occasionally dulness on percussion over the upper portion of the sternum has existed, and this has been attributed to persistence of the thymus gland. Rheumatic or neuralgic pains are not infrequent; and the tongue, lips, and nose may show striking increase in size, while the nails are small in proportion to the great growth of the bones in the hands and feet. Acromegaly may affect only one side of the body.

In *myxœdema* the tumefaction is not confined to the extremities, but is very general, and depends not upon changes in the bones, but upon a peculiar infiltration of the connective tissues. The skin is thickened and adherent to the subjacent tissues, and not pliable as in acromegaly. The face is "moon-shaped," and the jaws and malar bones are not projecting.

In the condition known as *osteitis deformans*, or *Paget's disease*, the changes in the bones of the face give to this the appearance of an inverted triangle. Besides, the disease attacks the long bones of the body, which undergo deformity.

Certain chronic diseases of the lungs and pleura are attended with enlargement of the terminal phalanges of the fingers and toes, and of the distal epiphyses of the bones of the legs and forearms. The finger-nails are curved, and the vertebral column is often bent. The changes are, however, usually restricted to these parts, and the disorder is not likely to be mistaken for acromegaly.

### Diseases characterized by Paroxysmal Pain.

There is a group of nervous disorders characterized solely by pain, confined ordinarily to one nerve. These nervous pains bear the generic name of *neuralgia*. Indeed, in all neuralgias the chief symptoms of the disorder resolve themselves into one symptom,—the symptom of pain. The pains are acute, follow the course of a nervous branch, and come on in paroxysms having distinct exacerbations, succeeded by distinct intermissions. In some cases these intermissions are long, in others short; in some they are complete, in others the pain is lasting and becomes from time to time exalted,—rather remissions, therefore, than intermissions. When the pain is severe it may be attended with muscular twitching. Sometimes, too, there is pallor followed by redness and swelling, though swelling is rare. Tenderness is present only when the neuralgia is of long continuance; at least there is not tenderness along the aching nerve, though we may find certain sensitive spots, as where a nerve-trunk emerges from a bony canal, or passes over a hard surface, or through fascia to become superficial, or at the point of division of a nerve-trunk, or of anastomosis of two nerve-trunks.

The pain of neuralgia is of a purely nervous character, and exists independently of inflammation, or of any recognizable textural change of the nervous centres or nervous trunks. All fixed pain and persistent early tenderness and evidences of trophic changes in the skin or muscles, and cutaneous eruptions in the course of the affected nerve, bespeak *neuritis*, and not neuralgia; and it is only when, after a minute search, we can detect no definite organic cause for the local pain, that we may set down our patient as laboring under neuralgia. Among other points of difference, too, we observe that the pain of neuralgia is often relieved by deep pressure, while that of neuritis is thus increased. Changes in the hair may take place as the consequence of either neuritis or of neuralgia, should the latter have existed for a considerable time.

From the characteristics of the pain just mentioned, it is evident that it is not likely to be confounded with that of ordinary local inflammation. But there is a kind of local pain for which neuralgia is often mistaken: the pain of *subacute* or of *chronic*

*rheumatism*. Yet this is in reality very dissimilar. The rheumatic pain is attended with soreness, is aggravated by movement or by pressure, is more diffuse and irregular, much more constant, much more influenced by alternations of temperature, but not acute or paroxysmal, and, finally, not limited anatomically to the course of one nerve, but scattered over parts supplied by several. Except as regards the influence of the weather, the pain of *myalgia* presents much the same points of difference, in addition often to the history of a muscular strain.

The source of the neuralgia should always be determined as closely as possible, on account both of the prognosis and of the treatment. In many cases it will be found to be connected with *anæmia*; in others, with the poison of rheumatism, of lithæmia or gout, of malaria, of syphilis, or of uræmia or ptomaines, or to be due to injuries to nerves, by contusion or wounds. It may be owing to emotional disturbances, or to exposure to cold. It is often reflex, the pain being far away from the seat of the disease, and due to irritation reflected through the nervous centres. For instance, an affection of the digestive apparatus, of the liver, or of the kidneys, may give rise to neuralgia in parts quite remote from them. It is evident that if such be the origin of the disorder, and if the malady which lies at its root and excites it can be controlled, the neuralgia will simultaneously disappear. Yet it must be confessed that we cannot always detect the cause, and we have often to treat the neuralgia by employing those agents which are suitable to the greatest number of cases.

Neuralgia may occur in any portion of the body. It may shift rapidly from one part to another, as in that peculiar neuralgia described by Putegnat,\* excited by a desire to pass water and by the act of micturition, beginning with numbness and acute burning or lancinating pain along the urinary passages, then affecting particularly the nerves of the forearm, especially the ulnar, and disappearing completely after micturition. The most frequent seat of neuralgia is about the head; and we shall here notice chiefly a few of its most common kinds. Most of the other varieties of the disorder will be elsewhere alluded to.

Neuralgia long continued becomes associated with the signs of

---

\* Gazette Hebdom. de Méd. et Chir., April, 1864.

nervous weakness, so-called *neurasthenia*. Indeed, in this state neuralgic symptoms are common, as are also abnormal sensations in the head, and many hysterical manifestations. Hysteria is, in truth, interwoven with most of the marked cases in women; and the history of excesses or of great mental or bodily strain usually accompanies the disorder in men.

**Facial Neuralgia.**—The sensory branches of the fifth pair are often the site of agonizing pain. But all the branches of the nerve are not equally liable: the lowermost of them is rarely affected. When the supra-orbital division is the seat of the ailment, the pain shoots to the forehead, the eyebrow, and the eyeball, which is apt to become injected. There are tender points just above the supra-orbital notch or foramen, in the outer part of the upper eyelid, at the lower edge of the nasal bone, and sometimes within the eyeball. If the infra-orbital nerve be disturbed, the pain darts to the upper lip, to the upper row of teeth and the posterior nares, and the cheek reddens and tingles, or the eyelids twitch. Tender points are found at the infra-orbital foramen, at the side of the nose, over the malar bone, and on the gums of the upper jaw. When the pain occurs in the inferior branch, it radiates to the lower lip and the chin, and is frequently accompanied by a flow of saliva. Tenderness exists at the inferior dental foramen, on the temple, and over the parietal eminence. Sometimes only one, at other times two, at other times all of the branches of the fifth are implicated in the complaint, or they may be seized upon alternately. There is often also pain at the vertex.

The disease is one of those belonging to advancing years; one of the neuralgias of bodily decay on which Anstie dwells. It has the same general causes as any other form of neuralgia. Sometimes it is associated with decayed teeth, or with an abnormal state of the bones of the head or face, such as thickening of the frontal, ethmoid, and sphenoid bones. Many of these cases terminate, after months or years of excruciating agony, in apoplexy.\* When from decayed teeth, the pain finally localizes itself in the dental arch, and there is persistent discomfort in addition to the neuralgic exacerbations.†

---

\* Sir Henry Hallford's *Essays and Orations*, p. 37 *et seq.*

† An interesting collection of cases is given in an essay by Brubaker on Reflex Neurosis associated with Dental Pathology.



The intervals between the paroxysms of neuralgia are of varying length. They may be of six months', or even a year's, duration; but so long an intermission is uncommon. Seasons in which sudden changes of weather are frequent generally excite attacks.

The malady is easily recognized. It may be mistaken for, or rather there may be mistaken for it, a disease of the bones of the face. But the local signs of this are different, and the pain is not paroxysmal. *Painful anæsthesia of the fifth nerve* is discriminated by the insensibility of the painful portions to touch, or indeed to any irritation. *Spasm of the face* is distinguished by the absence of pain from the convulsive twitchings of reflex origin which sometimes take place in facial neuralgia or "tic douloureux."

The *epileptiform neuralgia* described by Trousseau is dissimilar in these peculiarities: whether simple or combined with rapid convulsive movements of the muscles on one side of the face, it is quickly over; it lasts but ten or twenty seconds at a time, never more than a minute. Yet during the short duration of the seizures the pain reaches an intensity greater than in ordinary neuralgia. Moreover, in some persons who suffer from this terrible malady—the attacks of which may happen in quick succession by day as well as by night, and then perhaps remit for weeks or months—vertiginous sensations or epileptic fits occur, and thus the diagnosis is facilitated by the history of the case.

**Hemicrania.**—The pain here is limited to one side of the head, but it may extend to the other side, and be bilateral. It is intensified by sound of any kind, by movement and by light, and is often preceded by disorder of sight,\* sometimes of hearing and taste, by numbness and tingling in the limbs, by transient aphasia, by a sense of weight, though rarely by muscular weakness. Nausea generally attends the headache, but is most pronounced when this has reached its height, and is, as a rule, followed by vomiting or retching; the nausea and vomiting of the "sick-headache" are usually, indeed, prominent features of the parox-

---

\* There may be obliteration of objects in the field of view, or a curious glimmering attended with colored outline near the outside corner of the field of vision. These ophthalmic migraines have been described by Charcot (vol. iii. of his Clinical Lectures) as being at times among the forerunners of general paralysis.

ysm, hardly less prominent than the pain. During the attack there is commonly pallor of the face, which at the close gives place to flushing; slight mental change may also be noticeable. The attack lasts for hours or days; often it is severe for half a day. It may end with free diaphoresis or diuresis. At its termination, the patient feels exhausted, yet soon recovers his usual health, and may remain free from a seizure for a long time. But, as the disorder most commonly happens in women, and usually at their menstrual periods, the interval is not apt to extend beyond four weeks. At times the sensory phenomena of migraine occur with little or no headache.

Hemicrania, migraine, or megrim, has been explained as a neurosis of the sympathetic; or as a discharge of nerve-force, a "nerve-storm," from centric disorder. It is a stubborn affection, the tendency to which diminishes after middle age, but which, as Liveing\* clearly demonstrates, has an hereditary character.

Hemicrania must be carefully separated from the pain in the head which accompanies an *organic cerebral affection*. The main points of distinction are, that the neuralgic malady is paroxysmal, is attended with the same group of symptoms during each attack, and produces no nervous derangement in the intervals between the seizures.

*Rheumatism of the scalp* differs from hemicrania in the pain being continuous, dull, and superficial; in occupying generally both sides of the head; in being augmented by moving the affected muscles, and relieved by warmth. Moreover, there is almost always other evidence of rheumatism, and the pain is intensified by pressure; whereas in hemicrania, although the hair may be sensitive to the touch, strong pressure on the forehead, and even on the hairy part of the scalp, does not increase the pain, may indeed afford relief.

In *periostitis* affecting the bones of the head, particularly when syphilitic, we may find the same violent pain as in hemicrania. But there is considerable tenderness on pressure, the parts attacked are swollen and less elastic than the healthy portions, and the pain is especially severe at night.

**Sciatica** —This is sometimes a neuralgia following the course

---

\* On Megrim, London, 1873.

of the sciatic nerve, but often it is neuritis. The seat of the greatest suffering is generally the lateral surface of the thigh; thence the pains extend to the popliteal space, and in some instances along the anterior part of the leg. Often, too, the patient complains of an aching near the sciatic notch and in the loins. The pain is more or less steady; but it has its periods of fierce exacerbation, and damp, cold, and pressure augment it. When the nerve is inflamed there is tenderness on pressure over the course of the nerve. Pressure on localized points always develops pain, and the points that are most marked are on the lower end of the sacrum, on the side of the trochanter opposite the emergence of the great and small sciatic nerves, various points on the posterior aspect of the thigh, one at the head of the fibula, and one behind the outer ankle.

The disease is obstinate, and lasts for weeks or months. It interferes with locomotion, because of the distress which movements of the leg and foot occasion. It is a very rare disease in children. Generally it depends upon exposure to cold, or upon the rheumatic diathesis, or upon a neuralgic predisposition, or upon an irritation affecting the nerve before it leaves the pelvis, the result not unusually of sexual disorder, or of pressure from a gravid womb, or from an accumulation of *fæces* in the lower bowel. In many instances it is connected with gout, in others with anæmia, with syphilis, with disease of the hip-joints, and it may be, although it very rarely is, symptomatic of cerebral disease. Occasionally it is due to reflex excitation of the nerve. Sometimes it occurs after forced marches or long rides; probably in the majority of these cases, however, the sciatica is rheumatic. It is seldom double, except when of diabetic origin, or when due to compression from a growing tumor in the pelvis or from enlarging cancerous vertebrae.

Sciatica, when of long duration, leads to loss of motor power in the leg, to tingling, and to anæsthesia; and certain nutritive changes are observed in the limb, which is found to have decidedly dwindled, or there may be œdema. In many instances, it may be said truly in most, the disorder is the result of neuritis, and then there is generally much more, and persistent, tenderness,—in truth, in pure neuralgia there is not much,—and movement and position have but little influence on the pain. Further, the history of the case in pure neuralgia, the spontaneous pain,

the frequent anæmia, and the coexistence with other neuralgias, are very significant. Occasionally the neuritis ascends to the cord.

It is often a very essential matter to determine whether or not an effusion has taken place within the sheath of the nerve. In the main, what Fuller tells us is correct, that the presence of fluid within the nerve-sheath may be inferred when a patient who is suffering from sciatica complains of a dull aching or a benumbing pain in the limb, causing it to feel swollen, and when this sense of numbness and increased bulk has succeeded to pain of greater intensity, accompanied by cramps and startings and more or less inability to move the limb.

The disorders which are most likely to be confounded with sciatica are: *rheumatism of the muscles* and fibrous sheaths around the hip-joint; affections of the joint; and pains caused by irritation of the kidney. The first is very readily distinguished. It is generally, what sciatica is rarely, double-sided; and the pain is dull, diffuse, not paroxysmal, not limited to the sciatic nerve and its area of distribution, nor as much increased on pressure as that of sciatica. But, practically speaking, this kind of rheumatism is seldom seen unless associated with rheumatic inflammation of the sciatic nerve.

In *affections of the hip-joint* the suffering is increased by standing with the weight of the body thrown on the diseased leg. Moreover, the pain is usually limited to the hip- and knee-joints; does not descend in the course of the sciatic; is not associated with tenderness of the nerve; the aspect of the limb points to the disorganization that is going on; the leg shortens. Yet, before admitting this as a mark of difference, it must be ascertained by careful measurement; for, in consequence of muscular contractions, the affected limb in sciatica may appear to be shorter than it is. The main points of distinction between sciatica and a nervous affection of the hip-joint are the usual combination of the latter with hysteria, the very superficial tenderness, and the fact that the pain is apt to extend over the whole thigh.

*Irritation of the kidney* causes pain shooting down the thigh. The distress exists, however, in the course of the anterior crural nerve, is therefore not localized in the sciatic, is unattended with tenderness, but is accompanied by a frequent desire to pass water, and by other signs of disorder of the urinary functions.



*Sciatica* is sometimes *feigned*, especially by soldiers. But the copy is rarely a very accurate one. Impostors complain of pain on pressure and on motion, but are ignorant that the pain is prone to exacerbate after intervals of comparative quiet, and to increase in violence as night approaches. Their fancied torment is constant, but does not prevent them from sleeping; they wince when the muscles of the thigh are touched, yet, if their attention be diverted, the hand may be pressed along the sciatic nerve without any sign of tenderness being manifested.

## CHAPTER III.

### DISEASES OF THE UPPER AIR-PASSAGES.

#### SECTION I.

##### DISEASES OF THE NOSE AND ASSOCIATE ORGANS.

THE naso-pharynx, the larynx, and the trachea constitute the upper air-passages. The nose and naso-pharynx, or the space between the posterior nares and a horizontal line drawn through the lower end of the soft palate, first demand consideration. But as these disorders largely belong to a class which require surgical treatment, a brief review will be given in this place of those only that have features of medical interest. The *frontal sinuses* are in direct connection with the nasal chambers, and, in case of occlusion of their normal outlet, there is an accumulation of secretion, which may cause *headache*. This is likely to occur especially in the peculiar form of catarrhal inflammation attending influenza, in which the headache may continue for weeks until the inflammation subsides, or the patency of the outlet is restored. The headache may be associated with vertigo. Tumors may develop from the mucous membrane lining the frontal sinuses, of the same character as those of the nasal chambers; and the larvæ of insects, or mature forms, such as centipedes, find their way at times into these cavities and there cause pain and irritation.

*Frontal sinus diseases* are to be distinguished from *supra-orbital neuralgia*, *migraine*, and *cerebral disease* by careful inspection of the interior of the nose, examination with the curved probe to test the patency of the canal, by the presence of tenderness and other local signs of inflammation, such as swelling or discoloration, and by the history of the case.

The affections of the antrum Highmorianum, or maxillary

sinus, are similar to those of the frontal sinuses, and are more surgical than medical. Many of these cases are attended by a pain above the corresponding eye, and may be mistaken for *migraine* and the *neuralgia of frontal sinus disease*.

The diseases of the nasal chambers may be divided into acute and chronic. Among the former are coryza, acute rhinitis, glanders, hay-fever, hemorrhage or epistaxis, hydrorrhœa, mycosis, and abscess. Prominent among the second class are rhinitis, hypertrophic and atrophic, cirrhosis of mucous membrane, ethmoiditis, thickening and deviation of septum, rhinoscleroma, new growths, specific destruction by tuberculosis, syphilis, lupus, and malignant disease.

Another division might be made into local affections and those occurring in eruptive fevers and other acute diseases. For instance, in scarlatina, smallpox, typhoid fever, and diphtheria, swelling with increase of secretion and ulceration may occur, while in rheumatism and influenza changes are noted that may be regarded as almost characteristic. Many, probably most, of the instances of marked deviation of the septum and associated abnormalities are not evidences of disease, but are due to heredity. Bryson Delavan has found asymmetry in the nasal chambers to be the rule rather than the exception.

As regards the diagnosis of affections of the nose which are usually attended by mucous or purulent discharge and more or less obstructed breathing, it is well to bear in mind the possibility in children of there being a *foreign body in the nose*, and in older patients *rhinoliths* are sometimes detected upon rhinoscopic examination, of which Harrison Allen has reported examples. In one remarkable case described by Warren,\* the breech pin of a gun was discovered embedded in the right nasal fossa, where it had been driven by the explosion of a gun, several months previously, and its presence in the nose had not been suspected.

*Coryza*, or *acute catarrh*, is a general affection, which manifests itself by inflammation of the mucous membranes of the nose especially, but other mucous surfaces of the air-passages may be previously or subsequently affected. It often follows exposure to cold and dampness, and attacks those principally of lowered

---

\* Surg. Obs., Boston, 1867.

vitality. *Rhinitis* may be due to local irritation, as rough manipulation, operations upon the nose, or strong applications. It may be confined to one nostril. Upon inspection, the mucous surface is of a red color in certain locations, the turbinated erectile tissue is engorged, shows slight abrasions, and bleeds readily. The fossæ may be obstructed to a greater or less degree by inflammatory swelling of the mucous membrane, and the secretions vary in density from a clear serous fluid to a caseous or fibrinous exudation. To the latter the names of *rhinitis caseosa* and *rhinitis fibrinosa* have been applied. Cases with marked œdema have been reported under the title of *rhinitis œdematosa*.\* *Diphtheria of the nose* may occur independently, but is usually a concomitant or sequel of *faucial diphtheria*. At the present day the presence of the Klebs-Loeffler bacilli is relied upon to decide the diagnosis of diphtheria, although this micro-organism has occasionally been found in cases of other affections of the mouth, throat, and nose, which gave no clinical evidence of being diphtheritic. An acute catarrh is sometimes an early symptom of some specific disease, and, if attended by conjunctival inflammation, *measles* is suggested. Nasal catarrh accompanies *erysipelas*, and is one of the initial symptoms of *influenza*. Nasal catarrh may be also produced by the administration of remedies, as of the iodides, or in certain persons by the inhalation of drugs, such as *ipecacuanha*. At certain times of the year, when the air is filled with pollen from the *artemisia absinthifolium*, or ragweed, the *ailanthus* tree, or from roses or grasses, many persons suffer from what is called hay-fever, hay-asthma, rose-cold or hyperæsthetic rhinitis, which has for its most marked sign the reappearance of the symptoms upon the same day each year, suggesting a strong neurotic element. Daly, Roe, and Sajous have shown the dependence of many of these cases upon nasal abnormalities, the correction of which afforded comparative or complete relief.

*Nasal hydrorrhœa* is distinguished from *coryza* by the excessive flow of a serous fluid from the mucous membranes, especially over the turbinated bodies, which are pale and sodden. The affection is usually unilateral, but it may be bilateral. It happens at all seasons, thus distinguishing itself from hay-fever, though

---

\* J. C. Mulhall, Trans. Am. Laryng. Assoc., 1893.



like the latter it is found in neurotic subjects. It may occur in paralysis of the trifacial nerve, or as a result of head injury, and may be associated with polypi, or myxoma of the nose; it is at times of months' duration.\*

*Nasal hemorrhage* may indicate a general condition of the vascular system, a degeneration of the arterial coats, with or without increased tension. Occurring after the middle period of life, in a person otherwise apparently healthy, it suggests the likelihood of apoplexy or contracted kidney. It is important to distinguish between those cases in which the blood only passes through the nose, and those in which the blood comes from the nose. A rhinoscopic examination, both anterior and posterior, is essential to an exact diagnosis. In nasal hemorrhage, the blood very frequently comes from the septum low down, where it can be easily inspected.

*Post-nasal catarrh* has for its prominent symptom the dropping of mucus from the soft palate into the throat, and the expulsion at times, usually in the morning, of masses of gelatinous mucus or of hardened crusts from the naso-pharynx, giving rise to the unpleasant habit of hawking and spitting. In such cases rhinoscopic examination usually reveals inflammation of mucous membrane in the vault of the pharynx, and the glandular tissue, or so-called "pharyngeal tonsil," may be hypertrophied and form polypoid excrescences, or even adenoid tumors that may entirely occlude the posterior nasal openings. Digital exploration is a valuable means of diagnosis, especially in children. Deafness may be caused by occlusion of the Eustachian tubes, and mouth breathing is a necessary sequence. A form of post-nasal catarrh, of less severity, attends posterior hypertrophies and other abnormalities of the nose attended by increase of secretion. The diagnosis is made by rhinoscopy.

*Glanders* is manifested in the nose and throat by the evidences of inflammation of the mucous membrane, from which there is a free discharge of a viscid, muco-purulent or sanious fluid; pustules and spreading ulcers may appear on the surface of the body, farcy. The general symptoms are those of acute blood-poisoning, great prostration, and, in the human subject, a speedily fatal

---

\* As in a case reported by C. E. Bean before the Laryngological Association in 1891.

result. An erysipelatous rash appears on the nose, and spreads to the cheeks and forehead, vesicles soon appear upon the inflamed surface, intermingled with the characteristic pustules, which also occur on the mucous membrane of the throat and nasal passages; ulceration and gangrene follow. As the larynx becomes invaded, the voice is lost and the muco-purulent expectoration becomes more abundant and more bloody. Though the typhoid condition is a marked feature, the case cannot be mistaken for one of *typhoid fever*, on account of the absence of the rose-spots and of the characteristic temperature record. Glanders bears a close resemblance to *pyæmia*, but its pus-collections are external, the rigors are less marked and may be entirely absent, and it does not have the history of traumatism or surgical operation. There is no danger of confounding glanders with *purulent* or *serofulous rhinitis* in which there is no fever or constitutional disturbance. A very severe purulent rhinitis of acute form may be caused by accidental *gonorrhæal infection of the nose*; the history of the case gives the explanation.

The chronic forms of rhinitis attended by catarrhal thickening of the septum or of the turbinate bodies, atrophy or cirrhosis of the mucous membrane, and the development of cysts, polypi, or papillary fibromata, are recognized by careful rhinoscopic examination, and belong to surgery rather than to medicine. In former times ozena was regarded as a disease, but it now is recognized as an attendant upon chronic atrophic rhinitis, in which the secretions dry into crusts which undergo putrefactive changes, and thus produce the offensive odor. In cirrhosis there are contraction of the mucous membrane and evidences of atrophy, without decided catarrhal symptoms. Hypertrophy of the mucous membrane is considered as a preliminary stage to atrophy; and the production of polypi, or of myxoma, or, more rarely, of fibroma, is not an uncommon result. Hypertrophies are distinguished from new growths by their situation, color, density, and immobility; new growths being in abnormal situations, of peculiar color, and pedunculated; for instance, *polypi* are white and glistening, *cysts* are white, *papilloma* are rare and may look like a small bunch of grapes.\* The diagnosis of papillary hyper-

---

\* Mulhall's case, Proc. Amer. Laryng. Asso., New York, 1890, p. 54.

trophy from true papilloma, papillary fibroma, depends upon its location, appearance, and the microscopic details, which also distinguish the latter from epithelioma and sarcoma. The distinction of myxoma from cysts is given further on.

In making the diagnosis between benign and malignant growths, the rapid development and general appearances of the latter are usually depended upon as conclusive. At the same time the difficulty is enhanced by the danger of benign growths becoming transformed in their character. Bosworth reports a case in which sarcoma developed after polypi had been operated upon rather harshly by means of forceps, and a similar case is narrated by Heyman.\* Traumatism has been also observed to result in fibro-sarcoma and other fatal malignant growths of the nasal chambers.

The reflex disorders arising from naso-pharyngeal obstruction, by hypertrophies or new growths, have been studied by a multitude of clinical observers. Obstinate headache, asthenopia, earache, persistent cough, and vertigo are symptoms of special cases of nasal disorder. Voltolini first directed attention to the fact that nasal polypi may be the cause of *asthma*, and a deflected septum and hypertrophic rhinitis are also claimed as causes by Bosworth. Weber showed that diseases in the upper air-passages readily produce turgescence and swelling of the bronchial mucous membrane. Glycosuria has been known to originate in nasal obstruction, and to disappear when this was removed.†

*Nasal polypi* apparently may cause spasmodic stricture and difficulty in urinating, as in a case reported by Mulhall;‡ the spasmodic stricture was at once relieved by the removal of the nasal polypi.

*Rhinoscleroma* is a form of new growth allied to round-celled sarcoma, first observed by Hebra, characterized by the appearance of flat, slightly raised patches which are smooth on the surface and of ivory-like hardness, and which first appear at the edges of the nostril and spread to the upper lip. It is essentially due to infiltration of the lower layer of the integument by small round hyaline cells, which, Cornil tells us, do not contain micrococci. The skin is not discolored; the patches are slightly tender, but not painful, and they do not ulcerate. The lesion may

---

\* Revue Mensuelle de Laryngologie, 1888, p. 24.

† Bayer, Revue de Laryng., 1894, xv. 19.

‡ American Laryngological Association, 1892, p. 42.

gradually extend into the tissues of the mouth and nose and to the larynx, producing alteration of the voice and narrowing of the glottis. Rhinoscleroma is to be distinguished from syphilis, epithelial cancer, and keloid. It differs from venereal disease, according to Morell Mackenzie, mainly in its very chronic course, the absence of softening or ulceration, and its absolute intractability under every kind of medication. From epithelioma it can be discriminated by its smooth glistening surface, its hardness, the absence of bleeding or ulceration, and its persistently local character. The history and cause of the case and its general appearance serve to distinguish it from keloid, which has the puckered, white, and irregular outline of scar-tissue.

---

## SECTION II.

### DISEASES OF THE LARYNX AND TRACHEA.

Of these affections those of the larynx are far the most frequent and the most readily recognized. There are, indeed, symptoms in laryngeal diseases which at once direct attention to the seat of the malady. The larynx is the organ of speech: hence changes in the *voice* constitute the most striking manifestations of disorder. These changes vary in degree. The voice may be merely hoarse, or completely lost. In young children the different tone of the cry corresponds to the altered voice of adults. The alteration of the voice depends almost wholly upon an affection of the vocal cords, and this may be organic, such as inflammation, œdema, ulceration, cicatrices, and morbid growths; or it may proceed from perverted or impaired innervation. Very often the hoarseness or loss of voice is caused by diminished tension and want of certain and prompt action of the vocal cords, whether connected with structural change or not. The same cause gives rise, for the most part, to the modifications of the voice which show themselves as huskiness in speaking, or in the loss of certain notes in singing.

Next to the voice in diagnostic importance stand the character of the breathing and the cough. The *breathing* is labored and difficult, and is frequently perceived to be noisy, and coarse or shrill,—the so-called *laryngeal stridor*: a sign encountered whenever the orifice through which the air has to pass is narrowed,



either temporarily by a spasm, or more permanently by any state which gives rise to a constriction of the parts ; for instance, by swelling of the mucous membrane, or diphtheritic deposit.

The difficulty in breathing is in some diseases slight ; in others great. One of the peculiarities of laryngeal dyspnœa is its tendency to recur in paroxysms, during which the patient appears to be in imminent danger of strangling. These fits of suffocation are produced mostly by a spasm of the intrinsic muscles, particularly the adductors of the larynx. They occur in pure spasm of the glottis ; in croup ; in œdema of the larynx ; in ulceration and in polypi of the larynx.

The *cough* of laryngeal affections presents frequently the same peculiarity as the dyspnœa,—it happens in paroxysms. Another peculiarity, although not one so constant, is its harsh and ringing tone. The cough is often short and dry ; sometimes it is followed by a muco-purulent expectoration of roundish shape, or by a blood-streaked sputum, or by the spitting up of false membrane. It is readily excited by the act of swallowing, its seat is referred by the patient himself to the windpipe, and it is apt to be especially troublesome at night.

*Pain* is not so usual a symptom of laryngeal disease as either cough or changed breathing. In chronic affections it may be, indeed, wanting. It is rarely severe ; often more a sensation of tickling, of burning, or of uneasiness than of actual pain. It is apt to extend down the trachea to the upper part of the sternum. Sometimes it is increased on pressure, as in acute laryngitis and in ulceration of the mucous membrane ; and it may be also augmented by the act of swallowing.


By the symptoms, then, of altered voice, cough, dyspnœa, and, in some cases, of local pain and difficulty in deglutition, we recognize a laryngeal affection. The stethoscope is occasionally of service ; yet, on the whole, it furnishes little information. But inspection of the larynx by the aid of the *laryngoscope* has revolutionized our knowledge of laryngeal diseases. Indeed, any one who attempts a positive diagnosis of laryngeal diseases without the laryngoscope attempts to do without the only means which renders the diagnosis at all trustworthy, and is guilty of neglect. The instrument consists of a small mirror fixed on a long stem. It may be either circular, square, or oval. The circular mirror

occasions least irritation. It may vary in size from half an inch to an inch and a quarter in diameter. The larger the mirror we can employ, the better is the image.

The mirror is in some cases all that is necessary to practise laryngoscopy. It is heated in warm water or over a lamp, and then introduced into the back of the mouth in the manner presently to be described; the person to be examined having been placed with his face toward the sunlight, so that its rays may strike the laryngeal mirror.

But examinations by direct light are practicable only on some days and at certain periods of the day. Usually we require a second mirror to illuminate the throat and the laryngoscope. This mirror, when sunlight is employed, has a plane surface; when artificial light is used, it is better that the reflector be slightly concave. One of circular form, about three inches and a half in diameter, and with a focus of from ten to fourteen inches, answers best. It may be either attached to the head by means of a band, or worn on a pair of spectacle-frames, or placed on a movable stand, or affixed to a lamp. The spectacle attachment introduced by Semeleder is convenient. When this or the frontal band is made use of, the observer may either place the mirror opposite to one of his eyes, and look through the central perforation, or adopt the easier method of wearing the reflector on his forehead.


FIG. 13.



Laryngoscopes of various shape;  
not quite natural size.

The French have recourse for the most part to lenses, and concentrate the light directly into the throat. But a better arrangement is obtained in Mackenzie's rack-movement bracket and bull's eye condenser; or by a combination of lenses attached to a metallic frame which is fastened to a lamp, as in the well-known apparatus of Tobold. The best light to employ is coal oil; the most convenient, an argand gas-burner. The incandescent gas-burner yields a steady, white, and brilliant light which is especially favorable for office use. I have used the electric light very satisfactorily. A portable electric light is obtained by the aid of a small storage battery, which can be used at the bedside.

FIG. 14.



Laryngoscopic examination, as made with the means ordinarily employed.


To examine the larynx by artificial light, we should proceed thus. The patient, sitting in an upright position, with his head inclined slightly backward, is placed near a lamp, burning with a

steady, brilliant light, and the flame of which is behind and about on a level with his eyes. He is directed to open his mouth widely, to put out his tongue, and to hold between two fingers its point enveloped in a soft napkin or handkerchief. If he cannot accomplish this readily, the observer must hold the protruded tongue, or a tongue-depressor must be employed. The observer now seats himself directly in front of the patient, and nearly a foot from the mouth. Putting on his spectacles or frontal band, he throws a disk of light into the back part of the mouth; he then rapidly introduces the laryngeal mirror, previously heated in warm water or over a lamp and its proper temperature ascertained by touching his own hand or cheek. The mirror, great care being taken not to bring it in contact with the tongue, is placed with its back against the uvula, which, with the soft palate, is pressed backward and upward; the lower surface of the laryngoscope should be firmly applied to, or, if this be found to occasion too much irritation, should be held near, the posterior wall of the pharynx. The inclination of the mirror varies with the position of the patient and the parts we wish the more particularly to explore. As a general rule, it may rest at an angle of about  $45^{\circ}$ .

When the mirror is stationary, as for instance in the Tobold laryngoscopic lamp,—a less portable but far easier mode of illuminating,—the reflector is attached to the lamp by a freely movable brass rod, and the light is thus thrown into the mouth, leaving

the examiner unembarrassed. When the mirror has been introduced in the manner described, the laryngeal image is readily perceived. We see the epiglottis, the glottis, the cartilages, the true vocal cords, the superior thyro-arytenoid ligaments or ventricular bands, and in some cases even the rings of the trachea. We may be able to discern each portion

FIG. 15.



Laryngeal image, as seen in the laryngoscope under favorable circumstances.

of the laryngeal aperture with distinctness, or it may take several examinations to do so.

In health, the color of the various parts is very different.



Stoerck has well described it in likening that of the epiglottis, the interior of the larynx below the glottis, and of the cricoid cartilage, to the coloration of the conjunctiva of the eyelid; and the hue of the aryepiglottidean folds and the prominences of the arytenoid cartilages to that of the gums. The mucous membrane of the trachea between the rings is of a pale pink color; the vocal cords have a white, glistening look. Mackenzie takes special notice of the whole of the under surface of the epiglottis being in some cases of a bright-red hue; and Gibb points out that in negroes the cartilages of Wrisberg have a yellowish tinge.

The laryngeal image in the mirror bears this relation to the real position of the parts: the right vocal cord of the person who is examined is seen on the left side of the mirror, and the left vocal cord on the right; or, to state the matter in a form easy to be remembered, the cord which corresponds to the right hand of the patient is the right, that seen toward his left hand is the left. The epiglottis appears in the laryngoscope at the upper portion and behind; so do the other structures that lie in front. The arytenoid cartilages show at its lower portion, and toward the front.

To judge of the movements of the vocal cords, we tell the patient alternately to inspire deeply and to utter, as a prolonged high note, a sound like "ah." During this the vocal cords are closely approximated and stretched, and the epiglottis, in fact the whole larynx, is somewhat elevated; while during a full inspiration the cords are far apart, and hence the glottis is wide open. To obtain a satisfactory sight of the deeper-seated parts, we must bear in mind that the more horizontally the surface of the mirror is placed, the more distinctly they come into view. For the exploration of these structures, and particularly of the trachea, the light must be thrown from below upward upon the laryngoscope. To elevate the larynx decidedly, and especially to bring the epiglottis fully into view, the patient should in a high pitch pronounce *ee* as in the word *see*.

In some, laryngoscopy is easy; a conclusive examination may be made at the first attempt. In others, a course of training is required to subdue the sensibility of the fauces, which may be general, or be limited to a very small spot. As a means of overcoming the difficulty, sucking small pieces of ice, or the previous

administration of bromide of potassium, or the local use of a solution of cocaine from two to five per cent., is useful. But the best means is skill in the use of the instrument,—its rapid and decisive handling. The administration of an anæsthetic may be, however, necessary. This is especially the case in refractory children suffering with papilloma or other conditions demanding ocular inspection for diagnosis or treatment. To overcome pharyngeal and laryngeal reflexes, Seanes Spicer\* recommends the cautious use of a ten per cent. spray of solution of cocaine, and, for removal of salivary secretions, the free use of dry mops of absorbent cotton-wool. Five assistants are required: one to give chloroform, maintaining moderate general anæsthesia; the second holds the child in the usual position for laryngoscopy; the third stands behind the child, keeping its head erect and square to the operator; the fourth stands behind and to his right, holding the mouth-gag; the fifth holds the tongue out with forceps during the operation.

In some persons with very irritable throats, I have obtained good views by pressing the instrument against the roof of the mouth, instead of passing it back into the pharynx, and by altering the position of the head a little, tilting it more backward. The epiglottis, and the structures at the entrance of the windpipe, are thus readily enough brought into view: with the deeper parts we do not succeed so well; but in many cases we get sufficient guide for topical applications. There are further obstacles, such as a rising up of the tongue, greatly-enlarged tonsils, a long uvula, a pendent epiglottis, all of which at times interfere with our investigations. But in any case we should not endeavor to make the view more satisfactory by constantly altering the position of the mirror. It is better to introduce it repeatedly, than to shift it often when introduced, or to keep it for any length of time in the patient's mouth. Digital examination of the larynx with the index finger is an expedient of value in children and others who will not permit laryngoscopy. It is practised for diagnostic purposes in laryngeal œdema, and in new growths, such as papilloma.

If the mirror be passed behind the uvula, and the reflecting surface directed upward, the posterior nares may be examined.

---

\* Journal of Laryngology, Rhinology, and Otology, London, Oct. 1894.

To practise *rhinoscopy*, however, the mirror should be small and fixed to the shaft at a right angle. The patient is directed to keep his head erect, or bend it slightly forward, and while his mouth is wide open a strong light is thrown to the back of the throat. But before the rhinal mirror is placed in position, a tongue-depressor is applied, with which the back of the tongue is well pressed down, and which may be given to the patient to hold. Yet a difficulty remains,—namely, to get the uvula out of the way. This is not easily accomplished without a palate-hook, by which means the uvula, with a portion of the soft palate, is gently drawn forward and upward, the handle of the hook being held to one side of the mouth: Voltolini's palate-hook widens the pharyngo-nasal space satisfactorily, or Sajous' soft palate-elevator may be employed. But by instructing the patient to breathe through the nose and to breathe heavily while the mouth is open, we obtain relaxation of the muscles of the soft palate, and in most cases, after a little training, may dispense for diagnostic purposes with the palate retractor. The mirror, with its reflecting surface upward, is now passed along the tongue-depressor, until it reaches the posterior wall of the pharynx. By then raising somewhat the handle of the mirror, we obtain a view of the septum; and by slanting the mirror first toward one side and then toward the other, the posterior nares and the orifices of the Eustachian tubes may be inspected.

The art of rhinoscopy is more difficult than that of laryngoscopy, and, though the rhinal mirror aids us in detecting morbid appearances which would otherwise escape observation, it does so neither as readily nor as completely as the laryngoscope.

Let us now look at the chief diseases of the larynx. Grouped in accordance with their main features, and without classifying them in strict obedience to laryngoscopic inquiries, they may be arranged as follows:

#### ACUTE ORGANIC DISEASES.

Congestion, or hyperæmia.

Inflammation of the mucous membrane of the larynx—Acute laryngitis.

Œdema of the larynx.

Acute affections of the larynx and trachea as met with in children.	}	Spasmodic and pseudomembranous laryngitis —False and true croup.
---	---	---

Specific affections—Syphilis, tuberculosis, lepra, diphtheria, erysipelas, typhoid, etc.

## CHRONIC ORGANIC DISEASES.

Inflammation of the mucous membrane of a part, or of the whole—Chronic laryngitis in its various forms—Abscess.

Destruction of the cartilages.

Growths and tumors of various kinds.

Ulcers, simple and specific.

## AFFECTIONS OF THE NERVES.

Spasm of the larynx. (Laryngismus stridulus.)

Nervous aphonia. { Functional, or purely nervous aphonia.  
{ Paralysis of the muscles of the vocal cords.

Chorea of the larynx.

## Acute Laryngeal Affections.

**Acute Laryngitis.**—In its mild form, acute laryngitis is neither an uncommon nor a dangerous disease. In its severe form it is much more uncommon, and very much more dangerous. When it is slight, it occasions simply hoarseness; a feeling of tickling and irritation in or near the larynx; a trifling, though annoying, cough, or rather a constant disposition to clear the throat, more than a cough; and, owing in a great measure to a coexisting inflammation of the fauces, some difficulty in swallowing. The disorder passes off in the course of a few days.

When the inflammation is violent, and especially when it involves the submucous tissues, the symptoms are much aggravated, and life is in peril. The respiration becomes seriously impeded; with each breath a wheezing or whistling noise is heard. There is but little expectoration; and the cough is distressing and painful, and has a harsh sound. The voice is hoarse, or sinks into a scarcely audible whisper. The patient knows the seat of his disease: he feels that it lies in the windpipe, and complains of this being tender when pressed, and of a feeling of constriction in the throat. There is difficulty in swallowing, and fever, with a full pulse and flushed face. If the case advance unchecked, the countenance becomes distressed and pale, the lips bluish, the pulse irregular, and death sets in with all the signs of deficient aeration of the blood and of strangulation.

The disease in its graver form runs a very rapid course. If in a few days after its commencement no improvement show itself,



life does not last long. Sometimes death takes place on the first day of the attack. It rarely waits for the sixth. (Edema of the laryngeal mucous membrane is often the consequence of the inflammation and the cause of the danger.

The marked symptoms of the perilous complaint prevent it from being overlooked, and render its discrimination easy. There is fever with dyspnœa in the *acute pulmonary affections*; but the voice remains unaltered, and they exhibit physical signs which acute laryngitis does not,—they show rales, or abnormal respiration-sounds; while in laryngitis the murmur of the lungs is that of health, although it is sometimes enfeebled by the impediment in breathing, or obscured by the shrill sound which issues from the larynx. We find difficulty in swallowing and some hinderance in breathing in *tonsillitis*; but inspection of the oral cavity immediately detects the source of the disorder. There is difficulty in swallowing in *pharyngitis*, but there is not embarrassed breathing, or a peculiar voice, or cough, and the fauces appear dusky and injected, while they are but slightly affected in laryngitis, unless the inflammation of the larynx have supervened upon that of the throat. *Croup* resembles acute idiopathic laryngitis most nearly; but it is as rare in the adult as acute laryngitis is in the child, and, as we shall presently see, obvious differences in the symptoms exist.

An affection of the larynx cooccurring only in winter, *laryngitis hiemalis*, has been described by Mulhall, in which the secretions form adhesive crusts, producing difficulty in speaking, or more often aphonia. This is to be diagnosticated from *laryngitis sicca*, which is a part of a general process, and follows pharyngitis sicca and atrophic rhinitis.

There is a peculiar form of inflammation of the larynx, *diffuse cellular laryngitis*, a diffuse inflammation of the cellular tissue, with lymph or pus infiltrated in the submucous tissue, to which attention has been called by Henry Gray.\* It is a formidable affection, which bears a strong likeness to erysipelatous laryngitis, but, what is not by any means constantly the case in this disorder, the symptoms begin in the fauces and larynx; and, wholly unlike erysipelatous laryngitis, the submucous tissue is primarily attacked,

---

\* Holmes's System of Surgery, vol. iv.

and the neck becomes greatly swollen from the effused products around the larynx, trachea, and œsophagus filling its cellular tissue. The disease begins with chills, soreness of throat, and fever, soon succeeded by a hacking cough, by dyspnoea, by a dusky hue of the fauces, by enlargement of the tonsils and of the glands in the neighborhood of the jaw, and by great difficulty in swallowing. As the complaint proceeds, the neck increases greatly in size, the fever assumes a low type, and the patient either sinks gradually or dies asphyxiated, perishing sometimes rapidly from a speedy increase of the laryngeal tumefaction.

A form of œdema of the laryngeal mucous membrane exists, due to submucous effusion of a myxoid substance which occurs in the course of influenza, and to which Seiler has applied the term *myxoid œdema*; there are aphonia, swelling, and severe localized pain. The diagnosis is chiefly made by considering the history of the case, and the symptoms, which are not so acute as in ordinary œdema.

Other forms of inflammation of the larynx to which attention has of late years been called are *hemorrhagic laryngitis*, an acute catarrh of the larynx, attended by bleeding from the inflamed membrane, and *laryngeal rheumatism*. This generally happens in persons of rheumatic diathesis, is attended with considerable pain, and may or may not be associated with other signs of rheumatism.\* There are cases in which laryngeal symptoms are marked, and cases without them. Roos reports† several instances of rheumatic angina that terminated in attacks of general rheumatic arthritis. The principal features of rheumatic angina are excessively painful deglutition, redness and swelling of one or both tonsils; the disease is of slow development, and occurs with or without abscess-formation. It has been suggested that the joint affections are really secondary manifestations, pseudo-rheumatic in character, and that the polyarthritis belongs to the category of attenuated pyæmic infections. H. L. Wagner has found articular rheumatic affections following follicular amygdalitis, in which bacterial investigation showed that the synovial fluid obtained by tapping the joint

---

\* Archambault, Thèse de Paris, 1886.

† Revue de Laryngologie, etc., 1895.

contained the same micro-organisms as were found in the diseased tonsil.\*

Following inflammation or ulceration of the larynx, various irregularities may occur as the result of cicatricial contraction, or adhesions between the cords, which may be studied with the aid of the laryngoscope. There are usually alterations in the voice, with attacks of dyspnœa simulating asthma, and impairment of general nutrition.

**Œdema of the Larynx.**—The danger of acute laryngitis of any kind is much aggravated by the precise seat of the disease. When the inflammation takes place immediately around the glottis, and causes a serous fluid to transude, *œdematous laryngitis*, the peril is greatly increased. The inspiration is audible, noisy, hissing, and labored; there is a distressing sensation of constriction or obstruction in the windpipe, and the patient makes repeated efforts, by swallowing or by hawking, to clear his throat of the substance which seems to be clogging it. His difficulty of breathing is intense, and occurs in frightful paroxysms, sometimes of a quarter of an hour's duration, in which strangulation appears to be imminent; and, indeed, often he does perish by strangulation.

This grave disease, *œdema of the larynx*, sometimes follows an extension of the peculiar inflammation of the throat in the exanthemata, or is of erysipelatous origin, and it occasions death quickly, and amidst great suffering. But the œdema may arise without preceding acute inflammation, whether this be specific or not. It may result from long-continued pressure on the trachea or larynx, or in exceptional instances occur in connection with Bright's disease. Again, an effusion of serum may cause death suddenly in a person who has been laboring under a chronic laryngeal disorder.† Such cases of œdema of the larynx are distinguished from those produced by active laryngeal inflammation by the absence of fever, of local tenderness, and of marked difficulty of deglutition. It is true that, if the œdematous affection

---

\* Rheumatic Affections of the Body due to Tonsillar Disease, Trans. Amer. Laryngol. Assoc., 1894.

† As in tubercular laryngitis, which may be complicated both with acute and, more frequently, with chronic œdema. See an interesting paper on the connection in Archives de Physiologie, No. 6, 1882.

ensue upon a chronic inflammation of the larynx, tenderness and an impediment in swallowing may be observed. But the history of the malady and the non-existence of fever leave little room for error.

The diagnostic sign proposed for œdema of the larynx—the swelling of the epiglottis, as ascertained by the touch—cannot be relied upon, because this swelling does not always exist to an obvious degree, and, even when it does exist, is not readily determined by the finger. In the acute cases of œdematous laryngitis the laryngoscope shows a bright-red mucous membrane; sometimes the tumid epiglottis presents the appearance of two round red swellings. It is generally erect and tense. The œdema, in rare instances, may be altogether below the glottis.

**Croup.**—Croup is inflammation of the larynx and trachea; but it is something more. It is a spasmodic action of the muscles of the larynx, which spasmodic action gives rise to much of the peculiar cough, the stridor, and the paroxysms of dyspnoea, so characteristic of the disease. As croup is thus an affection composed, as it were, of several distinct elements, it differs somewhat according as one or the other of these elements preponderates. Thus, the inflammation may be comparatively slight, yet the spasm play a very prominent part; or the inflammation may be very severe, and result in the formation of a false membrane. To the first class belongs the disorder known as false croup, catarrhal croup, spasmodic croup, spasmodic laryngitis; to the second, the true or membranous croup, and diphtheritic croup.

*False or catarrhal croup.*—This is one of the most common diseases of childhood. Its seizures happen chiefly at night; and the child that has gone to bed well, or perhaps fretful from teething, or with a slight catarrh, wakes up suddenly in a state of alarm, breathing with difficulty. It coughs with violence at short intervals, and the cough is noticed to be loud and ringing and hoarse; and so are the voice and the cry. Each inspiration is attended with that shrill, “croupy” sound which, once heard, is never forgotten. The face is flushed, the pulse frequent, and the temperature but little above the normal. The paroxysm continues in this manner for about an hour; the breathing then becomes quiet, the child falls asleep, and rests well until toward



morning, when the attack is apt to be renewed. The little patient may, however, escape this altogether, and keep well; or else the paroxysm recurs the next night, or for several nights in succession. In the intervals the voice and respiration are natural, there is little or no fever, little or no cough. Yet sometimes a cough occurs, during the day, which has every now and then a croupal sound; the voice, too, is slightly hoarse.

Catarrhal, or false croup, most frequently follows exposure. It is very rarely fatal. The laryngoscope shows marked congestion with swelling of the mucous membrane and copious muco-purulent secretion. Cases in which the inflammation is extensive and severe, without having led to a plastic exudation, and in which the inflammation is apt to be chiefly subglottic, approach in their persistency and in the character of their symptoms very closely to true croup. Indeed, one form of the complaint may run into the other, warranting the assumption that they are not two diseases, but only two forms of the same disease. Spasmodic croup may be a symptom of abnormalities, such as of hypertrophies or of adenoid growths in the nose or pharynx, and, if persistent, should suggest a digital or rhinoscopic examination.

The main element in the production of the symptoms of false croup is undoubtedly *spasm of the glottis*. But *laryngismus stridulus*, as laryngeal spasm or spasm of the glottis is called by many, is a neurosis which, while it may complicate any affection of the larynx and trachea, may also exist independently, from central, or direct, or reflex, causes of irritation. The laryngeal spasm may, therefore, form a distinct disorder, which differs from catarrhal croup by the absence of all inflammation and by several circumstances which proclaim its non-identity, such as its usual connection with rickets, and its frequent association with other convulsive symptoms,—with distortion of the face, rolling up of the eyes, spasmodic contraction of the hands and feet, and general convulsions. Laryngismus and tetany are often associated; indeed, by many laryngismus is looked upon as the laryngeal expression of tetany. The Trousseau sign of tetany—pressure upon the large arteries and nerves of a limb developing a paroxysm of tetany—is said to be never absent in the laryngo-spasm.\*

---

\* Escherich, address before the Tenth International Congress.

Some cases of supposed purely nervous laryngeal spasm in children are undoubtedly symptomatic of laryngeal growths, or of paralysis of intrinsic muscles, and are really attacks of dyspnoea due to laryngeal obstruction. Laryngoscopic examination should be made in severe cases, even though an anæsthetic be required, as in Scanes Spicer's method. Laryngismus stridulus is an affection of children under two years of age. Crying may bring on the attacks, the child dying of suffocation or during convulsions. In some cases mentioned by Mackenzie, the attack assumes the form of a sudden, almost soundless, spasm that does not relax until life is extinct.

As in croup, the seizures are apt to take place at night. Generally the child has been fretful from teething, or from gastric or intestinal irritation, when suddenly an attack of difficult breathing occurs, accompanied by several loud, crowing inspirations, and by an appearance of the most manifest distress and of threatening suffocation; yet the paroxysm is not associated either with cough, or with fever, or with an altered voice or a materially changed cry. A fit of this kind may be repeated twenty or thirty times a day. It may terminate fatally in a short time; usually, however, the paroxysms are spread over weeks, or even over a longer period.

In addition to the frequent combination with other convulsive symptoms, the protracted duration of the disease, and the absence of febrile disturbance, of hoarseness, and of cough, point out the distinction between laryngeal spasm and catarrhal or spasmodic laryngitis. From bilateral palsy of the vibrators of the glottis, laryngismus is readily distinguished by the great and persistent difficulty of breathing in this affection. Laryngeal spasm also occurs in the laryngeal crises of tabes; the absent knee-jerk and the ataxia tell us its meaning.

In the adult, glottic spasm produces symptoms to which the name of *laryngeal vertigo* has been given; the attack comes on suddenly, the patient gasps for breath and becomes unconscious and asphyxiated. In such cases attendant disease of the pharynx often affords the explanation of the symptom.

*True or membranous croup.*—True croup is a formidable affection, in which there is inflammation that results in the formation of a false membrane. The plastic exudation is found lining the

larynx, extending at times into the trachea or down into the bronchial tubes.

The symptoms of this dangerous malady are : the same brazen cough, the same stridulous breathing, as in false croup, a decided change in the voice, dyspnœa, and fever. But all these symptoms do not show themselves at once, and generally only attain their intensity after ordinary catarrhal croup has existed for a few days. High fever—the temperature ranging from  $103^{\circ}$  to  $104^{\circ}$ , or upwards—and difficulty in breathing are manifest, and, although they exacerbate and remit, only cease when the disease ceases. There is much thirst, no appetite ; but what is taken is readily enough swallowed. The voice, changed almost from the onset, becomes hoarse and whispering, and, as the disease advances, often totally suppressed.

The child remains in this condition for several days : restless, with its head thrown back, its respiration labored, and the croupal sound never completely disappearing. The great efforts at getting air into the lungs through the obstructed windpipe give rise to inspiratory recessions, seen at the supra-sternal notch, above the clavicles, and at the epigastrium. Sometimes solid masses of membrane are coughed up. Finally, the cough stops altogether ; the intervals between the paroxysms of dyspnœa are effaced ; the countenance becomes livid ; the skin loses its sensibility ; the extremities grow cold ; and, unless relief be afforded, either by medicinal means or by an operation, the little sufferer dies comatose or suffocated. The fatal termination is not unfrequently hastened by an intervening attack of bronchitis or of pneumonia, —a fact which teaches us not to neglect examining the lungs in cases of croup, so as to be sure that no disease is there silently running its course with its symptoms masked by the tracheal malady.

The application of a stethoscope to the larynx or trachea does not give us much information as to the exact seat and the extent of the affection of the windpipe. Still it is not without value. It may enable us to judge of the position of the exudation, for we may occasionally hear a vibrating sound, as if a membrane were being tossed to and fro by a current of air. In a case that came under my notice some years ago, this sign was perceived with great distinctness at the lower part of the trachea and toward the

commencement of the left bronchial tube ; and, at the autopsy, at exactly this point was found a thick layer of membrane lying unattached in the tube. Laryngoscopic examinations in croup are difficult of accomplishment ; but, when successful, they demonstrate deep-red infiltrated subglottic tissue with muco-purulent incrustations ; or exudation upon and immobility of the vocal cords, and the arytenoid cartilages held together by false membrane in the interarytenoid space.\* Auscultation of the lungs, by showing to what extent the air is still capable of entering them, furnishes us with a clue to the degree of the laryngeal obstruction.

Membranous croup is a disease not apt to be mistaken. Yet we must be cautious not to attach too much weight to any one of the symptoms ; we ought rather to judge of the existence of the disorder by their grouping. But when we take the symptoms collectively,—the ringing cough, the peculiar respiration, the dyspnœa aggravated in paroxysms, the changed voice, the fever, the expectoration ; when we regard the comparatively short duration of the disease,—there is, with the exception of the ever-present question of the diphtheritic origin, but one interpretation of the phenomena possible.

It is, of course, of the utmost consequence to distinguish between spasmodic laryngitis or *false croup* and membranous croup. The main difference consists in this : in the former, the invasion is usually more sudden ; there is little fever, or this disappears with the paroxysm ; and so do the croupal breathing, and, to a great extent, the hoarse voice and the loud, barking cough. The disorder lasts rarely more than two or three days, the attack usually occurring at night ; whereas in true croup the duration is seldom less than from four to six days, the disease progresses steadily, and the voice and respiration show at all times the nature of the affection. Then in the latter we may find expectoration of false membrane. This is, indeed, the most absolute proof ; yet the absence of membrane in what is coughed up or vomited is not a positive sign that the case is not one of membranous croup. The membrane may be retained in the larynx ; and we meet, indeed, with instances in which it is impossible without a laryngo-

---

\* Pieniazek, Arch. f. Kinderh., x. 5.



scopic examination to say whether the inflammation has or has not produced a plastic exudation; whether, in other words, the case is a severe one of false croup, or one of membranous croup.

The disorders, excluding diphtherias, which, next to false croup, are most likely to be mistaken for the formidable malady under consideration, are: acute laryngitis, œdema of the larynx, retropharyngeal and retrolaryngeal abscesses.

*Acute laryngitis* in its ordinary form, such as we see in adults, is a very rare disease in children. Acute catarrhal laryngitis is in them closely connected with the phenomena of spasmodic croup; and the croupy symptoms, the changed voice, the barking cough, the paroxysmal dyspnœa, the slight or absent difficulty in swallowing, tell us what we are dealing with. In membranous croup these signs also are intensified, and we are apt to have high fever. A form of laryngitis, however, happens in children, which is very liable to be considered as croup: it is the *secondary laryngitis of the exanthemata*, especially of variola. Attention to the history of the case, and to the circumstance of the inflammation having spread from the throat downward, will go a great way toward forming a correct opinion of the disease. Yet the diagnosis is sometimes one of extreme difficulty.

*Edema of the larynx* resembles croup, in its severe or its membranous form, in the dyspnœa, the fits of suffocation and of coughing, the altered voice, and the noisy inspiration. It resembles it further in the fact that most of the symptoms do not disappear in the intervals between the paroxysms. Here is certainly a strong likeness. But the cough has not the croupal, brazen sound; expiration is comparatively unembarrassed; there is no fever, unless the œdema occur in the course of an acute affection; and, above all, œdema of the glottis is a disease of adults. Again, the history of the case often guards against error, for œdema of the larynx happens frequently, perhaps most frequently, in those who have been long laboring under chronic, or ulcerative laryngitis; it is also seen from the toxic effects of iodide of potassium. In cases in which we are able to use the laryngeal mirror, the peculiar œdematous look of the parts is readily recognized.

*Retropharyngeal abscesses* share with croup the dyspnœa, the

stridulous respiration, and the altered voice. They do not, however, share with it the peculiar cough ; and, further, in croup there is not the difficulty in swallowing, or the evident tumefaction and stiffness of the neck, nor can a tumor be recognized by the touch, as it can be when an abscess is seated behind the walls of the pharynx. Moreover, the dyspnœa and the voice present somewhat different characteristics. In the case of abscess, the former is greatly augmented or paroxysms of it are brought on by attempts at deglutition ; it is always preceded by dysphagia, is increased by pressure against the larynx, and is aggravated by the horizontal position. In croup, the patient seeks relief by throwing his head back, and, although he loses his voice and speaks in a hardly audible whisper, still the words are sufficiently distinct ; while an abscess gives a nasal or guttural tone to the voice, that makes it impossible to understand what is being said.

*Retrolaryngeal abscesses* following inflammation of the areolar tissue of the retrolaryngeal space present dyspnœa, attacks of suffocation, and cough like those of croup, and run, moreover, generally an acute course ; but they also present dysphagia and severe pain, occasioned by pressing on the thyroid cartilage.\*

*Abscess of the larynx* bears a strong resemblance to retropharyngeal abscess, and may be, like it, mistaken for croup. Abscess of the larynx in its acute and primary form is not a frequent disease : rare in adults, it is still rarer in children. No swelling can be detected in the pharynx to account for the pain, the cough, the difficult breathing and impeded swallowing ; but on close observation it is found that the larynx projects, and that there is induration at the posterior margin of the thyroid cartilage. The neck is not markedly swollen, as in diffuse inflammation of the cellular tissue. With the laryngoscope, we observe a circumscribed swelling, red at its base, and often yellowish at its apex. We do not find, as we so commonly observe in croup, that both inspiration and expiration are interfered with ; the latter, indeed, may be both unembarrassed and noiseless.

Abscess of the larynx may have unsuspected causes. Poli †

---

\* Goix, Archives Générales de Médecine, Oct. 1882.

† Gazzetta degli Ospitali, Naples, May 14, 1894.

reported a case in the discharge from which the sulphur-yellow granulations of *actinomyces* were detected. Watson Williams\* found the Gaffky typhoid bacillus at the base of ulcers and in the structures of the larynx.

Further, croup may be mistaken for tonsillitis, for capillary bronchitis, for whooping-cough, or for the presence of foreign bodies in the larynx or trachea. But the points of distinction are evident. In *tonsillitis* or in *tonsillar abscesses* the breathing is not at all, or but very slightly, impaired; and a glance into the mouth is sufficient to reveal the real nature of the malady. So it is in *peritonsillar abscess*, where otherwise the suffocative attacks that are prone to happen might be misleading. In *capillary bronchitis*, there is dyspnoea, as in croup; but the dyspnoea is unremitting, and associated with fine rales in the lungs, and not with a ringing cough, a harsh tracheal breathing, a hoarse voice. In *whooping-cough*, paroxysms of coughing and of obstructed respiration occur; but then follows the distinctive hoop; and there is no fever, the voice is not husky, and the child does not suffer between the spells. *Foreign bodies in the windpipe* give rise to stridulous breathing and to cough, but they do not often mimic croup closely enough to deceive; and the absence of the peculiar cough and of fever, and the history of the case, prevent error; so also does attention to the fact that the signs vary as the foreign body shifts its position. Furthermore, as Gross † in his elaborate work points out, the embarrassed breathing caused by a foreign body is chiefly found in expiration.

The diagnosis of membranous croup has been considered connectedly because it is convenient and practically useful to so consider it, and because I am still of the belief that there is such a disease as a membranous laryngitis which is not diphtheria. It is, undoubtedly, comparatively rare; undoubtedly all the older descriptions of membranous croup must be with our modern eyes viewed with suspicion. But I believe it as much an error to deny its existence as it was formerly to hold to the comparative restriction of laryngeal diphtheria. The strong points in the diagnosis of non-diphtheritic membranous croup are: the gradual

---

\* Journal of Laryngology and Otology, Oct. 1894.

† On Foreign Bodies in the Air-Passages.

origin and the slow deepening of the symptoms ; the fact that no membranes appear in other localities ; that the disease has a laryngeal onset,—though this may happen also in diphtheria,—and the absence of the Klebs-Löffler bacillus in any shreds of membrane in the expectoration. In discussing laryngeal diphtheria the matter is further examined into.

### Chronic Laryngeal Affections.

Of the chronic diseases of the larynx, chronic inflammation of the mucous membrane, and the changes produced in it by inflammation, thickening and ulceration, are the most common.

**Chronic Laryngitis.**—Alteration of the voice, cough, and an uneasy feeling in the larynx are the main symptoms. The cough is at first dry, but when of any standing is followed by a yellowish opaque expectoration. It either presents nothing peculiar in its tone or else is harsh and barking. The breathing is little, if at all, embarrassed, except when the mucous textures are greatly thickened or ulcerated. In that case there is dyspnoea, the respiration is apt to be noisy and the voice completely lost, because the vocal cords have also suffered. There is, moreover, considerable pain on pressure ; the sputum is muco-purulent, or else purulent and streaked with blood ; and sometimes, if the cartilages also be involved, fragments of them are expectorated, and by the touch we recognize the changed state of the tube.

The symptoms of chronic laryngitis are mostly not purely local. Chronic laryngitis is frequently, indeed, found to be connected with a broken constitution, because the inflammation of the larynx, both in its simple and in its ulcerated form, is often combined with tuberculosis, or with syphilis. In every patient, therefore, suffering from chronic laryngitis, we must endeavor to ascertain whether either of these morbid conditions is present. Chronic laryngitis frequently turns out, on thorough examination, to be laryngitis linked to a serious pulmonary difficulty ; or we detect ulcers in the pharynx associated with those in the larynx and cicatrices, and are enabled to trace clearly the ravages of constitutional syphilis.

As seen with the laryngoscope in chronic laryngitis, hyperæmia, general or partial, is present, associated in cases of long standing with considerable and uniform swelling of the mucous membrane ;



the vocal cords are often uneven at their edges, and there may be, chiefly between the arytenoid cartilages, superficial ulcers. Papillary growths upon the edges of the vocal bands may follow inflammation or repeated attacks of hyperemia.

Chronic laryngitis is liable to be mistaken for an aneurism of the aorta, or, more strictly speaking, an aneurism of the aorta is liable to be regarded and treated as a case of chronic laryngitis. The distinction, as will hereafter be shown, is mainly made by attention to the physical signs; often the paralysis of a vocal cord is of great significance.

Cases of functional or *nervous aphonia*, too, are sometimes confounded with chronic laryngitis; and it is by no means always easy to avoid this error. The loss of voice may be either partial or complete. It not unfrequently comes on without any previous warning; and this fact aids us greatly in diagnosis. So does the absence of cough, of expectoration, of local pain, and of all difficulty in breathing; for none of these symptoms are commonly observed in aphonia which is solely nervous. One of the causes of the disorder is overstimulation of the vocal nerves, by straining the voice in singing or in speaking. We also meet with it as occasioned by narcotics or by lead poisoning, and perhaps most frequently as a reflex manifestation, due to irritation of the intestines by worms, or to a disorder of the uterine system. In these instances of nervous aphonia the voice suddenly disappears and as suddenly reappears, a phenomenon not unusual in the aphonia of hysteria; and we may have from impaired but not wholly lost power the voice absent only for some hours daily. It is evident that in all cases of nervous aphonia the laryngoscope will assist us greatly, as it will show the true condition of the parts, as regards both their structure and their mobility. It also aids us in distinguishing these laryngeal disorders from cases of *aphonia* due to want of strength in breathing,—to want of power in expiration.

Enlarged bronchial and cervical glands, or an aneurism which compresses the laryngeal nerves, also produce hoarseness, and ultimately complete loss of voice. Under such circumstances there is a short cough, attended often with loud tracheal rales; and we observe attacks of dyspnoea, with a noisy, hissing respiration. The practical lesson which all such cases teach, is to remember that the symptom considered most characteristic of

chronic laryngeal inflammation—the altered voice—may occur when no laryngitis exists ; also to examine with the laryngoscope, and to note the effect of palsy of the muscles the result of nerve-pressure. In thoracic aneurism, pressure symptoms, such as the dyspnœa and the altered voice, with paralysis of laryngeal muscles, may be produced either by pressure upon the recurrent laryngeal nerve, which on the left side passes around the arch of the aorta, or upon the vagus. Pressure upon the vagus will give rise to abductor paralysis of the corresponding side, with adductor spasm of the laryngeal muscles of the opposite side, the spasmodic movements being intermittent. Pressure upon the one recurrent nerve causes one-sided abductor paralysis, the cord being at first in the middle line, but later it assumes the position due to adductor paralysis. Pressure upon one vagus, inducing double adductor spasm, produces serious dyspnœa and difficult phonation ; but pressure on one of the recurrent nerves may occasion intermittent dyspnœa that is usually not very troublesome, and scarcely, if at all, affects phonation. Major's \* researches have given us much of this definite knowledge.

Now, in the nervous forms of aphonia just mentioned, with the exception of those caused by pressure, the loss of voice is due to deficient power, and the cords move sluggishly or not at all. When the disorder reaches a high degree, we perceive, on looking into the laryngeal mirror, that the vocal cords do not approximate as the patient attempts to say *a* or *o*. But, besides these cases, owing to general want of force, we find cases of *spasm of the tensors of the vocal cords* with most peculiar, partially interrupted voice, and of *absolute paralysis of individual muscles*, as of one adductor of a cord ; or of one or both posterior crico-arytenoids, or abductors ; or of the crico-thyroids, or tensors. In some of these there is considerable dyspnœa, with noisy breathing ; in all the laryngoscope affords the only means of diagnosis. In paralysis of the external tensors of the vocal cords, the crico-thyroid muscles, there is inability to use the higher notes with any freedom ; the voice is rough or entirely lost, and viewed with the mirror we find in phonation a want of longitudinal tension, as indicated by a wavy outline of the glottis, the convexity of the

---

\* Transactions Amer. Laryng. Assoc., New York, 1891, p. 66.

upper surface of vocal bands on expiration and phonation, and concavity on forcible inspiration. The contraction of the muscles which can be felt externally in the healthy subject during phonation is completely absent. This form of disorder most frequently results from overstraining the voice; it may be caused by cold, and is apt to be bilateral. Geo. W. Major\* reports a case caused by exposure to cold. Palsy of the thyro-epiglottic muscles has its usual origin in diphtheria. The epiglottis stands erect, and does not move during attempts at deglutition. In palsy of the relaxors of the vocal cords, the thyro-arytenoid muscles, the deep tones are nearly gone. It is often unilateral, and comes mostly from overexertion of the voice during catarrhal laryngitis. Viewed in the laryngeal mirror, the edges of the cords do not approach in the median line, and the edges seem excavated. In paralysis of the posterior crico-arytenoid muscles, we see in the mirror the glottis merely as a narrow slit, becoming still narrower during inspiration. There is no disturbance of voice, and scarcely any sign of laryngeal catarrh, but there is most marked and noisy laryngeal dyspnoea. This paralysis of the abductors may happen from compression of the recurrent nerves by an organic stricture of the œsophagus.† Bilateral paralysis of the adductors is a common disorder, occurring in connection with locomotor ataxia and affections of the brain and of the medulla.

Unilateral paralysis of the adductors is more rare; it accompanies malignant disease of the œsophagus, aneurism of aorta, and, exceptionally, metallic poisoning, as lead and arsenic. It sometimes follows exposure to cold, or attends rheumatism or phthisis. When met with in connection with paralysis of the same side of palate or tongue, it is centric, at times bulbar. E. Fletcher Ingals has described cases thought to be hysterical in origin. We also encounter sensory neuroses of the larynx, and among these hyperæsthesia is common.

Chronic laryngitis, or rather its chief symptom, loss of voice, is at times *feigned*; and the deception may be kept up for an indefinite period. Yet we possess, in the use of anæsthetics, the means of detecting the fraud at any moment. Just before the impostor

---

\* Proceed. Amer. Laryng. Assoc., 1892, p. 10

† Case of Dujardin, Annales des Maladies de l'Oreille, 1887.

falls into the deep sleep produced by ether, or as he is recovering from the insensibility it occasions, his will no longer controls his voice, and he speaks in his natural tone, or even screams violently.

Now, under the term chronic laryngitis, which formerly for want of more precise knowledge was made to embrace most kinds of chronic diseases of the larynx, many different morbid processes are embraced, the exact nature and seat of which we may discriminate by the laryngoscope. Thus, the disorder may be wholly, or almost wholly, confined to the *epiglottis*. We may find this structure highly congested and enlarged; we may be able to note that it is pendent, nearly completely covering the glottis; and it is frequently the seat of ulceration. The attending symptoms in any case are those regarded as characteristic of a greater or less degree of laryngeal inflammation. In instances of ulceration, there is soreness with pain in swallowing, hoarseness and irritative cough, followed at times by blood-streaked expectoration. The ulceration may terminate in total destruction of the epiglottis. A turban-shaped swollen epiglottis is often met with in phthisis associated with pyriform swelling of the arytenoids. Pallor of these structures, indeed of the whole larynx, is one of the early signs of pulmonary tuberculosis, as Cohen has pointed out.

When the *vocal cords* are affected, we recognize in the laryngeal mirror either their reddening in part or entirely, or their induration and thickening, or we observe œdematous swelling in and around them, or their ulceration; and we can usually detect during breathing and phonation their impaired action. The inflammatory redness may be only in one cord. Small collections of mucus are often found adhering to different parts of the laryngeal membrane. Now, all these conditions are generally combined with marked aphonia; the voice, indeed, may be reduced to the merest whisper. Venous congestion of the larynx is so rare an affection that Mackenzie has met with but four cases of it.\* In making our diagnosis we must always be careful to find out if the laryngeal phenomena be not secondary, forming part of a general morbid state, such as dropsy, tuberculosis, syphilis, or changes in the blood. Chronic hypertrophy of the ventricular bands is the

---

\* Disease of the Throat and Nose, vol. i, 1880.



result of inflammatory thickening, and, as Tauber \* proves, occurs mostly in those who use the voice much in their professional vocations. Türek has given the name of "chorditis tuberosa" to a condition of the vocal bands in singers, having these characters. Midway in the upper plane of the bands there is a peculiar uneven surface, and, in addition to redness and swelling, some white, opaque spots as large as poppy-seeds.†

Paralysis of one vocal cord may exist, with immobility of one side of the larynx, and yet voice may be restored and be absolutely normal; the healthy cord, as in cases narrated by Bosworth, swinging over to the paralyzed side, so as to make up for the loss of power on that side. Voice may even exist, to a restricted extent, not only without vocal cords, but after entire extirpation of the larynx, as in the remarkable case of Hickey, reported by J. Solis Cohen,‡ in which the larynx was removed for malignant growth, and the trachea permanently fixed in the neck. After swallowing air, the patient was able to talk, and even to sing, by skilfully using his pharyngeal muscles.

Alteration of the voice, mumbling speech, as though there were some difficulty in closing the glottis, while the movements of the vocal cords appeared normal as seen with the laryngoscope, without true aphasia, is mentioned by John N. Mackenzie as a symptom in a case of bulbar disease.§

*Diseases of the cartilages and of the perichondrium* are most frequently encountered in connection with tuberculosis, syphilis, and low forms of fever. The affection often begins in the sub-mucous tissue, and the ulceration spreads until the cartilaginous parts of the larynx are involved. The arytenoid cartilages are generally first attacked; and portions of these cartilages may be thrown off and expelled. At times pus is formed which gives rise to swellings that can be recognized by the aid of the laryngeal mirror; sometimes a displacement of the cartilages takes place, before any portion of them is completely separated, and the most distressing and dangerous attacks of suffocation result; or the

\* Cincin. Lancet, 1887.

† Klinik der Krankheiten des Kehlkopfes, Wien, 1866.

‡ Pharyngeal Voice, Transactions of the Amer. Laryng. Assoc., 1894.

§ Transactions Amer. Laryng. Assoc., New York, 1891, p. 6.

perichondritis may lead to the development of bone-substance and a constriction of the tube. In some instances the purulent collection presses on a vocal cord, which, with the laryngoscope, may be seen to be immovable.

This instrument reveals very generally the ravages the disease has committed ; and we are thus enabled to form an opinion as to how far the destruction has progressed, and which of the soft parts as well as of the cartilages are involved. Leaving out the frequent perichondritis and caries of the cartilages which follow the deposition of *tubercle*, we find in laryngeal phthisis considerable swelling and ulceration of the epiglottis, and often semi-solid pyriform swellings of the aryepiglottic folds. The thickening is more regular and uniform than that of syphilis, and the tubercular ulcers not large and solid as in this affection, but small and numerous, and both vocal cords are involved ; while in this as in every other respect syphilis is more apt to be local and unilateral. Tubercle bacilli are found in the discharge from the laryngeal ulcer, and in catarrhal ulceration the ulcers are generally very superficial and on the vocal cords. The symptoms of laryngeal phthisis are difficulty in breathing and in swallowing, local pain and soreness, a greatly-altered or a lost voice, and a distressing, harsh cough, which is followed at times by purulent expectoration. Besides, we find the manifestations of disease of the lungs. But it occasionally happens that we encounter cases of tuberculous ulcers with abundant bacilli, in which no lung disease exists ;\* and it is not uncommon to find the tubercular disease of the larynx preceding that of the lungs. At times we note syphilitic and tubercular ulcers in combination. We may also meet with catarrhal ulcers where there is tubercular disease of the lungs.

The diagnosis between pachydermia of the larynx and the inter-arytenoid tumor of phthisis is that in the latter the swelling is distinctly a tumor, with a more or less well-defined margin. The color is usually red or pink ; in pachydermia it is whitish-gray or only slightly pink, and the surface may be smooth or coarsely papillary, while in the former it is smooth or finely granular, with sometimes a furrow or cleft. These features are

---

\* Canadian Practitioner, 1887.

specially relied upon by McBride.\* Ulceration occurs in pachydermia only exceptionally and as a complication ; it is common in phthisis of the larynx.

As the result of disease of the cartilage and of the perichondrium, especially as the result of the process of cicatrization, we may have stricture of the larynx and trachea ; for this is, in truth, the most common origin of *laryngeal stenosis*. The inspiration is prolonged and noisy ; the voice is generally, although not of necessity, affected. There is dyspnœa, and with the laryngoscope we can see how greatly the calibre of the tube has been encroached upon. Cicatrization is common after syphilis, but Cohen's case † proves that it may occur spontaneously also in tubercular ulcerations. Adhesions may be congenital, a web-like membrane uniting the vocal cords through a part of their extent, as in a case of Morell Mackenzie's. According to Paltauf,‡ primary stenosis of the larynx may be caused by scleroma, which may develop early in the larynx. The diagnosis depends upon the detection of the characteristic minute structures. Lepra occurs in the larynx only by extension from the pharynx, and is accompanied by manifestations upon the surface of the body, which clearly explain its character.

*Ulcers* in the posterior walls of the larynx give rise, as a rule, to distressing cough. Respecting *tumors* of the larynx and polypoid growths in its interior, we cannot distinguish them, by their symptoms alone, from chronic laryngitis. Their most trustworthy signs, irrespective of the cough and the altered voice, are a steadily increasing difficulty in breathing, and attacks of suffocation for which nothing in the lungs or heart or great vessels accounts.

New growths may occur in the larynx, of the benign form. Papilloma, papillary fibroma, is probably the most common ; myxoma is rare ; fibro-myxoma and fibroma unusual. Malignant disease in various forms may affect the structures of the larynx. Very frequently the diagnosis can be positively made only with the aid of the microscope.

The detection, at the seat of the larynx, of a growing tumor,

---

\* Edinburgh Med. Journal, April, 1893.

† Amer. Journ. Med. Sci., Dec. 1888.

‡ Sajous, Annual of Univer. Med. Sci., 1893.

accompanied by a severe cough, by a sanious sputum, by signs of destruction of tissue, as seen with the laryngeal mirror and as found in the expectoration, by perichondritis and exfoliation of the laryngeal cartilages, by hemorrhages, and by emaciation, would, in addition to the symptoms just enumerated, warrant the diagnosis of *cancer*, whether or not much pain were present. In some instances, too, gangrenous pneumonia occurs. If the tumor be flat, with a broad infiltrated base, rather soft in consistence, and color not much changed from normal, and there be a peculiar musty odor to the breath and expectoration, a probable diagnosis of carcinoma may be made. This may be confirmed by the subsequent rapid development of the malignant disease, associated with distress in swallowing, bloody expectoration at times, and quickly appearing cachexia. *Polypi* in the larynx may sometimes be seen by depressing and dragging forward the tongue until the epiglottis is brought into view; at least they have been thus discovered, and even successfully operated upon. But as regards polypi, or, indeed, any form of morbid growth, we possess in the laryngoscope the most certain, usually the only certain, means of detecting them, and even of aiding us in removing them, as is now being constantly done. These laryngeal growths vary much in size and in color; they are often seated at the anterior free edges of the true cords, or still more generally just above or just below their origin, and are, as a rule, readily discerned. Sometimes they may exist for years, merely producing changes in the voice and some cough, but no very great distress; or they may lead to fits of strangulation and to sudden death. It is impossible to be sure of their nature without repeatedly examining portions of them. *Papillomas* are usually cauliflower-like or in bunches; they occupy most frequently the vocal cords, while *sarcomas* are oftenest found at the anterior portion of the larynx. *Cysts* of the vocal cords are much rarer than other forms of growths; they sometimes rupture spontaneously, and the hoarse voice quickly clears.\* Myxomata of the larynx and the epiglottis, according to Van der Poel,† may be manifestations of pernicious anæmia. They

---

\* Heinze, Archives of Laryngology, New York, 1880.

† Amer. Laryng. Assoc., 1890.



differ from cysts in being a pure, gelatinous growth characterized by stellate fusiform cells, often anastomosing, embedded in a homogeneous, or finely fibrillated, soft, gelatinous basement substance. Many cases that are classed as cysts would properly come under the head of myxoma if the aid of the microscope had been sought.

Before concluding these remarks on diseases of the larynx, it may be thought necessary to point out the differences between them and diseases of the *trachea*. But affections of the trachea need not be separately considered. Lying between the larynx and the bronchi, the trachea commonly shares in their disorders. Thus, we have seen croup to be a malady in which both larynx and trachea are involved. Slight inflammation of the trachea occurs constantly in slight attacks of laryngitis or of bronchitis. *Ulcers* in the trachea may exist without ulceration of the larynx; but then they usually escape detection. Sometimes, however, they reveal themselves by a constant pain at the lower portion of the neck and the upper part of the sternum, joined to all the symptoms of ulceration of the larynx except the impaired voice. *Morbid growths*, too, occur in the trachea,—cancer, carcinoma, syphilitic growths,—as they do in the larynx, and the tube may be altered in form and in structure. Vegetations also form in the trachea after tracheotomy.\* We can make use of the laryngoscope to assist us in the diagnosis of any of the forms of tracheal disease referred to. Yet the instrument is not always available; for it is only under favorable circumstances that the entire extent of the trachea can be seen.

In *narrowing* of the trachea the bronchial tubes are also at the same time often narrowed. The stenosis may be caused by external compression, as from a goitre, from an aneurism, or from a mediastinal tumor; or the constriction may be due to some cause, such as new formations, in the walls of the tubes. The chief symptoms are the same in either case; and they are, long-drawn-out respiratory acts, noisy breathing, especially in paroxysms, dyspnoea, particularly marked in inspiration, epigastric retraction, feebleness or absence of vesicular murmur, with clear pulmonary

---

\* See cases collected by Petel, *Des Polypes de la Trachée*, Paris, 1879.

resonance, loud wheezing heard with the stethoscope at or near the place of constriction, and voice slightly, if at all, impaired. This, the normal appearance of the larynx as shown by the laryngoscope, and the almost imperceptible motion of the windpipe during breathing,\* are of great value in distinguishing a tracheal from a laryngeal stenosis. A bronchial stenosis is chiefly discriminated by the signs of the constriction being one-sided, and attended with marked thrill of the thoracic wall of the affected side, and with loud sounds issuing from it, loud enough to be heard at a distance. Subglottic œdema may be detected by the laryngoscope on deep inspiration.

---

\* Gerhardt; also Riegel, in Ziemssen's *Cyclopædia*.

## CHAPTER IV.

### DISEASES OF THE CHEST.

AN examination of the diseases of the chest must be prefaced by a description of the methods of investigation which have given to their diagnosis such certainty. The same methods may be applied in the study of the maladies of other parts of the body, but they are of special service in the recognition of thoracic disorders, and will be here, therefore, most appropriately considered.

The discrimination of disease by the eye, the ear, the touch, in fact by the direct aid of the senses, is called *physical diagnosis*; the signs thus ascertained are connected with perceptible alterations in the material properties or physical nature of structures,—such as alterations in their form, their density, or their sounds,—and are known as *physical signs*.

Physical signs are, then, the exponents of physical conditions, and of nothing more. But as the same physical conditions may occur in various diseases, so may the same physical signs occur in various diseases. An isolated sign is, therefore, not diagnostic of any particular malady. It reveals usually an anatomical change; but it does not determine the disorder occasioning this change. The tendency to ascribe to each thoracic affection, and even to each stage of an affection, a pathognomonic sign, has greatly retarded the usefulness of physical exploration. By presenting a never-ending list of specific signs, it has frightened many from attempting to become acquainted with the most serviceable of all the means of diagnosis, and many more, by the unnecessary complications introduced, have been disheartened at the very threshold of their studies. The subject may be much simplified by laying less stress on individual signs, and by grouping them together according as their association becomes distinctive of certain well-marked physical states. Morbid anatomy then steps in with its teachings, and tells us in what diseases these states are commonly found. It is in conformity with these views that I shall at-

tempt, in the following pages, to delineate the signs of thoracic affections.

But physical signs cannot be acquired from books; they must be learned at the bedside. Their value can be ascertained by reading; yet to distinguish them with readiness requires constant cultivation of the eye, of the ear, and of the sense of touch. And it is of great importance to have clear ideas regarding the structure of the parts to be investigated, and of their action in health.

For the sake of convenience, the surface of the chest has been mapped out into regions. Various arrangements of these have been made by different authors. The simplest division of the chest is into anterior, posterior, and lateral surfaces. The regions into which the anterior surface may, for practical uses, be subdivided, are an upper region, extending from just above the clavicle to the fourth rib, and a lower region, from the fourth rib downward. Posteriorly, also, there are an upper and a lower part of the chest to be specially examined. It is hardly necessary to say that all these regions are double,—the same on each side of the chest. Many more divisions are usually made; but they are perplexing to the student, and of doubtful value. The artificial boundaries generally laid down are, indeed, too minute and yet not minute enough; they are too minute for ordinary purposes, not minute enough when it is desirable to localize a physical sign. Whenever this is requisite, instead of resorting to the names of the regions usually employed, I think it preferable to designate the seat of the sign with reference to some fixed anatomical point. This may be done for the anterior part of the chest by indicating the distance above or below the clavicle, or near what part of the sternum, or at which rib, or spreading over how many intercostal spaces, the sign in question is perceived. At the posterior part of the chest, the spinous ridge of the scapula, its lower angle, and the spinal column, serve as landmarks. For most clinical purposes, it is only needed to study the region above the spinous process of the scapula, as separate from the space below. But in some instances it may be necessary to notice the region between the scapulæ, inter-scapular, or that extending from the lower angle of the bone to the limits of the chest, infra-scapular.

Let us now examine the different methods of physical diagnosis, and particularly in their relation to pulmonary diseases.



## SECTION I.

## DISEASES OF THE LUNGS.

The Different Methods of Physical Diagnosis, and the  
Physical Signs of Pulmonary Diseases.

## INSPECTION.

If the chest be examined with the eye, we obtain an idea of its form, size, and movements. In health this inspection shows us that the two sides of the chest are, to a great extent, symmetrical in form, as well as in size and in movement. Both sides rise equally during inspiration and sink equally during expiration. On both sides the motion of inspiration is longer than that of expiration, and the pause between them extremely slight.

This *respiratory movement* is visible over the whole thorax. In males it is most distinct at the lower portions of the chest; in females it is most discernible at the upper. This difference in the two sexes becomes the more manifest, the more hurried the breathing. In healthy adults the lungs expand with regularity from sixteen to twenty times in a minute. In certain pulmonary affections, especially in pneumonia, the number of respirations often exceeds fifty in a minute. But hurried breathing and changed movements of the thorax occur independently of diseases of the lung; the heaving of the chest in an hysterical paroxysm proves this. Where the diaphragm does not descend, as in consequence of peritonitis or of abdominal dropsy or of tumors, the breathing is much more rapid, and is perceptible at the upper parts of the chest. Again, the thoracic movements may be distinct on one side and hardly noticeable on the other, as in pleurisy or in pneumothorax. Lastly, as happens in some cerebral lesions, the motions of the chest may be very slow and labored, or irregular, or they may have apparently ceased, and the breathing be altogether abdominal.

The *form* of the chest is sometimes strikingly altered by disease. Congenital malformations, imperfect development, and curvatures of the spine modify it; so do intra-thoracic affections.

Frequently the chest presents a retracted or an expanded look. Retraction denotes diminished size of the lung, and, if one-sided, is usually indicative either of chronic changes in the lung-tissue, as in chronic pneumonia or in the forms of phthisis, or of false membranes which bind down the lung; or it is found in a very marked manner in empyema with external opening. Expansion of the chest is met with in emphysema, in pneumothorax, and in pleuritic effusion. A local or partial expansion, or bulging, may be encountered in the latter disease, or it may depend on thoracic tumors, on pericardial effusions, or on hypertrophy of the heart.

The *size* of the chest can be only approximatively judged of by the eye. Where accuracy is necessary, measurements must be resorted to.

#### MENSURATION.

To measure the circumference of the chest or of the abdomen, or to ascertain the distance from one portion of the surface to the other, a graduated tape is all that is required. To attain the former object, the spinous process of a vertebra is chosen as a fixed point, and the tape is thence passed round the body to the median line, first on one side, then on the other, taking care that it be applied evenly to the skin, and that the level of the measurement be the same on both sides. This level, if the examination be recorded, should always be noted, that we may have a uniform standard of comparison. And for the same reason it is best to adopt the plan of making our measurements as nearly as possible on the same line: for example, in determining the circular width of the thorax, we can, as a rule, select a line immediately above the nipple, or draw the tape around the chest toward the sixth costo-sternal joint, and, therefore, on the level of the sixth rib near its attachment to the cartilage. We measure thus the width of the chest; if we wish to obtain the longitudinal diameter, the line from the clavicle to the base of the chest is taken. Where the chest is deformed, Woillez's cyrtometer, a chain with links, may be used in place of the tape.

In estimating the size of the chest in disease, it must be borne in mind that even in health its two sides vary widely. The half-circle on the right side is, in right-handed persons, at least half an

inch larger than the half-circle on the left. But the measurements, to be trusted, must be performed while the patient is holding his breath in expiration. In inspiration the girth of the chest is increased fully three inches. In well-developed men it measures at the upper part about thirty-three to thirty-four inches during expiration.

If it be desirable to ascertain in how far the respiratory acts modify the dimensions of the chest or of the abdomen, this may be readily effected by the ingenious "chest-measurer" of Sibson, or by the "stethometer" of Quain or of Carroll,\* all of which instruments register accurately the movements of breathing; or the respiratory curves can be traced and studied by the atmograph of Burdon Sanderson, or by the anapnograph, an instrument made use of by Bergeon and Kastus, and similar to the sphygmograph,† or by Riegel's double stethograph; or the curves of the respiratory movements may be seen in the tracings of the pneumograph applied to the chest. Ransome has called attention to the value of recording the exact extent of the respiratory movements by stethometry as a means of prognosis in chest disease.‡

The transverse diameter—the breadth—of the chest may be determined by means of a pair of callipers, arranged specially for the purpose; and the curves or flatness of the surface may be ascertained, should it be necessary, by Alison's stetho-goniometer (Fig. 17); but it is rarely necessary. In fact, these minute measurements, however interesting to the physiologist, have, as yet, not been made available to the physician. Inspection teaches us the same as mensuration. What it teaches with less precision can be learned for purposes of diagnosis with a graduated tape.

FIG. 16.



The stethometer of Quain. The box is placed on the sternum, and the string carried around the chest. One revolution of the index, which is moved by a rack attached to the string, indicates an inch of motion in the chest.


\* New York Medical Journal, 1868.

† Gazette Hebdomadaire, Sér. 2, v., 1868.

‡ Medico-Chirurgical Transactions, vol. xlv., 1881.

Mensuration may be employed not only to judge of the size of the chest and of its movements, but also to ascertain the amount

FIG. 17.



The stetho-goniometer of Scott Alison.

of air which is received into the lungs. The instrument used for this object is the *spirometer*, an invention of Dr. John Hutchinson, and since his time numerous modifications of the instrument have been made; for instance, the ordinary dry and the wet gas-meter have been adopted to the purposes of spirometry, and an instrument small enough to be carried in the pocket has been suggested. The results the spirometer has yielded are of value from a physiological point of view; from a clinical, there are too many sources of fallacy and too many drawbacks to render them of great importance; and not the least of these drawbacks is, that it takes much practice to learn how to blow. The spirometer may indicate that a large quantity of air enters the lungs, and thus become a rough test of their normal condition. But when less air passes into the organ than the spirometric standard requires, this leads in itself to no conclusions; certainly not to any concerning the disease which occasions the diminished vital capacity. In estimating results arrived at by the spirometer, it must be remembered that sex, weight, age, and height have to be taken into account. To the latter Hutchinson assigns much importance, since he enunciates the law that for every inch above five feet, eight cubic inches are to be added to the healthy standard. For the height of five feet, the breathing volume is one hundred and seventy-four cubic inches. But these calculations are not exact; they only approximate the truth. Moreover, the vital capacity may be increased by practice, with the spirometer or by the use of pneumatic instruments designed to breathe in compressed air or to breathe out into rarefied air.



To determine both the expiratory and the inspiratory power, the hæmadynamometer may be employed. Hammond\* recommends the use of the instrument in the examination of recruits. According to his observations, healthy men of five feet eight inches raise the column of mercury about two inches by inspiration, and about three inches by expiration.

Waldenburg measures the force in respiration by a special apparatus, and has introduced *pneumatometry* as a means of diagnosis. The power exerted in expiration is greater than in inspiration. In some affections the expiratory pressure is largely diminished, as in emphysema and asthma, while in the forms of phthisis the force of inspiration is much lessened.

#### PALPATION.

Palpation, or the application of the hand, confirms the results obtained by inspection and mensuration as to size, form, and movements. It may, in addition, be employed to determine spots of soreness, the density and condition of tumors, the state of the thoracic walls, the frequency of the breathing, and the action of the heart. The hand may further be of service as a means of distinguishing vibrations produced by rhonchi, *rhonchal fremitus*, or by the voice, *vocal fremitus*; or it may detect fluid by the sense of fluctuation it imparts, or a roughened serous membrane by the friction fremitus. When both fluid and air are present in a large hollow space, by shaking the patient a distinct vibration of the parietes is felt, accompanied by a splashing sound, known as the Hippocratic or succussion sound.

Palpation is to be practised by applying the palmar surface of one or of several fingers evenly, and without too much pressure, on the part to be examined.

#### PERCUSSION.

By percussing or striking bodies we elicit sounds by which we judge of their composition. Percussion was first practised by striking directly with the hand over the organs to be explored; a


---

\* Treatise on Hygiene, Philadelphia, 1863.

method which, although serviceable to ascertain marked differences, or to obtain an idea of the general resonance of a part, is inferior to mediate percussion. The media used to receive the blow are various: a disk or plate of ivory, of wood, or of leather; a piece of india-rubber; or the middle finger of the left hand. The finger answers best for percussion of the chest; for abdominal percussion a pleximeter is preferable.

When the finger is employed, it ought to be applied with its palmar surface firmly pressed against the chest, and as parallel as possible to the ribs. One or two fingers of the other hand may then be used to tap with,—for the finger is, for ordinary purposes, better than any of the percussion hammers invented,—the greatest attention being paid to the circumstance that

FIG. 18.



The pleximeter: about natural size. It may be conveniently made of hard rubber.

the percussing finger strikes perpendicularly, whatever pleximeter be used, and not slantingly, as is too generally the case. The whole movement should proceed from the wrist, and only from the wrist, and ought not to be too rapid, or unequal, or of great force. If all of these apparently unimportant points are attended to, the results obtained may be relied upon; if not, the want of manual dexterity invalidates the conclusions. No other fault is so often committed by the beginner as that of raising the finger used as a pleximeter from the surface,—thus obtaining the sound of the finger, and not that of the organ he wishes to percuss,—unless it be the fault of striking with great force, as if the object were to break into the cavity of the chest. Forceful percussion is of use only when the sound of deep-seated organs is to be brought out.

The main sounds elicited by percussion may be designated as dull, clear, and tympanitic. Of course, these, like all other sounds, may differ in strength, in duration, and in pitch.

A *dull* sound denotes absence of air. It is the sound both of fluids and of solids. It is, thus, the sound sent forth from the airless viscera,—from the liver, spleen, and heart. When it


takes the place of the pulmonary sound, it bespeaks consolidation, from whatever cause induced, or the presence of something which checks the normal vibrations of the lung-texture. Dulness is always associated with an increased sense of resistance to the percussing finger, and over parts emitting it the vibrations of the tuning-fork, which Bass has introduced into diagnosis, are weak, while they are loud over normal pulmonary structure.

A *clear* sound is produced by a series of marked and unhindered vibrations which are emitted from a substance containing air. As thus defined, a clear sound evidently is yielded by percussing any air-containing organ. But custom has restricted the employment of the term clear to denote the peculiar resonance obtained by striking over pulmonary tissue. When, therefore, a clear sound is spoken of, it means a sound having the nature of that of the lungs, or of normal vesicular or pulmonary resonance. A resonance analogous to the vesicular resonance may be obtained, Flint points out, by percussing a loaf of bread.

A *tympanitic* sound, on the other hand, is a non-vesicular sound, having the character of that of the intestine. Wherever heard, it indicates the presence of quantities of air in conditions similar to that contained in the intestine,—namely, enclosed in walls which are yielding, but neither tense nor very thick. When elicited over the chest, it may be only the transmitted sound of a distended stomach or colon. But generally a tympanitic sound over the seat of the lungs is expressive of emphysema, or of pneumothorax, or sometimes of a cavity or of œdema of the lungs. Again, as Skoda has taught us, it occurs in moderate pleural effusions above the level of the liquid. Many find difficulty in distinguishing between the clear sound of the pulmonary tissue and the tympanitic sound. The more ringing character of the latter, and its higher pitch, constitute its essential properties.

If the cavity communicate with a large column of air in the bronchial tube, the note on percussion varies, as pointed out by Wintrich, accordingly as the patient opens or closes his mouth. It is more markedly tympanitic and higher in pitch when the mouth is wide open. Altering the position from a sitting to a horizontal one, when the cavity is partially filled with fluid, Gerhardt has shown changes the tympanitic percussion note, and I have

FIG. 19.



observed it to be markedly altered—indeed, to disappear—on full held inspiration.\*

As modifications of the tympanitic sound may be viewed the *amphoric* or *metallic* sound, and the *cracked-pot* or *cracked-metal* sound. The first of these is a concentrated tympanitic sound of raised pitch, and denotes a large cavity with firm, elastic walls. The second is not unfrequently found associated with it. It requires for its development a strong, abrupt blow of the percussing finger while the patient keeps his mouth open. The condition that usually occasions the sound is a cavity communicating with a bronchial tube. It is also met with uncombined with an excavation, as in the bronchitis of children, in pleurisy above the seat of effusion, near a pericardial exudation, in emphysema, and in certain instances of pneumothorax. Indeed, any disorder in which the chest-walls remain very yielding, and in which a certain amount of air contained in the lung or pleura is, by sudden percussion, forced into a bronchial tube, will occasion this cracked-metal sound.

In addition to the character of all these sounds, we study their *degree*, or amount of fulness : such changes as are

FIG 19.—A serviceable model of a percussion hammer; not quite natural size. The india-rubber is screwed to the ring, which has a diameter of from five-eighths to three-quarters of an inch. The metallic ring is attached to a steel stem with a very decided spring. The pointed portion of the india-rubber is used to strike with on the pleximeter.

\* Amer. Journ. of Med. Sci., July, 1875.



expressed by "more or less," "diminished or increased." Thus, a clear sound may be increased, owing to stronger vibrations and a larger quantity of air, yet not lose its distinctive pulmonary character, as happens often, for instance, when the air-cells are dilated; the sound of the large intestine is fuller, more tympanitic, than that of the small intestine, and so forth.

With changes in fulness or volume of sound go hand in hand changes in its *pitch*. Increased volume is linked to lowered pitch, diminished volume to higher pitch; but so is increased tension.

To sum up the chief results of percussion, as above described:

#### QUALITY, OR CHARACTER OF SOUND.

**CLEAR:**—Presence of air,—as in the lung-tissue.

**DULL:**—Solidification or compression.

**TYMPANITIC:**—Certain amount of air enclosed in a structure or cavity the walls of which are not too tense.

*Metallic:*—Large hollow space, with firm but elastic walls.

*Cracked-metal sound:*—Usually a cavity communicating with a bronchus.

#### DEGREE, OR INTENSITY.

Any of the sounds mentioned may be *diminished* or *increased* in intensity as the conditions which produce them are modified.

#### PITCH.

Heightened or lowered as amount of air or as tension is altered.

If it be desirable to obtain a more distinct idea of the sound than can be done by the ordinary method of practising percussion, it may be accomplished by resorting to *auscultatory percussion*,—a method which was introduced by Cammann and Clark, and which consists in listening, with a stethoscope applied to the parietes, to the sounds elicited by percussion. It is a means of determining with accuracy the boundaries of various organs, as of those of the lungs or heart, or of the liver or spleen, and yields particularly exact results when carried out with the double stethoscope.

The percussion sound will also be found to vary with the respiratory movement, and useful information may be obtained by

the appreciation of the note elicited by percussion while the breath is held after a full inspiration or in a prolonged expiration,—a method of diagnosis for which I have proposed the name of *respiratory percussion*.\*

As a standard for comparison in disease, the results of respiratory percussion in health must be carefully determined. It will be found that in the normal chest, anteriorly, a full held inspiration increases the resonance, makes the sound fuller, and raises the pitch; but, making allowance for the cardiac region, the resonance below the apices is relatively less increased on the left than on the right side.

Posteriorly, we find in the supra-spinous fossæ, and on a line toward the spine, that a full inspiration makes the percussion sound fuller and raises the pitch, especially on the right side. In the inter-scapular and infra-scapular regions the tone on gentle percussion is distinctly pulmonary and the pitch moderately high. On the left side an admixture of tympanitic resonance may be detected, particularly in the infra-scapular region. The pitch is somewhat lower in the left scapular and infra-scapular region than in the right. A full held inspiration elevates the pitch, increases the resonance very much, and makes the difference between the sides less apparent.

A held and complete expiration greatly lessens resonance and lowers the pitch on percussion.

The quality of the percussion note during an arrested respiratory movement is but little changed; perhaps it is somewhat less soft, corresponding to the marked resistance to the percussing finger. In a held inspiration, nevertheless, we obtain the idea of a greater mass of tone; in a held expiration, the reverse. Increase in volume of percussion note accompanies, contrary to our usual experience, heightened pitch; and this is more especially noticed in connection with the slight change in quality above mentioned. This anomaly is probably due to the altered tension of the structures, both lung-texture and chest-walls, during held respiratory movement.

These are the chief facts connected with a study of respiratory

---

\* Amer. Journ. Med. Sci., July, 1875; see also Friedreich, Deutsches Archiv für Klin. Med., Bd. xxvi.

percussion in health. The application to disease is manifold, as we shall find in the study of emphysema, of phthisis, of pleurisy, and of pneumothorax.

### *Percussion of the Healthy Chest.*

The sound elicited by striking a healthy chest differs in accordance with the part percussed. The *anterior* portion renders a clearer sound than the posterior, on account of the slighter thickness of the thoracic walls. But the pulmonary resonance is not, even anteriorly, alike at all parts. The portion of lung above the clavicle yields a sound which becomes somewhat tympanitic as the trachea is approached. Percussion is difficult in this region, as it is almost impossible to apply the finger or pleximeter properly to the surface; hence arise errors in diagnosis if too much value be attached to trifling differences between the two sides. Over the clavicle the sound sent forth is clear and pulmonary at the centre of the bone; at its scapular extremity it is duller; toward the sternum it becomes of higher pitch, and mixed with the sound of the bone. In the region bounded above by the clavicle, and below by the upper margin of the fourth rib, the resonance is very marked. In fact, the sound of this region may be taken as a type of the pulmonary sound: it is very clear and distinct, and but little resistance is offered to the percussing finger. Yet a slight disparity generally exists between the two sides. On the right side the sound is somewhat less clear, shorter, and of a higher pitch, than on the left. From the fourth rib downward, on the right side, the resonance of the lung, on strong percussion, is found to be slightly deadened; near the sixth rib the perfectly dull sound indicates that the liver has been reached. On the right side, during full inspiration, the liver is pushed downward for the space of an inch or more; and the dull sound on percussion begins, therefore, lower down, and on a line corresponding to the displacement of the organ.

On the left side the heart deadens the sound from the fourth to the sixth rib, and, in a transverse direction, from the sternum to the nipple. This dull sound is lessened in extent during inspiration, and in cases of emphysema; indeed, under any circumstances in which the lung more completely covers the heart. Lower down, owing to the liver reaching over to the left side, and to the pres-

ence of the spleen and a portion of the stomach, the sound rendered on percussion consists of a mixture of the dull sound of the solid viscera and of the clear sound of the lung with the tympanitic sound of the stomach. The latter character of sound predominates when the stomach is empty. Over the upper part of the sternum, to the third rib, the percussion sound is slightly tympanitic; at the lower part, the heart and liver cause this tympanitic or tubular character of sound to give way to a dull sound.

Position exerts some influence on the results of percussion. On exchanging the recumbent for the erect posture, the pitch of the sound on the front of the chest is raised.

At the *posterior* portion of the chest the sound varies materially according to the part percussed. Directly on the scapulæ the sound is duller than between the bones, or than below their inferior angles. Beneath the scapulæ a clear sound is emitted as far as the lower border of the tenth rib; here, on the right side, the dulness of the liver begins. Strong percussion, however, causes the dulness to become manifest higher up. On the left side, below the angle of the scapula, the percussion sound may be tympanitic if the intestine be distended; or it may be rendered slightly dull by the spleen. In and under the axilla the sound is very clear. But on the right side, at the lower border of the sixth rib, dulness becomes perceptible; at a corresponding situation on the left side, the sound is clear or tympanitic from distention of the stomach; and at the ninth or tenth rib, dulness and a sense of resistance to the finger disclose the presence of the spleen.

#### AUSCULTATION.

Auscultation, or listening to sounds, informs us of the play of organs, and furnishes us with the most trustworthy means of studying their action. It is of signal service in affections of the chest. Indeed, any one who reflects upon the certainty with which cases of thoracic disease, which would have set at defiance the skill of a Sydenham or a Cullen, are now capable of being detected, even by comparative tyros, will gladly acknowledge the heavy debt of gratitude we owe to the genius of Laennec.

The method he practised was the *mediate*, or by the stethoscope.



Another method has since his time grown up,—the *immediate*, or the direct application of the ear to the chest. For ordinary purposes, this is the best; but where it is desirable to analyze circumscribed sounds, as in diseases of the heart, the stethoscope is preferable.

Stethoscopes are made of various materials and of different shapes. One of moderate length, with an ear-piece which fits the pavilion of the ear, and with the extremity not too much expanded,

is to be preferred. The material is of less importance. I like best those of gun-metal, introduced by Hawksley. Of late years double stethoscopes have been much employed. The instrument invented by Cammann, of New York, consists of two tubes, the extremities of which are placed into the ears. It has since been modified by making the tubes attached to the ear-pieces of flexible rubber. It possesses the advantage of rendering sounds louder: its great draw-



back is that it indiscriminately intensifies all sounds, whether in the chest or not, and its use is, therefore, at first confusing. A similar kind of stethoscope is the differential stethoscope of Alison, by which each ear receives simultaneously the sound from a different region.

In auscultating, the following rules are to be borne in mind:

1st. Place yourself and your patient in a position which is the least constraining and permits of the most accurate application of the ear or stethoscope to the surface. Above all, avoid stooping, or having the head too low.

2d. Let the chest be bare, or, what is better, covered only with a towel or a thin shirt.


3d. If a stethoscope be employed, apply it closely to the surface, but abstain from pressing with it. This may be obviated by steadying the instrument, immediately above its expanded extremity, between the thumb and the index finger.

4th. Examine repeatedly the different portions of the chest,




and compare them with one another while the patient is breathing quietly. Making him cough or draw a full breath is, at times,

FIG. 22.



The double stethoscope.

FIG. 23.



Alison's differential stethoscope.

of service, especially the former, when he does not know how to breathe.

*Intrathoracic auscultation*, or ausculting the organs of the body from within the body, has been suggested by Richardson\* as a new departure in physical diagnosis. To a long œsophageal flexible tube a double stethoscope is attached, and the tube is passed down the œsophagus, and, if necessary, into the stomach. He thinks it will prove particularly valuable in diseases of the heart and in thoracic aneurism, and in determining strictures of the œsophagus.

---

\* *Lancet*, Nov. 5, 1892.

*Sounds of Respiration in Health and in Disease.*

The ear applied over the trachea of a healthy person, and subsequently over the lungs, discriminates two dissimilar sounds, which may be severally taken as starting-points.

The first is plainly blowing, both in inspiration and in expiration. It is heard over the larynx and trachea; and in a slightly modified form, as a less intense and hollow sound, at the upper part of the sternum; and sometimes, owing to the closeness of large bronchial tubes to the surface, it is perceived between the scapulæ, on a level with their ridges. It is occasioned by air passing through the tubes, and is known as the tubular or the *bronchial* sound.

The sound over the lung-tissue is different: it is much softer, more gradually formed, of lower pitch, mainly inspiratory, and almost immediately followed by a shorter and far less distinct expiration. This is the *vesicular* murmur,—produced in the finest bronchial tubes and air-cells by their expansion and contraction. The expansion gives rise to the distinct breezy inspiration; the noiseless contraction of the elastic walls of the vesicles and the passage of air back into the smaller bronchial tubes cause the short, indistinct, sometimes almost inaudible expiration. But the vesicular murmur is not exactly alike at different parts of the lungs. It is, as a rule, better marked over the upper lobes than over the lower, and more clearly defined anteriorly than posteriorly. Nor is the sound of the two lungs precisely the same; a disparity may generally be noticed at the apices. Most authors describe the vesicular murmur as more intense on the right side. Investigations instituted to determine this point lead me to agree with Flint that the reverse is the case. More expiration, a higher pitch, therefore more of the bronchial element, is presented by the upper portion of the right lung; but a stronger, more vesicular inspiration belongs to the left lung.

The murmur of the air-cells, then, is the sound which the ear encounters when it is placed over the greater part of the chest. Bronchial respiration is constantly engendered in the tubes of the lung; but, either because it is overpowered by the sounds of the myriads of expanding air-vesicles, or because the pulmonary tissue is a bad conductor for a deep-seated sound, or perhaps because the sound requires consolidated tissue for its perfect production,

bronchial breathing is not heard over the chest, except at the very limited space indicated, unless the action of the air-vesicles have been suppressed.

Disease, however, gives rise not only to changes as absolute as suppression of the vesicular murmur and its substitution by a bronchial respiration, but also to certain modifications of the murmur, which serve as valuable guides in diagnosis. Thus, the vesicular murmur may be abnormal in its intensity, or in its rhythm, or it may have lost some of the elements of its distinctive character, such as its softness.

**Changes in the Vesicular Murmur.**—The changes of the murmur which are of importance may be summed up as follows :

ALTERATION IN INTENSITY...	{ Increased, or puerile breathing ; Diminished, or feeble respiration ; Absent respiration.
ALTERATION IN RHYTHM.....	{ Divided and jerking respiration ; Alteration of length of expiration relatively to inspiration.
ALTERATION IN CHARACTER..	{ Harsh respiration.

*Intensity.*—An increase of the vesicular murmur is called *supplementary* respiration, or, from its resemblance to the breathing of children, *puerile* respiration. It depends upon an increased action of the air-vesicles ; more air, or air with greater force, entering them. The sound is simply a loud, distinctly vesicular respiration ; both inspiration and expiration being augmented in duration and loudness, but retaining their relative length.

Puerile breathing is not in itself a sign of any disease. It indicates rather greater activity and energy of the part over which it is heard, which activity makes up for the deficient action of other parts. In this manner effusions compressing one lung, one-sided deposits, or obstruction of the bronchial tubes by secretions, necessitate a supplementary respiration in the healthy portion of the same lung, or in the other.

A diminution of the vesicular murmur, or *feeble* respiration, consists in a lessening of the whole sound without change in its character. But the relation of inspiration to expiration does not remain the same as in health. In the large majority of instances the inspiration suffers most, and the expiration does not dimin-



ish in proportion : a circumstance explained by reference to the states which occasion the diminished vesicular murmur. These are varied ; but their causes may be reduced to four :

1st. Any cause which obstructs the passage of air and prevents it from fully reaching the pulmonary tissue. Foreign bodies lodged in the trachea or bronchi ; affections of the larynx ; considerable thickening of the mucous membrane of a bronchial tube ; its compression, or the accumulation in it of secretions, or its contraction by a spasm,—all diminish the quantity of the air and the force with which it reaches the vesicles, and hence reduce the strength of the murmur.

2d. Deficient respiratory action. This may arise either from general debility ; or from impairment of the nervous force, as in paralysis ; or from local pain, as in pleurisy or in pleurodynia.

3d. Causes which interfere mechanically with the free expansion of the air-cells. Pleuritic effusions, by compressing the lung-tissue, will of course diminish the vesicular murmur ; so, too, will morbid growths, or malformation of the chest. Comparatively slight deposits in the pulmonary tissue of tubercle or of lymph obliterate some air-cells, and prevent others from unfolding, and, by having impaired their elasticity, diminish their sound. The same loss of elasticity happens in emphysema : the overdistended cells cannot expand much more, they are rigid and more or less fixed ; the vesicular murmur is therefore feeble.

4th. The respiratory murmur may be imperfectly transmitted to the ear, owing to intervening fluids or solids. To this category belongs the enfeebled murmur so constantly met with in fat persons.

As so many conditions occa-

sion a feeble respiratory murmur, it is only by association with other phenomena that it acquires much importance. Taking the diseases in which the sound is most frequently found, it may be

FIG. 24.




Diagram illustrative of the main forms of feeble respiration. *a*, from distention of the cells in vesicular emphysema; *b*, from deposits in the pulmonary texture; *c*, from a solid body (*d*) lodged in a bronchial tube, which has led to partial or, in some spots, to complete collapse of the air-vesicles.

stated that, if a feeble murmur be combined with dulness on percussion, it signifies a tubercular deposit, or a pleuritic effusion : the former, if at the upper, the latter, if at the lower part of the lung. If it be connected with increased clearness on percussion, distention of the air-cells is its cause. A vesicular murmur, feeble throughout both lungs, with the percussion sound unaltered, arises from general debility, or from obstruction of the upper air-passages. Where the feebleness of the murmur is found to change from place to place, it is dependent upon a loose foreign body which is shifting its position in the bronchial tubes. Joined to unwillingness to expand the lung, on account of the pain thereby brought on, feeble respiration denotes pleurodynia or beginning pleurisy.

*An absence of the vesicular murmur* is produced by the same causes, carried a step further, which occasion feeble respiration. Complete obstruction of the tubes by foreign bodies, extensive deposits in the pulmonary tissue, or its compression by large pleuritic effusions, arrest the vesicular murmur. But, practically speaking, there is only one condition in which we are apt to find it entirely wanting, and that is when, associated with flatness on percussion, the presence of a large collection of fluid in the pleura is attested. Extensive deposits in the lung-tissue, tubercular or lymphous, also suppress the sound of the air-cells; but they do not suppress all sound. The noise of the tubes, the bronchial respiration, then takes the place of the vesicular murmur, and denotes the perfect consolidation of the pulmonary tissue.

*Rhythm.*—The inspiration and the expiration may be altered as regards their rhythm. The inspiration may be broken up into little puffs,—jerking respiration; or both inspiration and expiration may be lengthened or shortened. But neither lengthening nor shortening of the inspiratory murmur has a distinct clinical value; and *jerking inspiration*, met with as it is in spasmodic affections, in hysteria, in pleurodynia, and in tubercular infiltrations, is present under too many different circumstances to have by itself much diagnostic significance. But if limited to the apex, it may serve to excite, or aid in corroborating, a suspicion of tubercular deposit. One modification of the rhythm is, however, of decided importance,—a marked increase in the duration of the expiratory murmur while the patient is breathing quietly.

*Prolonged expiration* denotes that the air has difficulty in getting out of the lung. It is detained in consequence either of loss of elasticity of the cells, or of an obstruction in the bronchi. The former state may be occasioned by overdistention of the air-vesicles, as in emphysema, or by deposits which impair their contractile power. In the first case, the prolonged expiration is associated with augmented clearness on percussion; in the second, with impaired clearness. Where the prolonged expiration is met with at the apex of the lung, in connection with dulness, it is for the most part caused by a tubercular deposit.

But a prolonged expiration from tubercular or from any other kind of infiltration is not simply the pure, prolonged expiration of deficient elasticity of the air-cells. It is something more. The solid material conducts a portion of the sound of the bronchial tubes to the ear; and bronchial breathing is nearly always best and earliest perceived in expiration. Thus, a prolonged expiration, when joined to dulness on percussion and to an inspiration still vesicular, is a sound partly vesicular, partly bronchial, and may be interpreted as consolidation of the lung-tissue; consolidation not sufficient to have obliterated all the air-cells, but sufficient to have obliterated some, and to have impaired the contractile power of others.

The obstacle to the exit of the air may reside wholly in the bronchial tubes. Such is the source of the prolonged expiration when the mucous membrane of the bronchi is swollen. Not only does this condition cause the air to be retained longer in the air-cells, but the resistance to the exit of the column of air brings out more of the bronchial sound. On the whole, then, an accurate study of the expiration is of decided value; and it is of great importance to impress on young auscultators the advantage of inquiring into the expiration separately from the inspiration.

*Character.*—A distinctive character of the vesicular murmur is its softness. From the moment it loses this, it begins to pass into the bronchial sound. Respiration which is wanting in softness is termed *harsh* respiration, or, to modify slightly a term introduced by Flint, *vesiculo-bronchial*. Harsh breathing is, in truth, a union of the vesicular and bronchial sounds; it is a vesicular sound mixed with some of the qualities of a bronchial sound,—a rough inspiration devoid of all the softness of the normal respi-



ratory murmur, with a prolonged, somewhat blowing expiration. Any affection which, without destroying the murmur of the vesicles, causes the sound in the bronchial tubes to be produced with greater intensity, or to be better transmitted, will occasion harsh breathing. Thus, it exists when the bronchial membrane is swollen, as in bronchitis, and still more frequently in diseases which are attended with compression of the lung-tissue, or with partial condensation, such as some stages of the forms of phthisis or of pneumonia. Being a transition murmur to bronchial, harsh respiration shares the properties of the latter in having its expiration more developed than its inspiration. It is true, the inspiration alone may be harsh, and the expiration not be much changed; but this is uncommon.

Harsh respiration may be confounded with puerile respiration, with sonorous rales, and with bronchial breathing. From the first it varies by its higher pitch, its roughness, its more distinct and blowing expiration; from sonorous rales, with which, however, it often coexists, by the absence of all vibrating or musical character. From bronchial respiration harsh respiration differs merely in degree: it is mixed with more of the vesicular sound, is less blowing in inspiration, and, when produced by condensation, is not associated, owing to the smaller amount of deposit giving rise to it, with so much dulness on percussion.

**Bronchial Respiration.**—Purely bronchial respiration may exhibit the same modifications as the vesicular murmur in respect to rhythm and intensity. But neither its rhythm nor its intensity is of significance; its character is. To hear well-defined bronchial respiration is, in the majority of cases, to meet with complete consolidation of the pulmonary tissue. It is thus that in extensive infiltrations and in hepatization of the lung we find the bronchial or blowing breathing so marked; particularly so in the latter morbid state, for the most distinctly blowing or tubular respiration is heard in pneumonia.

The bronchial breathing encountered in disease resembles more that heard in health over the larynx or trachea, than that heard over the larger bronchial tubes. It entirely replaces the vesicular sound, which has for the time being ceased to exist. It differs from the normal vesicular murmur by its higher pitch; by its occurrence equally in inspiration and in expiration; by its blow-



ing character, especially in expiration ; and by the pause between inspiration and expiration. Harsh respiration resembles it most ; but this or vesiculo-bronchial respiration is, as already stated, a transition from vesicular to bronchial breathing.

Whether bronchial respiration be owing, as Laennec taught, to a better transmission of the sound of the tubes through the solid lung ; or whether it be produced, as Skoda declared, by consonance, is not of much consequence for diagnosis. The important practical fact connected with this form of respiration is, that it happens when the pulmonary tissue is condensed, which, in the large majority of cases, takes place from exudations or deposits ; in a small proportion only, from compression by growths or effusions. At times bronchial respiration is also met with in severe cases of asthma in which the air does not expand the air vesicles.

A variety of bronchial respiration, at least so far as the quality of the sound determines the point, is that significant sign, *cavernous* respiration. This is essentially a blowing sound ; yet it is not always distinct during both inspiration and expiration, being often only perceptible in the one, and mixed in the other with gurgling. The question whether it can always be distinguished from bronchial breathing has given rise to much dispute. That cavities may exist without cavernous respiration being perceived, or, on the other hand, that, owing to peculiar physical conditions, cavernous respiration may have been heard where no cavities are present, cannot be denied. But that a sound is met with which is less diffused, much more hollow, and, above all, of much lower pitch than ordinary bronchial respiration ; that connected with it other signs of a cavity are found ; and that, under such circumstances, a post-mortem examination proves an excavation to have existed at the spot where during life the sound was detected,—are facts which equally cannot be denied. The peculiar sound occurs, and may be discerned by the ear ; and no theory, however cautious it may make us in our conclusions, can put aside the evidence of the senses.

Cavernous respiration is, then, a blowing sound of low pitch, circumscribed, alternating with gurgling, and deriving its chief character from the cavity in which it is formed. Hollow spaces of any kind—from abscesses, from bronchial dilatation, from breaking-down cheesy degeneration, from softening tubercle—give

rise to it. How it is to be distinguished from bronchial respiration has already been indicated. A student learns this sooner than he does to discriminate between cavernous breathing and the vesicular murmur,—the best proof that the ear recognizes a difference between bronchial and cavernous respiration, since the latter, as a sound of lower pitch, is more like the vesicular murmur. It is only necessary to recall, with reference to the distinction from the sound of the air-cells, that this murmur is devoid of all blowing quality.

*Amphoric* respiration is a blowing respiration engendered in a large cavity with firm walls. Its peculiar character is owing to an echo from the walls of the cavity. It may be humming and of low pitch, or decidedly ringing and metallic. An imitation of the sound, though only an imperfect one, is effected by blowing into an empty jar.

Amphoric or metallic respiration is always indicative of a large cavity ; the sound is rarely met with in phthisis ; much oftener is it heard over the cavity which is formed between the layers of the pleura, by the entrance of air.

Another variety of breathing connected with a cavity is the so-called *metamorphosing breath sound*, to which Seitz has called attention. It occurs only in inspiration, and consists of a very harsh sound, which lasts for about one-third of the period of inspiration, when it is continued as blowing respiration, attended with metallic echo or ordinary rales. The cause of the phenomenon is the air having to enter through a narrow opening to reach the cavity. Flint \* regards this sign as a variety of what he calls *broncho-cavernous* respiration. The sound of expiration in broncho-cavernous breathing is bronchial, high in pitch, and indicates a cavity situated near a portion of consolidated lung. In *vesiculo-cavernous* respiration the cavity is surrounded by comparatively intact pulmonary tissue, and this gives an admixture of vesicular sound.

**New, or Adventitious Sounds.**—These consist of sounds which have no analogue in the healthy state, and which are not, therefore, modifications of the normal respiration. Of this kind are the rales ; crackling ; the friction sound.

---

\* Lectures on Physical Exploration of the Lungs, 1882.

Nearly all *rales*, or rhonchi, are sounds which are generated in the air-tubes by the passage of air through them when contracted or when containing fluid. In the first case are occasioned dry, in the second, moist rales. Rales may occur in inspiration or in expiration, or during both acts. They may obscure or entirely take the place of the natural murmurs. They may have their seat in the upper air-tubes, or in any division of the bronchi. When in the larynx or in the trachea, they are called tracheal rales; of these the death-rattle is an example. When in the bronchial tubes, they are designated bronchial rales; and, as this is their

FIG. 25.




Diagram illustrative of rales. The narrowing in one division of the tube gives rise to dry, the fluid in the other to moist rales. The rales at the termination of the tube and in the air-vesicles are the crepitant or vesicular rales.

most frequent situation, the term *rale* means a bronchial rale unless the location be specially indicated.

Dry rales are, for the most part, produced by the vibration of thick fluids which the air cannot break up, and which temporarily narrow the calibre of the tube. When this narrowing exists in the smaller bronchial tube, the sound which results is high-pitched, —*sibilant*; when in the larger, unless the calibre be much altered, it is low-pitched, more musical, —*sonorous*. A similar dif-

ference is observed with reference to the moist or bubbling sounds. When the fluid is thin, whether it be mucus, blood, or serum, and breaks up into large bubbles, large bubbling sounds are occasioned; when it separates into small bubbles, small bubbling sounds are the consequence. The latter, for obvious reasons, generally take place in the smaller tubes.

Neither dry nor moist rales are persistent, but vary in intensity, or shift their position, as the air drives the liquid which gives rise to them before it. Dry rales are particularly prone to be dislodged by coughing. When they are uninfluenced by the act of breathing or of coughing, they do not depend upon the presence of secretions, but upon a narrowing of the air-tubes from the pressure of surrounding tumors or from a fold of thickened mucous membrane, or by a spasm.

It has just been stated that rales are, for the most part, produced in the bronchi by the passage of air through fluids there contained. This is their most frequent seat; but they are not limited to the tubes. Similar conditions may give rise to rales in other places. We find liquids in cavities breaking up into large, sharply-defined, bubbling rales, the so-termed cavernous rale,—*gurgling*; or having in cavities of considerable size a ringing *metallic* character; and again, the presence of fluid in the air-cells occasions a minute rale,—the *crepitant*.

This vesicular rale, or *crepitation*, is a very fine sound, or rather a series of very fine uniform sounds, occurring in puffs, and limited to inspiration. It resembles the noise occasioned by throwing salt on the fire. Its name indicates its seat. It is caused by the agitation of fluid in the air-cells or in the finest extremities of the bronchial tubes; or, to adopt a view now held by many, by the forcing open during inspiration of the air-cells agglutinated by the exuded lymph. The first stage of acute pneumonia is the state in which this rale is mostly engendered.

The rales, including crackling, may be thus grouped:

BRONCHIAL RALES.	{	Dry or vibrating sounds.	{	Low-pitched (sonorous).
		Moist or bubbling sounds.		High-pitched (sibilant).
VESICULAR RALES.	{	Crepitation. Crackling?	{	Large bubbling (mucous).
				Small bubbling (subcrepitant).
RALE OF CAVITIES.	{	Hollow bubbling, or gurgling. Metallic rales.	{	



*Crackling* is a sign closely connected with rales, and, though its mechanism is undecided, it is regarded as a rale. It consists of a few fine and readily-discerned crackling sounds which happen generally in cases of pulmonary tubercle, and of which, therefore, they are considered as diagnostic.

The distinction between crackling and the crepitant rale is one most puzzling to a beginner. Nor is there, in reality, any difference, except in the number of the sounds. Crackling is a few fine sounds limited to inspiration, and heard commonly at the apex of the lung. Crepitation is a number of fine sounds limited to inspiration, but more diffused, and heard generally at the base of the lung. The sound is similar because the conditions giving rise to it are similar. Both depend upon tenacious fluid or semi-fluid matter in the ultimate structure of the lung: in the one case it is tubercle or cheesy degeneration, in the other usually the lymph of beginning inflammation. The crackling which indicates softening, as of tubercle,—called by some authors moist crackling, by others clicking,—is a succession of sounds like small moist rales, only less liquid than these, because breaking-up tubercle is not very fluid. The fine or dry crackling of the earlier stages of phthisis corresponds, then, to a vesicular rale; the coarser, or moist crackling, to the small bubbling sound. When the bubbles become larger and larger, and cavities form, and the fluid matter in them is agitated by the ingress and egress of air, the large, bubbling, ringing rale of cavities, or gurgling, is occasioned. Dry crackling, moist crackling, and gurgling accord then with the crepitant rale, small bubbling, and large bubbling sounds, and happen in the progressive stages of infiltration and softening of deposits, and generally in those of a tubercular nature.

*Pleural friction*, or the sound due to the rubbing together of roughened pleural surfaces, consists of a number of abrupt superficial noises heard in inspiration and expiration, rarely in either alone. Its seat is not usually extended, for it is, as a rule, only audible over portions of the lower part of one side of the chest. Sometimes it is so creaking and intense as to be distinctly perceptible to the hand as well as readily recognizable by the ear. But it may be so much like crepitation that even long practice in auscultation will not enable us to determine at once whether the fine sounds we hear are the friction of a roughened pleura, or the vesic-

ular rales of an inflamed lung. It is easy to lay down as the distinguishing mark that of greater superficiality ; but at the bedside the difficulty remains the same, and is removed only by attention to the physical signs and symptoms accompanying the doubtful sounds.

Nor is it, in some cases, less perplexing to discriminate between fine friction sounds and fine moist rales. By the sound alone it is often impossible ; concomitant phenomena must be taken into account. A friction sound is mostly confined to a smaller space, and is uninfluenced by cough ; while cough changes the position and the distinctness of rales. Yet even this rule is not absolute. A fine friction sound may be temporarily increased during the deep breathing which follows the act of coughing ; on the other hand, the influence which cough exerts on the small moist rale is not so great as on the larger bubbling sound. As for the more marked character of moisture which a rale is said to possess, that only aids us in some cases. Where the secretions are viscid, it would require a sense of hearing more delicate than belongs to the majority of mankind to judge, by the application of this test, whether the sound we perceive is formed in the lung or on its covering. As the result of investigations undertaken to ascertain whether there is any positive difference, so far as the ear can detect, between some of the finer kinds of friction and fine moist rales, I have come to the conclusion that frequently little or none exists ; and still less is there between crackling and the crackling variety of friction sound, or between this and the vesicular rale. The features most at variance are : that the friction phenomena are not strictly limited to inspiration as are the vesicular rales, are not seldom coarser in expiration than in inspiration ; that they are less uniform ; and that their seat is more circumscribed. Their production nearer to the ear may assist us, but does not always. The reason why some of the finer friction sounds resemble so closely fine moist rales or crepitation is apparent when we reflect that the irregularities in the pleura may be slight, and be surrounded by fluid which keeps them moistened. Bruen has called attention to the value of making the chest-walls immovable.\* When the chest is fixed, especially at the lower two-thirds, by the hand of an assistant, and the ear or the stethoscope is applied over

---

\* Physical Diagnosis.

the doubtful sounds, they will be found to have disappeared if of pleural origin, but to be still discernible if rales.

The creaking or grating varieties of friction are much easier of recognition than the finer forms. Their discrimination from rales is readily effected by noticing the rubbing and harsh character they possess.

### *Auscultation of the Voice.*

Attention to the voice, as heard over the chest, is by some regarded as very important in examinations of the lungs. Yet the information derived from a study of the thoracic voice is very slight unless confirmed by other physical signs.

When the ear is applied to the thorax of a healthy person who is speaking, a confused hum is perceived, most distinct in adults who are possessors of a deep voice, and tremulous in the aged. Now, the normal *vocal resonance*, for by that name the ill-defined vibrations are called, is more marked on the right than on the left side, and corresponds to the vesicular murmur. Over the bronchial tubes a more concentrated sound strikes the ear. This, termed *bronchophony*, accords with bronchial respiration, and, when detected over the lung, denotes, with rare exceptions hereafter to be referred to, the same as bronchial respiration,—increased density of pulmonary tissue caused by pressure or by deposit. Any normal vocal resonance which is augmented passes by degrees into bronchophony, and has a meaning similar to it.

Of the sound known as bronchophony there are several varieties: the *simple bronchophony* just explained,—observed in pneumonia, or in any form of consolidation; the hollow, *cavernous voice*, or pectoriloquy; and the bleating variety, or *ægophony*. The latter, indicative of a thin layer of fluid between compressed lung and the ear, is a sign generally too transitory to be of much diagnostic value; and pectoriloquy, if by this be understood what Laennec meant,—complete transmission of articulated words,—is of no special significance, as it may be met with where no cavity exists. But if the term be applied to a well-defined chest-voice, of hollow character, and heard as such over a comparatively limited space, pectoriloquy is a distinct physical sign, and really deserves the name of cavernous voice. This is particularly true of *whispering* pectoriloquy. Over large cavities the voice is peculiarly ringing and *metallic*. The conditions which produce



amphoric or metallic voice are the same as those which occasion any of the amphoric or metallic phenomena. Be the respiration metallic, be the voice metallic, be the rales metallic, they are all caused by a cavity large enough and with walls firm enough to reflect, to echo the sound.

Bronchophony and amphoric voice are instances of increase and change of character of the normal vocal resonance. A *diminished vocal resonance* occurs when the lung is compressed by air or fluid, as in pleuritic effusions, or in pneumothorax ; or when it is greatly distended with air, as in extreme cases of emphysema. Clinically speaking, the sign is oftenest encountered in pleuritic effusions.

The vibrations of the voice may be *felt* as well as heard. The vibration detected by placing the hand over the thorax when the patient speaks, the *vocal fremitus*, is, like the voice, increased by all consolidation of pulmonary tissue, and diminished by fluid or air in the pleura. Its relations to the voice are, however, not uniform ; and sometimes with increased density of the lung-tissue there is no increased fremitus, although there is increased chest-voice. In women the sign is valueless ; indeed, its main importance is derived from *its absence* in cases of pleuritic effusions. Just as the chest voice, it is most marked on the right side.

Rales, when extensive, sometimes cause a vibration to be transmitted to the chest-walls, as do the fluids in cavities. The former phenomenon is called the *bronchial fremitus*, the latter the *cavernous fremitus*. A friction sound that may be felt is designated as the *pleural fremitus*.

### The Combination of the Physical Signs, and the Examination of Patients affected with Disease of the Lungs.

In the preceding pages isolated physical signs have been discussed. But if in the investigation of disease we were to trust solely to isolated signs, our conclusions would be incomplete and unsatisfactory. All the methods of physical exploration must be employed, the results obtained compared with one another, and the attending symptoms carefully inquired into and brought into connection with the physical signs, before a diagnosis is made.

A patient presents himself for examination. After having obtained the history of the case, it is well to notice his general appearance ; to scan the expression of his countenance ; to feel the



skin and the pulse ; to inquire into the nature of the cough and of the expectoration ; and to determine the existence of pain. The character and frequency of the breathing are noted. Next we proceed to a physical exploration. The chest is examined ; its movements, its size, are inspected,—if necessary, measured. Percussion is employed, then auscultation.

The manner of investigating by these methods has been detailed ; it need not here be repeated. But what may be repeated is, that there are two lungs ; that it is incumbent always to explore both, and, as we proceed, to compare the action of one with that of the other. Nor, even when the pulmonary affection has been made out, ought the examination to be stopped. The state of other organs and of the system must be inquired into, so as not, in the pursuit of a few physical signs in the lung, to pass by accompanying disorders of the heart, or liver, or stomach ; so as not to overlook vital conditions, compared with which, as respects the treatment, the physical phenomena often sink into insignificance. There are acute and chronic diseases of the lung. The physical signs of both may be the same ; but the general symptoms and the constitutional state attending them are not always identical. In truth, these are at times, in the same malady, so different as to render a remedy which is of use in one case, useless or worse than useless in another.

As many of the signs elicited by the various methods of physical diagnosis depend on the same physical conditions, they may be studied in groups. The following will be usually found to be associated :

## ASSOCIATION OF PHYSICAL SIGNS.

PERCUSSION.	AUSCULTATION OF RESPIRATION.	AUSCULTATION OF VOICE.	VOCAL FREMITUS.	PHYSICAL CONDITION.
Clear .....	Vesicular murmur or its modification.	Normal vocal resonance.	Unimpaired.	Lung-tissue healthy or nearly so ; at any rate, no increased density of lung-tissue from deposit or from pressure.
Dull.....	Bronchial, or harsh respiration.	Bronchophony.	Increased.	Solidification of pulmonary structure.
	Absent respiration.	Absent voice.	Diminished or absent.	Effusion into pleural sac.
Tympanitic.....	Cavernous or feeble, according to cause.	Uncertain ; cavernous or diminished.	Uncertain ; mostly diminished.	Increased quantity of air within the chest, or air confined in particular points ; states commonly due to a cavity, or to overdistention of the air-cells.

Amphoric or metallic.....	Amphoric or metallic.	Amphoric or metallic.	Mostly dimin- ished.	Large cavity with elastic walls
Cracked-metal sound.....	Cavernous respiration.	Cavernous voice.	Uncertain.	Generally a cavity communi- cating with a bronchial tube.

In adults these phenomena are commonly combined. In children, however, their connection is not so constant nor so apparent. Owing to the extreme elasticity of the thoracic walls and the naturally clearer sound of the lungs, the relations of percussion to auscultation are not the same as in the adult. Dulness, even where the condition exists for its production, is rarely as marked; nor is comparison between the two sides of the chest as valuable, since most of the acute pulmonary affections of childhood are more often double than those of adolescence. Again, the diagnosis of the diseases of the lung in children requires some knowledge of the disorders to which they are peculiarly liable, and, above all, great care and patience.

Among some of the peculiarities of the respiratory function, before the age of puberty, may be mentioned the greater frequency of breathing. Infants between two months and two years breathe irregularly, and about thirty-five times in a minute. Between the ages of two and six years the average number of respirations in the same space of time is twenty-three. The breathing is also of a different type from that of the adult; it is abdominal, and can be more readily counted by noting the rising and sinking of the abdomen than by watching the slight movements of the chest.

Of the methods of physical exploration, auscultation is in children the most applicable. It is far more so than percussion, and is to be practised first, since percussion causes the child to cry. The voice as well as the breathing may be advantageously listened to; and although the fretful patient will not or cannot speak, it can and does cry. From the cry, when studied with the ear applied to the thoracic walls, we may obtain the same indications as from the vocal resonance. The back of the lungs should be invariably examined. It is there that the mischief is mostly seated. Fortunately, also, this investigation does not occasion the same fear or struggling on the part of the little sufferer: hence it is better not to place the ear to the anterior portion of the chest until the posterior has been listened to. The position, too, in which the child is auscultated should vary with its age. Very young children may be examined either in a lying or sitting pos-

ture on the lap of their nurses, or may be held in the arms of an attendant, who is directed to present the different parts of the thorax successively to the ear of the physician.

Before proceeding to the discussion of the symptoms of pulmonary diseases and of the diseases themselves, let us group the latter according to their anatomical seat.

## DISEASES OF THE LUNGS AND THEIR COVERINGS.

BRONCHIAL TUBES...	{	Inflammation, or Bronchitis;	{	Acute ...	{	Of large-sized tubes.
				Of capillary tubes.		
		Dilatation ; Narrowing ; Diseases of bronchial glands ; Spasm of muscular fibres, or asthma.	{	Chronic.	{	Ordinary chronic catarrhal form.
						Putrid bronchitis.
						Fibrinous bronchitis.
LUNG-TISSUE .....	{	Congestion ;				
		Hemorrhage ;				
		Apoplexy ;				
		Œdema ;				
		Collapse ;				
		Hypertrophy ;				
		Inflammation, or pneumonia, in varied forms ;				
		Induration ;				
		Abscess ;				
		Cirrhosis ;				
		Gangrene ;				
		Emphysema ;				
		Tubercle, chronic and acute ;				
		Phthisis of different kinds ;				
		Cancer ;				
Deposits, such as syphilitic, melanic, etc. ;						
Parasites.						
PLEURA.....	{	Inflammation, or pleurisy ;				
		Empyema ;				
		Hydrothorax ;				
		Hæmothorax ;				
		Tuberculosis ;				
	{	Malignant growths.				
PLEURA AND LUNG..	{	Pneumothorax ;				
		Perforations and fistulous openings.				
WALLS OF CHEST....	{	Pleurodynia ;				
		Intercostal neuralgia ;				
		Abscesses, etc.				

### The Principal Symptoms of Diseases of the Lungs.

After having described the physical signs of pulmonary affections, it is necessary to inquire into the more prominent symptoms they occasion. At the same time, several of the disorders which are mainly recognized by these symptoms, and the physical signs of which are comparatively unimportant, will be dwelt upon.

Yet of the symptoms about to be mentioned, not one belongs exclusively to pulmonary diseases. We have met with some of them in studying laryngeal complaints; we shall meet with them again in examining the affections of the heart. And in investigating them here we shall not view them simply with reference to morbid states of the lungs, but shall indicate their general relations to diseased conditions, even at the risk of discussing what might in part be more appropriately discussed elsewhere.

The symptoms which it is proposed more specially to sift are dyspnœa, cough, and hæmoptysis.

**Dyspnœa.**—Dyspnœa means difficulty of breathing. It is accompanied mostly by a sense of uneasiness and suffocation, and by increased frequency of the respiratory act. But, strictly speaking, it is not correct to apply the term dyspnœa to mere increased frequency of breathing, for accelerated respiration and difficult respiration do not of necessity go hand in hand. The breathing may be slower than natural, yet laborious; it may be quick, and not impeded. Pneumonia furnishes often an example of this.

Dyspnœa depends upon various causes. Feeble persons are sometimes troubled with it after the slightest exertion. It may be temporarily produced by any bodily or mental excitement. It is observed when the play of the diaphragm is interfered with, and the lung cramped in its expansion. This is its cause in ascites, in abdominal tumors, and in pregnancy. It may occur in perverted innervation, as in hysteria, or in connection with cerebral affections, from a want of power in the respiratory muscles, or it may be due to morbid blood conditions, as in anæmia, scurvy, uræmia, and septicæmia. It is, however, most frequently met with as a prominent symptom of the disorders of the larynx and trachea, or of the heart, and in the various diseases of the lung



and pleura, whether idiopathic or secondary. Being common to so many morbid states, it is not diagnostic of any.

Dyspnœa is usually aggravated by position. When the patient lies on his back, the respiration becomes more difficult. The form of dyspnœa in which the sufferer is obliged to remain in the erect posture in order to breathe, is termed *orthopnœa*. This is witnessed in hydrothorax, in œdema of the lung, and in affections of the mitral or tricuspid valves. In bronchitis of the finer tubes the difficulty in breathing is very great; so, too, is it in pneumothorax, in emphysema, and in pleurisy, if the lung be extensively compressed.

Dyspnœa may come on in paroxysms, and constitute the only, or certainly the main, symptom of disease. This is the case in asthma.

*Asthma*.—Asthma consists mainly in a spasmodic narrowing of the bronchial tubes, caused by contraction of their circular muscular fibres. Its chief symptom is distress in breathing, occurring in paroxysms, and attended with wheezing. These spasms may be preceded by a feeling of suffocation, or they may come on suddenly. The patient wakes up out of his sleep, finds himself wheezing and with a fit of the disease fully on him. He continues to respire with great difficulty, sits upright in bed, or walks about the room gasping for breath. His look is anxious, the face pale, and the color of the lips shows that the blood is not properly aerated. In spite of the struggle to get air into the lungs, the chest moves but little, and when the ear is placed on it, no vesicular murmur is heard,—simply the same loud wheezing that is perceptible to the by-standers; or bronchial breathing at the upper part of the chest, or sonorous and sibilant rales are detected, due for the most part to the narrowing of the bronchial tubes, and disappearing with the spasm. These dry rales are chiefly expiratory, and the lungs are very full of air, and displace the diaphragm downward by several intercostal spaces. At the end commonly of some hours the fit passes off with copious expectoration, and as suddenly as it came. But it may last for days, ameliorating in the daytime, exacerbating at night, and only ceasing gradually.

The exciting causes of these bronchial spasms are various. In some persons there is no apparent reason for the attack; in others

it is brought on by the inhalation of irritating fumes or of disagreeable vapors. In some it is preceded by digestive disorder, or by bronchial catarrh; in others, again, an interruption to the free circulation of blood in the lung, or a disturbance in the sexual organs or in the urinary secretions, seems to occasion it. It is not unusual to find, on closely questioning patients, that for some time prior to the asthmatic paroxysm they have passed a scanty, dark-colored urine. During the attacks Leyden\* found in the sputum peculiar crystals, further on depicted, which by some have been thought to be the cause of the attacks. Another interesting fact connected with the paroxysm has been pointed out by Curschmann,—the presence on the turgid, swollen mucous membrane of the bronchioles of a characteristic viscid exudation. This generally shows in the sputum in little pellets that have a spiral structure, very easily discerned by the microscope. These spirals come especially from the finest bronchial tubes; they will be presently described more particularly.

Now, whatever be the exciting agent that calls the bronchial spasm into existence, the symptoms of the attack of asthma are the result of the spasm. Yet asthma is not often a pure neurosis. The seizure itself is the expression of perverted nervous action. But there are generally permanent conditions present, such as disease of the brain or medulla, of the heart, of the lungs, of the ovaries, of the kidneys, of the stomach, or of the nose, such as polypi or hypertrophic rhinitis, which act as constantly predisposing causes to these seizures, and lead to attacks, either by direct irritation of the pneumogastric nerves or through the medium of the reflex system. Emphysema especially is a fruitful source of spasmodic asthma. Asthma has been noticed to replace other neurotic affections, such as epilepsy.†

The detection of the causes inducing an asthmatic fit may be difficult; but the diagnosis of the fit itself is not so. No disease of the lungs or bronchial tubes is likely to be mistaken for it, because no disease of either gives rise to the same symptoms. The dyspnoea of pleurisy or bronchitis is not paroxysmal, nor is

---

\* Virchow's Archiv, 1872.

† Lancet, June 10, 1893.

it attended with wheezing. Some of the affections of the larynx and trachea bear a nearer resemblance; yet they, too, announce themselves by different symptoms. Asthma may be distinguished from *croup* by the entire absence of fever, and by its lacking the peculiar hoarse voice and cough which appertain to both forms of this malady. The age of the patient is also very different: asthma is as rare in a child as croup is in an adult. *Edema and spasm of the glottis* differ from asthma by the much more markedly paroxysmal nature of the difficulty of breathing, by the shorter duration of the seizures, and by the absence of the loud and continued wheezing. The sensations of the sufferer, further, indicate correctly the seat of the obstruction. And so they are apt to do in some of the *paralyses of the vocal apparatus*, where noisy dyspnœa happens, and is aggravated in paroxysms. Further, we are aided here by the aphonia, by the inspiratory character of the stridulous breathing, by the absence of chest rales, and by the obvious lesion seen in the laryngeal mirror. A large *goitre* pressing on the trachea may give rise to dyspnœa and to a noisy sound in breathing; but the cause of both is easily traced to the tumor in the neck.

The most deceptive condition is when the *glands of the neck* enlarge suddenly and press on the trachea. I had, some time since, a young man under my care for acute bronchitis. He was progressing favorably, when one day he presented himself, breathing with great difficulty, and each respiration attended with a noise like the wheeze of asthma. I should have regarded him as having been attacked with asthma had I not, in looking at his neck, detected the group of enlarged glands. Such cases are extremely rare, and belong to the curiosities of medical practice.

Marked dyspnœa may be occasioned by the pressure of an *aneurismal tumor*, or by an *organic disease of the heart*. But it is hardly necessary to enter here into a detailed description of the distinctive character of either of these forms of troubled breathing. The stridor and the persistent difficulty of respiration in the first, aggravated though it may become in paroxysms, and the constant want of breath in the second, are not likely to be mistaken for the wheezing and the paroxysmal dyspnœa of asthma. True asthmatic seizures may both produce and be produced by a disease of the heart. But what is called "cardiac asthma" is not



often a spasm of the bronchial tubes: it is usually only a temporary increase of the dyspnœa, dependent upon a decided obstruction to the circulation in the lungs, and not accompanied by wheezing.

So, too, *renal asthma* is only very rarely a true bronchial asthma, being usually an aggravated form of dyspnœa in connection with chronic Bright's disease. So-called *thymic asthma* is a severe dyspnœa connected with enlargement of the thymus glands. It is especially met with in children.

There is a peculiar form of difficulty of breathing connected with a *loss of power in the diaphragm*. The patient, when the disorder is fully developed, cannot make even the slightest effort without his being seized with a feeling of suffocation and his respiration being greatly accelerated. He cannot take a long breath, and often his voice is much enfeebled. But the most significant sign of the paralysis is, that during inspiration the epigastrium and the hypochondria are depressed, while the chest dilates; and the converse takes place during expiration. If there be merely a lessened power of the diaphragm, these phenomena are observed only during forced breathing; a paralysis of one-half of the muscle occasions them on one side alone. Duchenne adds another important diagnostic test of a paralyzed state of the diaphragm,—namely, that if the phrenic nerve be galvanized, the diaphragm acts again with proper strength, and during inspiration the abdomen rises simultaneously with the thoracic walls. To discriminate the cause of the impaired or lost muscular force,—whether this be due to a lesion of the nervous system, or to inflammation of the muscle or of the adjacent textures, whether produced by rheumatism or by lead poisoning, or originating in progressive muscular atrophy,—we have to rely chiefly upon the history of the case. In *rheumatism of the diaphragm*, an absence of the vesicular murmur over the lower portions of the chest; respiration effected by the upper ribs exclusively; tense, hard abdominal walls; want of power to strain so as to aid the bladder or intestines in expelling their contents, with darting, stabbing pain from the spine to the margin of the ribs on each effort to inspire,—have been particularly noticed.\* In *fatty degeneration*

---

\* Chapman, Boston Medical and Surgical Journal, July, 1864.



of the diaphragm, which often coexists with a fatty heart, we find, in its last stage, great distress and difficulty of breathing, and death may rapidly follow the embarrassed respiration.\*

Another form of dyspnoea is the so-called *Cheyne-Stokes respiration*. It consists in inspirations at first short, then deeper and more and more labored, till the paroxysm is at its height; then becoming shorter, and more and more shallow, until the breathing is suspended. The pause lasts from one-quarter of a minute to a minute, when the respiration begins again in the same manner, first faint, then a little stronger, then still stronger, until it reaches its height, when it again subsides in a descending scale, to end in the same stand-still. This kind of breathing is a very bad sign. It is apt to happen when from some cause the supply of arterial blood is cut off from the respiratory centre in the medulla, or from this and the adjacent vaso-motor centre. It is rare in diseases of the lungs, much more common in fatty heart, in disease of the aorta, in tubercular meningitis, in apoplexy and affections compressing the medulla, in uræmia, and in sunstroke. It may be found in cases that recover. Sometimes it is of long duration.†

**Cough.**—Cough is a sudden and violent expiration, having usually for its object the expulsion of some annoying substance from the air-passages. But it may be purely nervous, and unconnected with the presence of any irritating matter in the respiratory organs. There are several kinds of cough: according to the amount of expectoration, a cough is dry or moist; according to its origin, it is laryngeal, tracheal, bronchial, sympathetic, etc.

A *dry* cough is indicative of irritation. This is often seated in the larynx and the trachea, or in their vicinity, or in the bronchi, or in the lung itself. An elongated uvula, and many of the diseases of the nose or the pharynx, give rise to a dry cough: it happens, too, in pleurisy and in the earlier stages of phthisis. In disorders of the larynx and trachea the cough is attended with a peculiar shrill noise, or a hoarse sound. But the irritation may not be situated at all in the respiratory system. Affections of

---

\* Callender, *London Lancet*, Jan. 1867.

† As in the case of granular kidney, reported in the *Clin. Soc. Trans.*, vol. xxiii., 1890.

the liver, stomach, intestine, uterus, or brain will occasion an obstinate dry cough. It is also produced by dentition, by diseases of the ear, by the presence of worms in the intestinal canal, by disorders of the heart, and by thoracic aneurism. Again, it may be strictly nervous. The brazen cough of hysteria is dry ; indeed, nearly all sympathetic coughs possess a dry character.

A *moist* cough may succeed to a dry cough. The moist cough depends, for the most part, on the presence of fluid in the bronchial tubes or the lung-structure. It attends bronchitis with free secretion, œdema of the lung, the more advanced stages of all the forms of phthisis, and pneumonia when the exudation is breaking up. It is generally accompanied by a free expectoration, which varies in appearance and amount with the morbid state causing it.

Cough is frequently preceded by a sensation of tickling in the larynx, to which the patient is apt to refer his whole disorder. It is much affected by position. Lying down often increases its intensity. Sometimes a cough occurs in severe paroxysms. In various laryngeal affections, in abscess of the lung, in consumption, and in bronchial phthisis, such fits of coughing are observed. But in no complaint are they so constant as in whooping-cough.

*Whooping-cough.*—This is essentially a disease of childhood, and the result of an epidemic influence, or of contagion. The peculiar spasmodic cough succeeds to a catarrh of more than a week's duration. During the paroxysms the eyes fill with tears, the child's face is injected and anxious, and its whole appearance shows how it is suffering for want of breath. The air in the lungs is expelled by a series of abrupt spasmodic expirations, when a long-drawn inspiration, attended with a hoop, temporarily puts a stop to what appears to be threatening suffocation. The rest is, however, short. The cough recommences, and is again followed by the loud whooping inspiration. It continues in this manner until, after a copious expectoration of stringy mucus, or after vomiting, the paroxysm ceases, and a more lengthened calm ensues. These fits of coughing repeat themselves at varied intervals during the twenty-four hours. They are very frequent at night. Yet the child's health remains good, in spite of the violence of the attacks and the length of time they are spread over. The spasmodic cough lasts for weeks ; the hoop then ceases, the cough loses its ringing sound, and gradually leaves entirely. It is only in comparatively rare

instances that it persists, and is followed by the development of tubercles in the lungs ; just as it is only in exceptional instances, or in certain epidemics, that bleeding from the nose or lungs, petechiæ on the forehead, ecchymosis of the conjunctivæ, or convulsions happen during the violent coughing. In about one-half the cases the cough is violent enough to produce ulceration of or around the frænum linguæ, from the force with which the tongue is propelled against the teeth. Frequently the ulcer is covered with a grayish exudation ; it is never noticed before the paroxysmal stage is well established. Sugar is at times found in the urine. As an early symptom of whooping-cough, photophobia with dilatation of the pupils has been observed.\*

An affection of so long duration, marked by such a peculiar sign as a hoop, is easy of diagnosis. Yet there are certain conditions with which it may be confounded. In its first stage, before the characteristic cough sets in, it may be mistaken for *acute bronchitis*. There is, indeed, at this period, no means of distinguishing between the two disorders, except by taking into account the tendency to choke, to flushing of the face, and to vomiting in whooping-cough ; for it is only seldom that the cough possesses from the onset a decided ring. And bronchitis is in fact the most frequent complication, or, to state it more accurately, almost an essential element, of the malady. It is usually present in a mild form at the onset ; it outlasts the paroxysmal stage. At the height of this, a severe attack of acute bronchitis or of broncho-pneumonia may temporarily mask the special traits of pertussis. Again, occasionally acute bronchitis may exhibit paroxysms of spasmodic cough. But the want of the nervous element in the disease, the absence of the hoop and of the recurring flushing of the face as well as of the vomiting, the dyspnoea between the paroxysms, the decided fever, do not permit us to be long in doubt.

A disease less easy to discriminate from whooping-cough is tuberculation of the bronchial glands, or *bronchial phthisis*. It, too, produces a ringing paroxysmal cough. It, too, occurs in children. There is, however, this difference : the enlarged bronchial glands are apt to press on the surrounding parts. This becomes

---

\* Huguin, quoted in Brit. Med. Journ., Sept. 26, 1891.

manifest by the engorgement of the veins of the neck, by the lividity and puffiness of the skin, by the difficulty in breathing and in swallowing. The character of the voice, also, may change; and yet there may be no abnormal physical signs in the chest. But often there is dulness on percussion between the scapulæ, where the swollen bronchial glands lie, and impaired respiration in portions of the lung. The symptoms are those of pulmonary phthisis, with which the disease, indeed, may be associated: there are emaciation and the same loss of strength, the same sweating at night, the same hectic fever, the same tendency to diarrhœa. At times the affection of the glands induces a caseous pneumonia,\*—in reality tubercular. Now, when we compare these phenomena with those presented by whooping-cough, we miss the hoop, the vomiting accompanying the fits of coughing, the ulceration or tearing of the frænum of the tongue,—a symptom usual, at least, in decided cases,—the epidemic or contagious origin, and the distinct periods, first of catarrh, then of spasmodic cough, then of gradual decline. We see, on the contrary, an affection of more gradual and uniform progress, which often proves its existence by special signs, among which a venous hum, heard when the stethoscope is placed upon the upper bone of the sternum while the child bends back the head, has been particularly noticed.†

When emaciation, hectic fever, and marked cough are met with in the last stage of whooping-cough, it is always highly probable that this has been followed by a tubercular deposit. It is not likely that such cases will be mistaken for those instances of pulmonary consumption in which violent paroxysms of coughing occur. The age, the origin, the history, are different. Equally dissimilar are the history and the symptoms in other spasmodic coughs, such as that of hysteria, or of some laryngeal affections.

**The Sputum.**—The consistency of the expectoration varies very much. When it is viscid and tough, it contains a large amount of mucus or muco-pus, and depends generally upon inflammation or a high degree of irritation of the bronchial membrane or of the lung parenchyma. When it is less tenacious, it

---

\* Samuel Gee, St. Barth. Hosp. Rep., 1877.

† Eustace Smith, London Lancet, Aug. 1875.



has far less mucus, and a preponderance of pus. When fluid and full of air, it floats; when dense and without air, it sinks. Fluid sputum forms a homogeneous mass; dense sputum assumes a round or irregularly round shape. When these purulent masses float in a thinner expectoration, we have the coin-shaped or nummular sputum, so common in instances of pulmonary cavities.

The quantity of the expectoration varies greatly in different diseases of the lungs. In the most acute stages, or in spreading inflammations, it is usually small, and increases as the difficulty lessens. In bronchial dilatation, in pulmonary abscesses, especially when they burst, and in the voiding of a collection of pus in the pleura through the bronchial tubes, the amount discharged is very large, and consists almost entirely of pus.

The color of the sputum depends a great deal on its constituents. When mucous, it is white; when muco-purulent, yellowish or yellowish-green; when purulent, generally greenish or of a yellowish-green. It is also tinged by bile, by pigment, and by blood.

Sputum consists chiefly of water, serum, albumin, mucin, nuclein, and many salts, such as the chlorides of sodium and magnesium, the sulphates of sodium and calcium, and the phosphates and the carbonates of sodium, of magnesium, and of calcium. It has an alkaline reaction. In certain diseases, especially in putrid bronchitis and in gangrene of the lung, it contains ferment that acts like the pancreatic ferment.\*


Microscopically examined it exhibits pavement, columnar, and alveolar epithelium, leucocytes, blood-globules, various forms of crystals, such as the slender needles of the fatty acids, fibrinous coagula, bacteria, fungi, and elastic fibres. The alveolar epithelium is mostly of elliptic shape, and often shows little fat drops or pigment particles. The fatty acids and the elastic fibres are encountered in diseases involving disorganization of the lung-tissue. The fungi are most common in the sputum from cavities, in putrid bronchitis, and in gangrene. The leptothrix masses are readily recognized by their blue stain from a solution of iodine and iodide of potassium. Mould fungi, forms of leptothrix, and sarcinæ have been specially noticed. The latter are smaller than the sarcinæ ventriculi.

---

\* Stadelmann, *Zeitschr. f. Klin. Med.*, xvi., 1889.

*Elastic fibres* in the sputum are very significant; they indicate lung-destruction. They may be found as a bundle of fibres, or

FIG. 26.




Elastic fibres of pulmonary tissue, after treatment with caustic soda.

in the shape of the alveolar lung structure. In the latter instance they are even a more valuable sign than in the former. Elastic fibres are met with most frequently in tubercular lung-destruction, but they also occur in abscess and in gangrene of the lung, in cavities from bronchitis, and, according to Jaksch, in pneumonia, even where there is no abscess. The most certain way of finding the elastic fibres is to liquefy the sputum

by means of caustic soda, or to boil it in a solution of caustic soda or potassa; the particles that fall to the bottom of the vessel can be readily removed and placed under the microscope.

FIG. 27.



A spiral magnified.

*Spirals* are structures also possessing considerable significance. They have been studied especially by Curschmann, who traced them to an exudative inflammation of the bronchioles. They are most common in asthma, and bear a close relation to the Charcot-Leyden crystals which are often embedded among the coils. They have also been found in pneumonia. They consist chiefly of a substance allied to mucin, and are large enough to be detected in the sputum with the unassisted eye, though their peculiar structure and the central thread are recognized

clearly only with the microscope.

*Fibrinous coagula* are found in the sputum of pneumonia, of diphtheria that has extended into the lung, but especially in plas-

tic bronchitis, where they furnish a very valuable diagnostic sign. They are moulds of the ramifications of the finest bronchial tubes, whitish in color, of arborescent appearance, and consisting of fibrin. They are small in pneumonia, and do not generally occur in any number; should they be numerous the gravity of the case is greatly increased. They can be seen with the naked eye, or studied with a low magnifying power. In the description of plastic bronchitis a fibrinous mould is depicted.

Different *crystals* can be discriminated only with the microscope. We find cholesterin crystals in the sputum of tuberculosis and in abscess of the lung; the long, thin needles of margaric acid more especially in pulmonary gangrene and in putrid bronchitis. In the latter disease and in empyema breaking into the lung there have been also noticed by Leyden tyrosin crystals. Crystals of hæmatoidin follow a hemorrhage retained for a time in the bronchial tube. If the blood-crystals be conjoined to cells, they indicate, according to Jaksch, a previous hemorrhage; if any large number of them exist free, they point to a rupture of an abscess from neighboring parts into the lung structure. Uric acid crystals are encountered in the expectoration of gouty patients. The colorless, sharply pointed, octahedral or rhomboidal crystals described by Leyden and Charcot, and named after them, occur in various conditions, as in acute bronchitis; in the blood, intestinal tracts, and bone-marrow of leukæmia; but particularly in asthma. There they seem to have a direct connection with the attack. They are found in the sputum as little, round, yellowish bodies, but require a microscope for satisfactory study. They are soluble in warm water, in ammonia, in acetic acid, and in the mineral acids, and are supposed to be phosphate of ethylenimine.

Sputum very frequently contains *parasites*. In it may be found the scolices and free hooklets of echinococcus, the actinomyces fungus, the amœba coli, and others. Sputum full of amœbæ is thin and oily. They generally get into the lung from

FIG. 28.



Charcot-Leyden crystals.

hepatic abscess following amœbic dysentery. But the parasites of most consequence are the vegetable parasites, especially the bacilli and the cocci. Now, there are many bacilli and cocci that are not linked to any special morbid condition. But these have no particular diagnostic value; the pathogenic organisms are of the greatest importance, and most important is the tubercle bacillus, which is revealed by its significant action toward certain stains, an action that it shares only with the bacillus of leprosy. If the bacillus be exposed to an aniline dye dissolved in an alkali, unlike other pathogenic and non-pathogenic minute organisms, it retains the color on the subsequent addition of decolorizing reagents, such as acids and alcohol. There are many different tests based on this principle. The one of Koch, as modified by Ehrlich, is still, I think, the favorite; though the Ehrlich-Weigert method is also much employed. The Koch-Ehrlich method is as follows: A small drop of sputum is spread very thinly over the surface of a cover-glass, a second cover-glass is laid upon this, and the two are pressed together and then separated by sliding one over the other. The thin layer on the surface of the cover-glass we select to test is dried, by holding it over a gas or an alcohol flame, the side of the specimen being up. The dry sputum is now stained by letting the cover-glass lie for twenty-four hours at ordinary temperature in a saturated solution of aniline oil in water, made by adding the oil drop by drop to distilled water in a test-tube until the mixture becomes turbid, when it is filtered; a few drops of a saturated alcoholic solution of fuchsine, of gentian violet, or of methyl violet, are then added. At the end of this time all the component parts are stained, including the bacilli. The cover-glass is now lifted and immersed for a few seconds in a mixture of one part of nitric acid to three parts of water, until the preparation, previously red, becomes yellowish-green. The preparation is then placed in alcohol of seventy per cent. until no more color is given off; the color disappears, except that of the tubercle bacilli, which are red. We can then counter-stain the other parts blue by immersing the cover-glass for a few minutes in a two per cent. watery solution of methyl-blue or of malachite green, unless gentian or methyl-violet has been employed, when Bismarck brown must be used for the background. The cover-glass is then washed in absolute alcohol,



dried, and the preparation mounted in oil of cloves or in Canada balsam.

There have been many different processes proposed, among which those of Ziehl-Neelsen and of Gabbett, in which carbolic fuchsin instead of aniline water fuchsin is employed, are much used. The latter is excellent for rapid staining.\* The cover-slip is kept for from two to five minutes in a cold carbolic fuchsin solution, and then counterstained with a methylene-blue

FIG. 29.



Tubercle-bacilli in sputum, magnified about fifteen hundred diameters.—From a specimen prepared by Dr. Longstreth.


sulphuric acid solution, two grammes of methylene-blue to one hundred grammes of a twenty-five-per-cent. solution of sulphuric acid. The preparation is then rinsed off in water; the tubercle bacilli show the marked red stain.

As seen when stained, tubercle bacilli are fine rods, absolutely motionless, of the diameter of a human blood-corpuscle, and

\* See, for a clear description of the different processes, Schenk's *Manual of Bacteriology*, transl., London, 1893.

forming spores of oval outline. Their presence in any number is proof of the existence of tuberculosis ; when but few are found it

FIG. 30.



Tubercle bacilli.

is a question whether they may not have accidentally got into the air-passages.

Another valuable micro-organism in the sputum is the *pneumococcus*, especially the one described by Fraenkel as characteristic of pneumonia ; it has, however, also been detected in the saliva, in abscesses, in meningitis, and in empyema. Two cocci generally are found together. It is depicted in discussing pneumonia. Let us here examine only the process by which it is best discerned, which, moreover, is a most valuable one in the diagnosis of many micro-organisms, the process of Gram.

Gram's decolorizing method makes use of an aqueous solution of iodine and iodide of potassium : one part of iodine, two parts of iodide of potassium, and two hundred and fifty parts of water. The preparation is previously stained in aniline water solution of gentian violet, made in the usual way by shaking up in a test-tube filled with water one to two cubic centimetres of aniline until an emulsion is formed, which is filtered, and to which enough of a concentrated solution of gentian violet has been added to render the liquid of a dark color. The iodine solution is then washed out of the tissues ; the bacilli or cocci are easily isolated by the stain. The prepared section or cover-glass should be slowly but completely warmed in the aniline solution of the gentian

violet, either on the water-bath or over the flame, then laid for from one to two minutes in the aniline water solution of gentian violet, and subsequently placed in absolute alcohol until the color is discharged. The bacteria show the stain of gentian violet; the tissue may be double-stained red with picrocarmine or other dyes.

This method of Gram is of the greatest value in distinguishing micro-organisms. For instance, it separates the pneumococcus of Friedländer, which does not stain with it; and the bacilli of cholera, of typhoid fever, and of glanders do not retain the stain.

**Hæmoptysis.**—Sputa are streaked with blood in bronchitis, intimately admixed with blood in pneumonia; yet we do not call this hæmoptysis. It is only when a certain quantity of pure blood is expectorated that the complaint is regarded as hæmoptysis, or hemorrhage from the lungs. Now, a pulmonary hemorrhage may be an idiopathic affection; but it is not often so. It is mostly symptomatic of a grave disease of the lungs or of the heart. It is at times a discharge that takes the place of a suppressed flow of blood from another part of the body, as in vicarious menstruation. Among diseases of the heart, mitral disease is most generally connected with hæmoptysis. Tubercular disease of the lungs is by far the most common pulmonary affection in which hemorrhage happens. But it may also occur in gangrene, in bronchial dilatation, in abscess, in congestion of the larynx, and in the early stages of pneumonia. We also meet with it in purpura, in typhoid and typhus fevers, and in arthritic subjects.

When called to a person who has been spitting blood, we have first to solve the question, Where does the blood come from? It may issue from the nose or mouth; from the trachea; from the œsophagus or stomach; it may stream from an aneurism which has burst into the air-passages; or it may be that the lung is bleeding.

When in *epistaxis* the blood, instead of flowing out of the nostrils, flows backward, it is coughed up. But on the patient inclining forward, it will issue from the nose. The color of the blood is not florid; and it can be seen trickling down the pharynx. Inspection is of equal service when the blood comes from any part of the *oral* cavity; especially if it proceed from the gums. Their swollen state, their spongy appearance, and the readiness with which they bleed when pressed, point out at once the source of the hemorrhage.

Loss of blood from the *larynx* and the *trachea*, or from the *œsophagus*, is exceedingly rare ; and when it does occur, it is dependent upon some local lesion, such as an ulcer, or the presence of some foreign substance that has been swallowed. By attention to the history, then, we can recognize the cause and the seat of the hemorrhage. The blood itself furnishes no certain mark of distinction. Occasionally the hemorrhage takes place into the interior of the larynx, and only a very small quantity of blood is expectorated. Cases of hemorrhagic laryngitis are usually connected with catarrhal inflammation of the windpipe, with or without ulceration ; they are accompanied by severe dyspnœa, and with the laryngeal mirror the blood can be seen trickling down the windpipe.

When blood is vomited from the *stomach*, it is preceded by a feeling of weight and uneasiness in the epigastric region, and sometimes by decided nausea. The ejected matter consists of a dark grumous blood, thus altered by the gastric juice, and is often mixed with broken-down food. Its dark color is invariable, except where an artery has been laid bare by an ulcer, in which case a sudden discharge of florid blood takes place. There is not commonly more than one act of vomiting ; the blood which remains in the stomach passes into the intestines, and goes off with the stools. Hæmatemesis is attended with tenderness at the epigastrium. It is usually symptomatic of an organic affection of the stomach, liver, intestine, or spleen ; it may, however, depend upon the swallowing of irritating poisons ; or happen in fevers or in scurvy, or as a substitute for suppressed discharges.

The blood which gushes out of the mouth when an *aneurism* opens into the air-passages is red and arterial. It spurts out in jets, and the patient rarely long survives the hemorrhage. Should this not prove quickly fatal, we are seldom at a loss to determine the cause of the bleeding ; for we find the physical signs of the aneurismal tumor in the chest.

But when the blood comes from the *lungs*, it presents characters and is connected with symptoms totally different from any of those just mentioned. The bleeding is preceded by a sense of weight and of uneasiness in the chest. The patient perceives a saltish taste in the mouth and a tickling sensation in the larynx, when suddenly the mouth fills with blood, or after a slight cough



he expectorates a quantity of light-red and frothy blood. His anxiety becomes great; the skin is covered with a cold sweat; the pulse is quick and bounds under the finger. He spits up more blood, and this continues to come up at varying intervals and in changing quantities all day, or for several days, or even for a much longer period. It is at first pure blood, or mixed with the sputum; is red and not coagulated, and frothy, except when the hemorrhage is very profuse. But after one or two bleedings, the matter which is coughed up contains dark clots, being the blood which has been retained somewhere in the air-passages since the previous attack. The blood is never, at the onset of the hemorrhage, dark and grumous; yet in rare cases it has more of a venous than of an arterial hue. The amount which is brought up at one bleeding ranges from one to two drachms to as many pints; but the quantity that comes out of the mouth is by no means an index of the quantity extravasated. The blood may be effused into the pulmonary structure, and but little be expelled.

After the description above given, it is unnecessary to point out the marks of discrimination between blood ejected from the lungs and blood from other parts. The symptoms are different; the blood itself is different. And listening to the chest detects bubbling sounds in the air-tubes; still, to find these is not requisite for the diagnosis of pulmonary hemorrhage, and indeed, while the bleeding is going on, the patient's welfare forbids an extended thoracic examination. But as soon as circumstances permit, that examination becomes of immense value by showing us with what morbid state the hemorrhage is connected, and whether the bleeding is symptomatic of a disease of the heart or the lungs, or does not depend upon either. It is mostly owing to an affection of the heart or the lungs, and is exceedingly prone to be repeated.

Yet the lungs may bleed frequently without there being an organic lesion within the chest to account for the hemorrhage. I had, some years ago, a patient under my care who had been spitting blood daily for five years. Although enfeebled by the loss of blood, his general health remained good. His lungs and heart appeared to be sound. Another patient had pulmonary hemorrhages at varying intervals for eighteen months. He finally died of exhaustion; but he never presented any physical signs of tho-

racic disease. An examination of the body was, unfortunately, not permitted. But in the case of a gentleman that I had watched for years, the repeated hemorrhages were found at the autopsy to be unconnected with disease of the lungs. He died of an acute disease complicated with pleurisy.

In these instances the hemorrhages recurred often. But we meet with robust persons in whom the loss of blood follows active exercise or exertion and is not apt to be protracted or to be repeated. In such cases, of which I have seen a number in soldiers sent to hospitals after the fatigue of a long march or the excitement of a battle, simple congestion of the lungs is probably the cause of the disorder.

Except under the circumstances mentioned, hæmoptysis is a grave symptom. It is not dangerous as regards its immediate termination, but dangerous because it is, for the most part, the indication of a serious malady. Few die as the direct consequence of the hemorrhage, but many die of the disorder of which the hemorrhage is the consequence.

### Diseases in which Clearness on Percussion is met with and constitutes a Valuable Sign.

Some of these ailments are acute, others chronic; and nearly all have as their prominent symptom a cough, and are affections, or follow affections, of the bronchial tubes.

**Acute Bronchitis.**—This is an acute catarrhal inflammation of the bronchial mucous membrane, which occurs idiopathically, or happens as a secondary complaint in the course of fevers, of rheumatism, and of cardiac disorders. Let us examine the manifestations of the idiopathic malady.

Bronchitis varies considerably according to the size of the tubes involved. When the smaller tubes are affected, a disease called capillary bronchitis, or suffocative catarrh, is established, the prognosis of which is very grave, and the diagnosis of which presents points for special consideration.

The symptoms of acute bronchitis of the *large and middle-sized tubes* are, a sensation of tickling in the throat, soreness or pain behind the sternum and along the lower ribs, a slight oppression in breathing, and a paroxysmal cough. Let us add to these pain in the limbs, coryza, and a fever of moderate intensity, rarely

reaching  $103^{\circ}$ , and we have the main phenomena met with during the onset and at the height of an attack of ordinary acute bronchitis. The fits of coughing in the earlier stages are followed by a clear, frothy expectoration, which, as the cough becomes looser and less fatiguing, changes from an almost transparent fluid to a yellowish or greenish sputum. This may be uniform or streaked with blood; it may be small in amount, or in considerable quantity; and it consists chiefly of pus-cells and of large, round, alveolar cells with some blood-corpuscles. The fever soon leaves; but long after it has ceased, the patient still has a cough and expectoration, both of which only gradually disappear.

The physical signs may be inferred from the lesions. As there is no condensation of pulmonary tissue, there is no dulness on percussion, the thickening and injection of the bronchial mucous membrane not being sufficient to modify materially the normal resonance. But these conditions must alter the respiratory murmur. They bring out more of the bronchial element of sound, hence more expiration with the coarser inspiration,—in other words, a harsh respiration; or the swelling obstructs the entrance of air into the air-vesicles, and enfeebles the vesicular murmur. Again, new sounds, the rales, are produced; first dry, then moist. This succession of the rales is, however, not absolute, and depends, to a great degree, on the density of the fluid in the bronchial tubes. Dry rales, mixed with moist, may be perceived even in the later stages of acute bronchitis, and long after the febrile signs have ceased. In fact, the tenacity alone of the exudation determines the nature of the rales, and even somewhat their exact character; for every dry rale is not precisely like every other dry rale, nor every moist rale equally moist. With reference to size, the sonorous rales and the large bubbling sounds prevail when the disorder attacks the larger tubes. Sometimes, when the bronchial inflammation is severe and extensive, we find a sound which seems to be neither a dry nor a bubbling rale, but rather a compound of both,—a dry sound, yet not continuous, giving the idea of being caused by the breaking up of fluid. Or, there may be a mixture of the sounds of respiration with the rales, occasioning a peculiar kind of breathing,—one in which we can recognize neither a distinctly vesicular nor a distinctly bronchial element, nor a well-defined rale. All these states are dependent upon the amount, and, above



all, upon the condition, of the exudation in the bronchial tubes. But they indicate nothing beyond the fact that there is an exudation present which is very large in quantity and tenacious in character. When the sounds are of the indeterminate nature just alluded to, the vibrations produced in the tubes are apt to be transmitted to the parietes of the chest, occasioning with each respiration a marked fremitus.

The diagnosis, then, of acute bronchitis is determined by the cough, the fever, the expectoration, and the signs of clearness on percussion, diffused rales, or harsh respiration. From all those diseases of the lung which result in the consolidation of the pulmonary tissue, such as *pneumonia* and *tuberculosis*, we distinguish bronchitis by the absence of dulness on percussion. Some cases of *acute tuberculosis*, on account of the sudden invasion of the malady and the general diffusion of the physical signs, are liable to be mistaken for acute bronchitis; but the different progress of the disorder usually clears up all doubt. Error in diagnosis is more likely to arise from the habit, when the signs of bronchitis have been made out, of not looking further; forgetting, in the attention to the disease within the thorax, the various morbid states which bronchitis may accompany, and particularly its frequent association with the eruptive fevers, such as measles and smallpox, with typhoid fever, and with malaria.

*Capillary Bronchitis*.—This is a disease of the aged and of young children. It begins with an acute inflammation of the larger bronchi; or the disorder may from the onset affect the smaller tubes. In either case, signs of obstructed circulation soon manifest themselves; there is lividity of the lips and cheeks, with hurried breathing, a rapid pulse, an anxious countenance, great restlessness, moderate fever temperature, and a cough, followed by viscid expectoration. As the malady advances, the color of the skin and the mucous membranes shows more and more the want of properly-aerated blood; the sputa cease with the failing strength; and in old persons delirium and coma, in young children convulsions, mark the closing struggle.

The physical signs are those of ordinary bronchitis, but modified by the seat of the malady. High-pitched whistling sounds, accompanied or superseded by very fine moist rales, denote the smaller size of the tubes involved. The resonance on percussion



is clear, or very slightly different from that of health. When materially duller, it indicates that the pulmonary tissue itself shares in the inflammation, or that it has been exhausted of its air and has collapsed.

The parts of the lung which the physical signs prove to bear the brunt of the disease are the lower lobes. In the upper there may be large rales and some fine ones; but it is low down and at the posterior portion of the chest that the fine sounds are most abundant. Yet when the inflammation is extensive, and the accumulation of secretions and morbid products great, quantities of small rales are heard at every part of the chest.

From this description of capillary bronchitis it will be apparent that it differs from *ordinary acute bronchitis* in the greater tendency to prostration and to suffocation, in the signs of imperfect aeration of the blood, and in the fineness of the rales.

Like the more usual kind of acute bronchial inflammation, capillary bronchitis is liable to be mistaken for *acute lobar pneumonia* and for *phthisis*. And in the majority of cases the same rules serve for its discrimination; the absence of percussion dullness and the diffusion of the morbid sounds are here again of the utmost value. The rapidity of the attack and the signs of suffocation might mislead into the supposition of the existence of œdema of the glottis, of laryngitis, or of croup; errors in diagnosis which the detection of fine chest rales will prevent.

Capillary bronchitis which really merits the name is a very rare disease, though I believe it to exist. What is called capillary bronchitis is for the most part *catarrhal pneumonia* or *broncho-pneumonia*, one of the most common, as it is one of the most fatal, of the diseases of childhood. Like capillary bronchitis, the disease affects both lungs. It is commonly observed in connection with measles, whooping-cough, influenza, or diphtheria; it is especially seen in children previously in impaired health or scrofulous or rachitic. It is apt to be attended by cerebral symptoms,—indeed, it may set in with these,—by rapid breathing and paroxysms of dyspnoea, and by high and irregular fever, ranging between 102° and 105°. As the inflammation is limited to the lobules, it yields but imperfect signs of consolidation. The bronchial breathing is rarely very marked; crepitant rale is not usually perceived, or can scarcely be distinguished from the small

bubbling sounds of fine bronchitis ; and, from the usual association with inflammation of the fine bronchial tubes, it is in individual cases often extremely difficult to say whether portions of the lung-tissue are consolidated. Theoretically, broncho-pneumonia may be distinguished from capillary bronchitis by the dulness on percussion ; practically, this aids but little. Dulness on percussion is in children difficult to elicit ; and, again, a dulness may be temporarily produced in capillary bronchitis by collapse of the pulmonary tissue.

Broncho-pneumonia may or may not be preceded by bronchitis of the fine tubes. We may suspect that the inflammation has infiltrated the lobules, if the breathing be very rapid, the fever severe, or the temperature, which is rarely above  $102^{\circ}$  in the preceding bronchitis of the finer tubes, rise suddenly by several degrees ; if the cough lessen as the pneumonia develops ; if laryngeal symptoms arise ; and if, in addition to rales, not very diffused, spots of dulness, which do not change their seat, and do not disappear under respiratory percussion, be discerned, and plastic pleurisy appear as a complication. On the other hand, when there are early most marked signs of deficient aeration of blood ; when the child seems to suffocate from want of power to expectorate ; when a multitude of fine dry and moist sounds are heard at every part of the chest, and little or no corresponding impairment of resonance on percussion is detected,—we know that the capillary bronchi are extensively filled with pus and morbid secretions, and that true suffocative catarrh is threatening life. Capillary bronchitis is a rapid disease ; catarrhal pneumonia runs a much slower course, lasting perhaps weeks, and showing a temperature record that is marked by great alternations between morning and evening.

**Chronic Bronchitis.**—The symptoms and signs of chronic bronchitis are not very different from those of the ordinary form of acute bronchitis. The duration of the complaint and the absence of fever, except during marked subacute or acute exacerbations, are the chief distinguishing elements. Yet the cough, although on the whole chronic, is far from being constant. It may disappear almost altogether, and then reappear with more than its previous severity ; and this state of things may go on for years, undue exposure and change of season aggravating the disorder.

The sputa vary, even more than in acute bronchitis, in tenacity and quantity. There may be merely a small quantity of yellowish matter expectorated in the morning, or an almost continued flow from the bronchial tubes,—*bronchorrhœa*. The physical signs differ accordingly. A harsh or feeble respiration, and few or many, either dry or moist, rales, are present, in conformity with the state of the bronchial mucous membrane and of its secretions. The sound on percussion is clear, and this, with the diffusion of the signs discerned on auscultation, is of great importance. Excessive secretions somewhat impair the pulmonary resonance, but only temporarily; for with the shifting secretions shifts the very slight dulness.

One of the most important points in the diagnosis of chronic bronchitis is to attend to the manner in which it arises. It may follow a seizure of acute bronchitis, or be the result of recurring attacks of subacute character; it may appear as a primary affection, or it may follow the exanthemata; or, again, it may complicate some previously existing disorder, as Bright's disease, rheumatism, lithæmia, gout, psoriasis, or eczema, and be directly traceable to the constitutional taints of these maladies; and its symptoms will vary and be influenced by those of the general malady to which it is subordinate.

In the ordinary idiopathic malady the general health, as a rule, suffers but little. In some instances, however, emaciation takes place, and the disease simulates phthisis. This is particularly the case in the bronchial affections among knife-grinders and coal-miners, also in those of granite-masons, of sandpaper-makers, of flax-dressers, and of potters. The resemblance becomes still greater when superadded bronchial dilatation and fibroid induration of the lung produce physical signs like those of pulmonary consumption.

A chronic *catarrhal inflammation of the mucous membrane of the nose* may be mistaken for chronic bronchitis, with which, indeed, it may coexist. But when occurring uncombined, there are no rales in the chest or altered breathing-sounds indicative of disorder there, though there may be a cough, from the throat being also affected. The secretion, too, from the nose is very copious and of muco-purulent character, the upper part of the nose looks somewhat flattened, and the sense of

smell is impaired,—not one of which signs is met with in chronic bronchitis.

It seems almost unnecessary to speak of the differential diagnosis between chronic bronchitis and *rose cold* and *hay asthma*. The coexistence of marked signs of irritation of the eyes, the nose, and the throat; the appearance of the distressing affections at a particular period of the year; the fixed time in which they run their course; their occurrence in those of neurotic constitution and having an irritable nasal mucous membrane; the almost instant relief on leaving the regions where the attack has been brought on and on reaching favorable localities; the depression of the nervous system; and, on the other hand, the less decided signs of bronchial affection,—clearly distinguish the maladies.

We meet occasionally with a form of bronchitis in which the expectorated matter is solid. This *plastic bronchitis* presents all the usual signs and symptoms of bronchial inflammation. It may be chronic, or it may be acute. It is most frequently chronic, with occasional acute or subacute exacerbations. The disease extends in this way over weeks, months, or even years, and is apt to end in complete recovery. But in its acute form it is a complaint of great danger and accompanied by much dyspnoea, and has led to death by suffocation.\* Males, as we find by looking at the cases which Peacock† has collected, are more often attacked than females. The same carefully-collated observations show that the disorder affects more commonly the upper than the lower part of the lungs. As regards the physical signs, Fuller,‡ who has met with a number of well-marked examples of the complaint, states that there is weakness or entire absence of breathing over the affected portions of the lungs, and that, from attending collapse, complete and rapidly-developed dulness on percussion may ensue. But the only absolutely diagnostic phenomenon is the peculiar membranous material expectorated. In form this may be either in thin shreds, or moulded into an accurate cast of a bronchial tube

---

\* Andral, Clinique Médicale.

† Transactions of the Pathological Society of London, vol. v.

‡ Diseases of the Chest.




and its ramifications. The expectoration of the firm bodies is sometimes attended with copious hæmoptysis. The casts consist of layers of fibrin in which leucocytes and alveolar epithelium are embedded. Leyden's crystals and Curschmann's spirals may also be detected. The disease is most apt to occur in the spring months, and the chronic cases are not infrequently associated with tuberculosis.

The little round solid pellets which consumptive patients or even some persons in good health cough up from time to time are the result of a plastic bronchitis on a very limited scale. A kindred disease to plastic bronchitis has been described as "bronchiolitis exudativa." The sputum is grayish and very tenacious, and full of spirals which come from the bronchioles. Gradually-increasing dyspnœa and attacks of asthma are prominent symptoms.\*

Another variety of chronic bronchitis is *putrid bronchitis*. This may happen in connection with bronchial dilatation or with chronic pneumonia, or without these conditions; occasionally it appears after a suppurative pleurisy which has broken into the lung. There is fever with irregular temperature; at times chills occur. The distressing cough is followed by a copious, half-liquid sputum, extremely offensive, and containing little yellowish plugs, the so-called Dittrich's plugs. The peculiar odor is thought to be due to a micro-organism, especially to a short, slightly-curved bacillus described by Lummitzer.† Cases of putrid bronchitis may be mistaken for gangrene of the lung; but the odor is different, and they lack the physical signs of lung destruction and elastic fibres in the sputum. We must, however, bear in mind that putrid bronchitis may terminate fatally

FIG. 31.



Cast from a case of plastic bronchitis.

\* Curschmann, Deutsch. Arch. für Klin. Med., Nov. 1882.

† Wien. Mediz. Presse, May, 1888.

by induced pneumonia or pulmonary gangrene. Sometimes it produces death by metastatic abscess of the brain.

**Emphysema.**—A distention of the air-cells is a frequent sequel of chronic bronchitis. It may happen in only one lung; but the air-vesicles of both are usually distended. The effect of this is to obliterate some of the capillaries, and to interfere with a flow of blood through the lungs. From this proceed, to a great extent, the feeling of constriction and the dyspnoea, the anxious look, the bluish lip, of emphysematous patients, and the tendency the disease has to produce dilatation or dilated hypertrophy of the right side of the heart.


Emphysema is essentially a chronic malady; but in its course subacute attacks of bronchitis occur which much augment the difficulty of respiration. The embarrassment in breathing is, indeed, the most prominent of the symptoms. It is not so much the difficulty of getting air into the lung, as it is of getting it out, which annoys the patient. He breathes as if he had no object but that of forcing the air out of the pulmonary tissue. And this task is often aggravated by spasmodic narrowing of the bronchial tubes: hence it is very common to meet with the loud wheezing of asthma in those whose air-cells are permanently dilated. In long-standing cases of the disease the patient looks cachectic, is cyanosed, the shoulders are rounded, the chest is barrel-shaped, and dropsy of the feet is noticed. There may be also a chronic cough, which may be dry and occur in paroxysms of marked intensity.

The physical signs of emphysema are easily deducible from the pathological conditions. The distention of the lung-tissue explains the great prominence and fulness of the chest, and the displacement of the liver or heart. The ringing clearness on percussion—at times almost tympanitic in its character—and the increased resistance to the finger have the same cause. Nor is it difficult to understand how the loss of elasticity in the dilated air-cells will give rise to an unchanged note on respiratory percussion, to prolonged expiration, and to a feeble inspiratory murmur. If bronchitis coexist, the signs on auscultation are necessarily somewhat altered. The respiration is harsh, or intermixed with dry and moist rales. The former especially assume great prominence, and are heard as sonorous, or still oftener as sibilant, rales,

during the prolonged and labored act of expiration. Occasionally a crackling sound is heard in emphysema.\* When the emphysema is partial, all these signs are limited; when it is more general, they are diffused.

If the upper lobe of the right lung or the lower lobe of the left, which, according to Louis,† are the parts most frequently

FIG. 32.



Appearance of the chest in a patient suffering from a high degree of emphysema. The heart is displaced. The other physical signs are extreme percussion clearness; a feeble, hardly audible inspiration; a very prolonged expiration.

affected, be emphysematous, the visible local bulging might mislead into the idea of the prominence being due to an aneurismal tumor, or to the presence of fluid in the pleural cavity. Any doubt will, however, be dispelled by a careful examination of the chest. The dulness over an *aneurismal tumor*, its pulsation, and its sounds, are different from the exaggerated clearness on percussion and the changed respiratory murmur of an emphysematous

\* Gerhardt, Berlin. Klin. Wochenschr., March 12, 1888.

† Mém. de la Soc. Méd. d'Observation, tome i.

lung. *Pleuritic effusions* produce a bulging at the lower part of the thorax. But, although there may be a very clear, or rather a tympanitic, sound above the fluid, the absolute dulness over it shows that the prominence of the chest is not caused by distended air-vesicles. When the emphysema is extended and general, there is little or no action of the diaphragm, and the complaint gives rise to displacement of the liver or heart; and this circumstance, taken in connection with the dilatation of the chest and the dyspnœa, brings the malady into a category of affections which will be examined hereafter. When considering this group, we shall return to emphysema, and point out its distinguishing marks from the disease for which it is most likely to be mistaken,—*pneumothorax*. We shall only here add that the affection of the heart, the torpid displaced liver, and the presence of albumen in the urine, in emphysematous patients, may call away attention from the primary pulmonary cause.

An effusion of air may take place into the areolar tissue uniting the lobules. There are no physical signs peculiar to this *interlobular emphysema*; they are exactly the same as those furnished by dilatation of the air-cells, except that a dry friction-sound and a large, dry crackling, both of which occur occasionally in vesicular emphysema, are much more common. Nor are there any general circumstances specially indicative of the disease, save its suddenness, and the external emphysema which follows. The latter is detected under the jaw, or at the base of the neck, and yields a peculiar crepitation. Yet the extravasation of air into the areolar tissue of the neck is not a constant attendant on the extravasation of air in the lung. Besides, the possibility of a crepitating swelling in the neck being due to a rupture of the bronchial tube or of the larynx must be borne in mind.

The rupture of the air-cells which gives rise to interlobular emphysema is brought about by any severe effort, by violent coughing, by laughing, or by the throes of parturition. It has also been known to happen in the course of pneumonia or of pulmonary hemorrhage and to have caused sudden death. Its most frequent association, however, is with hooping-cough.

A *compensatory emphysema* is met with when distention of the air-cells takes place in the unaffected lung or in an unaffected lobe. It generally occurs at the anterior margins, and is grad-



usually developed by the high tension in the air-vesicles of those portions of the lungs which have to do more duty. It is thus chiefly met with in cases of extensive pleural effusion, in pneumothorax, and sometimes in pneumonia. The physical signs are those of ordinary emphysema.

In all the disorders which have just been treated of, the resonance on percussion has been dwelt upon as a most valuable sign. Before proceeding to consider the diseases in which dulness is encountered, a few words may here find their place on a morbid condition in which clearness rapidly gives way to dulness, and dulness changes quickly back into clearness. As, moreover, the complaint to which I allude—*collapse of the lung*—bears a close connection with bronchitis and emphysema, and has been made to play an important part in the explanation of some of their symptoms and complications, its consideration is at this time fitting.

In noticing that dulness on percussion sometimes appears in the course of capillary bronchitis, it was remarked that this does not of necessity show that the inflammation has extended to the lobules; it may be owing to the air in the lung being exhausted and to the pulmonary tissue having collapsed in consequence of accumulations in the bronchial tubes. These accumulations occasion collapse by shutting up the tube through which the air reaches the air-vesicles. No air can enter; the residual air is gradually exhausted, and the disordered portion of lung is reduced to a state as if it had never breathed. But, although in the majority of instances this condition of things is brought about by catarrhal secretions in the bronchial tubes which cannot be expectorated, it would be a mistake to suppose that these are always present. Any want of power to fill the cells of the lung with air may lead to their collapsing. In some of the typhoid forms of acute and chronic diseases, in the pulmonary congestions of the aged and enfeebled, and in those occurring just prior to death, large portions of the lung-tissue may collapse simply from inability to breathe with sufficient force. We also meet with collapse of the lung in hooping-cough.

When we inquire whether the diagnostic signs of collapse of the lungs are so clearly defined that we can always make out the state of the pulmonary tissue, we have to admit that our

knowledge of the pathological phenomena as yet exceeds our power to recognize them in the living. The physical signs are not satisfactory; the symptoms vary with the conditions producing the disease. There is dulness as in the other forms of condensation, as in pneumonia, as in pleurisy. Neither voice nor respiration is characteristic. The most usual physical sign is dulness on percussion, with an absence of all respiration, or with a blowing sound, which is faint and not so distinct as in pneumonia. The dulness is not great, may be changed during respiratory percussion, and in cases dependent upon inspissated mucus may disappear suddenly when the obstructing cause is removed. Yet collapse of the lung is at times a state of long duration. Great stress is laid by some on the signs of emphysema which surround the dulness of the condensed tissue. Should a pneumonic process affect the collapsed portion, the dulness is stationary, and we are apt to find the high but variable temperature of broncho-pneumonia.

After collapse the breathing becomes very difficult. The patient makes intense efforts at inspiration; owing to the non-expansion of the lung during these efforts, the ribs move inward and recede, instead of moving outward as in ordinary breathing. This sign, the suddenly-increased dyspnoea, and the appearance of dulness unaccompanied by marked bronchial breathing, are, in a case of bronchitis, the most trustworthy indications that collapse of the lung-tissue has taken place. Yet where the collapsed lobules are small and scattered through the lung, these signs are not all present, and the diagnosis is uncertain. The dulness is wanting; and the peculiarity in inspiration may not be observed.

When collapse affects a large portion of lung, it much resembles *lobar pneumonia* and *pleurisy*, from both of which, however, it may often be distinguished by the phenomena indicated, and, still more positively, by the history and the absence of that group of symptoms and physical signs which characterizes inflammation of the lung or the pleura. How nearly it resembles *broncho-pneumonia* has already been stated. The diminution in volume of portions of the chest, the shifting character of the physical signs, and the speedy re-entrance of air into parts that had shown signs of condensation, are the most trustworthy points in diagnosis.

### Diseases in which Dulness on Percussion occurs.

The diseases of the lungs in which dulness on percussion is met with are all those in which compression or consolidation of the pulmonary tissue takes place. Especially do we find dulness, and the physical signs which accompany it, in the phthises, in pneumonia, and in pleurisy.

**Phthisis.**—Phthisis presents itself in a chronic and in an acute form. The chronic variety is by far the most frequent. It is essentially “the consumption,” which is such a scourge to the human race. In by far the greatest number of instances this consumption is linked to tubercular disease. And although we can recognize a non-tubercular form, I shall, unless it be otherwise specified, use the term phthisis as meaning tubercular disease.

Beginning usually with a short and insidious cough, with a feeling of lassitude, and a decline in general health; attended at times from its onset with a pain in the affected lung and a somewhat quickened circulation; or giving the first indications of its existence by the occurrence of a hemorrhage; or developing itself after severe bodily or mental fatigue; or traceable to some neglected cold,—the disease becomes fully established, with symptoms which hardly need a detailed description. The harassing cough by day and by night; the impaired appetite and disturbed digestion; the loss of blood from the lungs; the steadily-augmenting debility; the short breathing; the exhausting night-sweats; the hectic fever; the deceptive blush which this imparts to the cheek; the increased lustre of the eye; the singular hopefulness; the temporary improvements; the relapses; and the greater vividness of the imagination, so strongly contrasting with the waning frame,—are phenomena with which sad experience has made not only every physician, but many a fireside, familiar.

The most constant of all these symptoms are the hemorrhage, the cough, and the emaciation. The *cough* is at first dry, and followed by a frothy expectoration. As the disease advances, the sputa thicken. They become greenish in color, streaked with yellow, and “nummular,” consisting of large greenish masses of a rounded form, or sometimes rounded yet with jagged edges, which masses do not sink in the cup containing them, but float imperfectly in a thin serum. This expectoration is, however, by

no means pathognomonic of the malady ; it is occasionally encountered in chronic bronchitis. In the last stages of consumption the sputa are often homogeneous, and have a dirty-grayish, decidedly purulent aspect. Examined microscopically, they show alveolar epithelium, pus-cells, exudation corpuscles, and elastic tissue, the most distinctive of which is the elastic tissue of the alveolar walls. Yet the only absolute sign in the sputum is the bacillus tuberculosis. Its presence bespeaks tubercular disease, its absence is an almost conclusive argument against the existence of this affection. The numbers found in the sputum bear a direct relation to the extent and gravity of the complaint ; in arrested tubercle they become very few or disappear. In lung destruction from syphilis or from chronic pneumonia, in the non-bacillary form of fibroid phthisis, in cavities from bronchial dilatation, in gangrene of the lung, the bacillus is not observed. But failure to find the bacilli in the expectoration is not as valuable evidence as finding them ; for Koch himself could not detect them in a certain number of cases of consumption. It is also a question whether a few of the bacilli may not be found in the sputum from accidental lodgement in the air-passages.

In rare instances, the cough remains slight throughout the malady ; but generally it is a very distressing feature of the complaint, and is particularly worrying at night. Sometimes its violent paroxysms bring on vomiting. But vomiting and other gastric symptoms occur irrespective of paroxysms of coughing. In truth, anorexia, nausea and vomiting are often very prominent and early symptoms, and may exist where no obvious lesion of the gastric mucous membrane is found ; dilatation of the stomach attending the dyspeptic symptoms is not uncommon. Early anaemia, with increase of the blood-plates, is another frequent symptom.

Among the less constant symptoms of pulmonary consumption are a troublesome and rebellious diarrhoea connected with catarrhal inflammation or with fistula in ano, chronic laryngitis and chronic pharyngitis, and the red line around the border of the gum. In some persons this gingival line is a mere streak ; in others it is more than a line in breadth ; in none is it a certain indication. A sign which has a much more definite connection with tubercular disease of the lungs is the appearance of the



nails. The end of the finger is somewhat clubbed; the nail is curved, prominent in the centre, depressed at the sides, its surface slightly cracked, its appearance bluish. This peculiar condition of the nails is sometimes met with even in the earlier stages of the disease. A similar nail is, however, seen in chronic pleurisy and in diseases of the heart. The laryngeal symptoms are apt to be a very distressing complication, and mostly end, no matter how they begin, in tubercular laryngitis. This, and the laryngoscopic appearance of the ulcers which attend it, have been described when treating of laryngeal diseases.

A significant symptom of phthisis is the heightened temperature as ascertained by the thermometer. Indeed, the temperature may be greatly elevated for several weeks before we find physical signs indicative of the deposition of tubercle, or of an undoubted increase in the already existing deposition. Furthermore, the rise in the body heat closely corresponds to the activity of the deposition of tubercle. If the temperature be decidedly and permanently elevated throughout the day, there is active deposition. When the animal heat is normal, the deposition in the lungs has ceased, and the tubercular process is arrested or retrograding. It may also be normal or even subnormal in very chronic cases.

But these statements, as I know from repeatedly examining into the matter, do not aid us much in discriminating lingering lung complications in febrile states, or affections intercurrent in tubercular phthisis, from a spread of the disease, or certain forms of persistent non-tubercular consolidations. The morning temperature in tubercular phthisis is often higher than the evening temperature, though we frequently see the reverse. Very generally the maximum temperature is reached in the afternoon; sweats occur in the evening, and there is a drop of two or three degrees towards morning. The temperature chart may simulate that of a remittent or an intermittent fever, and the frequent occurrence of chills and the sweats makes the resemblance still closer. In the last days of the disease the temperature may fall greatly.

The thermometer has been made use of in another manner in the diagnosis of tubercular consumption. Peter \* calls attention

---

\* Clinique Médicale, tome ii., 1879.

to the advantage of local thermometry. A surface thermometer is applied firmly in front of the chest in the second intercostal space, and if the temperature is higher there than on the other side, or than normal, it is because there are tubercles underneath. In beginning tuberculosis the increased local heat is in proportion to the extent of the lesions. In health the temperature of the chest-walls is about  $36^{\circ}$  Cent. ( $96.8^{\circ}$  Fahr.); it may rise in tubercle to  $37^{\circ}$  Cent., or more, and in consumption with cheesy degeneration still higher, surpassing the general fever heat of the body.

The symptoms which precede a fatal termination are various. Patients may go on failing for years; or an intercurrent attack of acute tuberculosis, of pneumonia, of tubercular meningitis, or of an affection of the intestines, may at any time result in death.

But at no stage of the disease do we derive as exact knowledge from a study of its symptoms as we do from a study of its physical signs. Before explaining these, it is necessary to recall briefly some facts connected with the general laws governing tubercle, and the tendency of tubercular matter to soften and destroy the textures among which it is infiltrated. It may undergo, at any period in its course, a retrogressive development, by shrivelling up, or by passing into a calcareous state. When situated in the lungs, it seeks the apices by preference; it is rarely limited to one lung, although one lung is usually the most diseased, and often at the beginning of the malady is alone affected. It is not merely a local complaint, but stands in connection with a peculiar, tainted state of the constitution, whether this be produced by infection from the products of the bacilli or not; hence the symptoms of phthisis are not solely the expressions of the condition of the lungs.

These pathological facts are all of the greatest importance. They tell us where to seek for the earliest indications of a deposit. They explain to us its signs. They teach us to look further than the lungs, and prepare us for finding lesions in other organs.

In accordance with the laws affecting tubercular depositions, we have three stages of phthisis, which run, however, by almost imperceptible degrees into one another. They are:

1. Incipient stage, or beginning deposition;
2. More complete deposition, occasioning consolidation;

## 3. Stage of softening and of the formation of cavities.

1. A few scattered tubercles do not change the normal percussion resonance ; nor do they appreciably alter the natural breath


FIG. 33.

Slight percussion dulness.

Feeble or harsh respiration.....

Prolonged expiration.....

Exaggerated respiration...



Beginning infiltration ; masses of tubercle have accumulated, but the intervening lung-tissue is still healthy.

sounds. But as soon as the deposit is sufficient to impair the elasticity of the lung-tissue or to increase its density, a relative loss of clearness on percussion on one side, and modifications of the vesicular murmur, such as feeble or jerking inspiration, or a prolonged expiration, may be ascertained. The dulness is readily detected by percussing the patient with his mouth open and during a fixed expiration, or the difference between the two sides becomes very manifest during held inspiration ; in other words, respiratory percussion will aid us. To find the dulness at the upper part of the chest posteriorly, the position recommended by Corson,\* of crossing the arms and clasping the shoulders, is very advantageous. In a certain number of cases, with the slight dulness on percussion and the changed breathing is associated a

---

\* New York Journal of Medicine, March, 1859.

blowing sound in the subclavian or in the pulmonary artery. A murmur is, indeed, at times present in the pulmonary artery long before any other physical indication of tubercle is discernible. All these physical signs may be accompanied by rales of various kinds. What makes them significant is, that they occur at the upper portion of the lung, whether anteriorly or posteriorly. If, therefore, any modification of the vesicular murmur, or any adventitious sound limited to the apex, exist; if there be a slight dulness on percussion above or under the clavicle, or in the supraspinous fossa; if this coincide with flattening of the anterior surface of the chest, especially on one side, with defective expansion of the thorax and shortness of breath, with a cough, and falling off in general health,—the diagnosis of beginning tubercular disease is almost positive. But these signs possess now less value to us than formerly, for the detection of bacilli would be of greater import than any or all of them.


2. As the infiltration advances, the signs become more decidedly those of consolidation. Greater dulness on percussion at the upper portion of one or of both lungs, scarcely influenced by respiratory percussion; more resistance to the percussing finger; stronger vocal resonance; a sinking in of the side most affected, and often soreness to the touch over the diseased part; a very harsh murmur; or, when the infiltration surrounds the bronchial tubes, a distinct blowing respiration,—are all present in varying degree, and all denote consolidation. And chronic consolidation at the apex has, in the large majority of instances, but one interpretation; phthisis. In the second stage, as well as in the first, we often meet with superadded signs of bronchitis which occasionally mask the respiratory sounds, and with friction-sounds from local pleurisies, or with fine crackling.

3. The diseased organ now passes into a state of softening, or rather some portions of the lung begin to soften, while others remain indurated, and in yet others fresh infiltration takes place. Moist crackling or persistent moist rales indicate that softening has begun. The broken-down material may be expectorated, and the malady for a time be stayed; but such is not often the case. The area of the softened mass widens; cavities form; and in addition to the moist rales, to the physical phenomena of the second stage, and to the increasing debility, night-sweats, and



hectic, the signs indicative of a cavity are noticed. What these are, may be learned from the following engraving. Prominent among them are the cavernous voice, especially in whispering,

FIG. 34.



and the hollow breathing. But the hollow, cavernous respiration may be caught only in expiration, or it may be temporarily superseded by very large bubbling sounds,—gurgling. Again, over small or over deep-seated cavities none of these sounds may be perceived; and, in truth, even when they exist, their limitation to a particular locality is an element in the diagnosis of a cavity almost as important as their presence.

The results of percussion over an excavation are not always the same. They depend much on the thickness and the state of the walls of the cavity. If dense, percussion yields a dull sound; if thin, a tympanitic, or its varieties, a cracked-pot or a metallic sound. If only a certain amount of indurated tissue intervene between the cavity and the surface of the chest, a singular sound, a mixture of dull and tympanitic, is produced. If healthy lung-tissue form the walls of the excavation, the sound is clear, or nearly so. Moreover, in all cases the pitch and, to some extent, the character of the

sound are changed by percussing over the cavity while the mouth is kept open. When it is shut, the sound elicited is of lower pitch. On inspiratory percussion, the previously tympanitic or mixed sound becomes dull. Another sign by which we may judge of the existence of a cavity at the upper part of the lung is the extraordinary clearness with which the heart-sounds are heard at that point, or a waving impulse in the second intercostal space.

Such, then, are the physical signs which indicate the varied structural conditions of the lung in the three stages of phthisis. With these signs are associated, as symptoms, cough, purulent sputum full of elastic tissue, increasing quickness of breathing, chest pains, progressive debility, hectic fever, digestive disorders, and emaciation,—symptoms the occurrence and severity of which mark also, though not very accurately, the periods of the malady. Since our knowledge of the value of the bacillus of tubercle, all points of diagnosis with reference to a pretubercular stage are of little importance. For, unless we found the bacillus in instances supposed to be pretubercular on account of defects of temperature, lessened muscular power, vomiting, imperfect assimilation, emaciation, sore throat, slight, dry cough, and of limited physical signs, we should have no proof that the disease had or was likely to have anything to do with consumption.

An interesting contribution to our knowledge has been made by Fowler,\* which will help us in the clearer appreciation of the malady. It consists in watching the extension of the lesions in their “line of march,” which is found to take place in a regular manner. The primary lesion is not often in the extreme apex of the lung, but has its site from an inch to an inch and a half below the summit of the lung, and rather nearer to the posterior and external borders. Lesions in this position tend to spread backwards, and thus is explained why we may have the physical signs of deposit marked in the supra-spinous fossa while they are still uncertain in front. Another site of primary affection is at a spot corresponding on the chest-wall with the first and second interspaces below the outer third of the clavicle. The lower portion of the lung is usually involved before the apex of the opposite lung.

---

\* The Localization of the Lesions of Phthisis, London, 1888.

Let us now look at the disorders with which phthisis, in its various stages, is likely to be confounded, premising that in doubtful cases the diagnosis is always to be established by the presence of the tubercle bacillus. They are, to speak of thoracic affections only :

CHRONIC BRONCHITIS ;

CHRONIC PNEUMONIC CONSOLIDATION ;

CHRONIC PLEURISY ;

PULMONARY CANCER ;

SYPHILITIC DISEASE OF THE LUNGS ;

BRONCHIAL DILATATION ;

PULMONARY ABSCESS ;

PULMONARY GANGRENE.

ACTINOMYCOSIS.

*Chronic Bronchitis*.—The first stage of consumption is particularly prone to be mistaken for chronic bronchitis. Distinct dulness on percussion at the apex is of much aid in discrimination, especially if it be on the left side, and if alterations of the vesicular murmur correspond to it. When the dulness is not discernible, we have to depend, in our efforts at a separation of the two diseases, on the history of the case, especially as to family, likelihood of exposure to infection, the occurrence of blood-spitting, the limitation of the physical signs to the apex, and the proofs of increased activity of the surrounding lung. Cough and expectoration are common to both affections. But they are associated, in chronic bronchitis, with physical signs more or less diffused through both lungs, and unaccompanied by much constitutional disturbance ; while from the onset of phthisis the falling off in general health is out of proportion to the local lesions.

Where the deposition is at all extensive, an erroneous diagnosis of bronchitis is with ordinary care impossible, unless, as is always highly improbable, phthisis should be complicated with *emphysema*, or the tubercles be quiescent and so diffused as not to impair the resonance on percussion. Under the latter circumstances especially, the occasional tympanitic character of the sound over the seat of the tubercular deposition is liable to be misconstrued into increased clearness on percussion, and into a disproof of the existence of phthisis. When tubercle and emphysema coexist, the percussion note may really be pulmonary and like that of

healthy lung. We should then have to judge of the one disease following the other mainly by the respiratory sound, which becomes much feebler; generally, too, the dyspnoea is increased. The thermometer, as Ringer suggests, by showing a higher temperature than in pure emphysema, may assist us. But the most certain sign would be the bacilli in the sputum.

A difficult diagnosis may be at times the distinction between chronic bronchitis and the *phthisis of old people*. This, indeed, often happens in a latent form, and is very slow in its development; the temperature may be normal or subnormal. Besides the microscopic examination of the sputum, auscultation alone is of much value, since the chest remains resonant on percussion, owing to the dwindling of the muscles of the thorax, the ossification of the ribs, and the rarefaction of the lungs.

In the stage in which the signs of consolidation become well defined, phthisis may be mistaken for any of those conditions which occasion the physical signs indicative of greater density of the lung-tissue, and which are accompanied by cough and by loss of flesh. Such are particularly pneumonic consolidation, pleuritic effusion, and cancerous deposits.

*Chronic Pneumonic Consolidation.*—Chronic pneumonic consolidation, or chronic pneumonia, gives rise to many manifestations which simulate consumption. These are cough, emaciation, and the local signs of chronic condensation,—increased voice and fremitus, sinking in of the chest-wall, feeble inspiration and prolonged expiration, or a fully-developed bronchial respiration. But in pneumonic consolidation the history usually points to an antecedent acute affection; the health is not so much impaired; there has been no hemorrhage, although, owing to intervening acute bronchitis, the sputa at times may have been streaked with blood; and the dulness on percussion and the other physical signs of consolidation are, for the most part, perceived over the lower lobe of one lung. In many of these cases interstitial fibroid changes ultimately take place in the lung, and we thus have a chronic interstitial pneumonia, and this allies the cases closely to fibroid phthisis. Yet it is clinically convenient to keep them apart, as the consolidation may slowly disappear, and the retraction of the chest and other features of fibroid phthisis as ordinarily seen are not present.



This position of the physical signs is of great importance. Yet there are two sources of fallacy which may arise. On the one hand, tubercles may, by way of exception, be seated in the lower lobe; on the other, chronic pneumonic induration may affect the apex. When an infiltration of tubercle takes place in the lower lobe, its distinction from chronic pneumonic condensation is very difficult. Our surest guides are attention to the pathological law which teaches that consumption is not met with in an advanced state in one lung alone, and the examination of the sputum for bacilli; and by noting the want of those serious symptoms which go hand in hand with the physical signs of tubercular phthisis we may determine the real nature of the case when an inflammation of the upper lobe has resulted in its persistent induration. I adduce a few instances, by way of illustration :

A gentleman was under my care for years, in whom, after pulmonary inflammation, signs of condensation remained in the upper part of the right lung. He did not suffer at all, except from attacks of acute bronchitis, to which he was very liable. During these he lost flesh; but when they passed off he rapidly regained it. He had a chronic cough, but it was very slight. After the lapse of a number of years I lost sight of him.

In another case, with a similar history, I found dulness on percussion, prolonged expiration, and a friction-sound limited to the apex of the right lung. There had been a continuous cough, but very little constitutional disturbance, and no hemorrhage. The abnormal signs lasted for a year, and then almost disappeared under a succession of blisters, and the cough ceased.

In both cases the signs were confined to the summit of one lung. I had some time since under observation a patient affected much in the same manner, a man seventy-five years of age, in whom the dulness at the right apex had for years remained stationary. I might cite many more examples; but these are sufficient to justify the conclusions that can be drawn from the facts mentioned.

But to return to the points of difference between chronic induration of the lung and tubercular phthisis. They may be thus summed up: when the signs of consolidation, whether existing at the upper part of the lung or not, are out of proportion to the general symptoms, there is reason to believe that they are not the

result of tubercular infiltration. The non-occurrence of hemorrhage would tend to strengthen such an inference. Very important information is drawn from watching whether the physical signs undergo changes indicative of a deposit in the hitherto healthy portions of the pulmonary texture. To the presence or absence of the bacillus tuberculosis in the sputum great weight must be attached. But the presence is of far more value in diagnosis than the absence.

A great and complicating difficulty in the differential diagnosis remains to be mentioned. It grows out of the circumstance that tubercular disease may be developed in a lung which is in a state of chronic induration. Whatever the explanation, the fact cannot be disputed that we find persons who are without a trace of pulmonary disorder, seized with an inflammation of the lung, which is followed by persistent consolidation, and in the course of time by undoubted tubercular phthisis. Indeed, many of the reported cases of tubercle affecting primarily the lower lobe of the lung are, in reality, cases of tubercle following chronic pneumonic consolidation. The history is usually as follows. A person in all respects healthy is attacked with an acute pulmonary affection. He recovers from it, but with a trifling cough, with a persistent dulness on percussion, and with a feeble respiration, heard over one of his lungs. He continues ailing, yet is not positively ill, when, without any apparent cause, after a time varying from a few months to years, the pulse becomes frequent, his cough increases, the expectoration augments greatly in quantity and becomes decidedly purulent, the temperature rises, and he emaciates rapidly. Profuse night-sweats occur; and the physical signs, which have been stationary for a long time, now begin to change. The dulness extends; and, instead of the enfeebled respiration, a harsher, blowing respiration is perceived over the affected part, and moist crackling and the signs of a cavity follow. If doubt still exists as to the nature of the malady, the advance of the disease will clear it up. True to the laws of tubercle, a deposit takes place in the lung previously sound, and not at the lower portion, but at its apex.

Hemorrhage may or may not occur. In the patient from whose case the above description is drawn, it did not happen; and in others, too, it was wanting. Its presence is, therefore, strongly

in favor of the fact that tuberculosis has occurred ; its absence does not positively prove the contrary. At all stages a minute examination of the sputum will tell us when the bacillar infection takes place.

It is supposed by many that the tubercle bacilli have existed in the lung prior to the inflammatory disease, or may, indeed, have caused it. But this is not often borne out by the clinical history. It is more likely that the bacilli have formed, as a rule, in the damaged organ. Cases of the kind with the cheesy changes in the lung and the disintegrating products of the inflammation form the variety of phthisis that was not long since asserted to be a special disease, *pneumonic phthisis*, but which we no longer doubt to be only a clinically somewhat different variety of tubercular affection.

These remarks will apply almost equally whether the original seizure was an ordinary croupous pneumonia or a catarrhal pneumonia. In both we have the signs of consolidation remaining ; in both the same questions of diagnosis may arise, as to whether the lung is undergoing cheesy degeneration, and as to the occurrence of tubercle. Yet there are some points which the chronic consolidation attending a *chronic catarrhal pneumonia* exhibits, that I shall here refer to. In the first place, the history of a preceding acute catarrhal attack is clear, or there have been a series of attacks, after one of which the lung was left solid, and since which the patient has remained delicate, prone to take cold, and is easily put out of breath. Now, he may come under our observation in the midst of one of these broncho-pneumonic seizures, and we may watch him for five or six months with the signs of consolidation over portion of one lung, whether at base or apex, or with affected points, often symmetrical, in both ; further, there are night-sweats, fever with decided evening exacerbation, diarrhoea. Gradually these urgent symptoms yield ; he gets about, but a spot or spots of consolidation in one or both lungs do not pass away for many months ; or the chronic catarrhal pneumonia may remain as such, or pass into pneumonic phthisis, which means really tubercle.\* When this happens, great variation between morning and evening temperature, simulating a

---

\* See a paper of mine, *Phila. Med. Times*, June 19, 1880.

malarial fever, increasing cough and dyspnœa, marked sweats, decided emaciation, announce the event; while the physical signs show extending dulness, crackling and fine moist rales, over the affected spots or in parts not previously diseased, and ultimately cavities. At all stages repeated examinations of the sputum for tubercle bacilli are of decisive value.

*Chronic Pleurisy.*—A persistent cough attended with emaciation and with dulness on percussion is common to chronic pleurisy and to phthisis, and is a cause of many errors. But the seat of the dulness at the lower part of the thorax; its much more absolute character; the almost entire cessation of all breath-sound; the diminished or absent vibration of the chest-walls when the patient speaks; the dilatation of the affected side,—are in striking contrast with signs most manifest at the apex, with the distinctly-prolonged expiration, with the rales and the evidences of beginning softening. Nor are the symptoms of a pleuritic effusion as grave as those produced by phthisis. Even where the fluid filling the chest is pus, we do not find hectic fever so intense, emaciation so great, or night-sweats so constant and exhausting; and the patient coughs less, and never spits up blood. In those cases of chronic pleurisy in which the side, instead of being dilated, is retracted, the diagnosis is more difficult. Attention to the seat of dulness being at the lower part of the chest, to the diminished respiration, voice, and fremitus, and to the shrinking affecting only one side of the thorax, will, however, serve as the foundation for a correct conclusion.

Tubercle may complicate pleuritic effusions. We suspect this by the occurrence of hemorrhage, and by the marked emaciation and hectic. We can only be sure of it by finding signs of deposit on the non-affected side,—which deposit, in accordance with the custom of tubercular disease, will take place first at the apex,—and by bacilli in the sputum. Tubercular pleurisy may be a one-sided as well as a primary disease. It is not always accompanied by effusion. There may be only great and irregular thickening of the pleural membrane attended with variable fever, with coarse friction, with much pain, and with or without bacilli in the scanty expectoration. Chronic *double* pleurisy is apt to be associated with a tubercular affection of the lungs, but it may be rheumatic, or may occur without obvious cause.



*Pulmonary Cancer.*—Cancer of the lung has many symptoms which it shares with tubercle. Cough, night-sweats, hemorrhage, gradual wasting, belong to both diseases, as do the signs of pulmonary consolidation. But cancerous formations are usually limited to one lung. Only one side of the chest is, therefore, flattened or distended. Over the cancerous lung the percussion dulness is great. There is either loud, blowing respiration, or, if the mass have compressed or obliterated a bronchus, enfeebled or absent breathing and absent tactile fremitus. We find no rales; but all the signs of consolidation are more perfect than in tubercle. Owing to a cancerous deposit in the mediastinum, the dulness at times extends beyond the median line. Paroxysmal dyspnoea, enlargement of the clavicular lymph-glands, and prominence of the large veins on the chest and arms are common. Fever is very generally absent.

Cancer in the lung may soften; yet the signs of softening are rarely as manifest as they are in tubercle. The sputa are purulent, or like currant-jelly or prune-juice. Further, a cancerous tint of the skin may be present; and cancerous tumors in other parts of the body become almost absolute evidence in favor of a deposit in the lung being cancerous, since, with rare exceptions, cancer and tubercle do not coexist. The character of the pain must be also taken into account. In tubercle, it is transitory and shifting; in cancer, it is much more constant, and much more severe.

*Syphilitic Disease of the Lungs.*—Syphilis may occasion a specific form of bronchitis, preceding the syphilitic eruption; or produce gummata, which may soften and be eliminated, and which, according to Ricord, form in the lungs toward their periphery and base; or give rise to chronic interstitial pneumonia of the base. When syphilis manifests itself in the pulmonary structures, it produces most of the phenomena of phthisis. The chief differences are, that the nodules affect generally only one lung, most frequently the right, and principally the base or the lower part of the upper lobe; that they remain circumscribed, not spreading to the surrounding textures; and that they occasion, as a rule, neither hæmoptysis, nor fever, nor night-sweats, nor decided emaciation, nor marked cough or rales, but dyspnoea out of proportion to the local disease. The most common physical signs are dulness on

percussion, deficient fremitus, altered vesicular breath-sounds, and obvious sinking in of the supra- and infra-clavicular regions; in some instances signs of destruction of the lung are found. Still, the syphilitic affection can be distinguished with certainty only by the history of the case, and by the thickening of the periosteum of the head of one or both clavicles. Milroy,\* in his investigations on soldiers, also lays stress on the thickening of the perichondrium of one or more of the upper cartilages, with frequently a tumefaction of the soft parts between them and the skin. To these tests may be added that recognized by Broderick,† substernal tenderness, as a means of diagnosis of acquired syphilitic taint. In all cases, we must be careful that the thickening at the upper part of the chest-walls and the altered resonance thus occasioned be not looked upon as signs of a tubercular consolidation. And as regards the tenderness, pain on pressure, as has been correctly asserted, is met with at the lower part of the sternum in a large number of phthisical cases.

Syphilis of the lung may also be associated with syphilitic lesions in other organs, especially in the larynx, and we may find considerable cough, with emaciation, diarrhoea, and albuminuria. But even then there are no night-sweats and fever attending the emaciation, the great debility, and the marked dyspnoea. The diagnosis of syphilis has been made by microscopical examination of the sputum, finding nucleated granular cells, shrivelled nuclei, spindle-cells, and remnants of a finely-striated stroma.‡ To the absence of tubercle bacilli in doubtful cases great weight must be attached. Fibrous pleurisy and pleuritic effusions are comparatively frequent; even small cavities occur in the lung.§

The preceding diseases are most likely to be confounded with the stages of consumption prior to softening and the formation of cavities. Next let us review those affections which, like phthisis, occasion the signs of excavation, and which, therefore, may be mistaken for its third stage: they are, chiefly, bronchial dilatation, abscess, and gangrene of the lung.

---

\* British Army Medical Report, quoted in *Annals of Military and Naval Surgery*, vol. i., 1863.

† *Madras Medical Journal*, July, 1865.

‡ Sokolowsky, *Deutsche Medicinische Wochenschrift*, Sept. 12, 1883; Cube, also Guntz, quoted in *Schmidt's Jahrb.*, No. 6, 1882.

§ Satterthwaite, *Boston Med. and Surg. Journ.*, June, 1891.

*Bronchial Dilatation.*—A dilatation of the bronchial tubes takes place in two forms: either the tubes are uniformly dilated and like the fingers of a glove, or else they form cavities by undergoing a saccular enlargement. The former variety furnishes the symptoms and physical signs of a case of chronic bronchitis attended with copious expectoration. The percussion clearness may be slightly lessened, owing to the condensation of the surrounding pulmonary tissue; the respiration may be more strictly bronchial; but otherwise both symptoms and signs are those of chronic bronchial inflammation. In the globular form of dilatation we meet with all the sounds of tubercular excavations: the hollow, blowing respiration; the hollow, well-transmitted voice; gurgling; even metallic tinkling. In the acuter cases, Wilson Fox\* has observed the metallic quality of the rales to be very distinctive. Yet all these phenomena are in strange contrast with the almost unimpaired health, and with the non-occurrence of hemorrhage, of night-sweats, and of emaciation. Pain, Lebert has shown, is among the early manifestations of the disease.

Thus, when we find the signs of a cavity, and when the general symptoms do not indicate profound constitutional disturbance, we may suspect a bronchial dilatation. This suspicion becomes a certainty, if the cavity be at the middle or the lower portion of the lung, and if the resonance on percussion be but little impaired. For in bronchial dilatation the dulness over the seat of the disease is very slight; certainly not nearly so great as that yielded by the dense walls of a tubercular excavation. It is also true that the dulness on percussion is not increased by respiratory percussion, and, for the most part, follows, and does not precede, the auscultatory signs of a cavity. We find further evidence in the stationary character of the physical signs: for months they do not change; whereas in phthisis they continually alter with the advancing malady. The expectoration of bronchial dilatation, too, is more abundant than that of consumption, purulent, acid, and in very chronic cases fetid, suggesting, indeed, at times, the existence of gangrene. Nor does it look like the sputum of phthisis, for the bulk of it is much more fluid, and in the watery secretion float small masses of pus and detritus far

---

\* Treatise on Diseases of the Lungs and Pleura, London, 1891.

less compact than the nummular sputum of phthisis. It may contain elastic fibres, but not tubercle bacilli. As regards the cough of dilated bronchi, it is much more persistent, being constant by day and by night, and only at times relieved by expectoration, which then varies in copiousness according to the size of the sac.\*

Skoda† describes, as a peculiar physical sign present in sacculated bronchial dilatation, a large and coarse crackling, called by him the large bubbling, dry crepitant rale. In a case which came under my observation, the diagnosis was made by this auscultatory sign. The patient, a boy aged twelve years, had swallowed a bone, which lodged in a bronchial tube and gave rise to bronchitis and bronchial widening. He died subsequently of acute meningitis, and the bone was found firmly embedded on one side of the globularly-dilated bronchial tube.

*Pulmonary Abscesses.*—Abscesses of the lung may form in the course of acute pneumonia, but are not then likely to be mistaken for chronic phthisis. Different is it with abscesses which are developed three or four months after an attack of pneumonia, and where the lung-texture has remained partially consolidated. I have seen not a few examples of chronic induration of the lung terminating in this way. A man who was shot through the lung was seized, soon after the injury, with inflammation of that organ. Percussion dulness and blowing respiration continued at the lower part of the left lung. One day, after exertion, he suddenly expectorated a considerable amount of pus. The signs of a cavity were detected at once; but they subsequently disappeared, and perfect recovery took place. In another case of pneumonia, the disease in like manner lapsed into a chronic state. Five months after the acute attack, the evidences of an excavation became manifest at the edge of the right scapula, and existed there for two months; then, so far as physical signs could prove, the cavity closed. Instead of the hollow, blowing respiration and gurgling, only a somewhat roughened vesicular murmur was perceived.

Such is, however, not always the termination. The abscess may grow larger and larger, until the entire lung is destroyed;

---

\* Skoda, Allgem. Wien. Med. Zeitung, 1864, No. 26.

† Percussion and Auscultation.



amphoric percussion note, amphoric respiration, amphoric voice, and, at times, metallic rales, being the physical signs observed.

Lung abscesses differ from *bronchial dilatation* in not being permanent and fixed. They have this in common with *tubercular excavations*—they change. They increase like these ; but, further, they do what tubercular cavities do not—they decrease. Their physical signs are in every respect like those of all cavities, and vary with the size of the excavation. Sometimes metallic respiration and voice may be heard over it ; or perforation of the pleura produces the signs of pneumothorax with effusion. In fortunate instances the pus is expectorated, or the abscess opens externally, and a cure is thus established. But very large abscesses are apt to wear out the patient. Hectic fever, and occasional hemorrhage, attend them ; yet neither is as constant a symptom as it is in consumption. The sputa are usually copious, purulent, full of elastic tissue, and fetid, differing in this respect from the expectoration of phthisis, which is only temporarily fetid, if the secretions decompose in the cavities. Again, abscess of the lung may be distinguished from tubercular disease by being ordinarily situated at the base of the organ ; by its following—although there are exceptions to this rule, chiefly in septic conditions—pneumonic consolidation ; by the occurrence of copious expectoration being often, not constantly, sudden ; but especially by its limitation to one lung. The other lung remains perfectly healthy. It may enlarge, and its murmur be more distinct ; but the sounds denote its texture to be normal.

Abscess of the lung is not infrequent in suppurative diseases of the nose, or larynx, or œsophagus. It is still more common from embolic infection.

The small amount of constitutional disturbance which pulmonary abscesses sometimes entail is remarkable. In several patients, in whom I have noticed abscess of the lung consequent upon chronic pulmonary consolidation, the physical signs of a large cavity were in strange contrast with the regular pulse, the almost undisturbed breathing, the slight cough, and the healthy complexion.

Let us tabulate the differences between a tubercular excavation and a pulmonary abscess :

## PULMONARY ABSCESS.

Signs of cavity usually at the lower lobe.

Copious and purulent sputa, containing elastic fibres, but free from tubercle bacilli.

Comparatively small amount of constitutional disturbance.

One lung affected.

## CAVITY FROM PHTHISIS.

Signs in the upper lobe.

Sputa less copious, and at first nummular, containing tubercle bacilli, as well as elastic fibres.

Graver symptoms, and a different history.

Usually both lungs affected.

What has been called "dissecting pneumonia," a suppurative inflammation starting mostly in the peri-lobular and peri-bronchial tissues and dissecting the lobules, and subsequently destroying the parenchyma, leaving nothing but the bronchial ramifications and vessels, has symptoms that are in the main those of abscess, of which, indeed, it forms a variety. The absence of fetid breath and of fetid sputum distinguishes it from gangrene.\*

*Pulmonary Gangrene.*—Another disease which yields the signs of an excavation, and which, like phthisis, is attended with wasting of the body, here claims attention. Gangrene of the lung occurs either as diffused or as circumscribed gangrene, after pneumonia, after wounds of the lung, from blows on the chest, from poisoned blood, or from emboli in the pulmonary tissue. The physical signs are those of a cavity, seated usually in the lower portion of the lung. The symptoms are: great and increasing prostration, dyspnœa, a very pale face, a quick pulse, hemorrhage, emaciation, and a cough, followed by profuse purulent sputa of a greenish or brown color. But nearly all these symptoms happen also in phthisis. What is characteristic of gangrene is the extreme fetor of the expectoration and of the breath. The sickening odor is not perceived during each act of breathing, but mainly after coughing, and, as it were, in jets. It is the symptom by which, especially if taken in connection with the signs of breaking up of the pulmonary tissue and the sputum, gangrene is with certainty recognized. Some authors lay stress on the fact that a cavity is found in only one lung, and at its lower part. This is unquestionably of aid in discriminating between phthisis and gangrene; but it does not distinguish between a gangrenous excavation and a simple

---

\* See an elaborate paper by Hutinel and Proust, Arch. Gén. de Méd., Nov. 1882.

abscess of the lung. The only positive proof of gangrene of the lung is, as just stated, that the signs of breaking down of the pulmonary tissue are accompanied by a disgusting and more or less persistent fetor of the expectoration and of the breath; sometimes a sickening, faintly sweetish smell, sometimes fecal, oftener that of putrescence. I say persistent, because local gangrene, on a small scale, occurring around tubercular cavities or in bronchitis, may give rise to temporary extreme fetor of the breath. But it is only temporary, and therefore not liable to lead to fallacious inferences. The expectoration may be fetid in cases of bronchial dilatation or of abscess of the lung, but is never brownish, as is not uncommon in gangrene; and neither it nor the breath has that peculiar gangrenous odor which makes the patient as unbearable to himself as to his attendants. In rare instances pleurisy with fetid effusion may occasion a fecal smell of the expectoration and breath, which is gradually lost.\* The fetid sputum of fetid bronchitis is not associated with any signs of breaking down of the lung.

Yet in considering the diagnosis regarding *bronchial dilatation* we must not overlook the fact that, as Dittrich and Traube † have shown, this bears a marked relation to gangrene. Decomposition takes place in the secretions retained in the bronchial dilatation, and ulceration of the coats may ensue, leading to a gangrenous process in the surrounding tissue. Now, as just mentioned, the sputum even in bronchial dilatation may become fetid. As, moreover, it, like gangrenous sputum, may present a dirty greenish-yellow color, and separate on standing into three distinct strata, of which the uppermost is frothy though dense, the second serous, and the third dense, containing pure pus and detritus; as, further, we meet in both affections with little solid masses of particularly offensive odor full of fat and fine needle-shaped crystals of margaric acid,—we may have to depend, for a differential diagnosis, on finding with the microscope pigment grains and masses of elastic tissue. Bacteria and vibriones bespeak a similar pulmonary origin, and they and the substance in which they are embedded yield a purple or blue reaction with iodine.‡

*Pulmonary Actinomycosis.*—This rare disease resembles tuber-

\* As in the case reported by William Moore (Dubl. Quart. Journ., May, 1865).

† Gesammelte Abhandlungen.

‡ Leyden, Klinische Vorträge, No. 26, 1871.

cular disease of the lung in presenting cough, fever, wasting, and a muco-purulent expectoration. The attending fever is of irregular type, sometimes like that of typhoid fever, more generally like hectic fever. The physical signs are mostly those of tubercular deposit. The absolutely distinctive feature is finding the ray fungus in the sputum. Besides the lungs, other parts of the body may be involved, such as the jaw, the alimentary canal, and the subcutaneous tissues.

The diseases just considered exhibit, thus, points in which they are similar, and points in which they are dissimilar, to pulmonary consumption. Other affections might be added which are sometimes mistaken for this malady, such as intermittent fever, anæmia, dyspepsia, chronic diarrhoea, chronic laryngitis, chronic pharyngitis, and thoracic pains. But each of these, although it may accompany tubercular consumption and even mask some of its symptoms, lacks, when it is present as an idiopathic affection, those local evidences of deposition and softening, lacks that profound and persistent constitutional disturbance, which form as striking a part of phthisis as the disease in the lungs. An examination for bacilli is always of the greatest value.

In the remarks on the diagnosis of pulmonary consumption, the complaint has been assumed to be progressive; in rare instances it *retrogrades*. Now, before dismissing the subject of phthisis, the signs by which such retrogression can be discovered may be mentioned. They are not very fixed. In those cases in which many tubercles undergo a cretaceous transformation, calcareous particles are coughed up; the signs of softening cease; fibroid changes take place in the affected lung; the apex flattens; and a feeble murmur, with prolonged expiration or a harsh respiration, with slight dulness on percussion, is all that remains to indicate that tubercular disease has existed. It is hardly necessary to say that the cough stops, and that flesh and strength return. These phenomena may be noted even when large cavities have existed. But, unfortunately, it is not very often that we have opportunities to make such observations.

We meet occasionally with instances in which the physical signs of an infiltration into the lung-tissue depart with tolerable rapidity. They occur in those who have a decidedly scrofulous aspect, enlargement of the glands of the neck, or a scrofulous inflammation



of the eyes. In accordance with the acknowledged identity of scrofula and tubercle, we are forced to admit that the disease in the lungs is tubercular. Yet the connection with the enlarged lymphatics; the circumstance that the diminution in size of the glands is often followed by increased pulmonary deposits; that these depositions are very beneficially influenced by treatment; that they disappear sometimes altogether, or only reappear months afterward,—all make it a question whether there be not a *scrofulous disease of the lung* independent of a tubercular, one pursuing more the course of an external scrofulous disease, one, moreover, which presents a much more favorable prognosis than ordinary consumption. Among scrofulous children cases like those mentioned are not uncommon. The disorder certainly differs from the common forms of pulmonary tuberculosis, and it is not bronchial phthisis. It does not present the paroxysmal cough, the signs of pressure on the trachea or the large bronchi, and the dull sound on percussion between the scapulæ, which are the common accompaniments of enlarged and tuberculous bronchial glands. Indeed, the bronchial glands are not of necessity involved.

Some years since, I had an opportunity of inspecting the lungs in one of these instances of supposed pulmonary scrofula. I was treating a little girl for this affection, when she received a severe injury which resulted in her death. She had, when first seen, an eruption on the scalp, sore eyes, and enlarged cervical glands. She was also much troubled by a cough; and marked dulness on percussion was discerned at the upper portion of the left lung. Here, as in fact throughout the whole of the left lung and the upper part of the right, the respiration was harsh. But for two weeks before her death the symptoms and signs had strikingly improved under cod-liver oil and iodide of iron. She was rapidly losing her cough and gaining strength. The dulness on percussion was diminishing, the respiration becoming less and less rough. At the autopsy the greater part of the left lung and a portion of the right were found to contain yellowish, cheesy deposits, which exhibited under the microscope a large quantity of granules and some shrivelled cells, without distinct nuclei. The examination was made before the days of the knowledge of the tubercle bacillus.

It would be out of place to pursue here this intricate subject. I shall only add that there are no phenomena which serve as

a foundation for an absolute diagnosis of a scrofulous, in distinction from a tuberculous, infiltration. But the rapid fluctuation in the physical signs, their occurrence in those who present a strongly scrofulous aspect, and the course of the disease, may furnish a clue by which to separate clinically, as far as they can be separated, cases of the disorder. Perhaps the absence of hæmoptysis from among the symptoms may be a matter of importance from a diagnostic point of view. Certainly hemorrhage did not happen in any of the cases of pulmonary scrofula which have come under my observation. As regards bacilli in the sputum, I do not know of any observations, but most likely the bacillus of tubercle is present.

*The Acute Affections of the Lungs accompanied by Dulness on Percussion.*

In continuing the consideration of the diseases in which dulness on percussion is a marked sign, let us glance at a group of *acute* affections, in the distinction of which dulness and the physical sounds which correspond to it hold an important part.

The acute diseases of the lungs are bronchitis, pneumonia, pleurisy, and acute phthisis. They have some signs and many symptoms in common. They all present fever; they are all associated with more or less dyspnœa and thoracic pain; they all occasion a cough. If, therefore, a physician meet with an acute disease of the chest, and find the heart healthy, he at once asks himself, Is the malady acute bronchitis? is it acute phthisis? is it acute pneumonia? is it acute pleurisy?

Now, the symptoms and signs of acute bronchitis have already been discussed. It has been pointed out that the want of intensity of the fever, and particularly the unimpaired resonance on percussion, separate bronchial inflammation from all affections which occasion consolidation or compression of the lung-tissue. We may then proceed to examine the other acute pulmonary affections.

**Acute Phthisis.**—When phthisis runs its course rapidly, it is known as acute phthisis, acute tuberculosis, or galloping consumption. This formidable complaint is met with at the close of other diseases, especially of fevers; but exposure, toil, and anxiety are also among its exciting causes.

Acute phthisis shows, more even than chronic pulmonary consumption, that the disease is not simply one of the lungs. The

lesions found by the knife of the pathological anatomist are indeed for the most part insufficient to account for the early exhaustion and the emaciation. The disorder often begins with a severe chill: fever follows; at first like any fever with anorexia, quickened pulse, and elevated temperature, but soon accompanied by exhausting night-sweats and rapid emaciation, which, in connection with the intense restlessness and prostration, the high temperature, and the supervention of delirium, may cause the febrile disturbance closely to resemble typhoid fever. The symptoms that point to the thoracic malady are the accelerated breathing, the cough, the copious expectoration, the pain in the chest, and the spitting up of florid blood.

The physical signs are not always the same. If the tubercles be scattered through the lungs, no signs are perceived but those of diffused *acute bronchitis*; indeed, the sputum is the same in composition, and tubercle-bacilli are not found,\* or are infrequent. More commonly the signs are like those of chronic pulmonary phthisis, and associated with the fever and prostration we find the percussion dulness of a deposit or the evidences of the destruction of the pulmonary tissue, furnished by coarse, moist rales, and cavernous breathing. Tubercle bacilli are then usual.

When the malady assumes the form resembling chronic pulmonary consumption, the diagnosis from bronchitis is not perplexing; but when its phenomena are similar to those of acute bronchitis, the recognition of the tubercular affection is often impossible. This remark applies particularly to the distinction of the miliary form, *acute miliary tuberculosis*, from bronchitis of the finer tubes; since the slight constitutional symptoms and the coarseness of the rales of ordinary bronchial inflammation are too unlike the phenomena of acute consumption to occasion difficulty in their discrimination. But from *bronchitis of the finer tubes* the diagnosis can be effected only by taking into account that repeated chills, rapid emaciation, and profuse sweats are wanting in the bronchial affection; that the temperature is not so high, nor so irregular; that the skin is more livid; that the rales are more abundant and more perceptible at the lower part of the chest; and that, perhaps, the breathing is not so hurried or so difficult.

---

\* Jaksch, *Klinische Diagnostik*, 1887.

Moreover, with the intense dyspnoea there are generally frequent and violent fits of coughing, and marked chest pains, in the acute tubercular malady. Yet none of these signs are convincing proofs. The presence of dulness on percussion, or the sinking in at the upper part of the chest, the occurrence of hemorrhage, the finding of the tubercle bacillus, if present, the eruption of miliary tubercles in other organs, and the longer duration of the case are alone conclusive evidence in favor of the acute tubercular disease. Hemorrhage is, however, by no means so constant in the acute as in the chronic form of the affection.

Much the same symptoms will enable us to distinguish between acute tuberculosis of the miliary form and *broncho-pneumonia*, except that we can draw no inference from the dulness on percussion, further than that its early occurrence, with the bronchial symptoms, points to the pneumonic malady, its later occurrence, after the grave symptoms, to the tubercular.

When the dulness on percussion is well defined, acute phthisis might be mistaken for ordinary pneumonia. But the signs of deposit and of softening in both lungs, and the seat of the lesions at the apices, show differences from a disease which in the large majority of instances is one-sided and at the lower part of the lung, which exhibits a characteristic sputum, and in which breaking up of the pulmonary tissue is so rare.

Yet there are cases of acute phthisis that display symptoms and signs very puzzling, and strongly simulating those of pneumonia.

A person in perfectly good health is seized, after exposure, with cough and fever. They are accompanied by dyspnoea, and soon we find signs of consolidation of the lower lobe, or of the entire lung. The dulness on percussion does not disappear under treatment; and a hollow, blowing respiration and gurgling, usually first perceptible at the angle of the scapula, gradually appear, and indicate the formation of a cavity. Emaciation, which began from the onset, progresses more rapidly, and goes hand in hand with extreme prostration and profuse perspirations. The sputa are copious and purulent, but at no time mixed with blood. The other lung is carefully examined; all its sounds are normal. The case remains in this condition for several weeks, the patient temporarily improving under stimulants, yet, on the whole, growing weaker and tormented with fever of very irregular type. A slight



roughening of the inspiratory murmur, or dry rales at the apex of the unaffected lung, attract attention, and dulness on percussion and the signs of deposition become there more and more manifest. A post-mortem examination exhibits nearly the whole of one lung converted into a uniform yellowish or grayish mass of tubercle, and containing one or several large excavations; not a vestige of healthy lung-structure is to be seen. Scattered tubercles are found in the other lung, and mainly at its apex.

The case just described is one of a group which every physician has met with. The beginning of the case as one of pneumonia or broncho-pneumonia, the persistent consolidation, the occurrence of rales and of subsequent dulness on percussion at the upper part of the previously unaffected side, the continuance of the disease, and the prostration and sweats which accompany it, permit us to foretell its nature and the probable fatal termination.

Such cases were not long since classed as *acute pneumonic phthisis*, and looked upon as inflammatory, with resulting caseous infiltration of the pulmonary tissues and disintegration of the cheesy infiltration. With our present knowledge of the bacillar origin of consumption, they are explained by supposing that the tubercle bacilli have fastened readily on the altered lung, or, more generally, that these have occasioned the attending inflammatory process.

Acute phthisis may simulate other affections besides those of the chest. It has at times the delirium and prostration, the dry tongue, and the bronchial rales of *typhoid fever*. The diarrhœa and the abdominal symptoms are, however, wanting. Yet simultaneous deposition of tubercles in the intestine may cause these; and in this case the only mark of difference from typhoid fever is the absence of an eruption; unless, even under these circumstances, we are aided by the fact pointed out by Fox, that, unlike the persistent high temperature of typhoid fever with its regular diminution when the disease declines, the thermometric record in acute phthisis shows great and sudden variations, bearing no relation to the number of respirations or to the beats of the pulse. The temperature may vary many times in the course of the disease to the extent of six or seven degrees. Acute phthisis lacks the eye phenomena, the gastric disturbance, the rigid muscles, the convulsions, of *meningitis*; else the active delirium it occasionally produces might be attributed to inflammation of the membranes of the brain.

Acute phthisis sometimes progresses with extreme rapidity. I have seen a case terminate in thirteen days. It is almost invariably fatal. Yet it has its periods of deceptive improvement: the disease may proceed speedily toward softening, and then remain for a time stationary. In some instances the termination in death is the result of complications, as of tubercular meningitis, or of erysipelas of the throat and the bronchial tubes.


**Acute Pneumonia.**—Inflammation of the lung, or “croupous pneumonia,” is the type of the acute pulmonary affections. The hot, dry skin, the flushed face, the quickened pulse, the extremely rapid breathing, the thoracic pain, the cough, and the peculiar expectoration, point out at once the acute nature of the attack and the organ which is disturbed. Beginning commonly with a chill, or with flushes of heat, the disease progresses with the symptoms indicated. A few of these require a detailed description.

The expectoration is characteristic. It consists at first of a glairy mucus; soon it becomes more viscid, and acquires the appearance dependent upon the admixture of blood with the mucus and exudation-matter, to which the term rusty-colored has been given. This rusty sputum is pathognomonic of pneumonia; yet cases run their course without it. The expectoration is sometimes like prune-juice, or it is purulent. Both augur badly: both indicate that destruction of the lung-tissue has begun.

The shortness, or increased frequency, of breathing is another marked symptom. The patient draws from forty to eighty breaths a minute; but the pulse, although rapid, does not quicken in proportion. Pneumonia, therefore, forms an exception to the rule that with greater frequency of breathing the pulse rises. This perverted *pulse respiration-ratio* may be made an important element in the diagnosis. The febrile symptoms are ordinarily severe; still, they are not associated with decided cerebral disturbance. Headache is common; delirium is rare, and, when it occurs, is indicative of great danger. In drunkards it may take the form of delirium tremens. The flush on the cheek is so decided that by this and the hurried breathing alone the disease may often be recognized. The flush on the cheek is not accidental. It is sometimes very dark, and, according to Bouillaud, is most obvious when the inflammation affects the apex of the lung. Herpes is also a common symptom.

The temperature rises abruptly, and on the first or second day attains  $103^{\circ}$  to  $105^{\circ}$  F. In children and in robust adults it is specially high. It shows little change, except an evening exacerbation and a marked morning remission of from  $1.5^{\circ}$  to  $2.5^{\circ}$  for five to nine days. Between these days, sometimes on the fifth, generally on the seventh day, it falls abruptly, and a

FIG. 35.



Temperature chart in pneumonia. The observation was begun on the first day of the disease. The crisis commenced toward the end of the fifth day, and continued through the sixth to the seventh, with a secondary rise on the sixth. The chart is typical, except that the fever temperature throughout was about a degree lower than is usual. There was a slight right-sided pleurisy, but no attending bronchitis.

true crisis occurs. The temperature may sink to the norm, or even below it, and then another, though not marked, rise take place. At times there happens on the fifth day a partial but decided drop, soon again followed by ascending temperature. This pseudo-crisis is apt to occur in cases that become prolonged. It is, too, in this class of cases with slow resolution that a gradual termination of the fever is often observed. Sometimes the course of the fever is marked by sudden elevations and striking remissions. This is more common in double than in single pneumonia, and seems to correspond with fresh invasions of lung-tissue.



The urine is high-colored, and that of fever. Nitrate of silver does not precipitate its chlorides. They commonly disappear during consolidation of the lung, and their reappearance shadows forth returning health. The vanishing of the chlorides from the urine happens also in other acute affections; but in pneumonia it is most absolute.

In studying the clinical history of pneumonia we must not forget how often it exists in combination with other maladies. We find it in association with meningitis, and we must therefore always examine any cerebral symptoms with care; we note it in connection with endocarditis, which may coexist with meningitis; while its association with pleurisy is so common that this can be hardly looked upon as a complication. Among the rarer symptoms are jaundice, parotitis, croupous colitis, and transitory aphasia, appearing on the second or third day of the attack.

The *physical signs* which denote that the lung-tissue has become the seat of acute inflammation vary with the effects of the inflammation. In the first stage, or that of engorgement and beginning exudation in the air-cells, into which, however, the air is still capable of entering, there is only a slight impairment of the normal resonance on percussion. The vesicular murmur is at first somewhat altered; it may be feebler or harsher. But soon are heard with each act of inspiration, and limited to the inspiration, numerous rapidly-evolved, very fine, crackling sounds, the "crepitant" or vesicular rales.

As the exudation becomes firmer, and the tissue of the lung solidifies by occlusion of the air-cells, the stage of red hepatization is before us. Now all the signs of complete consolidation are discerned. We find decided dulness on percussion, unchanged by full inspiration; blowing respiration in its purity, high-pitched and tubular-sounding; bronchophony; and increased vocal fremitus. Rales from the accompanying bronchitis are heard with extreme distinctness through the solidified tissue, Skoda's consonating rales; so are the sounds of the heart. A crepitant rale is still here and there perceptible, or the ear catches a pleural friction-sound.

When the exudation is reabsorbed or expectorated, the signs of consolidation become less and less perfect. A vesiculo-bronchial succeeds to the bronchial breathing. The dulness on percussion



lessens; crepitant rales—not, however, so fine as at the onset of the affection, and mixed with larger moist rales—return; the cough increases; the expectoration becomes more copious, loses its tenacity and rusty color; the dyspnœa diminishes,—all phenomena indicative of the breaking up of the exudation, and of the return of air into the vesicles. If, instead, the exudation be converted extensively into pus, and the lungs soften, the physical signs are the same as in the second stage. The rarity of excavations of sufficient size explains why gurgling and the signs of a cavity are not perceived. We suspect the mischief that is going on within the chest from the protracted dyspnœa, the increasing rapidity of pulse, the purulent or brownish sputa, the pinched features, the dry tongue, and the mental wandering. Recovery may take place even then. This third stage is indeed

FIG. 36.




Diagram illustrative of perfect pulmonary consolidation, such as happens in the second stage of pneumonia.

not so much an abrupt, suddenly-established process, as it is the extension and greater diffusion of a state that may be found in portions of the lung which to the eye have all the appearance of red hepatization. In every instance of red hepatization the microscope shows that in parts the lung-tissue is infiltrated with granules and is undergoing softening, and it is probable that this breaking

down occurs, even though on a small scale, in all cases of pneumonia which recover. It is often impossible to determine that the third stage has arrived; and death may take place long before the lung presents the condition which pathologists term gray hepatization. With reference to the diagnosis of this third stage, we may suspect, from the symptoms, that the pulmonary tissue is seriously damaged. But we can never know it, unless we find the physical signs of extensive softening; and in the large majority of cases this cannot be done.

The morbid phenomena, physical signs and symptoms of the malady correspond, then, usually in this manner :

#### PNEUMONIA.

I. Stage of engorgement and beginning exudation.	Crepitant rale; slight percussion dulness.	Cough; beginning dyspnoea and rapidly-developed fever heat.
II. Stage of solidification of lung-tissue (red hepatization).	Percussion dulness; bronchial respiration; bronchophony; often a pleural friction-sound.	Rusty-colored sputum; dyspnoea; cough; high fever, temperature generally above 103°, with marked evening exacerbations and morning remissions.
III. Stage of softening (gray hepatization).	The same physical signs as in the second stage; unless large abscesses have formed.	Chills; prostration, etc.; purulent or brownish sputum; generally high temperature, 104° to 105°, or upwards.

Here is a disease which presents such striking symptoms and signs in nearly all its phases, in which the sputa are so peculiar, the hurried breathing so evident, the physical signs so distinct, that error is, with ordinary care, difficult. It becomes still more so, if a few of the pathological peculiarities of pneumonia be borne in mind: the fact that it is rarely double; that it comparatively seldom affects the upper lobe of the lung, and that it is often accompanied by the signs of pleurisy or of bronchitis. In some instances sudden disturbance of the circulation takes place with the rapid development of cyanosis. These symptoms bespeak a heart-clot.

Let us now contrast pneumonia with the various diseases of the

lungs with which it may be confounded. In its first stage, on account of similar signs, the acute inflammatory disorder is sometimes mistaken for œdema of the lung, or for the pulmonary engorgement in some fevers, or for other kinds of congestion of the lungs; and still more frequently these morbid states are mistaken for it.

*Pulmonary Œdema.*—This consists in the transudation of serum into the air-vesicles. It may be acute, the result of sudden congestion, such as that following injuries of the brain or irritation of the par vagum; or it may arise at the termination of acute affections of the lungs. It is more usually, however, chronic, and is seen as a dropsy of the air-cells, associated with dropsies elsewhere, and in connection with organic disease of the liver, heart, or kidneys. The characteristic manifestations of œdema—be it acute or chronic—are embarrassed breathing, expectoration of frothy serum, and crepitating and fine bubbling sounds diffused over both lungs, and dependent upon the fluid in the air-cells and small bronchial tubes. It presents, thus, many points of similarity to the first stage of acute pneumonia. The dyspnoea, the crepitation in the lung, may well mislead; but we cannot err, if the frothy sputum, the general distribution of the rales, their somewhat coarser character, the bluish lip, the noisy breathing, and the absence of fever be taken into account. In acute œdema these phenomena are but the precursors of death. In chronic œdema the rales are persistent, and so is the difficulty of respiration. The patient has usually to be propped up with pillows, otherwise he cannot breathe.

*Pulmonary Engorgement in Fevers.*—In fever of low type a crepitant rale, which might be supposed to be a proof of beginning inflammation of the lung, is often heard at the back part of the chest. The sound is the consequence of pulmonary congestion, with probably slight effusion into the finest bronchial tubes and air-vesicles. It is perceived over both lungs; and this, taken in connection with the history of the case, with the absence of decided shortness of breath, and with the rale not being followed by dulness on percussion and blowing respiration, shows that it is not dependent on inflammation of the pulmonary tissue. It is necessary to be aware that these fine rales may occur in fevers without being due to a true pneumonia; as otherwise the patient is apt to be treated for a disease of the lung which has no existence.

*Pulmonary Congestion.*—Besides the lung congestion just referred to as occurring in fevers, we have other causes producing a marked congestion, or “hypostatic pneumonia.” We find it in enfeebled hearts and in mitral and tricuspid disease, in those whose blood is impoverished and who are for any length of time bedridden, in instances of acute rheumatism, and due to the pressure of tumors. In the dependent portions of the lungs the manifestations of congestion show themselves first; they are, besides the signs of impeded circulation and of deficient aeration of blood, slight expectoration, scarcely any fever, varying shortness of breath, somewhat impaired resonance on percussion at the lower part of the chest,—generally more over the right than over the left lung,—feebleness of respiratory murmur, and a few fine and coarse moist rales. The sputum contains numerous epithelial cells, and blood pigment in all stages of change.

The congestion in all the instances mentioned is passive, and either hypostatic or mechanical. An active congestion of the lungs is a rare condition, though it may come on after strenuous exertion, during mountain climbing, or subsequent to extreme heat or cold. The physical signs are the same as those of passive congestion; the sputum is apt to contain more blood. There is little, if any, fever; and the history of the case, the stationary character of the physical signs, and their double-sidedness, distinguish the congestive disorder from pneumonia.

In its second stage, owing to the cough and dyspnoea, and in part, also, to some similarity in the physical signs, acute pneumonia may be confounded with pulmonary apoplexy, acute pleurisy, acute phthisis, and acute bronchitis.

*Pulmonary Apoplexy.*—An effusion of blood into the texture of the lung is generally, although by no means invariably, accompanied by external hemorrhage and by great difficulty of breathing. Over the effused blood there is dulness on percussion, and the ear hears an enfeebled or bronchial respiration. Around the seat of the mishap it encounters moist rales. Now, here are signs bearing some resemblance to those of pneumonia. But we miss from among them the decided fever. We note, on the other hand, not blood intimately mixed with the expectoration, but pure blood, florid or sooty-looking, almost devoid of air, in not large amount, at times surrounded with muco-purulent matter, and ordinarily



voided for a number of days. On close scrutiny a grave disease of the heart is generally detected to explain why an extravasation of blood into the pulmonary structure has taken place. Then we most frequently find the branch of the pulmonary artery leading to the infarcted part plugged by an embolus, which has been formed in the right cavities of the heart or been washed in through the general venous system, and most commonly affects the right lung. Again, we have more pain than in pneumonia, and the dyspnoea is different. In pneumonia it augments up to the height of the malady. In pulmonary apoplexy it is greatest, and it is very great, when the blood is extravasated; after that it declines. Yet the two affections often coexist. The closure of the vessel produces a pneumonia from embolism, or the blood acts as a foreign body, and around it is lighted up an inflammation of the lung-structure, which is apt to have its seat in the posterior part of the lower lobe of the right lung; further, the inflammation may be the starting-point of caseous degeneration; or sloughing or gangrene may result.

Pneumonia from embolism may be also caused by a pyæmic condition, and the clots may have their origin in bed-sores, in ulcers, and in various forms of suppuration. The plugs are saturated with ichor, and metastatic abscesses supervene. The symptoms are the same, and we can make a diagnosis only by the history; there are the same circumscribed spots of consolidation, and the same kind of pain, which is also often found to be associated with a localized pleurisy, sometimes followed by effusion.

Pulmonary apoplexy is met with in connection with other than thoracic affections. Observations by Brown-Séquard and by Ollivier have proved its association with central nervous lesions, and have demonstrated its occurrence on the same side as the brain lesion; \* which is not the case with reference to the ordinary acute pulmonary diseases, for these Rosenbach † has shown to be much more frequent on the paralyzed side of the body, and therefore, generally, on the side opposite to the cerebral mischief. Pulmonary apoplexy, or "hemorrhagic infarct," is also met with in malignant fevers.

Of the other diseases mentioned which resemble pneumonia, the

---

\* Arch. Gén. de Méd., Aug. 1873.     † Centralblatt, No. 16, 1879.

distinguishing points need not be here fully described. *Acute pleurisy* will be farther on more particularly studied. With regard to *acute phthisis*, it is only necessary to repeat that cases are encountered, apparently of pneumonia, in which, after the symptoms of acute inflammation of the lung pass off, those of phthisis come into the foreground. With reference to *acute bronchitis*, I shall merely recall that the dyspnoea is not so great, and that no percussion dulness is yielded by an inflamed bronchial membrane.


Percussion is thus of signal value in the diagnosis of pneumonia. In fact, when bronchitis complicates pneumonia, and loud, dry rales take the place of the blowing respiration, it is our only trustworthy guide. A single tap on the chest which elicits an absolutely dull sound tells the difference between pure bronchitis and the inflammation of the bronchial mucous membrane which accompanies inflammation of the parenchymatous structure of the lung.

The form of pneumonia most liable to be mistaken for bronchitis is the pneumonia of childhood or of old age, *broncho-pneumonia* or catarrhal pneumonia. This affection has already been described in examining capillary bronchitis. But, as the disease may also occur in adults, and has special features, a few words more will not be out of place.

*Broncho-Pneumonia*.—It mostly supervenes upon acute bronchitis, except in instances in which it arises from inhaling irritating gases. The bronchial attack is usually severe, but it may be so slight as to be readily overlooked. The spread of the disease to the lung-texture is attended with rapid rise of temperature. When the disorder attacks adults, it is apt to seize upon those debilitated by previous disease; it much more commonly affects the upper lobes than does ordinary sthenic pneumonia, and is generally bilateral. As the broncho-pneumonia merely solidifies lobules, the signs of marked consolidation are wanting, or are perceptible over only a small space. Crepitation is not common, but small moist rales are; bronchial breathing and increased fremitus show only over limited points; and the sputum is not rusty and viscid, but catarrhal. Cough and expectoration, sometimes absent in croupous pneumonia, are always present in broncho-pneumonia.

Catarrhal pneumonia or broncho-pneumonia is often noticed as a

FIG. 37.





complication of the infectious fevers, especially measles and diphtheria. It is the form of pneumonia developed when particles of food pass into the larynx and bronchial tubes,—aspiration or deglutition pneumonia. Catarrhal pneumonia pursues a much slower course than croupous pneumonia, and generally yields only gradually. The consolidation may continue stationary for weeks, showing a fever with marked daily remissions and exacerbations, like a hectic fever, and then slowly disappear. As interstitial inflammation of the bronchi and alveolar walls is distinctive of the disease, and as the perivesicular structures are markedly involved, persistent local consolidation from interstitial pneumonia or fibroid phthisis often follows. On the other hand, caseous degeneration and breaking down of the lung-texture may follow, or extended tubercular infiltration become manifest. Whether the bacillus finds in the consolidated lung a ready lodging, or the broncho-pneumonia is originally excited by the bacillus, phthisis, in truth, is in adults a not uncommon termination; in children, too, this may happen, or rachitis may develop, or an ill-defined but persistent cachexia, with a great tendency to catch cold.

There is a form of broncho-pneumonia, described as tuberculous inspiration broncho-pneumonia, that follows hemorrhage from tubercular cavities. It is usually preceded by active physical effort, and its first manifestation is a hemorrhage.\*

Pneumonia often shows itself in an epidemic form, and is now generally looked upon as an infectious disease, a lung fever. The evidence of a micro-organism as its cause is very strong. The diplococcus pneumoniae, as it is usually called, was found independently by Pasteur and by Sternberg, and has been fully studied by Fraenkel, after whom it has been named. It is present in the buccal secretion of a certain number of healthy persons. Its association with catarrhal pneumonia is not so close as with croupous pneumonia. Indeed, it is thought that the bacillus of tubercle at times excites this, making a specific broncho-pneumonia from the start; the staphylococcus and the streptococcus pyogenes may also induce it, as Northrup's observations clearly prove.


---

\* Bäumler, Deutsch. Med. Wochenschr., No. 1, 1893.



The cocci are best stained in dilute alcoholic solutions of the aniline dyes, and are readily seen in preparations colored by Gram's method. In this respect they differ from the pneumobacillus of Friedlaender, which is also found in a certain proportion of pneumonic lungs, but does not retain the stain after

FIG. 38.



The diplococcus pneumoniae of Fraenkel; the cocci are stained dark blue, the capsules are unstained. (After Jaksch.)


going through the process. The Fraenkel coccus is elongated or round, enveloped in a capsule, and often found in pairs.

The micro-organism of pneumonia has been found in the blood; in the meningitis that at times attends pneumonia, and generally appears at the height of the malady; also in the accompanying pleurisy, and in the lung complication of ulcerative endocarditis. This disease has, indeed, a close association with pneumonia, whether coming on with it or developed in its course. Osler has brought this fact out very prominently in his Gulstonian Lectures.

There are some varieties of pneumonia that present clinical features of a peculiar kind. *Apex pneumonia* is one. It is more usual in children than in adults, and the frequency with which cerebral symptoms arise and draw away attention from the chest is a matter of common observation. The cases, as a rule, are severe, and the temperature is high. *Double pneumonia* differs in nothing from ordinary pneumonia except in the severity of the

symptoms. The cases, unless speedily fatal, are generally of longer duration, and the temperature is less characteristic, for the reason that it rarely happens that both lungs are affected at the same time. Double pneumonia is rare; what is called double

FIG. 39.



Pneumococcus (diplococcus) of Friedlaender, without the capsule, from a pure culture upon gelatin from the sputum in a case of croupous pneumonia at the Pennsylvania Hospital. Drawn by Dr. Joseph Leidy, Jr.

pneumonia is generally inflammation of one lung and heavy congestion of the other. *Latent* pneumonia is not often seen except in the aged. There is but little fever, and it is only by the physical signs that the disease can be recognized. *Migratory* pneumonia, a condition in which different parts of the lung are successively involved, is not a frequent disease. The temperature shows a tendency to sudden falls, with rapid rises whenever a fresh part of the lung is involved. Some of the older clinicians, especially Wunderlich and Trousseau, regarded the disease as having a close connection with erysipelas.

It is always very important to find out whether pneumonia is primary or intercurrent in some other malady, such as in rheumatism, Bright's disease, diabetes, the exanthemata, the typh fevers, or in septic states. At times it is distinctly noticed to

follow contusions of the chest. As has been already said, it may be epidemic. By the symptoms and physical signs we cannot distinguish the sporadic and simple cases from those of the infectious malady. Further bacteriological research may solve the matter.

There are two other forms of pneumonia which have not been elsewhere considered, and which, as they present somewhat peculiar symptoms, require to be noticed. They are typhoid pneumonia and bilious pneumonia.

*Typhoid Pneumonia.*—Inflammation of the lung may be from its onset attended with extreme prostration. This form of the disease has been made a matter of warm controversy, both as to the symptoms which characterize it and as to the relation it bears to other varieties of the malady. Now, any one who reads the dissimilar descriptions given of it will become convinced that under the term typhoid pneumonia very dissimilar disorders have been ranged together. On the one hand, it has been applied exclusively to the inflammation of the lung which may complicate typhus or typhoid fever; on the other hand, it has been made to include an idiopathic fever in which the affection of the respiratory organs is occasionally wanting. To neither of these maladies ought to belong the name typhoid pneumonia, since in both the inflammation of the lung is but an incidental accompaniment. Then under the name of pneumo-typhus a disease has been of late years described, especially by German clinicians, in which typhoid fever begins with a well-defined pneumonia, that for the time being throws the enteric symptoms into the shade.

Typhoid pneumonia is pneumonia with symptoms of a typhoid type, and marked by rapid failure of the vital powers. The inflammation of the lung arising in the course of typhus or typhoid fever will of course be apt to present this character; but the malady is also noticed as a consequence of phlebitis; as supervening in cases of erysipelas, of Bright's disease, and of delirium tremens; or as the sole apparent affection. It happens not unfrequently in epidemics, and is very often observed among negroes. Its ravages on the plantations of South Carolina and Georgia are sometimes frightful. It is, also, very fatal in jails, and among troops in the field, serving under unfavorable hygienic conditions.

The physical signs are those of the sthenic form of the disease, except, perhaps, that the crepitant rale is less frequent. Most of the same symptoms, too, show themselves: cough, short breathing, and pain in the chest. All of these may be very marked, or so trifling as hardly to direct attention to the lungs. There is, however, one symptom characteristic and constant, and but one, and that is the great tendency to sinking. As regards the expectoration, it may be rusty-colored; yet occasionally, even in the early stages of the complaint, it consists of pure blood. The pulse is always quick, but weak. Dark sordes often collect on the teeth and gums, as they do in typhoid fever. Pain is absent in some cases, and extremely acute and of a radiating character in others. Concerning delirium, we know that it is much more common than it is in the sthenic variety of pulmonary inflammation, except this affect the apex in children. Some authors mention an eruption. It is, however, questionable whether the cases which came under their notice were not typhus or typhoid fever, in the course of which pneumonia appeared. The flush on the face in the low type of the malady under consideration is usually of a dusky hue, but not invariably: a pink-colored blush, extending sometimes all over the body, seems to have specially attracted the attention of observers. The disease is always dangerous, and, as Stokes\* points out, resolution is extremely slow. Chronic hepatization, with or without a low hectic fever, or a lurking congestion, may continue for weeks.

The symptoms of typhoid pneumonia are at times strangely mixed up with those produced by other conditions. In many districts in which the complaint is prevalent, it bears the distinct impress of malaria. Again, articular symptoms seem to predominate in some regions of country, and in some epidemics. Gibbes† speaks of an acute pain in the back part of the eye, in the ears, or in the side of the neck, attended with stiffness of the muscles; and of a swelling of the tonsils, and of the submaxillary and sublingual glands, which he states to be of evil augury. Dickson,‡ drawing his description of the disease from cases observed in and around Charleston, portrays several forms, the most common of which exhibits a respiration hurried and irregular; heavy sigh-

---

\* Diseases of the Chest.

† Amer. Journ. Med. Sci., 1842.

‡ Elements of Medicine.



ing; a feeling of weight at the præcordial region, with nausea and vomiting; and a tongue clean, but red. Delirium is present from the beginning, and does not subside until recovery takes place. The duration of such attacks averages from six to ten days.

*Bilious Pneumonia.*—Jaundice and other indications of hepatic and gastric derangement are not usual in ordinary sthenic pneumonia. They may be occasionally caused by the inflammation spreading to the liver, or may be of blood origin. But in the pneumonia so general in the spring and the autumn in the miasmatic regions of some of the Southern and Western States of this country, hepatic symptoms are common, and mark a special type of the disease, known as malarial pneumonia or bilious pneumonia, or by the familiar name of “bilious pleurisy.”


This form of inflammation of the lung is simply pneumonia, sthenic or asthenic, on whose features the stamp of malaria is imprinted. The chill with which it begins is usually protracted, and is followed by pain in the side, by fever, by hurried breathing, by cough, and by tenacious, rusty-colored expectoration. The pain in the side, which depends upon accompanying pleurisy, is sharp and severe, and renders the respiration irregular. The sputum is at times rusty-colored, while at others a frothy and bloody serum or pure blood is expectorated. The fever shows the type of the disease. It is much more paroxysmal than in the other varieties of the malady. This peculiarity, and the obvious symptoms of hepatic and gastric disorder, are indeed the only absolutely distinguishing traits of bilious pneumonia. The febrile exacerbations are stated by Manson, of North Carolina, to be preceded, during the morning hours, by an insensible chill,—a coolness of the ends of the nose, fingers, and toes, which, in grave cases, extends over the entire extremities.

The physical signs are those of ordinary acute pneumonia. Bronchial breathing and bronchophony are said to be more often absent, or to appear and disappear rapidly. It is certain, if this be true, that in these instances the malady could not have been inflammation, but was more probably a collapse of the pulmonary tissue. Any one, indeed, who compares the various statements made with reference to the disease, must have been struck with

the fact that cases of congestive fever in which the lungs have become simply engorged, or perhaps collapsed, and cases of inflammation of the lung arising in the course of remittent fevers, are included in the description of bilious pneumonia.

**Acute Pleurisy.**—Acute pleurisy has been so often incidentally mentioned, that a description of its main points will here suffice. It comes on from cold or exposure, or from injuries to the chest; but a great many cases are secondary to some general or infectious malady. The first effect of the inflammation is to

FIG. 40.




Roughening of the pleura from inflammation; a small amount of fluid has begun to collect.

reddden the pleural membrane; an exudation of a soft, grayish, easily-detached lymph then takes place. This constitutes the first or dry stage of the disease; and if the two inflamed surfaces unite, the disorder does not pass beyond this stage. Often, however, along with the exudation of lymph occurs an effusion of serum, which produces a special train of phenomena, and gives rise to the second stage, or that of liquid effusion.

The physical signs of the *dry* stage are impaired movement of the chest, a feebler respiration, and a friction sound of varying extent and intensity. The first two signs are caused by the patient instinctively refraining from expanding the lung, because of the pain it occasions. The mechanism of the friction sound, its nature, its superficial character and want of uniformity, have been pointed out in a previous part of this chapter. In the stage of *effusion* the physical signs differ according to the amount

FIG. 41.



Examination of the posterior portion of the chest while a large effusion is occupying the left pleural cavity.

of fluid the pleural cavity contains. A moderate quantity of liquid only constricts the lung-texture, and leaves the bronchial tubes intact; a large accumulation compresses everything; it drives all air out of the lung, pushes it into a small space against the vertebral column, and displaces the liver or heart. Wherever the fluid accumulates there is dullness on percussion. When the patient is in the erect posture, the flat sound on striking the chest

and the sense of resistance to the finger are marked at the lower part of the thorax, since the fluid naturally settles there. The line of dulness is, however, not the same in front as it is behind. It is generally much higher behind, and alters, of course, with the changing quantity of effusion, and somewhat with the position of the patient. When he lies upon his face, the fluid gravitates, if not circumscribed by adhesions, toward the anterior chest-walls, and the percussion dulness posteriorly becomes far less perceptible. The peculiar curve of the percussion line often found has been specially described by Calvin Ellis, and is named by Garland the letter S curve.\*

Where the effusion is extensive, the intercostal spaces are widened and their depressions effaced. The side appears to the eye distended, fluctuation may be perceived, and, owing to the absolute compression of the lung, no sound is heard over the chest when the patient breathes, or speaks, or coughs. In more moderate collections of fluid, the cessation of sound is not so absolute. There is an ill-defined, deep-seated respiration, and the voice reaches the ear with tolerable distinctness, and occasionally with a peculiar bleating resonance attending it. But, as large collections of fluid are more common than small ones, the former set of phenomena are, at the height of the disease, more frequent than the latter.

Above the liquid there is mostly increased resonance on percussion, or a tympanitic sound, Skoda's sound. Various explanations have been given of this. It has been attributed to the complete compression of the lung; it has been thought to be due to its slight condensation. Whatever be the true explanation, the fact of its occurrence is undeniable. This tympanitic sound is more manifest at the upper part of the chest in front; it may, indeed, be found in front when it does not exist behind. In some cases the sound has an amphoric, in others a cracked-metal character. When the ear is applied above the line of percussion dulness, it recognizes occasionally a friction sound; and near the spinal column posteriorly, where the compressed lung lies, it perceives almost invariably distinct bronchial respiration and bronchophony.

---

\* *Pneumono-Dynamics*, 1878, and *New York Medical Journal*, Nov. 1879.



When the fluid begins to be absorbed, the voice becomes more audible over the seat of the effusion, the vocal vibrations may be felt by the fingers, and the respiration is again heard. But for a long time it continues enfeebled, and its character is indeterminate; it is neither vesicular nor purely bronchial. As more and more of the fluid disappears, the voice becomes more and more distinct; a friction sound finally shows that the roughened surfaces have come in contact; and the dulness on percussion is replaced by a far clearer sound. False membranes now unite the two pleuræ; the intercostal spaces resume their normal shape; and the chest is either restored to its natural size, or is left permanently somewhat contracted. The bronchial breathing near the vertebral column persists for a long time, since the compressed lung unfolds but slowly.

These physical signs have been discussed first because they are the most important elements in the diagnosis of pleurisy. The symptoms, indeed, often hardly attract attention; and if we trusted to them, we should be groping in the dark. Pleurisy mostly begins with a chill, followed by fever and by a dry, irritating cough. The most distinctive, though not a constant, symptom of the first stage is the sharp, acute pain, the "stitch in the side." It is commonly felt under the nipple or in the axilla, and is somewhat increased on pressure. Its seat by no means always corresponds to the seat of the friction sound. As the effusion takes place, the pain disappears, dyspnœa becomes evident, and the patient ordinarily lies on the affected side. The febrile symptoms and dry cough continue; yet neither is marked, and both disappear long before the fluid is entirely absorbed.

Pleurisy may be idiopathic, coming on generally after exposure to cold and damp; or it may be an attendant upon other diseases of the lungs, such as pneumonia or tuberculosis, or may accompany measles, scarlatina, typhoid and typhus fevers. It may also be caused by wounds of the thoracic walls, by rheumatism, gout, Bright's disease, diphtheria, pyæmia, cirrhosis of the liver, and other morbid states.

The malady with which acute pleurisy is most likely to be confounded is *acute pneumonia*. Both are affections occasioning dyspnœa; both are, in the majority of cases, one-sided; both present, in their most advanced stages, dulness on percus-

sion. But the dulness in the latter disease is far less absolute than in the former ; nor do we, save in very rare instances, meet with a tympanitic or an amphoric percussion sound in pneumonia, while in pleurisy, as we have just seen, it is far from unusual above the level of the fluid. In the few cases in which an amphoric or a tympanitic sound is perceived in pneumonia, the peculiar tone is most obvious over the consolidated tissue.

The other physical signs of the two diseases show still less similitude. The absence of respiration, of vocal resonance, and of thrill is in striking contrast with the loud blowing respiration, the strong chest-voice, and the increased vocal thrill of pneumonia. There are, however, exceptional cases of pleuritic effusion, in which bronchial breathing is heard all over one side of the chest. Especially does this happen if pneumonic consolidation accompany the effusion ; but even in simple compression of the lung, and where the collection of liquid is not extensive, bronchial respiration may be perceived. The difficulty of distinguishing from pneumonia such cases of pleurisy, in which probably the lung-tissue is compressed around the bronchial tubes but these are not encroached upon, is great. As aids in diagnosis, we seek for dilatation of the chest ; we note the peculiarities of the breathing, which, although blowing, is mostly fainter than, and is unlike, the high-pitched, brazen respiration of pneumonia ; we find that the percussion dulness over the upper part and where the bronchial respiration is most distinct is not very great, and, especially, that it disappears on respiratory percussion ; we observe that the voice is less strong and ringing, and has, perhaps, a bleating tone ; and we take into account the appearance of the sputum and the character of the fever. On the other hand, pneumonia may present itself in a form distinguishable from pleurisy in the stage of effusion with the greatest difficulty ; it is when the bronchial tubes as well as the lung structure are filled with a fibrinous exudation. In this *massive pneumonia* we do not find either tubular breathing or fremitus attending the flat percussion note, and it is only by noting the absence of displacement of the heart or the liver, the violent coughing spells, and observing the fragments of moulds of the bronchi in the expectoration that a conclusion can be arrived at. But generally the diagnosis between pleurisy and pneumonia is easy. It may be thus summed up :

## PLEURISY.

Sharp pain; friction sound; dry cough; impaired chest-motion.

In stage of effusion, obliteration of the intercostal spaces; enlargement of the side; displacement of several viscera.

In the large majority of cases, dullness, with enfeebled or absent respiration, voice, and fremitus.

Decubitus is often on the affected side.

Sputa frothy; rarely any rales in the chest.

Fever slight. Temperature record irregular, and not characteristic; rarely 103°.

## PNEUMONIA.

Dull pain; crepitant rale; cough, followed by expectoration.

In stage of hepatization, none of these signs are manifest.

Dulness, with marked bronchial respiration; distinct thoracic voice; increased vocal fremitus.

Decubitus not peculiar; sometimes on the sound side.

Sputa rusty-colored; rales from accompanying bronchial inflammation common.

Fever severe. Temperature record much more characteristic. Temperature rises rapidly soon after onset, then is continuous, with marked evening exacerbations and morning remissions. Often reaches 105°, especially in pneumonia of upper lobe. Toward end of disease generally rapid defervescence.

In the first stage of pleurisy the pain might cause the disease to be confounded with pleurodynia or intercostal neuralgia. In all three pain is the prominent symptom. Let us see how it differs in each:

*Pleurodynia.*—Pleurodynia is described as a form of muscular rheumatism. But frequently it is myalgia, or pleurisy which does not pass beyond the dry stage. Of this nature are most of the fugitive chest-pains from which phthisical patients suffer. Yet there are cases in which no signs whatever of pleurisy exist, but which are attended with as much pain as pleurisy. The pain of pleurodynia is, indeed, often excessively severe; the patient refrains from deep breathing, since every motion of his chest increases his suffering. The pain is augmented by movements of the arm and by pressure, and is generally associated with tenderness. Pleurodynia shares with pleurisy the feeble respiration and the want of action of the affected side. It differs from it

by the absence of friction sound and of fever; by the shifting pain, often double-sided; and by the greater tenderness of the chest-walls.

*Intercostal Neuralgia.*—In anæmic women and in consumptives acute thoracic pain is not uncommonly the result of an intercostal neuralgia. The same want of expansion of the chest and the same enfeebled breathing as in pleurodynia are here noted, also the same absence of fever and of pleural friction. The distinguishing marks of intercostal neuralgia are: its intermittent character; its frequent association with uterine disturbance, especially with leucorrhœa, and the limitation of the tenderness to special points in the course of the affected nerve. Valleix has drawn attention to three painful spots which are tender to the touch: one at the exit of the nerve from the spinal column, the second in the axillary region, and the third near the sternum or in the epigastric region. It is on the left side that we are most apt to find intercostal neuralgia, and between the sixth and ninth ribs that the painful places are usually detected.

Pain occurs also in diseases affecting the lung-texture. There is pain of a dull nature in pneumonia, of a more severe character in cancer. But the pain is so dissimilar, and the coexisting symptoms are so unlike, that the error of confounding these maladies with pleurisy, on account of the pain, is not likely to be committed.

### Diseases presenting Dilatation of the Chest, Displacement of the Liver or Heart, and Dyspnœa.

A group of diseases may here be studied, all of which occasion more or less dilatation and prominence of the chest, and all of which are attended with decided shortness of breath. In bronchitis and pneumonia a slight increase in the diameters of the chest may take place; but it is not a sign of any diagnostic importance. In the recognition of emphysema, pneumothorax, and pleuritic effusions, the dilatation of the thorax forms one of the main elements; moreover, it is often combined with marked dyspnœa and with displacement of the liver or heart. These affections, then, may be examined in the same connection, and compared with one another, and incidentally with several less common diseases which present similar manifestations.



The history and signs of emphysema were given when treating of the diseases accompanied by clearness on percussion. It was then mentioned that in many instances the prominence of the chest is circumscribed. Such cases cannot be mistaken: the bulging is too limited. But when the emphysema is more general, and an entire side of the chest or the whole chest becomes dilated, or when the inflated lung displaces the liver or heart, the affection comes into the group under consideration. A patient seeks advice for shortness of breath. His chest is inspected, and looks enlarged. The physical signs prove that the disease is not one of the heart, or an aneurism. What, then, is it? Is it an effusion into the pleura? is it an intra-thoracic tumor? is it pneumothorax? is it emphysema? A tap on the chest goes far toward showing whether it is the former. If the sound rendered be resonant, it is not liquid in the chest that is producing the disturbance, nor, except under rare circumstances, an intra-thoracic tumor: the disorder is either pneumothorax or emphysema.

**Pneumothorax.**—Of all thoracic maladies, pneumothorax is the one the similarity of which to extensive dilatation of the air-cells is the greatest. In both, the large quantity of air occasions increased clearness on percussion; in both, there is considerable and persistent difficulty of breathing; in both, the distention of the chest and the displacement of organs may be obvious. In pneumothorax, however, the symptoms and signs are associated with different conditions. Pneumothorax is an accumulation of air in the pleural cavity, but it is something more: the entrance of air is soon followed by the effusion of liquid.

Air is let into the cavity of the chest by the pleura being perforated by wounds, or through the diaphragm by malignant disease of the stomach or the colon, or, as is most common, by its partial destruction consequent upon disease of the lung. It is in this way pneumothorax originates in the course of tubercular softening, of gangrene, of pneumonia, or from the bursting of a distended air-vesicle or of a dilated bronchial tube.\* In the large majority of instances it occurs in tubercular patients.

When air passes from the lung into the pleura, it usually happens during a paroxysm of coughing. The pain which ensues


---

\* Case recorded by Taylor, *Prov. Med. Journ.*, vol. i., 1842.

is most intense; and the frightful, suddenly-developed dyspnœa, the anxious expression of the face, soon show how seriously respiration is interfered with. If death do not take place, symptoms of pleurisy with effusion manifest themselves; and, as in pleurisy, the patient lies ordinarily, but not invariably, on the affected side.

The distinctive marks of pneumothorax are furnished by its physical signs. The ingress of air into the pleural cavity widens the chest, effaces the depression of the intercostal spaces, and occasions an extremely clear, or, more correctly speaking, a tym-

FIG. 42.



Physical signs in pneumothorax on the right side. The heart is observed to be displaced toward the left, as actually happened in the case from which the outline was taken. The percussion resonance on the right side was tympanitic, extending somewhat over the left margin of the sternum; the fremitus was annulled; the voice metallic.

panitic, sound on percussion. The air prevents the lung from expanding: hence there is an enfeebled or absent respiration, except near the spinal column, where the compressed organ lies, and where the breathing is bronchial. The hand, if laid on any

other portion of the chest, feels, when the patient speaks, no thrill, and no vocal vibration is detected by the ear. When the perforation has not closed, and the air rushes into the artificial cavity produced by the separation of the two surfaces of the pleura, the respiration is amphoric, or it, the voice, and the rales are all accompanied by a distinct metallic ring; respiratory percussion, too, changes the sound elicited, rendering it duller. Drops of fluid falling into the cavity, or the bursting of bubbles on the surface of the liquid in the pleura, are echoed to the ear with a metallic sound, and are often heard as a silvery tinkle. A metallic echoing sound is also obtained if the ear is placed on the back over the affected side of the chest while a coin is tapped on another in front.

The presence of the fluid in the pleural cavity gives rise to a dull sound on percussion at the lower part of the chest, which changes readily with the position of the patient, and to a splash, perceptible to the ear and to the finger, when the thorax is suddenly shaken. This continues until the effusion increases, and until the opening in the membrane closes, the air disappears, and the case resolves itself into one of chronic pleurisy,—the most favorable termination of pneumothorax.

Now let us compare the physical signs with those produced by emphysema. The sound on percussion in both is very clear, or is tympanitic; more so, however, in pneumothorax, which, in addition, exhibits dulness at the lower part of the chest. The respiration in both is feeble. But it is feebler in pneumothorax, and not accompanied by a long, laborious expiration; besides, it is often amphoric, and attended with metallic voice and tinkling,—phenomena which dilated air-cells cannot occasion. Moreover, there can be no splashing sound in emphysema, and this always exists in pneumothorax, except in those rare instances in which there is no fluid in the pleural cavity; on the other hand, the displacement of the heart is generally much greater in pneumothorax, and the dilatation of the chest more apt to be one-sided. Yet too much stress has been laid on the latter point as a means of distinction; for emphysema may be one-sided, and, on the other hand, pneumothorax, as I know from meeting with a number of instances, may occur on both sides. In some cases we are aided in the discrimination by noticing that bulging is perceptible over

the displaced heart, and that a metallic echo follows the cardiac sounds.

The physical signs of the two diseases are thus very different; so, too, are many of the symptoms. Difficulty of breathing exists in both. But in emphysema it takes more the form of attacks of asthma; besides, it does not set in suddenly and with intensity, and remain intense. In pneumothorax the patient remembers to have been seized with a pain in his chest, since which period he has been continuously very short of breath.

Yet there are exceptions to this: there are cases in which the symptoms occasioned by perforation of the pleura are from the onset so slight as not to attract the least attention. Such cases cannot be recognized, save by their physical signs. Among these, dilatation of the chest, with the widened intercostal spaces, the displacement of the liver or heart, and the exaggerated and altered resonance on percussion are most valuable in preventing the disease from being confounded with some affections which in other respects give rise to many of the same phenomena. In large cavities, for instance, the respiration and voice may be metallic; metallic tinkling, nay, even succussion, may occur. But the prominent chest, the extremely clear, tympanitic, or metallic sound on percussion, bordered by the line of absolute dulness due to the effusion, are not met with. The history also is different, and the dyspnoea is not so great. The same dissimilarities will prevent us from mistaking for pneumothorax a pneumonia in which the percussion sound over the consolidated lung is tympanitic. And a study of the physical signs, too, will at once enable us to discern whether the difficulty in breathing, though it be suddenly developed, and apparently under circumstances which make the swallowing of a foreign body seem likely, be due to this cause, or to perforation of the pleura and pneumothorax.\*

There is, however, a morbid condition which exhibits nearly all the signs and many of the symptoms of pneumothorax, and which, were it more frequent, would be the source of constant errors of diagnosis,—*diaphragmatic hernia*.

Of this rare affection we know but little. Yet, thanks to Bowditch,† what we do know of it teaches us that a protrusion of the

---

\* As in a case of the disease communicated to me by Dr. Walter F. Atlee.

† Buffalo Med. Journ., June and July, 1853.



abdominal organs through the diaphragm will generally dilate one side of the chest, compress the lung, and displace the heart. It will do more: it results in dyspnoea; and, as the stomach or intestines are, for the most part, the viscera which find their way into the chest, metallic tinkling and a tympanitic sound on percussion are detected. These are also signs of pneumothorax. There is, indeed, no mode of separating the two diseases, except by attention to the history of the case, by noting that the dyspnoea of the former suddenly appears and as suddenly disappears, that it has often existed from birth, and that the metallic tinkling happens when the patient is not breathing, and is mixed up with the rumbling sound arising in the stomach or intestine.

It has been made a question whether we can distinguish ordinary cases of pneumothorax from these very rare ones which are supposed to occur *without perforation*. Now, even admitting that such really happen as a sequence, for instance, of decomposition in pleuritic effusions, there are no signs by which we can recognize them with certainty. It has been claimed for them that there is no antecedent history of a chronic pulmonary affection, particularly of phthisis, that there is not that suddenly-occurring severe pain and extreme dyspnoea, that the sputum and breath are never offensive, that metallic tinkling is absent, or rare and inconstant, and that the amphoric breathing is not so well developed or so clearly defined. If in a case of perforation, however, the opening have closed, the physical signs, it is granted, are the same.\*

**Chronic Pleurisy.**—Chronic pleurisy is the third of the group of more usual affections which are characterized by dilatation of the chest, by displacement of the intra-thoracic viscera, and by shortness of breath. It is true that acute pleurisy in the stage of effusion would, strictly speaking, find here a place; but the acute symptoms bring it into another class, with which it has been more conveniently described.

Chronic pleurisy is established if the fluid, after an acute attack, be not absorbed, or if an accumulation of liquid take place gradually, in consequence of subacute inflammation of the pleura. It is also found, especially in its purulent form, in a number of

---

\* Boisseau, Arch. Gén. de Méd., vol. ii., 1867.

infectious diseases, particularly scarlet fever and typhoid fever. This form is also seen to follow pleuro-pneumonia and perforation of the pleura by softening tubercle. Chronic pleurisy has no constant symptoms, and is often remarkably latent: the patient frequently does not remember to have had acute pleurisy. He is not commonly troubled with much cough, nor is the want of breath so great as might be expected; he is not capable of talking for any length of time, or in a loud voice, but he does not really suffer from dyspnœa. His general health may remain good, and no emaciation occur. In some persons, on the other hand, the loss of flesh, the quickened pulse, the sweats, the paroxysms of hectic fever, are so marked as to produce a close resemblance to the last stages of tubercular consumption.

While the differing symptoms rather hide the pleurisy from detection, the physical signs render it easy of recognition. These signs have been studied in describing the effusion in acute pleurisy. It is only necessary to recall that the most significant are absent respiration and voice, a flat sound on percussion, with a vesiculobronchial or a bronchial respiration above the seat of the liquid. The intercostal spaces are obviously widened; their depressions are effaced. They are, indeed, sometimes convex, and the finger pressed on them detects a distinct fluctuation. During the act of breathing, the diseased side is almost motionless, presenting a strong contrast to the obvious play of the healthy side. The lung which is not disturbed increases in size. Its murmur is more intense, sometimes harsher; and the percussion sound over it is exceedingly clear. In some cases it becomes emphysematous. The heart or liver is displaced. A lateral curvature of the spinal column is apt to take place, and the shoulder remains fixed and stiff during the respiratory acts. To distinguish whether the fluid is collected in one cavity or in several, in other words, whether unilocular or multilocular, is generally impossible. Jaccoud\* has, however, called attention to some points which aid in arriving at a conclusion. If we have a zone in the dulness where vocal vibrations are preserved, as at the posterior part of the chest from along the vertebral column toward the sternum, and beyond this zone no vibrations are perceived, we may infer that the effusion is

---

\* Bulletin de l'Académie de Médecine, 1879.

divided by a band of pleural adhesion ; if the voice and fremitus be preserved, although weakened, over the whole extent of the dulness, except in a zone of a few finger-breadths at the lower part of the chest behind, while no tympanitic sound is elicited under the clavicle, we are to conclude that the pleurisy is multilocular. When adhesions to the diaphragm exist, the normal movements during respiration at the epigastrium and hypochondrium are reversed, and at each inspiration a marked depression of the inferior intercostal spaces is perceptible.

Effusions into the pleural sac may last for a long time, and lead to death by progressive exhaustion ; or the patient may recover by the fluid being absorbed, or by its finding a vent through the bronchial tubes or the thoracic walls. But the chest is rarely restored to its former state. The lung was too much compressed, or is still bound down by too firm adhesions, to resume its full share in the function of respiration. The walls of the chest sink in around it, and the side is flattened, sounds duller on percussion, and presents a feebler breathing than the other lung, which remains somewhat enlarged. The heart generally returns to its normal position, but the shoulder on the affected side is apt to show a permanent depression.

Notwithstanding the decided character of the physical signs, chronic pleurisy is frequently overlooked. The only explanation of this is, that too little attention is paid to the physical signs. Were the chest more often carefully explored, we should cease to hear of patients whose pleural cavity is filled with pus being pronounced incurable consumptives, because they are emaciating and have hectic fever and clubbed nails ; or being treated for disease of the heart, on account of the displacement of that organ, and of dyspnoea and œdema ; or being dosed with mercury, for an imaginary disorder of the liver ; or being subjected to long courses of quinine and arsenic, to check a rebellious ague which the chilly sensations and paroxysms of fever at times simulate.

These physical signs are the same whether the fluid be serum or pus. The character of the fluid produces, indeed, no distinctive changes either in the signs or in the symptoms. We suspect empyema if the emaciation be great and accompanied by a quick pulse, high temperature, and hectic fever ; but I have known pus

in the chest with a temperature scarcely above the norm, and, on the other hand, the accumulation not to be purulent with a temperature of  $103^{\circ}$ . Baccelli has proposed a new and simple test to determine the character of the fluid, which, on the whole, I believe to be of use. It consists in ascertaining accurately how the voice penetrates, especially the whispered voice. If easily and thoroughly transmitted, the liquid is serous and homogeneous; if with difficulty, it is fibrinous or purulent; if not at all, it is most apt to be the latter. In cases of much doubt I have long been in the habit of using a hypodermic syringe and removing with it enough of the fluid for microscopical examination.

The *microscopic* and *bacteriological* examination of the exudation will give us valuable information. In rare instances the fluid consists of fat-globules and of masses of cholesterine.\* In cases of hemorrhagic pleurisy the hæmoglobinometer will inform us accurately as to the amount of blood in the exudation.† We find it, indeed, full of blood and altered blood constituents in hemorrhagic pleurisy, a form which pleurisy may assume in low fevers, but which is more frequently found in cancerous, and sometimes in tubercular pleurisy. In the latter disease, contrary to expectation, tubercle bacilli, as we know from the observations of Ehrlich, are frequently absent. There is a group of cases in which, either in a serous or a purulent exudate, we detect the diplococcus pneumoniae; here there may or may not have been a preceding pneumonia. Cases in which the diplococcus pneumoniae is met with are generally of favorable prognosis. In septic pleurisy the streptococcus is found, and especially, as Koplik has shown, the streptococcus pyogenes; staphylococci are also met with. These are much more serious cases, both as to duration and as to recovery.

When we come to inquire into the thoracic diseases with which chronic pleurisy is likely to be confounded, we shall find that, although many have some signs in common, few, if any, present the same association of signs. Leaving out the malady which is most commonly mistaken for it,—pulmonary consumption,—since the points of difference have already been discussed, the affections

---

\* Debove, Soc. Méd. des Hôpitaux de Paris, tome xviii., 1881.

† Henry, Medical News, April 14, 1888.



with which chronic pleurisy, while the pleura is full of liquid and the chest enlarged, is liable to be confounded, are :

EMPHYSEMA AND PNEUMOTHORAX ;  
INTRA-THORACIC TUMOR ;  
ENLARGEMENT OF THE LIVER ;  
ENLARGEMENT OF THE SPLEEN ;  
ABSCESS IN THE THORACIC WALLS ;  
PERICARDIAL EFFUSION ;  
HYDROTHORAX.

*Emphysema and Pneumothorax.*—These, although such different diseases, are grouped together because they give rise, like chronic pleurisy, to a dilated chest, and to displacement of the liver or heart. But the other signs above pointed out, which indicate the presence of air, are so striking, that an error in diagnosis can only be the result of carelessness.

*Intra-thoracic Tumor.*—A tumor within the chest may occasion the same distention of its walls, the same displacement of organs, the same dulness on percussion, and the same absent respiration, as an effusion of liquid into the pleura ; yet the signs are not exactly alike. There is no fluctuation in the bulging intercostal spaces ; the vocal fremitus is not so constantly abolished ; and the level of the dulness is not changed by altering the patient's position. Nor is the flat sound so uniform or so strictly limited as that produced by fluid : amid the dulness may be detected here and there a spot yielding on percussion a clear sound. A tumor in the chest, moreover, presses on the nerves, or bronchial tubes, or great vessels, and thus gives rise to severe pain, and to dyspnœa and signs of interrupted circulation far more evident than are caused by a pleuritic effusion. It not infrequently grows into the mediastinum, and then leads to prominence of the sternum, and to dilatation of both sides of the chest. These phenomena are found, whatever be the nature of the morbid growth. As most of the thoracic tumors are cancerous, we are often assisted in our diagnosis by discovering a cancer in other parts of the body, as well as enlarged cervical glands, and by noting the severe pain in the chest, the harassing cough, and the expectoration of blood or of a peculiar jelly-like substance. Yet these evidences, while they aid us in establishing the fact of a new growth in the thoracic cavity, do not by any means determine its situation. We cannot go a step

further, and say with certainty whether the abnormal formation is situated exclusively in the lung, or in the pleura, or whether it affects both. When the tumor occupies the mediastinal spaces, and is not cancerous, it is most likely a sarcoma. Lymphadenomata come next in frequency.\* In children, however, sarcoma is a more frequent neoplasm than carcinoma.†

In those cases in which an effusion into the pleura complicates an intra-thoracic tumor, attention to the history and to the signs of pressure alone apprises us of its presence. Yet both signs and symptoms may be so closely like those of chronic pleurisy as to render a differential diagnosis impossible. Nay, friction sounds, a stitch in the side, and fever may be produced by a *cancer of the pleura*, and be apparently so rapidly developed as to cause the disease to be regarded as an acute or a subacute inflammation of that membrane. Cancer of the pleura, like tubercle of this structure, has, therefore, no pathognomonic signs. The most certain sign of cancer of the pleura is probably the one mentioned by Trousseau,—namely, that the fluid which is evacuated by paracentesis consists of a bloody serum; yet, though of great significance when present, its absence is not so valuable a test; since Moutard Martin found hemorrhagic effusion in only twelve per cent. of the cases he analyzed. Ehrlich ‡ has published seven cases, in three of which he detected special cellular elements in the fluid, and was thus enabled to come to a correct conclusion. In some instances there is no fluid in a greatly-thickened cancerous pleura.§

It is at times equally impossible to distinguish a *circumscribed pleurisy* from a tumor in the chest. In those rare cases in which adhesions bound the liquid effusion and encyst it, we observe all the marks of a tumor,—a restricted bulging and percussion dullness, and absent respiration and tactile fremitus, though this may be retained over the lines of the adhesions. Several cysts may form as the result of successive attacks of pleurisy, and may exist in any portion of the chest. The fluid may be collected in the mediastinum, or between the lobes of the lung, or anywhere between the surfaces of the pleural membrane. The purulent

---

\* Hobart A. Hare, *Affections of the Mediastinum*, 1889.

† Edwards, *Archives of Pediatrics*, July, 1889.

‡ Charité Annalen, 1882.

§ Purjesz, *Deutsches Archiv f. Klin. Med.*, Aug. 1883.

contents of the sac sometimes find their way into the bronchial tubes, and are expectorated, or give rise to a distinct fluctuation in the intercostal spaces, and then discharge through the thoracic parietes. In such cases the diagnosis is not difficult. But where these phenomena are not present, the dissimilar history of the case and the absence of symptoms of pressure are the only means of distinction from a tumor in the chest. Fortunately, encysted pleurisy is a rare disease; were it frequent, it would be a fruitful source of error. The same remark applies to hydatid cysts, which may occasion all the signs of a circumscribed pleurisy.\* An examination of the fluid obtained by an exploratory puncture, in which echinococci are found, is the only positive test.

*Enlargement of the Liver.*—An enlarged liver usually descends into the abdominal cavity; yet it may be forced upward as far as the fourth rib, and, by encroaching upon the lung, may give rise to many of the physical signs of a pleuritic effusion. The surest diagnostic test is, that during full inspiration and expiration the line of dulness descends and ascends; while the flat sound of a pleuritic effusion is not affected by the play of the lungs. This test will be applicable except where the liver is firmly adherent to the walls of the abdomen. As circumstances to assist in discriminating between the enlargement of the abdominal organ and the presence of liquid in the chest, may be mentioned that the heart, if at all displaced, is pushed upward, and not toward the side; and that the dulness of an enlarged liver extends higher up anteriorly than posteriorly, which is the reverse of what takes place in a pleuritic effusion. Moreover, the respiration at the lower portion of the lung posteriorly, although enfeebled, is still audible.

*Enlargement of the Spleen.*—An enlarged spleen is attended with prominence and with dulness on percussion at the lower part of the chest on the left side, and might, therefore, mislead into the idea of a pleuritic effusion. Error in diagnosis is prevented by attention to the fact that the dulness extends also

---

\* See the observations of Vigla, Arch. Gén. de Méd., Sept. and Nov. 1855; Lebert's Klinik der Brustkrankheiten, Bd. ii.; P. Kidd, Transact. Pathol. Soc., London, 1884-85, xxxvi.; C. Hochsinger, Wien. Med. Blätt., 1887, x.; J. D. Thomas, Australasian Medical Gazette, 1887-88, vii.; L. Bard and R. Chabannes, Rev. de Méd., Paris, 1888, viii.; Netter, La Semaine Méd., Aug. 1892.

downward and toward the median line. Again, the heart is not laterally displaced, but tilted upward; the respiration is feeble, but not absent; and the vocal vibrations are mostly unimpaired.

*Abscess in the Thoracic Walls.*—This, too, leads to local tumefaction and fluctuation; but we can ascertain whether a fluctuating tumor in the intercostal spaces communicates with the pleural cavity or not—whether, in other words, it is or is not the result of an effusion which is pointing externally—by watching how pressure and the acts of respiration affect it. For, unless the diaphragm has become immovable from the extent of the effusion, a bulging which is in connection with the pleura is diminished during a full inspiration, and becomes more prominent when the diaphragm ascends in expiration. The swelling, moreover, can be made to disappear to some extent by pressure. It is not so with an abscess seated in the walls of the chest. It is not reducible, and does not recede during inspiration.

*Pericardial Effusion.*—An effusion into the pericardium ought not to be mistaken for an effusion into the pleura. The first induces prominence and increased dulness on percussion over the region of the heart; the second, dulness and prominence over the back part as well as over the front of the lung. A few cases are, however, recorded in which an enormously-distended pericardial sac produced a flat sound posteriorly, and gave rise to signs of compression of the lung. But in these attention to the feeble impulse of the heart and its muffled sounds permitted it to be told that fluid had accumulated in the pericardium, and not in the pleura.

*Hydrothorax.*—A dropsy having its seat in the pleural cavity is called hydrothorax, or water on the chest. The term is, in truth, sufficiently significant, the fluid which is poured out being very thin and watery. The physical signs of hydrothorax are the same as those of an effusion due to inflammation; but, as the dropsy results from an organic disease of the liver, heart, or kidneys, the serum collects in both pleural sacs. Now, an effusion caused by an inflammation of the pleura is nearly always one-sided. Even where both pleuræ are filled with fluid,—a rare condition, except in tubercular pleurisy,—one is affected before the other. This does not happen in hydrothorax. Thus the double-sided effusion, and its usual association with dropsies in



other parts of the body, are matters of much significance. Besides, in forming a diagnosis of hydrothorax we may lay stress on the absence of friction sounds; on the smaller quantity of fluid; on the history of the malady; and especially on the presence of a structural lesion of the liver, kidneys, or heart.

These, then, are the diseases with which chronic pleurisy, when it produces dilatation of the chest, may be confounded. Indeed, in view of the frequency of the operation of aspiration or of paracentesis, it is important to know what affections besides chronic pleurisy may lead to prominence of the chest and to compression of the lung; and tapping the chest has in itself certain diagnostic bearings which may be here alluded to. One of these is an albuminous expectoration that follows, which may be looked upon as a passing albuminuria due to circulatory disturbances. It is not an unfavorable event; on the contrary, in cases in which it happens, retraction of the thoracic parietes is less likely to occur.\*

#### Diseases in which Retraction of the Chest occurs.

**Chronic Pleurisy.**—We may here continue the description of chronic pleurisy in the stage of absorption, since it is under these circumstances that the most marked retraction of the walls of the chest takes place. This shrinking of the thoracic parietes is not a sudden, but a gradual act, and instances are therefore constantly met with in which the upper part of the chest is flattened and the lower, owing to its still containing fluid, bulges. The contraction of one side of the thorax attains its highest degree when the effusion in the pleura is discharged through the chest-walls and external fistulous openings are established.

The symptoms in the stage of retraction are those of chronic pleurisy with dilatation of the chest, and present, therefore, the same variableness. But œdema of the affected side, which is sometimes so striking a symptom of chronic pleurisy when the effusion is considerable, is here not noticed. The physical signs alter somewhat, according to the presence or absence of fluid in the pleural sac. When none exists, respiration is heard all over the lung as a feeble inspiration with prolonged expiration, or as an indistinct blowing; and now and then a friction sound may be

---

\* Legroux, Arch. Gén. de Méd., Aug. 1873.

caught. When the pleura still contains liquid, these signs occur at the upper portion of the chest, and a much more absolute dulness on percussion, an absent voice and vocal fremitus at the lower part denote that fluid has there accumulated. The heart is found either in its normal position or still displaced. The force with which contraction takes place may pull it over to the side on which the shrinking is going on. Wasting of the muscles of the shoulder and sensory changes on the affected side of the chest have been observed as a result of chronic adhesive pleurisy.\*

Now, it is evident that chronic pleurisy, when leading to retraction of one side of the chest, cannot be mistaken for diseases attended with thoracic distention; but it may be mistaken for affections like pulmonary cancer, tubercle, and chronic consolidation, which also occasion a flattening of the chest-walls.

From *cancer* we distinguish it by the absence of the peculiar expectoration, and of hemorrhage; by the want of signs of perfect consolidation; and by the dissimilar history. We distinguish it from *tubercle* by the diminution of the chest in the latter not being confined to one side; by the physical signs indicative of deposit and softening at the upper portion of the lungs; by the presence of rales; by the occurrence of hemorrhage; by the greater emaciation; and by the tubercle bacilli in the sputum.

Chronic interstitial pneumonia presents, on the whole, most points of resemblance. But there is this difference: the shrinking of the side in this disease is less marked, and is confined to the part involved,—usually the lower lobe of the lung. The retraction is much more general in chronic pleurisy; or where it is partial, it is the upper segment of one side of the chest which is flattened,—the lower is prominent, and sounds very dull on percussion, shows no change on respiratory percussion, and yields the ordinary physical evidence of a fluid. In the former malady the blowing respiration, or the enfeebled inspiration and prolonged expiration, and the distinct voice are heard only over the consolidated lobe; in the other lobes the breathing is plainly vesicular. In chronic pleurisy the same abnormal signs, except perhaps the increased voice, are either manifest over an entire

---

\* Thévenet, Lyon Méd., 1894, No. 5.

side, or are perceived over the narrowed portion of the chest, and at the lower part the respiration, voice, and fremitus are abolished.

In that form of chronic pulmonary induration attended commonly with dilatation of the bronchial tubes, to which the name of *cirrhosis* of the lung,\* or *fibroid phthisis*, has been given, the flattening of the affected side is as obvious as it is in pleurisy. In truth, the two disorders bear a strong relation to each other. The increased formation of connective tissue in the pleuritic adhesions passes on into the lung, occasioning an interstitial pneumonia,—though the fibroid change oftener begins in the lung; as this progresses and the lung shrinks, bronchial dilatations usually follow. We distinguish cirrhosis of the lung by the copious and peculiar sputum which attends the bronchial affection; by the rales; by the harsh or bronchial or tubular or feeble respiration; by the dulness on percussion with an occasional tympanitic note; by the marked resistance of the chest-walls; by the increased vocal resonance; by the narrowing of the intercostal spaces; and by the displaced apex beat,—forced up, if the disorder be on the left side, one or several intercostal spaces, or so covered by the expanded left lung, if the disorder be on the right, as to be imperceptible, unless the shrinking of the affected lung be considerable, when the heart may be found drawn over on the diseased side. Further signs of the complaint, when the malady is left-sided, are that in the second intercostal space to the left of the sternum a double beat of the pulmonary artery is perceptible, and that whichever side is diseased shows the diaphragm greatly displaced upward, and a marked vesicular resonance in a line along the edge of the sternum caused by the overlapping of the healthy lung, and in strong contrast with the line of dulness of the cirrhotic organ.† The affection is a very chronic one, and unattended with fever or laryngeal symptoms. Loss of flesh and of strength is very gradual, and night-sweats are slight or inconstant. Dilatation, or hypertrophy with dilatation, of the right side of the heart, and dropsy, are not infrequent, and hæmoptysis is still oftener met with. It is a mistake to suppose that it occurs

---

\* Corrigan, Dublin Quart. Journ., vol. xiii.

† Nothnagel, Sammlung Klinischer Vorträge, 1874.

only when tubercles are present, or in what is called the bacillary variety of fibroid phthisis.\*

The disease has among its causes the inhalation of fine particles, such as of steel, of coal-dust, of iron-dust, of cotton. It may have an obscure beginning, or it may clearly date from an acute pneumonia, especially an acute or a subacute broncho-pneumonia, or a plastic pleurisy. It may become complicated with tubercle, and then tubercle bacilli are found in the sputum. The fibroid condition of the lungs—also called by some fibroid phthisis—in old tubercular lungs or around cavities is an evidence of a disposition toward healing, a local fibroid change, and is not fibroid phthisis, though this term is limited by some to cases of indurated lung of tubercular origin. Pulmonary cirrhosis often proves fatal from an acute affection, a pneumonia or a broncho-pneumonia, of the previously healthy lung. In very rare instances it is double.†

The connection of pleurisy with the cirrhotic lung has just been mentioned; and, though the origin of interstitial pneumonia from invasion through the pleura is in dispute, I hold the view to be correct. We must, however, not forget that primitive dry pleurisy is found also under other circumstances, and may give rise to retraction of the chest. Firm fibrous bands may result from organization in the pleura after a dry pleurisy, or after absorption of the effusion; or the dry pleurisy may be of tuberculous origin. It is then usually double-sided. Osler‡ mentions some remarkable vaso-motor phenomena when these primitive dry pleurisies affect the apex and probably involve the first thoracic ganglion, such as flushing or sweating of one cheek or dilatation of the pupil.

A *collapsed* state of the lung, resulting from a plug of mucus in the bronchial tubes, may yield the manifestations of chronic pleurisy with partial retraction. No signs distinguish such cases, except the more limited depression; the absence of any disease above the flattened spot; the want of friction sound, and of tenderness on pressure; and the rapid disappearance of the physical

---

\* Sir Andrew Clark, *Lancet*, July, 1892.

† McCollom, *New York State Med. Assoc.*, 1885.

‡ *Practice of Medicine*, 1892.



phenomena after an effort of coughing has removed the obstruction.\*

Where external *fistulous openings* exist, the shrinking of the side, as already stated, is carried to the highest degree. These fistulæ, whether produced artificially or by nature, may close after they have served the purpose of evacuating the fluid in the pleural cavity. But they often persist for months or years, and keep on discharging offensive, purulent matter. The patient emaciates under this continued drain, yet not so quickly as might be imagined. More or less troublesome cough annoys him, but it is not ordinarily accompanied by much expectoration. Every now and then, however, he discharges for days a quantity of fetid, purulent sputum. It is then very likely, as Traube has observed, that the pus has softened part of the pulmonary pleura sufficiently to soak through the lung into a bronchial tube. It seems certain, as far as physical signs can prove, that it is not the liquid in the pleura which is being voided through a distinct perforation of the pulmonary tissue, for the physical signs of pneumothorax are absent.

The clubbing of the nails is often extremely marked, and may exist to an extent far greater than in phthisis. The nail is rounded and bluish, and the whole end of the finger looks enlarged. This appearance is even more striking than the curve of the nail. The nails and last joints of the toes show the same alteration.

The fistulous opening is situated ordinarily in the intercostal space below the nipple. It may, however, be seated at the back of the chest, and communicate by a tortuous sinus with the intestine and other abdominal viscera. A pleuro-bronchial fistula may form; if the opening pass into the lung, the physical evidences of pneumothorax are present, but the side is still retracted, and striking the chest elicits a mixture of a dull and a tympanitic sound. Where merely an external opening exists, no signs of pneumothorax occur, because no air finds its way into the pleural cavity.

A fistulous opening into the pleura is not difficult of diagnosis.

---

\* An interesting instance of the kind is related by Prof. William Pepper the elder in the American Journal of the Medical Sciences for April, 1852.

It is easy to establish the fact that the fistula is not simply produced by caries of the rib ; for a probe may be run into the chest for two, three, or four inches.

I base these statements on a number of instances of chronic pleurisy attended with external fistula which have come under my notice. The seat of the opening near the nipple ; the peculiar nail ; the occasional flow for days of a most offensive sputum from the bronchial tubes, without any traces of pneumothorax ; the ease with which the fistula could be probed, and its depth ; the gradual emaciation ; and, I may add, the decided improvement under the persistent use of cod-liver oil and tonics,—belonged to them all, and justify the description given.



## SECTION II.


### DISEASES OF THE HEART.

The heart is kept from rolling about in the chest by the great vessels which spring from its base, and by the attachment to the diaphragm of its membranous covering,—the pericardium. It lies obliquely in this membrane, with its long axis directed downward and toward the left. Its base points backward and upward toward the right shoulder ; its under side rests upon the central tendon of the diaphragm. The interior of the heart is lined by a serous membrane,—the endocardium,—which is reflected over the valves. These valves all lie in close proximity to one another, and within a space of less than an inch square.

The relations the different parts of the organ bear to the chest-walls are as follows. The auricles are on a line with the third costal cartilages ; the right auricle extends across the sternum to the right side of the chest. The right ventricle is placed partly under the sternum, and partly to the left of it. Its inferior border is on a level with the sixth cartilage. The left ventricle lies within the nipple, between the third and fifth intercostal spaces. The apex is seated between the cartilages of the fifth and sixth ribs, to the inner side of, and from an inch and a half to two

inches below, the left nipple. The base of the heart corresponds posteriorly to the sixth and seventh dorsal vertebræ, from which it is separated by the aorta and œsophagus. The greater portion

FIG. 43.



Topography of the heart. The relations of each portion of the heart to the walls of the chest are shown. The dotted lines mark the lungs. The figure is based upon several careful dissections.

of the anterior surface of the heart is removed from the thoracic walls by the lungs. The right lung extends to the middle of the sternum. The left lung spreads out as far as the fourth cartilage, and covers the whole of the left ventricle, except the apex. The part of the heart which remains exposed consists thus mainly of the lower portion of the right ventricle; it presents the shape of a rude triangle.

The position of the valves can be learned by running needles into the chest before the viscus is taken out. In this manner it is ascertained that at the left border of the sternum, on a level with the third intercostal space, lies the mitral valve, and in front of

this, more directly under the sternum, and but a few lines lower, the tricuspid valve. The pulmonary orifice is seated opposite the junction of the cartilage of the third rib with the left edge of the sternum. Near it, very slightly lower, but placed more obliquely, are the aortic valves. The aorta then proceeds from left to right, and ascends to the upper border of the second costal cartilage on the right side; thence it crosses, under the sternum and in front of the trachea, to the left side. The pulmonary artery is found in the second intercostal space on the left side, enclosed in the pericardium, and passes to the cartilage of the second rib, where it bifurcates.

The size of the heart is about that of the closed fist. Its mean weight in adults is between eight and nine ounces. Only in very large persons does it exceed this.

The organ exhibits, when in action, a wonderfully perfect mechanism and regularity of movement. Its cavities contract on both sides at the same time, and distend on both sides at the same time. It then rests for a short period. The contraction of the ventricles occasions the impulse which is seen and felt in the fifth intercostal space. While the blood is flowing in and out of the heart, the valves are kept in constant motion. Their play makes itself known by two distinct sounds of unequal length, which are produced mainly by their opening and closing.

The first sound, long and dull, is caused by the forcible closure of the valves at the auriculo-ventricular openings. Yet it is not a purely valvular sound. The stroke of the heart against the walls of the chest, the muscular contraction itself, and the flow of blood into the aorta and the pulmonary artery aid in its formation. The first sound corresponds, therefore, to the closure of the auriculo-ventricular valves, to the impulse of the heart, to the opening of the valves at the orifice of the aorta and of the pulmonary artery, and to the passage of blood along the arteries. The second sound is short, abrupt, and ringing. It results from the sudden closure of the semilunar valves. During its occurrence the blood rushes through the opened mitral and tricuspid valves, and dilates the ventricles.



## Examination of the Heart by the Different Methods of Physical Diagnosis.

Before proceeding to examine the heart, we inquire into the history of the case, and into such symptoms as the expression of the face; the appearance of the eye; the condition of the capillary circulation; the presence or absence of dropsical swellings and of cough; the state of the breathing; the character of the pulse; and the frequency and violence of the palpitations. The cardiac region is then explored by the eye and by the hand; the size of the organ is estimated by percussion; and, lastly, its sounds are studied by the stethoscope. These different methods are most conveniently practised when the patient is in an easy position, leaning back in a chair or propped up with pillows in bed. To examine them more in detail:

### INSPECTION.

Inspection detects on the chest of some healthy persons a slight protrusion over the seat of the heart; yet this is far from being constant or even the general rule. When the heart is hypertrophied, or when fluid has accumulated in the pericardium, we perceive a marked prominence in the præcordial region. A depression at the lower part of this region may be natural; a very evident depression is almost always the result of an attack of pericardial inflammation.

Yet neither prominence nor depression is a very important sign. One much more so, which inspection shows, is the *impulse* of the heart. This is seen where the apex beats against the walls of the chest: between the fifth and sixth ribs, about an inch inward from the nipple and two inches downward. It is for the most part confined to this point, and appears as a brief raising of the integument, occurring with great regularity of succession. In lean persons it is very distinct; in fat persons it is generally not at all perceptible. Its seat, even in those who are in perfect health, is not always exactly the same. It is changed by different positions, and by the distention of the stomach after a full meal or by flatulence. It is most modified by the acts of respiration. During a long-drawn inspiration the heart descends somewhat and the

expanded lung sweeps it inward, and the impulse becomes discernible in the epigastrium. During a fixed expiration the beat moves upward, and appears more extended and weightier. The changes produced in its situation by disease, both thoracic and abdominal, are many. It is tilted upward and outward by the left lobe of an enlarged liver. It is displaced by diverse affections of the lungs and pleura. It is forced up by a pericardial effusion. It is visible lower down and over a larger surface in enlargements of the heart; but even then it is most distinct at the apex. The apex beat lies without the line of the nipple in most children up to the fourth year.\*

The alterations in the character and force of the impulse are as diversified as those of its seat. But they are more readily appreciated by the hand than by the eye.

#### PALPATION.

Palpation is, so far as the exploration of the heart is concerned, much preferable to inspection. Many an impulse can be felt which cannot be seen. The rhythm of the motion is changed by a large number of cardiac affections, both functional and organic. So are the extent and force of the beat. Both are temporarily increased by powerful excitement; both are permanently augmented by hypertrophy. In dilatation and pericardial effusion, the extent over which the stroke is felt is greater than in health; but the impulse is feeble, and in the latter disease irregular and wavy. Softening of the texture of the heart, diseases of the brain, some morbid states of the blood, and a low condition of the system will also enfeeble the beat.

The hand, when laid on the præcordial region, perceives at times *two* impulses. This double impulse is often recognizable in health, especially in thin persons. It becomes still more evident in hypertrophy with dilatation of the ventricles. One of the beats is systolic; the other corresponds to the diastole. Bouillaud cites examples in which the diastolic stroke was double. The systolic beat is occasionally split into several parts when the pericardium adheres to the heart.

---

\* J. Mitchell Bruce, Enlargement of the Heart, in Keating's Cyclopædia of the Diseases of Children, vol. ii.

All these modifications of the impulse stand in direct connection with the action of the ventricles. The auricles, save in some rare instances in which they are dilated and their walls thickened, give rise to no perceptible movement in the chest-wall.

Besides the impulse of the heart, other phenomena may be studied by placing the hand over the cardiac region. The sounds of the heart can be analyzed by means of the touch. They will be felt, the one as a long and dull, the other as a short and distinct, vibration. The motion is due to the play of the valves, and disappears with their destruction. The fingers applied over the heart perceive at times a peculiar thrill, or a rubbing movement. The first—called by Laennec, from its resemblance to the purr of a cat, the purring tremor—is nearly always indicative of a valvular lesion. The second is caused by the to-and-fro motion of a roughened pericardium.

A more accurate means of studying the varying impulse than is afforded by the fingers has been sought to be attained by instruments which record the beat of the heart. The cardioscope of Alison and the cardiograph of Marey and of Galabin have been used for the close analysis of the cardiac impulse, as have the sphygmographs of Pond and of Brondgeest. But as yet these instruments have not proved to be of marked diagnostic value.

#### PERCUSSION.

Percussion affords the readiest means of judging of the size of the heart. The patient is placed in a recumbent position; then, by a series of moderately strong taps, we proceed downward from near the middle of the left clavicle, until a dull sound, accompanied by decided resistance, tells that we are striking over a solid organ. The point at which this dull sound begins is over, or immediately at the lower border of, the fourth cartilage. It corresponds to the upper limit of the portion of the heart which is left uncovered by the lung.

The superior border of the dulness having been thus ascertained, we next percuss on the right side of the sternum, on about a level with the fifth rib, and progress across the bone. At, or very near to, its left edge we find marked resistance and a duller sound. Here we draw our second line, and continue to strike straight across the cardiac region up to the point at which a

clear, full note demonstrates that the pulmonary tissue is resounding. This determines the transverse diameter of the heart,—at least so far as it can be mapped out on the chest. The apex of the organ and its inferior surface remain to be fixed. The first is readily done by advancing in an oblique direction from the already ascertained right border. But we can save ourselves this trouble by feeling for the impulse or by listening for it with a stethoscope.

The inferior surface can be circumscribed by prolonging the line of the dulness on percussion of the upper border of the liver, and then judging by the greater amount of resistance and the fall in pitch that the heart has been reached. These are not easy to appreciate; nor is it indeed often necessary to define the contiguous edges of the left lobe of the liver and of the heart. If the other boundaries have been correctly drawn, the size of the heart can be accurately estimated,—accurately enough, at least, for any practical purpose. The dulness elicited by percussing the cardiac region is not so absolute as that of the liver or of some other solids. It is mixed with the sound of the lung-tissue, or with the resonance of the sternum. Nor is it a representation of the size of the entire organ. It simply portrays the more superficial portion, which is uncovered by the lungs.

In women it is particularly difficult to define these limits. It can be done only by having the mammary gland drawn to one side while percussing. It is equally difficult in children, as the space over which the dulness is perceived is very small. In adults the dulness ordinarily spreads over two, or nearly two, intercostal spaces. Its transverse diameter in a grown person of medium size is about two inches and a half. In tall, broad-chested men it is upwards of three inches. Such, at all events, is the result of measurements I have made.

The range of the dulness is changed by a number of causes, physiological as well as pathological. A full inspiration alters it materially, by bringing the lung down over the heart, and by displacing the organ itself. The upper border of the percussion dulness shifts to the extent of an intercostal space. Below the nipple, between the fifth and six ribs, the sound becomes clear; but over the dislodged lower part of the heart, the beat of which is distinctly seen under the cartilages of the ribs, at a



point varying from three-fourths to one and a fourth inch from the median line, there is dulness with resistance to the finger. A full expiration produces, for the most part, converse phenomena. It enlarges the boundaries, especially in an upward and transverse direction. The dulness reaches nearly, or even entirely, across the sternum. Auscultatory percussion enables us to fix the percussion limits more closely. Sansom's pleximeter also conduces to greater accuracy in cardiac percussion.\*

The area of dulness is diminished in emphysema. It is increased by a shrinking of the left lung, and by diseases of the heart and of its membranes. Prominent among these are hypertrophy, dilatation, and an effusion into the pericardial sac.

#### AUSCULTATION.

When the ear or a stethoscope is applied over a healthy heart, it detects two sounds of very dissimilar character: the first is long, dull, heavy, and corresponds to the impulse against the walls of the chest; the second is short and flapping, and occurs after the impulse. These sounds are audible at all parts of the præcordial region, but not everywhere with equal distinctness. The first, being more ventricular in origin, is best heard over the lower part of the heart; the second, a more strictly valvular sound, is more defined at the base.

It has been already stated that these sounds are, to a great extent, produced by the play of the valves. Each of these forms a separate sound, or at least a portion of one. Now, experience teaches that there are points at which the sounds of the several parts of the heart may be isolated. Some of these points accord with the anatomical seat of the valves; others do not. None do so very closely; and the proximity of the valves to one another is such as to make it desirable that the localities selected for listening to them should be some distance apart.

Clinical observation sanctions the following: the sounds of the aorta are to be studied at the right edge of the sternum, in the second intercostal space; from there the stethoscope may be carried to the second costal cartilage of the right side, the "aortic

---

\* See paper by William Ewart, *Lancet*, Aug. 1891.

cartilage," and down to the left edge of the sternum opposite the third intercostal space; that is, not far from the seat of the aortic valves. The pulmonary orifice lies very close to them; but the artery itself ascends to the second costal cartilage on the left side.

FIG. 44.




Diagram showing the points at which the separate valves may be listened to.

Its sound may, therefore, be isolated in the second intercostal space, near to the left edge of the sternum. The mitral is listened to immediately above the beat of the apex. The sounds of the tricuspid and of the right ventricle may be sought for in the vicinity of and somewhat above the ensiform cartilage.

Both sounds are discerned at each of these points. But the

same sound varies in different situations. The first sound over the left ventricle near the apex of the heart is dull, heavy, and prolonged ; that over the right ventricle is clearer, shorter, and of higher pitch. The second sound heard there presents no constant and appreciable variance from that over the left ventricle ; yet it is less ringing and distinct than the second sound of the pulmonary artery and aorta. Even these two are not precisely alike. The second sound of the latter, when compared with that of the former, is found to be sharper and more accentuated. The first sound, however, does not differ materially from that of the pulmonary artery. But the first sound of both does differ most materially from that over the ventricles. Compared with the first sound over the right ventricle, the first sound of the pulmonary artery is much duller, more indistinct and like a vibration, and not of so high a pitch. Compared with the first sound at the apex, the first sound of the aorta lacks the weighty, prolonged character which belongs to the ventricular sound.

These statements are based on a series of observations made, some with an ordinary, some with a double, stethoscope. They certainly seem to favor the view of Skoda, that the first sound, as heard over the great vessels, is not merely a transmitted sound, but is one which is partly, if not entirely, generated by the arteries themselves when the blood rushes into them.

The sounds just considered undergo various modifications, both when the heart is affected and when it is free from disease. They may be audible over a larger space of the chest than usual ; they may be changed in character and in rhythm. Their transmission over a larger space is an unimportant sign. They are undoubtedly perceived over a more extended surface when the heart is enlarged, or when the surrounding tissues are condensed ; even in perfect health their range is very diversified.

During a full inspiration, the sounds at the interspace between the second and third costal cartilages on the left side disappear almost entirely, and become faint at the aortic cartilage. The first sound at the apex lessens also very much in distinctness, but it is better heard at a new point of impulse, visible toward the median line and just below the cartilages of the ribs. During a full expiration, the extent over which the heart-sounds are perceived is increased.

The sounds grow in loudness in any functional disturbance of the heart. When the organ is palpitating violently under strong nervous excitement, they may become short and sharp, and sometimes so loud and ringing as to be audible to the by-standers. They are often permanently louder than in health, and are shorter and more clearly defined when the walls of the heart are thinned. This is particularly the case with the *first* sound. When the walls of the heart are thick, the first sound over the hypertrophied portion is apt to be dull and prolonged. The first sound is weakened if the structure of the heart be softened : hence it is feeble in some low fevers, and in fatty degeneration of the organ. It is also less distinct when there is a want of tone in the muscle, or when the mitral and tricuspid valves are thickened.

To determine whether a dull first sound at the apex be due to an injured mitral valve, or to an alteration of the muscular power of the heart, Flint advises to place the stethoscope over the apex of the heart, and then on the outside of the left nipple to isolate the element of impulsion, which unites with the valvular element to form the complex first sound. If there be a marked impulsion over the apex, but if by means of the stethoscope placed to the left we perceive no sound at all which possesses a valvular character, or hear a sound which is but faintly valvular, we infer that the mitral valves are more or less damaged.

The *second* sound is not so liable to be changed as the first. It is rendered somewhat duller by a thickening of the semilunar valves ; on the other hand, it is more ringing when they are thin, and in great functional excitement of the heart, and in altered blood conditions, as in lithæmia or in gout. The sound, indeed, always becomes more distinctly accentuated if the column of blood closes the valves forcibly. This occurs not unfrequently in hypertrophy of the ventricles, especially the left, and in the increased tension of the vessels in contracted kidney ; it affects the second aortic sound. Accentuation of the second sound also takes place where a decided obstruction exists to the passage of blood through the lungs, and in mitral valvular disease. In both the latter conditions it is over the pulmonary artery alone that this accentuated second sound is audible.

Both the sounds are occasionally obscure and seem to arrive at the ear from a distance. This happens when fluid has ac-



cumulated in the pericardium. The sounds may be changed in their relative proportion to each other, and the pauses between them be lengthened or shortened, or else the sounds may intermit from time to time. From this perverted rhythm we do not derive any definite instruction as to the condition causing it. It serves only to show that the heart is acting irregularly, and thus directs our attention to the state of the organ. It may be associated with organic disease or exist without it. The same may be said of reduplication of the sounds of the heart. The second sound is the one which is generally split. Yet both of them may be doubled, or one may be doubled over one part of the heart and not over another; so that four or three sounds are counted to each beat of the pulse. The cause of the reduplication is the want of synchronous action of the two sides of the heart. The direct value for diagnosis of the altered movement is not great; but indirectly it teaches a most important lesson: it tells us that each side of the heart forms its own sounds, and that, to arrive at accurate conclusions, each side has to be separately examined. Yet there is some diagnostic value to be attached to the changed rhythm. Thus, the peculiar alteration of the sounds, which causes us to hear three sounds during the action of the heart, two of them in the diastole, producing the rhythm that has been likened to the gallop of a horse, is often found in contracted kidney and in arterio-sclerosis. It is particularly heard over the mitral and the tricuspid region. Fraentzel \* has noted the frequent occurrence of this cantering rhythm in typhoid fever and in croupous pneumonia, and looks upon it as a sign of grave cardiac weakness; it is also a sign of serious import in chronic Bright's disease.

Such, then, are the modifications which the healthy sounds present. At times we meet with sounds which do not in the least resemble those naturally heard, and which overshadow them or take their place. They are called *murmurs*, and are mainly produced either within the heart or on its surface.

Those murmurs that are *endocardial* have a common quality: they are more or less blowing. Yet the sound is not always of

---

\* *Krankheiten des Herzens*, Berlin, 1889; see also Cuffer and Barbillion, *Arch. Gén. de Méd.*, 1887.

the same character or pitch. It may be low-toned, it may be high-pitched ; it may be soft, it may be harsh ; it may resemble the blowing of a bellows, it may be musical ; it may be filing, or rasping, or sawing. The ingenuity of every listener exerts itself in tracing a similarity to some familiar noise ; but to little practical purpose. These different sounds have not been proved to have a decided significance beyond that of a blowing sound. They teach us nothing certain as to its source. They are, moreover, not at all times the same in the same case, since the heart when excited may emit a sound different from that which it does when it is beating quietly.

A blowing sound originates in the altered relation of the blood to the part over which it moves. This general statement opens the way to the consideration of the specially acting elements, both in the blood and in the heart itself.

Usually a cardiac murmur springs from a change at one of the orifices. This may be either a narrowing or a roughening, which interposes a local obstruction to the flow of the blood ; or it may be an insufficiency to close the opening. In the latter case the blood regurgitates, and a murmur is occasioned by the deviation of the direction of the current and the establishment of another. This subversion of the course of the circulating fluid, added to its increased velocity and force, is the chief source of those temporary blowing sounds not unfrequently perceived when a heart is greatly excited, while both its valvular apparatus and its muscular texture are healthy. But we meet every now and then with instances where none of these causes are present, and where altered blood is the foundation of the murmur.

Thus, to sum up the subject, we find murmurs that depend upon organic change, and murmurs that are unconnected with any structural alteration ; and these inorganic murmurs are due either to an unnatural condition of the blood or to temporarily perverted action of the heart.

The murmurs, however caused, have different effects on the sounds of the heart. They either accompany the sound throughout the whole or a part of its duration, and thus obscure it, or else they take its place and hinder it from being generated. In time of their occurrence they correspond to the contraction or to the dilatation of the heart, and therefore to the first or to the

second sound ; at least, they do so practically. It is true, they may immediately precede or succeed either sound, or fill mainly the intervals of silence between them ; but attention to such minute divisions is irksome, and, for most purposes, unnecessary. In point of fact, it is often difficult enough to say whether the murmur we hear is systolic or diastolic. The readiest method of judging of the time of the production of a murmur is to feel with the finger for the impulse while listening with the stethoscope. The blowing sound which agrees with the beat of the heart is systolic ; the one between the beats is diastolic.

When a murmur is once established, it attends each motion of the heart that can give rise to it ; but it is not always equally perceptible. It may become very faint, or disappear entirely, by the patient changing his position. It is sometimes manifest only when the heart is acting strongly. Indeed, it always requires a certain force and velocity in the passage of the blood to generate a murmur. Yet overaction of the heart may be as destructive of its distinctness as diminished action. This is, however, a matter that, should it be desirable for diagnosis, we can control by the administration of medicines like digitalis, aconite, or veratrum viride, provided their use be not contra-indicated.

A murmur is sometimes heard by the patient himself, or is audible before the ear is placed over the heart. It may be perceived as an abrupt blowing sound, apparently coming out of the mouth. A gentleman, whose mitral valves permitted of regurgitation, was under my charge. When he held his breath and kept his mouth open, he, as well as I, could detect an abrupt blowing sound issuing from the oral cavity. This sound, when the heart's action was at all excited, accompanied regularly each impulse.

Posture exerts a decided effect upon murmurs. A blowing sound distinct in the recumbent position may become very faint or disappear when the patient stands erect ; and the reverse holds good, although less common. The nature of the murmur—whether organic or inorganic—does not seem to influence the readiness with which it is affected by change of posture, though anæmic murmurs are thought to be more intense in the recumbent position.\* Pressure, too, has an influence upon the abnormal

---

\* James H. Hutchinson, Amer. Journ. Med. Sci., April, 1872.

cardiac sound ; it notably augments it, and often raises its pitch. Yet pressing the stethoscope against the chest does not occasion as much alteration in endocardial as it does in pericardial sounds.

A murmur may be obscured by the respiratory sound ; but this is not apt to be a cause of error in diagnosis. It is not nearly so fruitful a source of mistake as considering the natural sounds of the lungs to be blowing sounds in the heart. Certainly the resemblance is often great ; but blunders may be readily avoided by listening to the heart while the patient suspends his breathing.

Having ascertained positively the existence and the time of occurrence of an endocardial murmur, the next thing is to determine its exact seat, and, if possible, its immediate cause. The *seat* of the murmur is judged of by the place of its greatest intensity, and by the relation this bears to one of the four points for the clinical examination of the heart above described. If it be most distinct at or near the apex of the heart, it is produced at the mitral orifice ; if immediately above or at the ensiform cartilage, it is generated in the right ventricle and at the tricuspid opening. If we hear it most plainly at the sternum, somewhat toward its left border on a level with the third intercostal space or even the fourth rib, and with equal or nearly equal distinctness at the second costal cartilage on the right side, we are enabled to decide that it is developed at the origin of the aorta. The pulmonary artery is not often the seat of a murmur. When it is, this is clearly perceptible in the second intercostal space on the left side, and extends, if the valves be diseased, to the junction of the third left cartilage with the sternum ; although we must bear in mind that occasionally in mitral affections the murmur is loudest in the pulmonary area, or, as Naunyn has shown, not exactly over the artery, but rather an inch and a half or more from the left edge of the sternum in the second interspace.

Any of these situations may be the site of a distinct murmur occupying only one sound of the heart, or being produced in both, —one murmur taking place with, the other against, the current of blood. Yet it rarely happens that the murmur is strictly limited to one of these positions : it will mostly extend in various directions from its point of intensity, growing fainter and fainter as this is left. A blowing murmur thus transmitted may drown the natural sounds of the heart at the parts not diseased. But when



one orifice only is affected, we can usually hear the sounds at the other valves. They may be obscured, but still they exist ; and it is a vast aid when they are heard, since they set the limits to the disease. How important is it, then, to examine each portion of the heart separately, as much for the purpose of saying what is not as what is deranged !

If satisfied as to the seat of the murmur, we naturally turn to inquire into its origin. Is it caused by an alteration of the valves ? Is it unconnected with any appreciable change of structure in the heart ? There is nothing in the murmur itself which will tell us positively. As a rule, it is true that a harsh murmur results from organic disease, and a soft murmur is inorganic ; but we judge with much more certainty by the time of the occurrence of the blowing sound and by the accompanying phenomena. A murmur attending the distention of the ventricles shows that the orifices are injured. A systolic murmur may be either organic, or it may indicate simply a change in the state of the blood, or of the force and velocity with which it is circulating. In the latter case, however, the abnormal sound is temporary, and disappears with the excitement. If arising from an impoverished state of the blood, it is generally soft, of low pitch, is perceived over the base of the heart, and is accompanied by a humming sound in the veins of the neck. It may be heard over the right base, or on the left side over the pulmonary artery ; although Balfour maintains that it is not really over the pulmonary artery, but about half an inch or more to the left of the pulmonary area, and is not an arterial, but an auricular sound.

Throughout the consideration of the endocardial murmurs, they have been treated as originating at the seat of the valves. In truth, it is there that they are formed. Still, they are occasionally due to morbid states in the body of the ventricle, or in the auricle. But in either case, then, they are clinical curiosities. As regards the auricles, they yield no appreciable sound in health, nor are they in disease except rarely the source either of sound or of murmur.

A blowing sound is not of necessity limited to the heart : it may be transmitted all over the arterial system. Yet it would be a great mistake to suppose that every murmur heard over the arteries is connected with a disease of the heart. It is often but

the sign of impoverished blood, or a sound dependent upon local roughening or narrowing of the tube. The latter may be temporarily produced by the pressure of a stethoscope,—a fact of which it is well to be aware. It is even stated that pressure over a healthy heart may generate a murmur; but I have never been able to satisfy myself of the truth of this statement. It is certainly incorrect as a general rule, and the murmur depends, when it happens, more likely upon the condition of the blood and the force with which it circulates.

Let us now examine the sounds which originate on the outside of the heart. These *pericardial murmurs* have all a common source: they all result from irregularities on the membrane. Like the pleura, the smooth serous covering of the heart moves noiselessly in health; but when it is roughened by a deposit of any kind, the friction of its surfaces gives rise to a sound which may be single, but which is usually double. The character of this sound is variable. It may be a to-and-fro rubbing murmur, or it may be grazing, or scratching, or creaking, or whistling, or clicking and resembling the valvular sounds. It has but one quality which is constant, and that is its superficiality. By this superficiality; by the strict limitation of the sound to the region of the heart; by its altering from time to time its precise seat; by its greater extent and intensity when the patient bends forward; by its occasional increase, and even change of character, on external pressure; by its following, rather than occurring with, the movements of the heart; and by the sensation of friction which it communicates to the finger,—we know that the sound heard is produced on the surface of the heart. Yet, in spite of this array of points of difference, it is often difficult to distinguish a pericardial from an endocardial murmur.

An error not easy at times to avoid is the failure to discriminate between the presystolic apex murmur, regarded as characteristic of mitral constriction, and a pericardial friction localized near the apex. The only trustworthy points of distinction are that the pericardial sound changes in its quality and loudness, that it is rendered stronger and changed in pitch by pressure exerted with the stethoscope, and that the second sound at the left base is unaltered.

A friction sound is prone to mask the natural sounds of the heart. At times, although heard over the cardiac region, it is not

due to inflammation of the pericardium. The exudation may be on the surface of the pleura adjacent to the pericardium, and the murmurs be caused solely by the movements of the heart, with the rhythm of which they coincide. Sometimes, again, the sound heard in the cardiac region is in reality the rubbing of an inflamed pleura. If any doubt exist, let the patient be told to suspend his breathing. As this is stopped, the pleural sound ceases.

Such is a brief description of the different physical signs met with in examining the heart, both in health and in disease. Their importance for diagnosis it is difficult to overestimate. A knowledge of the physical signs is the solid foundation, without which any structure that may be raised will soon tumble to pieces.

### The General and Local Symptoms of Diseases of the Heart.

It is not easy to say what are and what are not the symptoms that belong to diseases of the heart. There are vital manifestations directing attention to the heart which are not associated with any change in its structure; and most serious changes in its structure may occur without any of these vital manifestations. Yet we often find a significant group of symptoms which accompany an affection of the heart. Some of these attest directly the organ disturbed, such as pain in the cardiac region, and palpitation. Others are the indirect and more remote expressions of its derangement, such as cough, dyspnoea, hemorrhages, dropsy, disorders of the brain and nervous system, engorgement of the abdominal viscera, a peculiar state of the arteries and veins, and the aspect of the face. It is unnecessary to do more than mention some of these, since several have been already described in connection with pulmonary complaints, and there is nothing in the cough or in the shortness of breath by which we can absolutely determine it to be caused by a disease of the heart. The same with respect to the hemorrhage: there is nothing characteristic about it. It simply proves the efforts of the blood-vessels to relieve themselves of the strain which the disturbance in the flow of the blood has put on them. The capillaries and the smaller blood-vessels give way first; partly from the reason just assigned, and partly from the altered state of their coats, a common associate of cardiac disease. These hemorrhages

are prone to happen from the bronchial tubes and the lungs, and the blood is expectorated; but they may also take place directly into the pulmonary tissue, or into or from any part of the body. Their danger is in proportion to the amount, to the importance of the function of the structures into which the blood is effused, and to the possibility of its finding an outlet. The peril is greatest when the blood is poured out into the brain.

**Cardiac Dropsy.**—The dropsy caused by a disease of the heart is met with in different situations: in the cellular tissues, in the peritoneal and pleural cavities, in the pericardium, in the ventricles of the brain and under the arachnoid, in the air-cells of the lungs,—in fact, in any part where fluid can exude, and where there is a space which can receive.

In anasarca dependent upon a cardiac lesion, the dropsical swelling begins about the ankles and feet; hence œdema starting in this situation is regarded as among the surest of the symptoms of a disease of the heart. The accumulation is much influenced by position: the feet are more puffy toward evening, when the patient has been all day in the erect posture, and least so when he gets up in the morning.

What the condition of the heart is that gives rise to dropsy, has been made a matter of much dispute. It has been held to be uniformly connected with dilatation of the right side of the heart. It has been taught to be invariably linked to a valvular affection. Clinical experience shows us that it may or may not exist where these states are present. The dropsy is most constantly found to be associated with an impediment to, or disturbance in, the flow of the venous blood, and therefore with disorder of the right side of the heart, particularly with a dilatation of the cavities. It may be permanent or not. Its extent certainly does not bear a constant relation to the extent of the cardiac disease. It bears a more constant relation to the amount of venous congestion, and to the impoverishment of the blood.

**Derangement of the Circulation.**—Unmistakable evidence of the obstruction to the flow of the blood through the veins is afforded by their prominence in different portions of the body. This is especially manifest in the superficial veins of the neck, which, moreover, when the tricuspid orifice is permanently open, exhibit a distinct pulsation with each beat of the heart.



The turgid condition of the venous system is rendered equally obvious by the livid tinge of the skin and the bluish color of the lip, and by ramifications of fine bluish vessels on the surface. But the arterial system may also be gorged, and we may find the capillaries and the smaller arteries seemingly ready to burst. The conjunctiva is then highly injected, and the cheek has a coarse, red look. This change in the color and appearance of the face, the thickening of the eyelids, and the prominent eye, make up the peculiar physiognomy of a chronic cardiac malady. The state of the larger arteries is very variable, and mainly according to the nature of the disorder. The pulse may be small and tense; it may be full; it may be rebounding; it may be very irregular; and it is often out of all proportion to the forcible action of the heart.

The derangement of the circulation of individual parts manifests itself by special symptoms. It shows itself in the brain by attacks of cerebral congestion; by vertigo; by violent headache, occurring in spells, or, less acute, in dull persistent ache, increased on exertion,—a form especially met with in children. We see evidences of the congestion of the nervous system in the disturbed dreams; in the sudden starting up from sleep; in the irregular action of certain muscles; in the spots which float before the eye. It is possible that the strange sense of insecurity, and the irritability, of which patients afflicted with a cardiac malady complain, are produced by the same cause. At any rate, whether produced thus or not, they are remarkable symptoms. There is no disease which unnerves more than a disease of the heart. Indeed, the mere fear of its presence gives rise to restlessness and gloom, and breeds timidity in those who would look any external danger boldly in the face.

The disordered flow of blood through the abdominal viscera occasions organic changes and a disturbance of the functions of the several organs. Thus, the liver increases in size, or undergoes other alterations which interfere more or less seriously with the elimination of the bile; or the kidneys no longer secrete as in health, but become much engorged and drain off the albumen of the blood; or the spleen sustains textural transformations. These states all tend to give rise to more and more dropsy, and hence to more and more suffering.

The symptoms which point most directly to the heart itself are palpitation and irregularity of action, and pain. These symptoms imply that the function of the organ is disturbed, or that its innervation is in some manner deranged; but they imply nothing more. They are, therefore, common to functional derangement which occurs associated with structural changes in the heart, and to purely functional derangement.

**Cardiac Pain.**—Pain in or over the heart is met with both in acute and in chronic diseases; yet it is not a regular or well-defined symptom of either. When we reflect that the heart may be pinched, may be torn, without exciting any suffering, it will be readily understood why its disorders do not occasion much pain. Indeed, many a case of enormous enlargement of the heart, or of profound textural alteration of its walls or valvular apparatus, is unaccompanied by pain. Still, we meet with instances in which distress at the heart and various uneasy sensations are among the more marked symptoms of a chronic cardiac lesion; and we even find persons complaining of a persistent pain in the heart, which extends to the left side of the neck and arm, in whom this symptom has preceded the signs of a disease of the heart or of its great vessels.

In the acute cardiac affections pain is a not inconstant symptom. Uneasy sensations, not amounting perhaps to absolute pain, are complained of in endocarditis. Actual pain is among the vital manifestations of inflammation of the substance of the heart, and of the pericardium. In the latter disorder it is usually increased by pressure, and is frequently very severe. But no suffering is so harrowing as that which happens in the obscure malady termed *angina pectoris*.

*Angina Pectoris.*—Although the nature of this complaint may be hidden, the symptoms are obvious enough. We do not know what the precise cause of this angina is; but we do know that the disease occasions paroxysms of intolerable anguish. These paroxysms come on suddenly, and pass off as suddenly. Their main feature is an agonizing pain in the præcordia, as if the heart were being firmly grasped by an invisible hand, or as if it were being torn to pieces. The pain is, however, not limited to the cardiac region; it radiates in various directions, shooting to the back, to the neck, and especially down the left arm. But this is not all: worse than the pain are the intense anxiety and the

feeling of impending death. The heart palpitates during the fit ; yet, if we judge by the character of the pulse, its movements are not always materially disturbed. The beat of the artery at the wrist may be small, may be weak, may be irregular, may be accelerated ; but it may also be full, strong, regular, yet not increased in frequency ; very generally the arterial tension is high. Again, there may be a decided difference between the pulses, the left being almost or quite imperceptible.\* The face is generally pale. Difficulty in breathing, contrary to what might be expected, is not a prominent symptom, and is, in fact, often wanting, while sometimes the breathing is irregular and of the Cheyne-Stokes variety. Giddiness, spasmodic seizures, temporary coma, perverted sensibility, occasionally attend or follow the cardiac attack, and so does pericarditis.†

The duration of the fits is as uncertain as are the causes which excite them. They may cease in a few minutes ; they may last upward of an hour. They come on rapidly, without any assignable reason ; they are reproduced by bodily ailment, by exertion, by fatigue, by exposure to cold, or by mental irritation. However provoked, they are always dangerous. The heart may stop beating during the paroxysm. "My life is in the hands of any rascal who chooses to annoy and tease me," was a saying of John Hunter. And in truth, after he had suffered for years from these seizures, his ungovernable temper brought on one in which he expired. It happens sometimes that the second attack follows at a short interval the one by which the disease first declares itself, and proves fatal. Latham‡ narrates the history of two cases of this kind. In one, life ceased in a fortnight after the first seizure ; in the other, in ten days. Nay, it may be cut short even in the midst of the first manifestation of the malady. Such was the death of Arnold of Rugby.§ On the other hand, I have had a patient under my care who for weeks at a time has five or six attacks daily, kept in check, but not wholly averted, by nitrite of amyl ; and in another patient as many as forty occurred in one day.

---

\* Hamilton Osgood, *Amer. Journ. Med. Sci.*, Oct. 1875.

† *Clin. Soc. Transact.*, vol xvii p. 82.

‡ *Lectures on Diseases of the Heart*, vol. ii.

§ Stanley, *Life and Correspondence of Thomas Arnold*.

The immediate conditions on which the symptoms of the attack depend are veiled in obscurity. Whether they be or be not produced by temporary increase of weakness in an already-enfeebled organ; whether a cardiac spasm occur or do not occur; whether the pain and the sensation of approaching death be or be not caused by an acute distention of the heart with blood,—we do not know. All we do know positively is, that the excessive pain abruptly appearing and disappearing points to deranged innervation. Yet we can go a step further; we can say with certainty that angina pectoris is not often an uncomplicated neuralgia. Modern research has taught us that these outbreaks of a cardiac neurosis are frequently linked to some structural change. This structural change, so far as we can now see, is, however, not at all times the same. The list of disorders of the heart and arteries which angina pectoris may accompany is, indeed, very long. There is hardly an affection of the walls or cavities of the heart, scarcely a morbid condition of the arteries that nourish it or spring from it, with which the distressing malady has not been observed to be associated. It has been found as an attendant on ossification of the coronary artery; on every form of valvular disease; on thinning of the parietes of the heart; on their fatty softening; on fungoid growths springing from the apex of the organ.\* It has been thought that combined with all of these states is fatty degeneration, which thus would be at the root of the angina.† Whether this view be correct or not, it is certain that fatty degeneration is more frequently conjoined with angina than is any other organic disease. Yet fatty degeneration occurs often without angina, as does disease of the coronary arteries, and we are thus forced to admit that, however frequent the association, in either instance some unknown element is still the determining cause. During the attack, as Brunton has shown, there is a vaso-motor spasm of the smaller vessels, with a rise in blood-pressure and increased tension in the arteries. Angina pectoris is now very generally ranked among the vaso-motor neuroses. But evidence is not wanting, as Peter's cases prove,‡

---

\* B. Travers, *Medico-Chirurgical Transactions*, vol. xvii.

† Quain, *Medico-Chirurgical Transactions*, vol. xxxiii.

‡ *La Semaine Méd.*, March, 1892.



that neuritis of the cardiac plexus, the neuritis itself being consecutive to aortitis, is the cause of a certain number of cases.

Angina pectoris is easy of recognition. The points to ascertain in diagnosis are, whether it is linked to an organic cause, and to what organic cause, or whether it is a pure neurosis, either primary or reflected. It may be a question whether those *severe pains in the region of the heart*, which occur in feeble anæmic persons after unaccustomed exertion, or which are brought on by the excessive use of tobacco,\* or which happen in rheumatic or gouty subjects, especially while suffering from indigestion, are real angina, or whether they may be separated from this affection. They differ from it, irrespective of being far less violent and less radiating, by the circumstances leading to an attack, and by their constant association with palpitation. *Intercostal neuralgia* with palpitation might be mistaken for angina; but the painful spots in the course of the affected nerve, and the comparatively slight suffering, distinguish it. In truth, it is a complaint seated only in the thoracic walls, and referred by the patient to the heart. *Great irritability of the heart*, attended with faintness, with sensations of sinking, with flushing alternating with pallor, and with pain, due most likely to a neurosis of the cardiac plexus, is discriminated from true angina by the palpitations, and by their connection with pain which never rises to the anguish of angina pectoris. Often, too, this apparent or false angina is found in persons who are hysterical, or are subject to neuralgia, or are laboring under a disorder of one of the abdominal viscera, and is then clearly reflex. It must, however, be admitted that the distinction between true and false angina is one of degree rather than of kind; for the cardiac plexus is precisely the point particularly involved in angina, and it is now generally thought that the disturbance of the heart in this painful malady occurs mainly through the influence of the sympathetic fibres which meet in the plexus.

Another complaint that may be confounded with angina is what may be called *cardiac epilepsy*. In this rare affection intense pain in the region of the heart happens in paroxysms. But unconsciousness, however temporary, occurs also, and the pain

---

\* Beau, *Journal de Médecine et Chirurgie*, July, 1862; Eulenberg, "Angina," in Ziemssen's *Cyclopædia*.

is apt to follow rather than to precede the unconsciousness. Yet it may outlast it, and become associated with twitching of the muscles of the face and with other spasmodic movements. These, the unconsciousness, and the time at which the pain happens, distinguish the malady from those instances of angina in which, owing to the severity of the pain, the patient passes into a protracted faint.

**Palpitation.**—This arises in various diseases of the heart. It happens at the beginning of acute affections; it is an unfailing accompaniment of some chronic lesions. It is especially distressing when the cavities are dilated and the walls of the organ thinned. But it bears no positive relation to any special cardiac malady, and is therefore not diagnostic of any. So, too, with irregular rhythm of the heart's action, with which palpitation is in truth often combined. It tells us nothing more than that the regular movements of the heart are disarranged. Frequently this disarrangement is due to a serious change in the valves or in the muscular structure. But palpitation, with or without irregular rhythm, may take place in a perfectly sound heart,—sound, at least, so far as our means of investigation enable us to determine.

Often the pulsations of the heart become stronger, more extensive, and more perceptible, from mere nervous excitement. But it is not necessary to detail the symptoms of a purely nervous palpitation. Every one has experienced them. Every one knows that there is a feeling of slight constriction about the chest, with a hurried breathing, and a strange sensation as if the heart were leaping from its place. Every one is also aware that the organ is felt thumping against the walls of the chest, and with a force which shakes them. The popular notion, that the heart is the seat of the emotions, is based on these striking evidences of its disturbed action, and poets have seized upon and delineated with accuracy some of the even more strictly physical phenomena of the extended impulse under strong nervous excitement.\*

---

\* Thus, Shakespeare, in the "Rape of Lucrece:"

"His hand, that yet remains upon her breast  
(Rude ram to batter such an ivory wall!),  
May feel her heart, poor citizen, distressed,  
Wounding itself to death, rise up and fall,  
Beating her bulk, that his hand shakes withal."

But, apart from the increase of the beat by mere temporary agitation, a heart may act overfrequently and overstrongly and its action become sensible to the person, in other words, it may palpitate, from some more unremitting excitement dependent upon perverted innervation. This is the main cause of the altered impulse of the heart in the so-called functional disorders.

Persistent rapidity of cardiac action, or *tachycardia*, may happen without obvious cause in persons apparently healthy. It is very common in irritable hearts and in exophthalmic goitre. Spender\* has called attention to its occurrence among the earlier signs of rheumatoid arthritis. The extreme frequency of the action of the heart is in some instances remarkable. I have known it to beat over two hundred times in the minute. The disorder may occur in paroxysms, described as "cardiac nerve storms" by H. C. Wood.† Great rapidity may be joined to a condition in which the two sounds are precisely alike, and the pauses of equal length. This foetal rhythm, or *embryo-cardia*, is a sign of heart weakness.

On the other hand, the deranged innervation may lead to very retarded movement, and the heart beat less than thirty times in the minute. We may find this slow action, *bradycardia*, both in functional and in organic maladies, though it is most likely that the nerve-centres are in both affected in the same way.‡ *Bradycardia* is often associated with atheroma of the aorta or of the coronary arteries. It is also met with in a number of instances of fatty heart and in old-standing valvular disease. Its association with jaundice, with feeble heart action during convalescence from fevers, with epilepsy, and with melancholia, is well known.

### FUNCTIONAL DISORDERS OF THE HEART.

It has just been stated that the direct symptoms of a cardiac disorder—pain, palpitation, irregular action—are met with when no recognizable structural change has taken place. Under such circumstances the affection of the heart is termed functional, and its symptoms are those already mentioned, variously combined, sometimes the one predominating, sometimes the other. These

---

\* On Osteo-Arthritis, London, 1889.

† University Medical Magazine, March, 1891.

‡ See a very interesting analysis of ninety-one cases, by Prentiss, Transact. Assoc. Amer. Phys., vol. iv., 1889.

functional disorders are very much more frequent than the organic. They are, for the most part, produced by direct excitement of the heart, or by its being sympathetically disturbed by some source of irritation existing remote from it, or in the system at large. The symptoms may be said to constitute the disease.

### Functional Disorders characterized by Palpitation, associated or not with Change of Rhythm.

We have already briefly mentioned the causes of augmented action which are associated with organic changes, and those which occasion temporary disturbance of the heart. A more lasting form of palpitation is engendered when the organ is kept constantly excited by a deranged condition of some viscus remote from it; by the use of stimulating substances; or by some general morbid states. Thus, a disordered stomach or liver leads to a reflex disturbance of the heart, which ceases if the disorder of the stomach or liver be remedied. In gouty, lithæmic, and rheumatic persons the heart frequently pulsates with increased quickness, and sometimes with marked irregularity. Special articles of diet, especially tea or coffee, produce palpitation; so does the inordinate use of tobacco. Immoderate dancing, masturbation, and excessive sexual indulgence, but particularly masturbation, are prolific sources of continued palpitation. We see also those affected with it who, addicted to laborious studies, grudge themselves time for food, sleep, and exercise. Women who are hysterical, or whose uterine functions are disordered, suffer, or fancy that they suffer, from palpitation. So do so-called nervous people invariably complain of the beating at the heart.

In those whose blood is much impoverished, the palpitations are often severe and constant, and they imagine themselves to be laboring under an incurable disease. There is, indeed, from the strong resemblance to an organic affection, apparent cause for alarm. The heart strikes sharply and abruptly against the walls of the chest; its action is frequent; the breathing becomes hurried on the slightest exertion. Nay, even the physical signs may be those of a structural lesion. The altered blood gives rise to a blowing sound in the heart, which is transmitted into the carotid and subclavian arteries. The difficulty of diagnosis is at



times considerable. The age; the sex; the anæmic look; the presence of a continuous humming sound in the veins of the neck; the strict synchronism of the murmur with the impulse; its want of harshness; its seat commonly at the base of the heart,—furnish a clue to the nature of the case. Still, we have often to judge as much or more by the absence of the signs of cardiac enlargement, and of impediment to the flow of the blood, whether the heart be affected in its valvular apparatus, or whether it be simply functionally disturbed and circulating watery blood.

A troublesome kind of palpitation is that attended with marked *irregularity* of the action of the heart, displaying itself by the beat being now slow, now fast, or occasionally intermitting. Sufferers from lithæmia or gout, or old persons with feeble digestion, are particularly liable to it. This form of palpitation is not without danger. It is very prone to be associated with an alteration in the structure of the heart, such as flabbiness of the walls, which may not be sufficient to yield any distinctive physical signs, but which is nevertheless sufficient to be a source of apprehension.

Some who experience fits of palpitation faint away during them. But the almost complete suspension of the movements of the heart which characterizes an attack of syncope has no definite connection with any form of palpitation, nor, indeed, with any form of cardiac disorder, organic or functional. In those who are subject to attacks of palpitation or to irregular action of the heart, the organ may finally become enlarged.

A peculiar irregular action of the heart has been much discussed under the name of *hemisystole*. Leyden pointed out that there were cases in which with every two beats of the heart only one beat of the pulse was felt, and attributed this to the right ventricle contracting alternately with the left. Different explanations have been given of the fact, but the observations of Riegel and Lachmann, while they do not strictly confirm the alternate action of the ventricles as the cause of the phenomenon, point to irregular contraction of the muscles of the heart as the cause.\*

We sometimes meet with a singular form of functional disturbance of the heart which leads to textural changes, and to which

---

\* Virchow's Archiv, Bd. xlv.; Deutsches Arch. f. Klin. Med., Bd. xxvii. p. 393.

Graves called particular attention. It consists in a long-continued excitement of the organ, as evidenced by its increased force and rapid and irregular action, which is followed by a swelling of the thyroid gland, pulsation of the arteries of the neck, and prominence of the eyeballs. This disease, *exophthalmic goitre*, is most commonly observed in women, and connected with hysteria, neuralgia, or uterine disturbance; it has in some instances an evident origin in shock. It is generally considered to be owing to an affection of the cervical sympathetic nerve. But its cause is far from certain. There are those who hold it to be a neurosis of the nerve-centres, especially of the vagus centre; and the detection of ptomaines in the urine is thought to be a proof that this apparent neuro-cardial malady is really consequent upon secondary disturbance of the nervous system due to poisonous products from the thyroid.

The most characteristic sign, the greatly accelerated heart's action, varies much in extent. All the signs may remit or may become aggravated from time to time, and especially during a severe attack of palpitation. The turgescence of the thyroid gland arises quite independently of the usual exciting causes of bronchocele. It is accompanied by a pulsating thrill similar to that of an aneurismal varix, and by a distinct throb. At an advanced period of the complaint, these signs subside, and the gland becomes more solid. Indeed, the whole affection may disappear, and the gland, the eyes, the beat of the carotids, the action of the heart, may be all brought back to a normal condition. On the other hand, hypertrophy and dilatation may result from the cardiac palpitations. And the malady may be noticed in association with valvular disease, under circumstances which make it a question whether this has followed it or is a mere concomitant.

The protrusion of the eyeball is often combined with a symptom that Graefe particularly observed,—a want of agreement between the movement of the lid and the raising or depressing of the glance. The spasm of the elevator of the upper eyelid is held by Abadie \* to be pathognomonic. Another symptom of importance is trembling of the hands. This tremor, Charcot points out, affects the whole hand, but not the individual fingers. There

---

\* La France Médicale, vol. ii., 1881.

is also, as Charcot shows, greatly-lessened bodily resistance to the galvanic current; but this sign is not of much value, as Cardew \* has found the electric resistance to diminish greatly when the skin is moist. Other symptoms are cramps, usually at night, epistaxis, œdema of the legs and eyelids, lessened respiratory expansion, moderate elevation of temperature, sensation of heat, increased sweating, paroxysmal attacks of diarrhœa, atony of the large intestine, intermittent swelling and pain in the joints, pigmentation, bulimia without gain in flesh, glycosuria, migraine, rheumatic symptoms, and mental derangement. All the physical manifestations of the disease are double-sided; and this, with the unchanged state of the pupils, serves to distinguish it from those rare cases described by Eulenberg,† where a thyroid growth pressing on the sympathetic on one side produces symptoms of exophthalmic goitre, including the palpitations.

In the distinction from ordinary goitre, the absence of eye and heart symptoms is of most value. There is also no murmur heard over the enlarged thyroid gland; whereas in Graves' disease a continuous murmur there is most common, and is, indeed, looked upon by Guttman as of the greatest diagnostic importance, especially aiding us in those doubtful cases in which the exophthalmos is wanting. My own experience confirms this statement.

There is another form of functional disorder of the heart, so peculiar as to demand a special notice. It is the curious cardiac malady of which we saw so many examples in soldiers during our civil war, to which I gave the name of "*irritable heart*," and which we also find occurring in private life. Its main symptoms are habitual frequency of the action of the heart, constantly-recurring attacks of palpitation, and pain referred to the lower portion of the præcordial region. The palpitations come on chiefly during exercise, but may also take place when the patient is quiet, and in many cases happen most often, or indeed entirely, at night, thus interfering with sleep. Those who are subject to the disorder complain much of headache and of dizziness, and especially of being thus affected when suffering from palpitation. The pain

---

\* Lancet, Feb. 1891.

† Ziemssen's Cyclopædia.

is generally dull and constant, but is often also described as shooting, and as taking place only in paroxysms. Its chief seat is near the apex, and it is combined commonly with excessive cutaneous sensibility. Often there is pain nowhere else in the body; but in some instances the cardiac distress is associated with pain in the back, which itself is not unusually connected with the excretion of oxalate of lime by the kidneys.

The action of the heart is very rapid, and in many instances its rhythm is irregular. The impulse is slightly extended, but not forcible, like that of hypertrophy: it is rather abrupt and jerky. As a rule, to which I have met with but few exceptions, the sounds of the heart are modified as follows: the first sound is short, sometimes sharp, resembling the second sound; at other times it is extremely deficient and hardly recognizable; the distinctness of the second sound is much heightened. We either hear no murmurs in the heart or in the neck, or they are inconstant. The area of percussion dulness does not appear to be augmented. The pulse is almost always easily compressible; it may or may not share the character of the impulse. It is usually very much influenced by position, falling rapidly twenty beats or more when the erect posture is exchanged for the recumbent. The increased frequency of beat is not connected with increased frequency of respiration, for often with a pulse of one hundred the respirations scarcely exceed twenty in the minute. The disorder is very obstinate, and improvement comes but slowly.

The cause of the morbid cardiac impressibility is difficult to ascertain. It seems in many instances to have followed fatiguing marches; in some it occurred after fevers or diarrhoea; it was not connected with scurvy, or with the abuse of tobacco. That it was not due to anæmia, was proved by the general aspect of the men, which was often that of ruddy health. Similar conditions of the heart occur from excessive dancing, excessive smoking, and certain occupations, such as glass-blowing. For a fuller consideration of the subject I refer to observations elsewhere detailed.\*

---

\* Medical Memoirs of the U. S. Sanitary Commission, 1867; American Journal of the Medical Sciences, January, 1871; and the Third Toner Lecture, Smithsonian Institution, 1874, "On Strain and Overaction of the Heart," where also the forms of irritable heart occurring in civil life are described.



Yet another form of functional cardiac disorder is the one which I have described under the name of *cardiac asthenia*, or heart exhaustion. It shows essentially the signs of a weak heart, and follows long-continued worry and overwork. There is rapidity of cardiac movement with very feeble action, and a great tendency to faintness. The breathing is singularly undisturbed. The impulse of the heart is weak, the first sound short, valvular, the capillary circulation defective. The duration of the cases is a long one, and recovery comes but gradually. In the cases that are not purely nervous, but in which the heart-muscle is enfeebled, shortness of breath and functional dynamic apex murmurs are often noticed.\*

These, then, are the principal varieties of functional disorder of the heart. It is hardly necessary again to state that the physical signs present the most certain, if not the only, means of distinguishing the functional from the structural affection. They show us that neither the size of the organ, nor its sounds, with the exceptions above mentioned, are materially different from what they are in health.

The irritable heart just described, as indeed other forms of functional heart disorder, may pass into organic cardiac disease by the constant overaction of the heart. And *overaction* or *strain* may also, as I have proved in the publications just referred to, lead to valvular affection, sometimes by preceding hypertrophy, at other times by a slow process of inflammation or disorganization engendered at or near the seat of the valve. Of this I published several instances in the "Memoirs of the Sanitary Commission." Others have been brought forward by Allbutt † which happened among persons engaged in vocations requiring sustained and oft-repeated muscular effort,—such as lifters, smiths, sawyers. And in his elaborate monograph Seitz ‡ has detailed several fatal cases in which the symptoms of a fatigued heart, due to strain, were followed by extensive dilatation without valvular disease. Leyden, too, has added to our accurate knowledge of the subject.§

---

\* Amer. Journ. Med. Sci., April, 1894.

† St. George's Hospital Reports, 1872.

‡ Die Ueberanstrengung des Herzens, 1875.

§ Die Herzkrankheiten in Folge von Ueberanstrengung, Berlin, 1886.

## ORGANIC DISEASES OF THE HEART.

Organic diseases of the heart may be classified as follows :

## ORGANIC DISEASES OF THE HEART.

Diseases affecting the walls of the heart, and mostly changing the size of the cavities.	{	Hypertrophy. Dilatation. Atrophy.
	{	Fatty degeneration. Parenchymatous degeneration. Fibroid heart, cardio-sclerosis, etc. Malformations.
Diseases affecting chiefly the walls alone.	{	Rupture of the heart. Injuries and wounds. Aneurism of the heart. New growths and parasites.
Inflammations.....	{ of membranes. of muscular structure.	{ Endocarditis. Pericarditis. Myocarditis (Carditis).
Diseases of the valvular apparatus.....	{	Valvular diseases.
	{	Chronic pericarditis. Hydropericardium. Hæmopericardium.
Diseases affecting the pericardium.....	{	Pneumo-hydropericardium. New formations on pericardium : cancer, tubercle, etc.
	{	Abnormal positions. Closure of openings of right heart.
Congenital diseases.....	{	Opening between the ventricles. Narrowing and closure of pul- monary artery, etc.

These are the organic diseases of the heart, save the rarest. But let us study the cardiac maladies according to their symptoms and signs rather than according to their anatomical classification.

**Acute Diseases presenting Pain in the Cardiac Region; the Symptoms of a Disturbed Circulation; and a Change in the Sounds of the Heart, or their Replacement by Murmurs.**

All the acute affections of the heart come under this head. In all, the sounds are either changed in their character or are replaced by murmurs. This is certainly true of the only acute diseases of which we have an accurate knowledge,—endocarditis and peri-

carditis. All the acute disorders give rise, further, to more or less pain, and to anxiety of expression ; in all there is fever ; all are prone to occur in connection with other morbid conditions, and especially with a contaminated state of the blood. In all, moreover, the symptoms of a disturbed circulation are met with : palpitation, irregular action of the heart, deranged flow of blood through the capillaries of different organs, and a tendency to dropsical accumulations. That these symptoms are not so clearly defined as in some of the chronic cardiac maladies, is owing to the shorter time the complaint lasts.

**Acute Endocarditis.**—Acute inflammation of the lining membrane of the heart arises from exposure to cold, or without any cause being discoverable. It sometimes results from violent efforts, or from blows or other injuries to the chest. It is often connected with an acute infective process or a vitiated condition of the blood, as in puerperal fever, in pneumonia, in chorea, in cancer, in scarlet fever, in pyæmia, in Bright's disease, or in diabetes. But its most frequent association is with acute articular rheumatism.

The chief source of danger in endocarditis is the tendency the inflammation has to limit itself. It is confined to, or is most strikingly developed at, a part which bears least of all any impairment,—at the valves,—and often leaves behind it some permanent disorganization of their delicate structure. But it does not generally affect the entire valvular apparatus: that of the left side is usually alone the seat of disease. What morbid anatomy thus teaches, explains the occurrence and situation of the principal sign by which endocarditis is recognized. The roughness of the surface over which the blood flows, or the lymph deposited on or in the neighborhood of the valves, interfering with their function, occasions a distinct murmur, which is mostly confined to the mitral and aortic openings, and may be preceded by an altered character of the first sound.

Besides this blowing sound, there are other signs worthy of note. It is true, they do not form so leading a feature of the disease ; still, they aid in its correct appreciation. The excited heart beats with augmented force, and sometimes with great irregularity, as the not unusual doubling of the second sound at the base proves. The size of the organ is not notably increased, except in those cases in which its cavities are choked with blood

or fibrin-clots. The pulse corresponds to the action of the heart; yet not so closely as might be expected. It is, for the most part, frequent and strong, and rather forcible at first; sometimes it is small and frequent. It becomes irregular, one beat being strong, the next weak, if the circulation through the heart be seriously obstructed. But it may be feeble while the heart is thumping with violence against the walls of the chest. Occasionally at the onset of the attack it has been observed to be slower than natural.

The general symptoms are not uniform. There is usually a sense of uneasiness around the heart, with a fever showing a temperature ranging from  $101^{\circ}$  to  $103^{\circ}$ , a short cough, palpitation and some irregularity of cardiac action, difficulty of breathing, and anxiety depicted on the countenance. To these are not uncommonly added turgescence of the face, headache, slight delirium, gastric irritability, diarrhœa, and rigors, followed by sensations of heat. Pain in the heart is rare, and is not likely to happen unless the pericardium or the muscular walls be implicated. In some cases an eruption of subcutaneous fibrous nodules occurs, especially in the rheumatic endocarditis of children.

Now, where these symptoms are present; where they manifest themselves in one whose system is in a state in which endocarditis is apt to take place; and where they are accompanied by a blowing sound recently and rather suddenly developed,—we are certain that inflammation is working its changes in the lining membrane of the heart. Yet some circumspection is requisite before arriving at this conclusion, and before the patient is subjected to energetic treatment with the view of saving him from the supposed damage which his heart is about to undergo. A murmur may be attended with febrile signs and not be dependent upon acute endocarditis. The sound may be of organic origin; or it may be engendered in the course of an idiopathic fever, and the lining membrane of the heart be unaltered.

In the first instance the murmur is *old*, and results from some chronic injury to the valve, the attending fever being an accidental complication. Here is undoubtedly a difficult case for diagnosis. We see the patient for the first time; he has fever; his heart is acting strongly; a distinct blowing sound is perceived over it. How are we to tell that his complaint is not acute endocarditis? We have no absolute means of deciding that it is not. Yet by



careful inquiry we can usually come to a knowledge of the truth. If the patient do not recollect to have suffered previously from dyspnœa, palpitation, or other signs of an affection of the heart; if the cardiac excitement and irritation be well defined; if the face denote distress; if the accompanying symptoms indicate a state that is prone to be complicated with endocardial inflammation,—it is this disease under which he is laboring. I may add another and very important element of distinction deduced from the study of the blowing sound, to wit, that the murmur of endocarditis is not so rough, is not often heard during the distention of the heart, and may be changeable in its seat, which an old-standing murmur never is. Besides, it is not associated with those signs of enlargement which are invariably found when the valves have been for any length of time affected, unless the acute inflammation occur in a heart the valves of which have been previously spoiled. Under such circumstances, we can only conjecture what is going on within the organ from its increased excitement, and, if I may take my own experience as the general rule, from the character of the blowing sound being altered. It is rendered often less distinct, nay, it is even entirely muffled, by the products of the recent inflammation.

But how are we to distinguish between the soft murmur arising in the *course of fevers*, and that resulting from effused lymph? It, too, is not rough. It, too, happens with the impulse. It, too, is preceded, as some cases of endocarditis are, by a lengthening of the first sound. Here is assuredly a strong resemblance; yet by no means an identity. The blowing sound in fevers does not exist until the blood is profoundly altered. In endocarditis it takes place almost as soon as the disease begins,—certainly as soon as we are able to recognize positively its beginning. The heart in fevers may be softened, but it is not so directly disturbed in its action. We do not find those symptoms, local as well as general, which show that the circulation is obstructed. The blowing sound is rarely at the apex, but more over the body of the heart. To this some weight may be attached, since the murmur of endocarditis is very apt to be heard at the apex. But to no fact ought as much weight to be attached as to the one first mentioned, that the murmur takes place early and not late in the disease.

Throughout this description of endocarditis, only simple, un-

complicated cases have been kept in view ; yet it is not often that the malady is seen in so pure a type. It is more generally accompanied by the friction sounds and other signs of acute pericarditis, and by the swollen joints, the painful movements, the acid perspirations, of acute rheumatism ; or by the characteristic appearances on the skin of erythema marginatum ; or by tonsillitis ; or by the kidney symptoms of Bright's disease ; or by the evidences of chorea, or of gonorrhœa, pyæmia, or septicæmia.

Nor is a murmur in endocarditis invariable. If the question be asked, "Can endocarditis occur without a blowing sound?" it must be answered in the affirmative. When the seat of the inflammation is not near the valves, no murmur is generated. There may be also none if no vegetations exist on the valves, and perhaps in states of the exudation with which we are at present unacquainted. We cannot, under such circumstances, detect an attack of endocarditis. Yet it may be even then strongly suspected to be present if great excitement or irregularity of the heart manifest itself in a person who is laboring under a disease which predisposes to endocardial inflammation, such as rheumatism. Cases of this nature are, however, exceptional. They do not happen sufficiently often to invalidate the statement that the development of a murmur is the sign indicative of endocarditis. Still, they happen sufficiently often to impress upon us that our knowledge of endocarditis is not complete.

The clinical study of endocarditis is, in truth, a comparatively recent study. There are some points about it which are as yet unknown, and others which have not been long cleared up. To this class belong the interesting researches on the formation of clots of fibrin in the heart, and on the effects produced when they or the vegetations which stud the valves are washed into the circulation. The *formation of clots in the cardiac cavities*, if at all extensive, announces itself by a sudden appearance or a sudden augmentation of the symptoms of obstructed circulation : the skin is cold, and the surface may be bluish ; there is a struggle for breath, the pulse is frequent and feeble, the action of the heart becomes exceedingly irregular, its sounds are indistinct, or a more or less distinct murmur is heard, and the extent of the præcordial percussion dulness is somewhat increased. Great anxiety of countenance, nausea, vomiting, excitement of the nervous system and

delirium, turgid veins in the neck, and fits of fainting, are also among the manifestations of the clogged flow of blood through the heart. Yet these phenomena are not absolutely distinctive; for Walshe records that the effects of a rupture of a sigmoid valve or of a tendinous cord, during the acute endocardial disease, will give rise to symptoms exactly similar to the obstruction of the circulation resulting from polypoid concretions in the heart.

Portions of the clots, or of the vegetations on the valves, are sometimes washed into the current, and the *embolism* occasions symptoms which, before we were aware of the damages to which the detached masses may give rise, appeared inexplicable. But now—when we see the circulation speedily diminished or arrested in a limb, and the limb becoming painful, swelling, or beginning to mortify; when we find that the flow of the blood through the brain has become suddenly disturbed, and the muscles of one side drop paralyzed; when the difficult breathing becomes rapidly still more difficult, while there are no signs of a superadded affection of the lung, nay, while the power fully to expand the lungs remains unimpaired, or while an effusion of fluid into the air-vesicles follows the dyspnœa—we know what has happened: we know that a broken-off piece of fibrin has been driven into the artery of the limb, or into the brain, or into the branches of the pulmonary artery, and, being too large to go any farther, has stuck fast, and has given rise to all these sudden and sad consequences. Sad indeed they are; for, even if the plugs do not lead to an immediately fatal result, they lay the groundwork for structural alterations in any tissue in which they become impacted.

But let it not be understood that the detachment of vegetations from the valves, or of fragments of clot formed in the cavities of the heart, happens in endocarditis only. Pieces may be separated from valves that are in a state of so-called ossification. And the blood *in the heart* may clot from any interference with the current, from heart palsy, or from changes in the vital fluid wholly unconnected with inflammation. But when it coagulates, from whatever cause, the symptoms are the same as those just described. A murmur, too, is not uncommonly produced, which is not distinguishable from that due to endocardial inflammation, but which is not of long duration, since death generally follows the impediment in the heart in a few days at farthest.

*Inflammation of the aorta* may occasion many of the symptoms of acute endocarditis ; at all events, it may do so when the upper part of the aorta is implicated. But it cannot be said that it is a condition which may be discriminated with certainty. The most significant signs are hurried respiration, a sharp, rapid pulse, tumultuous action of the heart, pain in the præcordial region, often greatly increased by movements, and also felt along the course of the spine, burning pain referred to the sternum, great anxiety. The history of the case points to gout, alcoholism, syphilis, or malaria. There may be paroxysms of pain such as occur in angina, and a loud systolic blowing sound. When the abdominal aorta is affected, we notice strong local pulsation, and a marked murmur will be heard with greatest distinctness at or near the seat of the inflammation. In some cases of aortitis, Bright \* observed an extremely high degree of morbid sensibility over all parts of the body, which caused the patient to scream with pain when his wrists were merely touched. The disorder is most apt to happen in cachectic persons ; and it has been repeatedly observed in those attacked with erysipelas, or after operations and injuries.†

Dissimilar causes may lead to different sites of endocardial inflammation. Thus, puerperal endocarditis is apt to localize itself in the right heart. It has pulmonary complications, and the progress of the disease is often slow ; it may last several months.‡

There is a form of endocarditis which may be here briefly mentioned,—*ulcerative endocarditis*. It is not common in this country, although I have seen a number of instances. It occurs mostly in connection with low forms of rheumatism or with blood-poisoning, and the symptoms of this, or of pyæmia or a low septic fever, are apparently the prominent features of the case, or it may happen subsequently to pneumonia.§ The ulceration perforates the valves, and may extend into the muscular structure of the heart ; pneumonia or pleurisy, embolic formations, and infarcts and metastatic abscesses in various parts of the body are among the common attendants. The perilous affection shows an endocarditis with the ordinary physical signs developing amidst

---

\* Guy's Hospital Reports, vol. i.

† Chevers, *ib.*, vol. vi., and 2d Series, vol. i. ; Osler, Gulstonian Lectures.

‡ Luzet et Ettlinger, Arch. Gén. de Méd., Jan. 1891.

§ Arch. de Physiol., August, 1886.



the symptoms of profound blood-poisoning and prostration, although these physical signs may be masked by a pericardial complication. Marked and recurring chills, like those of malarial fever, but coming on irregularly; a temperature of  $105^{\circ}$  to  $107^{\circ}$ ; an extremely rapid pulse, becoming suddenly much slower, though very irregular; profuse sweats; vertigo; delirium followed by stupor; dry tongue; vomiting and diarrhoea; jaundice; tenderness over liver and spleen; and scanty, albuminous urine,—are among the prominent features of the malady. As regards the thoracic symptoms, there may be oppression, dyspnoea, and pain, as ordinarily in endocarditis, yet these symptoms may be wholly wanting. In some instances a peculiar diffused rose rash, here and there mixed with papules and spots of ecchymosis, is noticed. By some, ulcerative endocarditis is looked upon as diphtheritic; indeed, when it has happened during puerperal fever diphtheritic exudations have been found on the mucous membrane of the vagina and uterus. It is certain that micro-organisms are constantly present, generally streptococci and staphylococci, and are found not only in the heart, but also in the infarcts in the spleen and liver which are common in the affection. Death is the common ending,—either from gradual exhaustion, or suddenly by the tearing away of the injured valves.

The disease is extremely rare in children. It is more often mistaken for typhoid fever than for any other disease. But it is also mistaken for typhoid pneumonia, for cerebro-spinal fever, and for hemorrhagic smallpox. When ulcerative endocarditis happens in connection with malarial poisoning, a not infrequent association in Africa, its seat of predilection is in the aortic valves.\* The most common type of the disease is the typhoid type. The malignant endocarditis may become engrafted on a chronic valve lesion. Its clinical association with a suppurative wound or puerperal disease is common. The cardiac symptoms may be very obscure, and the occurrence of embolism during a febrile process be the first sign to explain their meaning. Rigors are common, and are the cause of malignant endocarditis being frequently mistaken for malarial fevers. High fever is the rule, and is an important element in the diagnosis. But I have met with in-

---

\* Lancereaux, Arch. Gén. de Méd., April, 1881.

stances, proved such by the autopsy, in which fever was almost absent.

**Acute Pericarditis.**—Acute inflammation of the serous membrane of the exterior of the heart is very similar to that of its interior. It is developed under the same circumstances. It is found in rheumatism, in gout, in Bright's disease, in scurvy, in alcoholism, in scarlet fever, in septic processes, or as an extension of inflammation from pleuro-pneumonia. The pericardial malady exhibits the same frequent association with rheumatism as the endocardial malady; it presents the same symptoms. Nature has not, indeed, drawn a very strict line of demarcation between the two diseases. When one exists, the other is very apt to attend it. Yet we do meet with endocarditis without pericarditis, and more often still with pericarditis without endocarditis.

The anatomical effects of inflammation of the pericardium are like those of acute endocarditis, and resemble still more closely those which inflammation of the adjoining serous membrane—the pleura—occasions. The pericardium becomes injected and dry; plastic lymph accumulates on its surfaces, and especially on the surface which fits tightly around the heart. This stage of the disease corresponds to the dry stage of acute pleurisy. It may have the same termination by the two roughened surfaces adhering. But it is often followed by a stage similar to that of pleural effusion. The bag in which the heart lies is filled with fluid; the effusion may remain stationary or be absorbed, and the rugged portions of the membrane be placed again in apposition.

From a knowledge of the anatomical changes, the physical signs may be foretold. It is obvious that there must be at first a friction sound; that then the fluid which distends the pericardium will increase the area of percussion dulness over the heart, and prevent the sounds and the impulse from being distinctly perceived. Yet the friction sound is not always the same in extent or in character, because the deposited lymph is not always the same in extent or in character. The sound is like the crumpling of parchment, or the creaking of new leather, or it is grazing, or like a series of irregular clicks. It is a single or it is a double sound, and is prone to mask the natural sounds of the heart. But these are all points which have been already described: we shall merely add that when the friction develops itself under our

observation, and with signs of excitement of the heart, it is as distinctive of inflammation of the pericardium as a recent blowing sound is, under the same circumstances, distinctive of inflammation of the endocardium. When the pericardial effusion takes place, it ceases ; but only gradually, and not always completely ; and in any

FIG. 45.




Illustration of the position of the heart in pericarditis, and of the distention of the pericardium with fluid. The heart-sounds are indistinct, except above the effusion ; the impulse is feeble. The extent and shape of the percussion dulness may be judged of by the appearance of the distended sac.

case it is not uncommon for the ear still to recognize the murmur at the base of the heart and around the origin of the great vessels.

The percussion dulness due to the effusion is generally considerable ; and its contour is characteristic. As the fluid gravitates to the lower portion of the sac, this distends, of necessity, more than the part where the pericardium adheres to the vessels. The consequence is that the dulness, when the patient is in the erect posture, is pyramidal ; when he lies on his back, or changes from side to side, the outline of the flat sound is somewhat altered. Rotch,\* in an

\* Boston Med. and Surg. Journ., 1878, vol xcix. ; also article " Diseases of the Pericardium," in Keating's Cyclopædia of the Diseases of Children, vol. ii.

elaborate inquiry into the matter, points to the dulness in the fifth intercostal space to the right of the sternum as occurring even in small effusions, and as an available diagnostic sign ; and Roberts,\* in his excellent monograph, speaks of the valuable aid afforded by it to surgeons about to tap the pericardium. Another significant sign connected with the dulness is that, as Bamberger has taught us, an area of dulness near the angle of the scapula which coexists with bronchial breathing and increased fremitus, and which is perceived when the patient is erect, is greatly influenced by position. It disappears, and with it the other signs mentioned, as he leans forward, to return as the erect posture is resumed.

In cases of considerable effusion, the intercostal spaces of the cardiac region widen, the eye recognizes a distinct bulging, and the dulness on percussion reaches far upward, to the second, or even to the first, rib. Within the space of dulness is sometimes seen an irregular, wavy motion ; and what the eye detects the hand feels. But no movements, or only slight movements, may be perceptible in the præcordia. The heart, with its point pushed upward and outward by the accumulating liquid, has to struggle to reach the walls of the chest. Its contractions are irregular ; its impulse is feeble, or all appreciable impulse has ceased. The sounds heard through the mass of fluid seem distant and muffled. Yet the second sound over the upper part of the sternum and at the base of the heart retains its sharpness.

During the stage of absorption the apex returns to its natural position ; the dulness gradually disappears ; the sounds and the impulse regain more of their normal character ; the friction murmur reappears, and then ceases, leaving frequently the two surfaces of the pericardium adhering,—a condition which is not harmless, since it leads to dilated hypertrophy, or to dilatation.

We cannot foretell how long it will take the disease to run through its different stages. Death may occur in less than thirty hours, the heart being paralyzed by an enormous effusion ; on the other hand, the acute attack may last for as many days, and then leave serious traces. But whatever stage the malady be in, it can be recognized only by the physical signs : by the friction, the peculiar percussion dulness, the enfeebled impulse and heart-sounds.

---

\* *Paracentesis of the Pericardium*, Phila., 1880.



There are no general symptoms that prove a pericarditis to exist. There are symptoms by which we may infer that pericarditis is present; but there are none which absolutely belong to it and would prevent it from being overlooked. The symptoms usually met with are those of inflammation of the endocardium, but with more decided local evidence of disorder. We find the anxious expression; the fever, not generally high; the œdema; the same uncertain or irregular pulse. But there is more pain over the heart,—acute, severe pain, shooting to the left shoulder, augmented by movement, increased by pressure, and associated with epigastric tenderness; there is more dyspnoea, because the distended sac presses on the lung; a dry, irritative cough; and sometimes difficulty in swallowing. Yet every one of these symptoms may be absent. The pulse may be regular; the breathing not perceptibly accelerated or laborious; and even the symptom regarded as the most important of all—the pain—may be wanting from the beginning to the end of the disease.

When the action of the heart grows weaker and weaker, the circulation becomes more irregular. The beat of the artery at the wrist is feeble, and intermits; the veins of the neck are prominent; the skin is cold and pale; the extremities are œdematous. These are always symptoms of grave import.

If next we inquire with what complaints acute pericarditis is likely to be confounded, inflammation of the endocardium and inflammation of the pleura occur at once to the mind. To contrast the signs of the first two maladies, for the slight difference in their symptoms has already been mentioned:

## ENDOCARDITIS.

Blowing sound; excited action of the heart.  
Slight, if any, increase of percussion dulness.  
Impulse strong.  
Sounds normal or more distinct, except at site where murmur is heard.

## PERICARDITIS.

Friction sound; excited action of the heart.  
In stage of effusion, marked and extended percussion dulness.  
Impulse wavy and feeble.  
Sounds feeble and muffled, except at base; no blowing sound.

Such is the distinction of pure cases of each disease. Still, as already stated, the affections are often combined. It is not uncommon to hear with the friction sound a distinct endocardial murmur. But there is sometimes a difficulty of another kind in

the way of a precise diagnosis. The murmur produced on the outside of the heart may simulate so closely the murmur produced in its interior that it is almost impossible to discriminate between them. The former may completely possess the blowing characters of the latter. Mostly, however, it is rougher; more prone to be double; and each division is like the other, equally rough, equally superficial-sounding, equally lacking in strict correspondence to the systole or to the diastole. And, above all, the sound alters at times both in situation and in character with amazing rapidity. Perceived now as an ordinary bellows murmur on the left side, it is after the lapse of some hours heard as a rough rasping sound on the right. These changes have a high degree of value. But they are not of constant occurrence; and to say that it is sometimes impossible to tell a pericardial from an endocardial sound is to say no more than is borne out by every-day experience. In the stage of effusion pericarditis is not likely to be mistaken for endocarditis.

*Pleurisy* gives rise to some of the same symptoms and signs as pericarditis. It develops a friction sound: it occasions dulness on percussion, dyspnoea, and cough. But the physical signs are in different situations. In one disorder they are in the region of the heart, and are confined there; in the other they are spread over the whole side of the chest, and are most perceptible at the back. This is true of the dulness, and, for the most part, of the friction sound, which, when of pericardial origin, is rarely heard posteriorly.

At times, however, we meet with very puzzling cases. A friction sound discerned over the heart may be in reality produced *in the adjoining pleura*. The patient is directed to suspend his breathing; the friction sound does not stop. Now, the inference from this would be that the sound originates in the pericardium; and in the large majority of instances this is a correct inference. But it is not always so. The friction may have its seat in the pleura and be caused by the movements of the heart. There are no absolute means, besides the intermission of the sound during some of the beats of the heart, as well as during some of the acts of breathing, especially in expiration, of detecting in these rare cases the true seat of the disease.

To confound the dulness on percussion caused by liquid in the pericardium with that due to *liquid in the pleura*, is a mistake

more likely to happen, because the two serous membranes, and indeed the lung, are often implicated in the same inflammation. But a pericarditis uncomplicated with pleurisy or with pleuropneumonia does not change the clear sound at the back of the chest save in very rare cases of enormous accumulation of fluid. Effusion into the pleura gives rise to a flat sound anteriorly ; to a still more perceptible dulness at the inferior portion of the chest posteriorly ; and the sounds of the heart remain unaltered, unless its investing membrane contain fluid also.

These, then, are the diseases with which acute pericarditis is liable to be confounded. There are several chronic cardiac maladies which will occasion some of the same signs and symptoms : such are thinning of the ventricles with distention of the cavities, and a dropsy of the pericardium. But the history of these affections is different, and their signs, although similar, are not precisely the same. The *dropsy of the pericardium* is associated with dropsies elsewhere, and with some obvious cause accounting for the watery exudation, and at no stage of its existence does it exhibit a friction sound.

But there is another acute complaint of which pericarditis sometimes borrows the garb. The thoracic symptoms may be latent, but the disease may produce the symptoms of extreme *gastric irritation* or inflammation. There are nausea and vomiting, and tenderness on pressure in the epigastric region. All the remedies are directed to the stomach ; and at the post-mortem examination the physician stands amazed at finding this viscus healthy and the pericardium full of serum or pus. An inquiry into the state of the heart might have saved him from a serious blunder.

Another grave error which may be thus obviated is the mistaking of some cases of acute pericarditis, on account of the wild delirium they present, for acute *inflammation of the brain*. Now, both in endocarditis and in pericarditis this active delirium may throw all the other symptoms into the background. It is difficult to see why a pericardial inflammation should give rise to such violent disturbance of the brain. It is not at all unlikely that it has its origin, in part, at least, in the contaminated state of the blood which occurs in the affections, as rheumatism or Bright's disease, with which pericarditis is often associated. However occasioned, it is necessary to be aware that the cerebral symptoms arising in

inflammation of the membranes of the heart may entirely draw off attention from the serious lesions within the chest. A fixed delusion of having committed some crime appears to Flint\* to be a distinguishing feature of the mental wandering; while Sibson† in his exhaustive analysis points out, what I have known to happen in more than one instance, that the desponding and taciturn or, as he calls it, sombre delirium lasts from two or three weeks to as many months.

Can we by the symptoms or physical signs tell the character of the fluid in the sac? We cannot by the signs, and by the symptoms we can only suspect pus if there be recurring chills, and irregular but high temperature, and if the pericarditis have arisen in the course of a malady that makes the presence of pus likely. *Hemorrhagic pericarditis* can also only be distinguished as a probability by the history. It happens in scurvy and in purpura, and may be an attendant upon cancer of the pericardium. *Cancerous pericarditis*, however, produces also serous or purulent effusion. It is never a primary disease, and it has no characteristic symptom, except it be, in some cases, darting pain in the præcordial region attending the signs of pericarditis. It is by the history and by the evidence of deposit elsewhere that we have to judge. The same is true of *tubercular pericarditis*. Here the pericarditis is often dry, and the membrane much thickened. Yet an enormous effusion may occur, as happened in a case recorded by Musser.‡

Before dismissing the subject of pericarditis, let us inquire in how far one of its terminations—by adhesion or agglutination of the surfaces—can be recognized. In many of such cases, whether or not there be coexisting dilatation, or hypertrophy, or that rare condition, cardiac atrophy, or even probably when the heart is of normal size, we find changed rhythm and dyspnœa. Yet these are not special signs of pericardial adhesion. Nor is the “abrupt, jogging, or trembling motion” of the heart, described by Hope, pathognomonic; nor the extinction of the second sound, on which Aran dwells. For the pericardial surfaces may be found most thoroughly glued to each other where neither of these signs is present. But it must be admitted that the double jog, the rebound,

---

\* Diseases of the Heart.

† Article “Pericarditis” in Reynolds’s System of Medicine.

‡ Medical Diagnosis.



is often felt, especially if the enlargement of the heart be extensive; enfeeblement or absence of impulse, while it may happen, is much rarer. Yet there is not a single symptom or sign constant, or characteristic of pericardial adhesion. The most trustworthy signs are those given by Skoda:\* a drawing up of the heart's apex during the contraction of the ventricles, with a depression in the intercostal spaces becoming visible at the same time, and sometimes with a simultaneous sinking in at the lower half of the sternum; the limits of the dull percussion sound remaining unaffected during inspiration and expiration; and a confused instead of a distinct and punctated beat of the impulse against the finger. Gairdner,† too, lays stress upon the marked movement of the intercostal spaces over the heart; while Waishe‡ thinks that the systolic dimpling and the undulatory movements in the præcordia only happen if there be, in addition to the pericardial adhesions, pleuritic adhesions in front of the organ, or if the agglutination of the pericardium be combined with cardiac hypertrophy. In the latter case, too, jogging, trembling action of the heart may be highly developed. Friedreich§ has called attention to a rapid emptying of the veins of the neck during the diastole of the heart, while with the systole they swell up; and Riess|| has told us that, owing to the close bringing together of the heart, diaphragm, and stomach, the heart-sounds resound with a metallic ring. When the pericardial surfaces are very extensively and firmly united, the eye is struck by the evident depression of the præcordial region. When the pericardium is adherent to the sternum and bands pass off compressing the aorta,—“indurated mediastino-pericarditis,”—a pulse vanishing with each full inspiration—*pulsus paradoxus*—has been described by Kussmaul.¶ The same sign has been noticed by Irvine in cases of adherent pericardium and pleura, and by Traube\*\* in exudative pericarditis where the mediastinum was not implicated. Duroziez†† regards as the most certain sign of pericardial adhesion a visible sinking in of the surface, while the hand of the observer feels the shock of the impulse; the nipple

\* Zeitschr. der k. k. Gesellsch. der Aerzte zu Wien, April, 1852.

† Edinburgh Medical Journal, 1851, 1859, etc.

‡ Diseases of the Heart, 4th ed., p. 244. § Virchow's Archiv, Bd. xxix.

|| Berliner Klinische Wochenschrift. No. 51, 1878. ¶ Ibid., No. 37, 1873.

\*\* Charité Annalen, 1876.

†† Traité clinique des Maladies du Cœur.

may be in perpetual motion. Aran has proved the tendency to sudden death in complete pericardial adhesion.

Closely connected with the subject of inflammation of the pericardium is that rare affection in which air is present in the pericardial cavity, *pneumo-pericardium*, or, more strictly speaking, on account of the frequent association with fluid, *pneumo-hydro-pericardium*. It occurs as the result of injuries, of communication established by disease between the pericardium and the neighboring organs, and in very exceptional instances is due to decomposition of liquids in the sac. Its chief diagnostic features are abnormal resonance over the cardiac region, and a metallic character of the heart-sounds. The tympanitic resonance alters in a most marked manner with changes in the posture of the patient, and is limited by a distinct line of dulness caused by the fluid. The metallic sounds may at times be heard at a distance, and may be attended with sounds of most extraordinary kind, friction sounds mixed with splashing and gurgling, the so-called water-wheel sound, the *bruit de moulin*; generally an intermittent sound, at first metallic, which Reynier\* has informed us has not a bad prognostic meaning, except when the pericardium is not intact, as in cases of traumatic opening. The symptoms of pneumo-pericardium are vague, generally those of a pericarditis, with great difficulty in breathing and failing circulation. In point of diagnosis we must be careful to take all the symptoms and signs into account, and not be misled by the modification of the cardiac sounds and the splashing and metallic phenomena due to a dilated stomach. From *pneumothorax*, even when encapsulated near the heart, we distinguish pneumo-pericardium by the dulness on percussion to be found over the displaced heart in the former malady, and the amphoric or metallic respiratory sounds that are heard in addition to the metallic heart sounds.

The entrance of air may happen, as in the cases of Meigs† and of Müller,‡ by a rupture brought about by the pericardial exudation,—in the one case into the œsophagus, in the other into the lung. These cases of ulcerative perforation almost all end fatally.

**Myocarditis.**—The substance of the heart itself undergoes at times inflammation. Of this there are several varieties, two of

---

\* Arch. Gén. de Méd., May, 1880. † Amer. Journ. Med. Sci., Jan. 1875.

‡ Deutsches Archiv für Klinische Medizin, Bd. xxiv., 1879.

which are the most distinctive,—the acute inflammation of the muscular walls, and the chronic myocarditis or fibroid degeneration of the heart. The *acute* gives rise to infiltration among the fibres of the heart of blood-corpuscles, of granules of exudation, and of leucocytes, and local softening and circumscribed abscesses, and even gangrene and perforation of the ventricle, may result. But, though familiar with the post-mortem appearances, we are not enabled to foretell the state of the heart during life, mainly because the muscular structure is rarely affected without the endocardium, or still more frequently the pericardium, being implicated, and thus the manifestations of these disorders occur mixed up with those of the carditis. On analyzing the cases on record, I cannot, indeed, find either a symptom or a sign which can be considered as in the least pathognomonic. Extreme pain in the cardiac region is the most usual and the most prominent of the symptoms. It is sometimes excruciating and sharp, at other times dull, yet distressing and constant. The breathing is generally much oppressed; delirium is often present; the skin becomes cold; the heart fails in power; and the patient dies in a state of utter prostration or appears to suffocate. The pulse, as in endocarditis or in pericarditis, exhibits no uniform character. The statement that it is invariably intermittent, feeble, and quick, is not correct. It is so as the disease advances, but it has been reported to be full, and not above eighty, long after the distress in the chest was unbearable.\* The signs of cardiac failure are quickly developed. In purulent myocarditis the temperature shows marked remissions and exacerbations, and rigors and sweatings are usual.† Acute myocarditis may occur in rheumatism, but it is most common in pyæmia and in phlebitis. Its occasional association with gonorrhœa has been pointed out, and it may be found with or without gonorrhœal rheumatism.‡ In children there is a distinctly cerebral form.§

---

\* Salter, Medico-Chirurgical Transactions, vol. xxii. In several of the cases on record, for instance in the one mentioned by Graves in his Clinical Lectures, there was coexisting valvular disease, which, of course, invalidates the statements as regards the character of the pulse, and, indeed, as regards many of the other symptoms.

† Bramwell, Diseases of the Heart, Edinb., 1884.

‡ Councilman, Amer. Journ. Med. Sci., Sept. 1893.

§ Mitchell Bruce, Keating's Cyclopædia of the Diseases of Children, vol. ii.

Acute interstitial myocarditis and parenchymatous myocarditis, the muscular fibres in both being infiltrated with granules, have no distinctive symptoms. They occur in fevers, particularly in typhoid fever, yellow fever, and smallpox, and in pericarditis, and may be suspected under these circumstances from the feeble heart action.

*Chronic myocarditis*, or fibroid degeneration, often results from rheumatism, or attends pseudo-hypertrophic paralysis. A very common cause is disease of the coronary arteries, especially obliterating endarteritis of syphilitic origin. The disease is most common in men, and may lead to aneurism of the heart. The diagnosis of chronic myocarditis is as uncertain as that of the acute form. The symptoms are those of a feeble heart: œdema, great dyspnœa, cough, hemorrhages into different organs, venous congestions, have been especially noted. In some cases there is pain over the heart. The percussion dulness in the cardiac region is somewhat increased. The first sound is indistinct, the second over the aorta very weak. The most characteristic sign is a want of correspondence between the heart and the pulse-beats; these are unequal and irregular.\* Some stress may be laid on the signs of pericardial adhesions, if present.

#### Chronic Diseases attended with Increased Extent of Percussion Dulness, but with Normal or almost Normal Heart-Sounds.

We often meet with a group of affections which present the phenomena of extended dulness on percussion in the cardiac region, associated with sounds like those heard in health: they may be louder or less loud, better defined or less well defined, still they are the natural sounds of the heart, and no cardiac murmur is detected, unless the disorder be no longer uncomplicated.

To this group belong those diseases which affect the walls of the heart or its cavities, without having involved the valvular apparatus, such as hypertrophy and dilatation,—types of the two different states of force and of weakness, but both exhibiting an extent of percussion dulness greater than in health, and heart-sounds not materially changed.

**Hypertrophy.**—Hypertrophy of the heart is an overgrowth of its walls, and usually also of its cavities; for, although we

---

\* Ruhle, Archiv für Klin. Med., 1878.




may have the muscle thickening without the cavity's enlarging, nay, even with its diminishing in size, neither this simple nor the concentric hypertrophy occurs, save in rare instances. It is evident that any one of the chambers of the heart may alone become hypertrophied. But, practically, the state we mean when speaking of cardiac hypertrophy is an increase of the ventricles, and especially of the left ventricle, in its wall and cavity, with a similar, although much slighter, expansion of the right side. Whether the auricles be enlarged or not, is a matter always more of conjecture than susceptible of proof.

The physical and vital manifestations of the heart having outgrown its natural dimensions are these. The pulse is full and strong, and somewhat tense. The face is florid, or else it is pale; and the mucous membranes of the lips and eyelids are injected. The eyes are bright, and apt to be prominent. The carotids pulsate forcibly under the least excitement. Some persons suffer from headache and giddiness; in fact, all the symptoms denote a circulation actively, too actively, carried on. Yet the symptoms directly referable to the heart are not marked. There is, as a rule, no pain or irregular action of the heart, nor do violent fits of palpitation occur. What the patient comes to consult his physician about, are rushes of blood to the head; or a ringing in the ears; or a feeling of weight in the epigastrium which troubles him after a full meal; or shortness of breath; or in consequence of the powerful action of the heart, when lying in bed, attracting his attention; or because he is alarmed about a dry cough, and believes himself the victim of pulmonary consumption.

The physical signs are more uniform than the symptoms. We observe a fulness or arching of the præcordial region, and an impulse, strong, heaving, and extended over several intercostal spaces. The apex does not strike the chest-walls between the fifth and sixth ribs, but its beat is perceived lower down, usually an inch or more to the outside of the nipple line. The extent of percussion dulness increases, both longitudinally and transversely; and particularly in the latter direction, if the right ventricle be much enlarged. This peculiarity in the expansion of the area of dulness on percussion forms, in truth, with the greater dyspnoea, and with an impulse more directly perceived over the right side of the heart, near the pit of the stomach, and often out of propor-

tion to the compressible and rather small radial beat, and with the increased distinctness of the second sound of the pulmonary

FIG. 46.



An hypertrophied heart lying in its position in the chest. The cause of the lowered apex beat, and of the extension of the impulse, as well as of the somewhat squarer outline of the increased dullness over the enlarged organ, is obvious from the shape and position of the heart.

artery, the sign that hypertrophy with dilatation has principally affected the right side.

The first sound of an hypertrophied heart is duller than in health, but prolonged and weighty. The second sound is not particularly changed. There are no murmurs, except under rare circumstances, which will be mentioned in discussing valvular diseases. Thus, the greatest value of auscultation is that, by showing the sounds but little altered, it enables us positively to exclude a lesion of the valves; just as the chief service of percussion, with reference to an enlarged heart, consists in permitting us to distinguish the excited motions of the simply disturbed organ from the action of a heart the walls of which are thickened; and as the main use in noting the impulse is that it serves as a means

of discrimination between hypertrophy and those affections in which the beat is weakened, such as dilatation or a pericardial effusion, or between the dulness in the præcordial region due to hypertrophy and that caused by deposits in the pleura, in the mediastinum, or in the lung. Where there is contraction of the left lung, as from pleurisy or fibroid change, more of the heart is exposed, and the dulness on percussion in the cardiac region is increased, as well as the impulse, which is felt over a larger space and to the left; but the cardiac sounds are unchanged, and deep inspiration alters the extent of cardiac dulness but little.

Hypertrophy may be combined with decided dilatation of the heart. This kind of hypertrophy presents a less dull, prolonged first sound, and the pulse, though full, is likely to be more compressible. Hypertrophy may affect specially any part of the constituents of the muscular walls. Thus, the connective tissue, as Quain has particularly called attention to, may be alone concerned in the morbid action. Hypertrophy of the heart is found much more frequently among males than among females. Its causes are various. It is common in Bright's disease and in general arterial sclerosis; continued functional excitement produces it; so does any kind of strain and overaction, and perhaps excessive nourishment. It is found to be common among inordinate beer-drinkers. But the main cause is an obstruction to the circulation, either in the heart or in other organs. It is for this reason that the complaint is so often met with in connection with diseases of the valves or of the large arteries, and that the right side of the heart enlarges when the pulmonary air-vesicles are overdistended. We also, as we have seen, encounter hypertrophy of the heart as a consequence of the obliteration of the pericardial sac by its two surfaces adhering. In the hypertrophy of Bright's disease reduplication of the first sound of the heart is often noticed.

There is a form of hypertrophy of the heart to which attention has been particularly called by Fothergill's description,—the so-called *gouty heart*. Generally, although not always, there is coexisting disease of the kidney of the chronic contracting form. In the first stage of the affection, when well marked, we find decided hypertrophy with accentuation or booming of the second aortic sound, high blood-pressure, tense pulse, hardened arteries, and the passage of large amounts of pale urine of low specific

gravity. The renal changes may or may not be evident ; we may or may not detect albumen in the urine. In a subsequent stage of the malady there is failure of the circulation, and with the signs of the heart-failure, very often going hand in hand with fatty degeneration, the bulk of urine diminishes and the renal affection becomes more marked. The cardio-vascular phenomena are early made perceptible by the sphygmograph. The full, tense pulse gives a full up-stroke, a broad summit, and a retarded down-stroke ; the "square-headed tracing" formed is very characteristic of the malady, and bespeaks the fibroid change in the kidney, whether or not albumen be found. In some instances there is considerable dilatation as well as hypertrophy, and then severe palpitations result. The high blood-pressure is due to the waste-laden blood ; and the defective nutrition is apt to show itself also in atheromatous arteries, which in part account for the sphygmographic tracings. The skin oftens exhibits little twigs of dilated vessels ; the ear is usually deep red, with a large glistening lobe ; or in spare persons the lobe looks withered ; the teeth become blunt and worn down in time ; the hair is apt to be iron-gray. There is the history of gout, acquired or hereditary, but there may have been no active outbreak of gout, rather the condition of imperfect assimilation and increased uric acid, known as lithæmia.

**Dilatation.**—Dilatation of the heart is the reverse of hypertrophy. By this it is not meant that because the cavities are dilated the walls may not be increased. But it is meant that the morbid condition in which the cavities have been stretched out of all proportion to the thickness of the muscular walls is the reverse of the condition in which the walls are stronger, firmer, and more powerful than in health ; in other words, the latter state is very different from the former, and when it predominates we call the affection hypertrophy ; when the former is in excess we speak of the disease as dilatation, no matter whether the walls be slightly thicker than normal, or of natural thickness, or thinner, and apparently hardly capable of supporting the weight of the blood.

From these almost opposite pathological states, almost opposite physical signs or symptoms might be expected. And so we find it. In dilatation we look in vain for the activity and power with which the blood is forced out of an hypertrophied heart. Every-




thing indicates inaction and stagnation. There is a strong tendency to venous congestions and to dropsies. The portal system is gorged. The liver increases in size. The bowels are constipated. The urinary secretion is interfered with, and sometimes albumen is passed. The hearing may become dull. The patient is languid and feeble, and his intellect obtuse. He suffers from chilly sensations, and from uneasiness in the cardiac region and palpitations. The pulse is small and irregular, and the veins of the surface are swollen. The skin around the ankles, and often at other parts of the body, pits on pressure. But, since it is the right side of the heart which is usually the most affected, the lungs show most plainly the effects of the venous stagnation. Difficulty in breathing, making itself at times manifest in paroxysms attended with wheezing respiration; a chronic cough; a collection of serum in the pulmonary structure,—all add to the misery which the perilous malady entails. And as it is commonly some obstructive disease in the lungs, such as emphysema, which has given rise to the dilatation of the right side of the heart, so this again augments the morbid state of the lungs, and aggravates the symptoms.

The physical signs are very unlike those of hypertrophy. The same extended dulness on percussion exists; but it is associated with a feeble and fluttering impulse, which is in strong contrast with the heaving, powerful blow of an hypertrophied left ventricle, and which at times may be seen, yet cannot be felt. The sounds are not always the same. When the walls are thin, they are clearer, sharper, and more ringing than in health: if, however, the muscular structure be at all disorganized, the first sound is faint and very ill defined. The second is often split, giving rise to the so-called gallop rhythm. But no murmurs are perceived, unless a watery state of the blood produces them, or unless it happens—and it does not unfrequently happen—that the dilatation of the heart is conjoined to valves incompetent, either temporarily or permanently, to prevent regurgitation.

Such is the description of cases of marked dilatation. All cases are not, however, so distinct, nor are they uncomplicated. Organic affections of the heart are, indeed, indefinitely blended, and dilatation is met with in different combinations and in every possible degree. Accordingly, its symptoms and signs are somewhat

dissimilar. But one constant feature it certainly preserves: it always holds up to view both the vital and the physical manifestations of a weak heart. Indeed, when an hypertrophied heart dilates, the signs of relative weakness become superadded, the impulse is not so strong as before in comparison with the percus-

FIG. 47.



A dilated heart, the right ventricle opened. In this case there was no valvular disease. Hence the characteristic physical signs; the increased dulness on percussion, the extended but weak impulse. The first sound was feeble, for the organ was soft as well as dilated.

sion dulness, and dropsy becomes a marked symptom. Pure dilatation is likely to be confounded with the diseases in which enfeebled action of the heart is encountered, and these are fatty degeneration and a pericardial effusion.

**Fatty Degeneration.**—This is one of those disorders with the anatomical characters of which we are far better acquainted than with their clinical history. The microscope has revealed to us that the soft flabby heart, which appears to the eye little changed from health, has had its muscular fibres atrophied and transformed into fat-granules and oil. It has thus explained to us why a heart seemingly so little altered should rupture, or why death should set in with all the evidences of failing circulation,

when nothing in the whole body can be found sufficiently diseased to account for the termination of the vital action. But our power to recognize the fatty change during life has not kept pace with our power to recognize it after death. There is as yet no sign discovered by which we can positively say that the dangerous disorganization of the muscular fibres of the heart is in progress. We may, however, suspect it, if the signs of weak action of the heart—feeble impulse and ill-defined sounds—coexist with oppression, with a tendency to coldness of the extremities, with a pulse permanently slow or permanently frequent and irregular, and be met with in a person who is the subject of gout or of a wasting disease, or is very intemperate, or has arrived at a time of life at which all the organs are prone to undergo decay. Something more than a suspicion is warranted if, in addition, there be proof of atheromatous change in the vessels, or of fatty degeneration elsewhere, such as an *arcus senilis*;\* or if it be ascertained that the patient suffers from pain across the upper part of the sternum and from paroxysms of severe pain in the heart; that he sighs frequently; that he is easily put out of breath; that his skin has a yellow, greasy look; that he is subject to syncope, or to seizures during which his respiration seems to come to a stand-still; and that he is liable to vertigo, or to be stricken down with repeated attacks having the character of apoplexy, save that they are not followed by paralysis.

Now, here are certainly a group of phenomena dissimilar to those of a dilated heart. Let us add that the extent of the cardiac percussion dulness remains unaltered, except in those instances in which *hypertrophy* or dilatation coexists, that dropsies and local congestions are not prominent symptoms, or indeed do not happen at all, and the dissimilarity becomes still greater. A differential diagnosis would, under such circumstances, be anything but difficult. But in point of fact the matter is generally not so easily decided, and there are several reasons why it is not. One is, that all the features described are rarely combined in the same case;

---

\* But the *arcus senilis* may be absent. Fothergill points out that there is a false as well as a true *arcus*. The latter alone is significant of fatty degeneration and tissue-decay. It is a ring around the cornea of yellowish hue with blurred out lines, and the cornea itself is cloudy.

indeed, some of the more marked, such as the seizures like apoplexy, the Cheyne-Stokes breathing, are uncommon rather than common, and the peculiar breathing occasionally occurs in other cardiac maladies. The second is, because non-fatty softening, the result of a granular infiltration, as met with, for instance, in fevers, may present much the same vital and physical manifestations. The third is, because a fatty heart has a tendency to become dilated, and the symptoms and signs of the former disease are then merged into the symptoms and signs of the latter, throwing us back for a diagnosis into the province of conjecture and probability. With the organ in such a condition, the practical value of a differential diagnosis is, however, not great; for both affections require that the power of the heart should be sustained. Decided dropsy would indicate that dilatation had happened.

The remarks about fatty heart apply particularly to that variety of the disorder in which the muscular structure in middle-aged or elderly persons has slowly undergone decay. But we also, although far less frequently, meet with fatty heart in young persons and in a more acute form; and we encounter it in chlorosis, in pernicious anæmia, after repeated hemorrhages, and after phosphorus poisoning. Poisonous doses of acids, such as nitric, sulphuric, oxalic, are said by Von Dusch also to give rise to the cardiac change.

Persons who have fatty hearts are subject to attacks of faintness, preceded or attended with *sensations of great coldness*, or a chill. Sometimes these attacks happen daily, or every few days, and in such a manner as to give rise to the impression that they are due to malaria. A number of instances of the kind have come under my observation, and I have met with them particularly at the end of fevers or other debilitating diseases happening in those affected with feeble hearts. The seizures, though bearing a resemblance to intermittent fever, are unlike it in being associated with signs of great weakness of the circulation or heart failure, sometimes joined to a sense of impending dissolution; in their irregular accession; and in their not being followed by fever. In doubtful cases the thermometer, by showing the absence of the great rise of temperature of the malarial disorder, will materially assist us in the diagnosis.



*Heart starvation*, to which Fothergill\* has called attention, has, in the feeble circulation, the cold extremities, the tendency to vertigo, and the pseudo-apoplectic attacks, symptoms common with those of fatty heart. But there is not a true areus, and the malady is not associated with disease of the arteries. It is often an attendant upon general ill nutrition, and worry, and long hours of work and short hours of sleep.

A fatty heart sometimes *ruptures*. Now, in spite of the care with which the physical signs of this mishap have been detailed, we know nothing positive about them; for death usually takes place far too rapidly to permit of any observations. The symptoms that are mostly noticed are these: the patient is suddenly attacked with intolerable anguish in the heart; he presses his hand to it, then faints, and soon expires. Or else he lives for a short time, suffering from faintness, cramps, and difficulty of breathing, and with death plainly written on his face.

Where there is *fatty accumulation on the heart*, without fatty change of its fibres,—a condition we sometimes find in persons whose internal viscera are loaded with fat,—the manifestations are those of a feeble heart, and different from fatty degeneration only in degree. The first sound of the heart is weak and toneless; the pulse is feeble, but, as Walshe tells us, regular. The percussion dulness in the cardiac region is somewhat increased. A sensation of oppression over the region of the heart, or even actual pain, is complained of. There is shortness of breath on taking exercise.

*Atrophy of the heart* is so rare a condition that its symptoms are scarcely understood. All we know is that at times in certain wasting diseases, such as tubercular phthisis and suppurating bone affections, the heart atrophies; it may also do so when the coronary arteries are calcified, or the pericardium is tightly adherent; and cardiac atrophy is said to happen occasionally after pregnancy and chlorosis. It has not a single symptom nor a single sign by which it can be recognized with certainty. Theoretically, the diminished percussion dulness, clear sounds, and feeble impulse should enlighten us; but, even in cases where there is no co-existing fatty change, they are too uncertain to be made a basis for diagnosis, or attending lung conditions throw doubt on several

---

\* Edinburgh Med. Journ., May, 1881.

of them. There is great tendency to palpitation, and the pulse, Hayden tells us, is quick, all but inappreciable, yet regular.

**Pericardial Effusion.**—Pericardial effusion also presents the signs of a weak heart with increased dulness on percussion in the cardiac region, and is very liable to be mistaken for a dilatation of the organ. Where the effusion forms part of a general dropsy, the detection of the cause of the latter, in connection with the different signs which fluid in the pericardium occasions, will prevent error. Where the liquid has remained after an inflammation of the membrane, both signs and symptoms are like those of the state of effusion in acute pericarditis, and, although there are points of resemblance to a dilated heart, there are also points of contrast, as the subjoined table shows :

DILATATION OF THE HEART.	CHRONIC PERICARDITIS WITH EFFUSION.
Percussion dulness increased in extent, but square in outline. Impulse in epigastrium.	Percussion dulness increased, but often of pyramidal shape. Impulse in third or fourth left interspace, apex tilted upwards.
Heart-sounds clear and sharp; sometimes, however, feeble.	Heart-sounds feeble and distant-sounding at the apex, but distinct near the upper part of the sternum.
No friction sound.	Often friction sound still heard at the base of the heart.
Dropsy; signs of venous stagnation; severe cough, and dyspnœa.	Neither dropsy nor venous stagnation is observed; or, if at all, only in a very limited degree. Cough and dyspnœa are not such prominent symptoms.
The history of the disease shows it to be gradually developed.	The history frequently points to the acute attack.

These, then, are the marks of distinction presented by a chronic pericardial effusion, a fatty heart, and cardiac dilatation; in other words, between the main morbid states which occasion the signs and symptoms of a feebly-acting heart. Before proceeding, let us glance at one more condition, fortunately infrequent, which may give rise to some of the same phenomena as those described,—*an accumulation of blood in the cavities of the heart*. Like dilatation, this increases the area of percussion dulness, and is often associated with perverted rhythm. The chief differences, as far as our limited knowledge of the subject permits us to define them, are

these: the impulse is generally much more labored and irregular, is sometimes strong, sometimes weak, not so almost uniformly indistinct or tremulous. There is much more venous congestion of the face, with greater dyspnoea, and we often find some acute malady, such as endocarditis or pneumonia, giving rise to the cardiac engorgement. But the matter is a very difficult one to determine; for many of the same states which lead to dilatation may produce an accumulation of blood in the heart; nay, dilatation itself predisposes to it. We must bear in mind also that an acute dilatation sometimes happens from shock or sudden fright.

**Diseases of the Heart exhibiting more or less of the Signs and Symptoms of Enlargement of the Organ, and accompanied by Endocardial Murmurs.**

**Valvular Affections.**—These, when not due to congenital malformations, are most commonly the result of rheumatic endocarditis, of slowly progressing sclerotic changes, or of heart-strain. A certain number of cases have their origin in some of the fevers, as in scarlet fever, and in septic conditions and blood-changes, as in ulcerative endocarditis. The different valves are not affected by these causes alike. Rheumatic endocarditis is the principal cause of disease of the mitral valve, especially of mitral insufficiency; but among prominent causes of this are also alterations in the muscular wall of the ventricle or in the tendinous cords. Aortic insufficiency is generally due to slow sclerotic changes in the valvulets, whether attended with atheroma or not, or to subacute or chronic endocarditis from heart-strain; it may be also owing to the sudden rupture of a valve previously damaged. Mitral constriction is mostly brought about by atheromatous or calcareous alteration, as is aortic constriction; but in mitral constriction we may have also a history of endocarditis in early childhood subsequent to rheumatism, an exanthematous fever, or chorea. In insufficiency of the tricuspid valve we can trace usually the result of overdistention of the right heart, such as follows pulmonary congestion caused by mitral disease, or of an obstructive disease of the lung, such as emphysema or cirrhosis. Tricuspid stenosis, and the other very rare valvular affections of

the heart,—those of the pulmonary artery,—are commonly congenital.

To find the sounds of the heart clearly and well defined, is to know that no disease of the valves exists. When the valvular apparatus is disordered, the mischief betrays itself, for the most part, by a murmur. If, therefore, a murmur of any permanence be met with in the heart, especially if it be associated with the signs of either hypertrophy or dilatation, the inference that valvular disease exists will in the vast majority of cases be a correct inference.

Yet it will not be so always; for there are other morbid states besides valvular affections which engender a murmur, which may be even accompanied by all the manifestations of enlargement of the heart. Malformations, such as communications between the auricles or between the ventricles, or between the great vessels near their origin, or impoverished blood, or a misdirected blood-current, may occasion a murmur.

Now, with reference to *malformations*, their presence in adults, or in children that have passed the days of infancy, is exceedingly rare. The most trustworthy symptom they present is that indicating the admixture of arterial and of venous blood; in other words, the symptom of cyanosis, the bluish discoloration of the skin. In addition, we may perceive the signs of disturbed circulation in the lungs, such as dyspnoea and cough, and irregular action of the heart, and a blowing sound in the cardiac region; hypertrophy of the heart, especially of the right heart, is also very generally present. Still, the recognition of these malformations is always more or less a matter of conjecture. With the aid of more such researches as those of Moreton Stillé,\* of Peacock,† and of Hochsinger,‡ we shall become more accurately acquainted with the different lesions, and perhaps be able ultimately to discern them with certainty during life. At present it is in their rarity that the safety against errors of diagnosis lies:

---

\* Amer. Journ. Med. Sci., July, 1844.

† Treatise on Malformations of the Heart; see also the valuable treatise of Keating and Edwards on Diseases of the Heart and Circulation in Infancy and Adolescence, 1888; Osler's article on "Congenital Affections of the Heart," in Keating's Cyclopaedia of the Diseases of Children, and Hirst's, in Starr's American Text-Book of Diseases of Children.

‡ Die Auscultation des Kindlichen Herzens, Wien, 1890.



As a few points of assistance, it may be mentioned that communication of the ventricles through the septum gives rise to a systolic murmur at or near the base of the heart not propagated into the arteries, but according to Roger and to Sansom also heard between the shoulders; that the passage of blood through an open foramen ovale very rarely engenders any sound, though presenting marked cyanosis; and that, whether coexisting with these lesions or not, the majority of instances of cardiac malformation, after the age of twelve, present signs of obstruction at the orifice of the pulmonary artery. In this instance either a systolic or a diastolic murmur may be there perceived; in the first case the second sound of the heart is weak or wanting in the second interspace on the left side. Mitral disease of congenital origin is very rare. Thrill over the præcordial region is seldom met with, except when congenital defect in the septum exists. Loud, vibratory systolic murmurs heard most distinctly over the upper third of the sternum without attending hypertrophy of the left ventricle point to persistence of the ductus Botalli. A curious result of cardiac malformation has been observed,—abscess of the brain without appreciable cause.\*

The resemblance borne by cases of *functional disturbance of the heart*, associated with impoverished blood, to valvular affections, has already engaged our attention. The age; the appearance of the patient; the seat of the blowing sound at the base of the heart, and its soft character; the venous hum; the fact that the cardiac murmur is followed by a sharp second sound,—are all points upon which some stress may be laid; yet not so much as upon the absence of the phenomena of an enlarged heart. But if the question be asked, Are the latter absolute demonstrations of the existence of an affection of the valves? cannot an hypertrophied or a dilated heart, with sound valves, be combined with a condition of blood capable of producing a murmur?—we are forced to answer that such is possible. Under these circumstances, the tact of the physician may help him to a well-judged decision; but the only proof of a well-judged decision is afforded by time, or by the result of treatment that restores the blood to its normal state.

---

\* Ballet, Archives Générales de Médecine, June, 1880.

A murmur caused, in violent excitement of the heart, by *misdirection of the current*, due chiefly to temporary interference with the closure of the valves, or perhaps owing to altered tension of the valves,—causes the exact working of which I have elsewhere inquired into,\*—may become a troublesome source of error in diagnosis, especially when heard over a heart in a state of dilated hypertrophy or of dilatation. Fortunately, a blowing sound of this origin and in this combination is comparatively rare, and we are enabled to discriminate it from an organic valvular murmur by its not being persistent. It is much more likely to be heard at the apex, or rather, according to my own observations, somewhat above the apex, than is a murmur owing to changes in the blood; and it differs from the systolic blowing sound of mitral disease partly by the peculiarity of seat just mentioned, partly by its non-diffusion, its usual absence at the back of the chest, the want of harshness in the inconstant murmur, and the low pitch. Murmurs of this kind are also caused by obstructive diseases of the lungs, without disease of the heart being present. They may be brought out, as John K. Mitchell has shown, by suddenly closing the hand tightly.†

At times a murmur is heard which is not dependent on a cardiac affection, but on *lung changes*. We find, for instance, in consolidation of the left apex, especially if the lung be also contracted, a murmur, almost invariably systolic, over the site of the pulmonary artery; or we may encounter over large cavities with thin walls situated in the neighborhood of the heart a systolic, cardio-pulmonary murmur, caused, most likely, by the agitation of the air in the cavity, the heart being quite sound.

These, then, are the causes which impair the value of the cardiac blowing sound as a sign of a valvular lesion. Yet they do not happen often enough to prevent us from regarding a persistent murmur as eminently indicative of an organic affection of the valves.

Let us suppose that we are convinced that the murmur is due to a structural lesion. Can we say what its precise nature is? Can we accurately foretell that the valve is merely roughened, or that it has undergone calcareous transformation, or that it has

---

\* On Functional Valvular Disorders, Amer. Journ. Med. Sci., July, 1869.

† Transact. College of Phys. of Phila., 1892.

been bound down, or that it is lacerated, or that vegetations spring from it, or that its muscular attachments are sound or unsound? No, assuredly not. The most we can do is to judge whether the orifices through which the current flows be narrowed, or whether, by the valves not closing, they permit of regurgitation; and to distinguish even this we have to take into account more the time of the occurrence of the sound than its particular character or pitch. Indeed, all distinctions based entirely on either of these are not borne out by clinical experience. Valves incompetent to close the openings at which they are seated may permit a murmur to be generated of any character and of any pitch. It is true that a harsh murmur, like that of a saw or of a rasp, is for the most part occasioned by a contracted orifice with rigid valves; but many contracted orifices with rigid valves exist without producing such a rough noise.

A cardiac sound which is rare, but which, when present, is most generally associated with a narrowed orifice, is a distinct *musical tone* heard at the mitral or aortic valves. It resembles the cooing of a pigeon; or the auscultator listens and listens again, and directs the patient again and again to suspend the respiration, before he becomes convinced that the sound is not a sibilant rale in the lung. It is sometimes perceived merely at the end of an ordinary bellows murmur, and disappears and reappears from time to time. Where this rare sound is met with, the valves after death are commonly found to be rigid and unyielding. Yet this is not always the case. Sometimes the musical note is produced by the vibrations of clots which impede the rush of blood through the apertures of the heart, or by the loose edge of a valve flapping to and fro in the current. Occasionally, too, we hit upon it in chlorosis; but only very occasionally, and never unless it be then equally or more marked in the arterial system. We have the authority of Stokes for the observation that it may be suddenly developed and precede the signs of structural alteration of the heart. Schroetter maintains that the musical murmur is due to the vibration of a fine fibrous band stretched across the ventricle or a valvular orifice.\*


It has been already stated that we judge best of the condition of

---

\* Wien. Med. Blätter, No. 1, 1883.

the orifices and of the valves by ascertaining the time at which the bellows sound occurs. To do this it is necessary to recall in what state the orifices are during the movements of the healthy heart. During the contraction of the ventricles, the valves at the auriculo-ventricular openings are closed, to prevent regurgitation into the auricles; and the valves of the aorta and

FIG. 48.



Narrowing of the aortic orifice by vegetations springing from the valves, the structure of which was indeed, to a great extent, destroyed. The engraving illustrates also the physical signs of aortic constriction.

pulmonary artery are open, so as to permit the blood to pass along the arterial trunks. During the dilatation of the heart the reverse takes place: the valves at the origin of the great arteries are shut, to keep the blood which has just been sent forth from regurgitating, and those valves the function of which is to act as



gates to the auriculo-ventricular apertures are swung back, to allow the stream to flow into the ventricles.


If thus a murmur occur with the contraction of the heart and the first sound, it is owing to the blood either regurgitating from the ventricles into the auricles, or meeting with difficulty in passing into the aorta or pulmonary artery; if it occur after the contraction of the heart, and correspond to the second sound, it is due to the blood passing through a narrowed mitral or tricuspid orifice, or streaming back into the ventricles through incompetent aortic or pulmonary valves. But can we distinguish at which valve the mischief lies? Generally we can. By attending to the site of greatest intensity of the murmur, we become aware of the seat of its production, provided it be borne in mind what are the points at which to listen to the different valves. It is, however, also necessary to recollect that, as the whole heart is somewhat lowered, these points are rather below what they are in a natural state of things.

Now, we cannot always say whether more than one valve is affected. A blowing sound in the heart, no matter where generated, is usually transmitted all over the organ. If it mask the natural sounds at other valves, it is very difficult, nay, it is often impossible, to tell positively how many of the valves are injured, unless several spots be detected at which the murmur is intense yet not alike in character. The valves that most frequently show coexisting disease are the mitral and the aortic, particularly insufficiency of both, or aortic narrowing with mitral insufficiency. Diseases of mitral and tricuspid are also found to coexist, whether the lesion be regurgitation or narrowing. In all instances the precise character of the murmur at the different sites of the heart is of the greatest significance.

Thus the blowing sound is the most conspicuous and most constant sign of a valvular lesion. The other signs and symptoms vary in individual cases. Where the valves are but slightly affected, let us say slightly roughened, as they sometimes are after an attack of rheumatic endocarditis, the heart does not undergo any decided change in size; the circulation is carried on regularly; and, in spite of the abnormal sound in the heart, the patient's health remains unimpaired, or it is only occasionally that he suffers from palpitations. An alteration of the valves of the heart

of any extent produces, however, an alteration either in the capacity of its cavities or in the thickness of its walls, and the symptoms of dilatation or hypertrophy make their appearance along with the physical signs of extended percussion dulness and feeble or heaving impulse. Ordinarily it is the latter we meet with, because the valves of the left side are so very much more frequently diseased, and their derangements lead to hypertrophy rather than to dilatation. Affections of the *tricuspid valves* are usually connected with dilatation of the organ; hence dropsy, venous turgescence, and albuminous urine are in them more especially observed; and Blakiston has taught us their frequent

FIG. 49.



Insufficient mitral valves permitting regurgitation of the blood. The position and time of occurrence of the most significant sign of the affection are indicated in the engraving.

association with engorgement of the vessels of the brain, and how this becomes the predisposing cause of cerebral apoplexy when in connection with cardiac disease. We also find in them, or rather in tricuspid insufficiency, what Mahot has more particularly called attention to,—a pulsation of the liver corresponding to each systole of the heart, perceived by gently depressing the abdominal parietes with the hand on the epigastrium. In combined tri-

cuspid and mitral narrowing we may have the signs of pulmonary-artery regurgitation.\* In high degrees of aortic insufficiency, a systolic apex murmur, as pointed out by Flint, is very often produced by dilatation of the mitral orifice. The murmur differs from that over the damaged aortic valves, and may be presystolic in time. In coexisting aortic and mitral insufficiency the compensatory hypertrophy is arrested. In some cases of mitral regurgitation the mitral murmur occupies only the middle or the latter part of the systole.† In instances of disease of three valves, both obstruction and insufficiency, as in the case reported by Shattuck,‡ double murmurs of dissimilar kind may be heard over the area of the different orifices.

All valvular lesions may be combined with pain in the præcordia, palpitations, restlessness, and disturbed dreams. And according as the deranged circulation through the heart interferes with the circulation in other parts, special symptoms show themselves prominently. Thus, we find those who labor under a *mitral disease* suffering most from cough, from dyspnœa, and from attacks of cardiac asthma, since it is the lung which has to bear the brunt of the embarrassed flow of the blood. If we examine this organ closely, the physical sounds afford direct proof of its disordered condition. Here and there are heard plentiful moist sounds from fluid which has leaked into the air-tubes; here and there the respiratory murmur is roughened, and percussion elicits impaired clearness. This loss of the natural resonance is at times very manifest at the upper part of the lung, and I have known it to lead to the suspicion of tubercular deposit in cases in which the autopsy showed the pulmonary tissue to be healthy, though in a state of extreme congestion. Respiratory percussion renders the sound again clear. Mitral insufficiency generally leads to hypertrophy of the heart; mitral stenosis not unusually becomes associated with dilatation, or there is only hypertrophy of the right ventricle.

When the *aortic valves* permit of regurgitation, this gives rise to effects which are perceptible along the track of the arteries.

---

\* Dyce Duckworth, Clin. Soc. Transact., Jan. 1888.

† Crozer Griffith, Amer. Journ. Med. Sci., Sept. 1892.

‡ Boston Med. and Surg. Journ., 1891.

These all look superficial, and beat with apparent violence, from the force with which the thickened left ventricle is driving the blood through the tubes. The pulsation of the vessels may be even seen in the retinal vessels with the aid of the ophthalmoscope. Yet, when the finger is applied to the artery at the wrist, the strength of the beat is not so great as is expected. A short, abrupt, jerking impulse is indeed communicated to the finger; but then the artery immediately recedes, proving that it was only imperfectly filled. This pulse is the only one which gives us any real information as to the state of the orifices of the heart. In general terms, it may be stated that the pulse is small and rather tense when the openings are narrowed. Still, no stress can be laid on this in a diagnostic point of view. The want of correspondence between its strength and the force with which the heart is acting is often amazing.


More information than by merely feeling the pulse can be obtained by studying it with the sphygmograph. But even with this, as thus far developed, we gather in valvular diseases rather corroborative evidence than knowledge which is not attainable by other means of diagnosis. Perhaps with further research the instrument may be made available to inform us with certainty of the degree of the valvular imperfection; and this would be a great step in advance. As regards the most distinctive graphical signs, we obtain them in aortic regurgitation,—a vertical line of ascent

FIG. 50.



Sphygmogram taken from a patient with aortic insufficiency. The line of ascent does not terminate in as sharp a point, nor is the descent as sudden, as we sometimes find it.

FIG. 51.



Sphygmogram taken from a patient presenting the signs of mitral regurgitation.

of great amplitude, a pointed summit, and a sudden descent, with comparatively little dicrotism. If there be also marked aortic obstruction, the line of ascent is oblique, or rather the first part is



vertical, and following the sharp point is a gradual curve-like rise ; if senile changes in the artery complicate the aortic insufficiency, the sharp-pointed process terminating the line of ascent passes into a more or less horizontal plateau. In instances of decided uncomplicated aortic obstruction there are sloping up-strokes and down-strokes.

In mitral regurgitation the pulse tracing is usually very irregular, such as is seen at times in aneurism ; the line of ascent is short and unequal, and the line of descent is disposed to be oblique and to present marked dirotism. In mitral constriction there is also, usually, irregularity ; it is asserted by Mahomed\* that the up-stroke is vertical, and that there is, especially after giving digitalis, a secondary and very characteristic contraction of the ventricle manifest in the dirotic wave. Sansom† agrees in the main with this observation.

But, instead of entering into a detailed description of the pulse, however studied, or of any separate symptoms of valvular disease, let us group them together with the physical signs, according to the combination in which we are wont to meet them :

TABLE OF VALVULAR DISEASES.

SEAT OF MURMUR.	SEAT OF DISEASE.	CHARACTER OF DISEASE.	CORRELATIVE PHYSICAL SIGNS AND SYMPTOMS.
Murmur most intense at or near apex of heart.	Mitral orifice.	With impulse, means insufficiency of valves, permitting of <i>regurgitation</i> ; after impulse, and running into or corresponding to the second sound, or, more accurately speaking, generally preceding the first sound, presystolic, means <i>narrowing</i> of the orifice.	In mitral disease the heart very commonly undergoes dilated hypertrophy, especially the right ventricle. When there is also hypertrophy of the left ventricle, it is not simply mitral narrowing. The second sound of the pulmonary artery, heard in the second left interspace, is sharp, accentuated. The cardiac murmur is often distinctly perceived posteriorly on the left side, near the angle of the scapula. Dyspnoea and dropsy are prominent symptoms, especially dyspnoea. Cough is not unusual, and the pulse is not unfrequently found to be feeble and irregular. In some forms of mitral narrowing, where the curtains are not too rigid, the murmur is always rough. This is the case usually with the presystolic murmur, which is pre-eminently regarded as the sign of mitral constriction. But in this affection all murmur may be absent,

\* Medical Times and Gazette, May, 1872.

† Diagnosis of Diseases of the Heart, 1892.

TABLE OF VALVULAR DISEASES—(*Continued.*)

SEAT OF MURMUR.	SEAT OF DISEASE.	CHARACTER OF DISEASE.	CORRELATIVE PHYSICAL SIGNS AND SYMPTOMS.
Murmur most intense at or near the middle of the sternum, or heard with equal distinctness close to the sternum in the second interspace on the right side, and thence propagated into the arterial system.	Aortic orifice.	With impulse, means <i>narrowing</i> , or obstruction; with diastole, and taking the place of the second sound, or occurring in both sounds, the first murmur short, means <i>regurgitation</i> .	either temporarily or permanently. In mitral narrowing a thrill in the cardiac region can be often felt. Mitral narrowing is frequently associated with contracted kidney.  Hypertrophy of left ventricle, often to a very great degree, the compensation being very decided. All the cardiac sounds may be normal, except at the aortic valve, although they are obscured by the murmur. This is distinct in the carotids, and is sometimes as well heard at the ensiform cartilage as over the sternum and on a line with the third intercostal space,—a fact necessary to be aware of, so as to avoid confounding the aortic lesion with one of the tricuspid valve. When the orifice is constricted, a purring thrill is frequently observed to attend the harsh or musical systolic murmur. The symptoms in aortic valve disease are often remarkably latent. There is very commonly neither dropsy nor dyspnoea. The pulse in regurgitation is abrupt and receding, and all the superficial arteries and the capillaries pulsate. It is not unusual to find a double aortic blowing sound attending aortic regurgitation, probably from slight coexisting obstruction of the orifice, though this is not always found; a double murmur is also heard in the carotids and femorals. A mitral apex murmur may be also noticeable.
Murmur most intense at or very near to the ensiform cartilage, and over the lower part of the right ventricle.	Tricuspid orifice.	With impulse, <i>regurgitation</i> ; with diastole, and taking therefore the place of the second sound, or preceding the first, <i>narrowing</i> .	Tricuspid regurgitation exists usually in combination with dilatation of the right ventricle, and therefore with the symptoms of this condition: with venous congestions, with dropsies, with difficulty in breathing. On account of the open state of the orifice, the cervical veins may pulsate during the movements of the heart; and in all cases they are distended. The pulsatile motion in the neck becomes especially visible when the breath is held in expiration. The cardiac murmur is ordinarily soft, of low pitch, is not transmitted into the arteries, and is not heard above the level of the third rib. In some cases it is so feeble as to be with difficulty discerned. In tricuspid narrowing, a very rare disease, there are presystolic murmur and thrill, cyanosis of the face and lips, and great dropsy.

TABLE OF VALVULAR DISEASES—(*Continued.*)

SEAT OF MURMUR.	SEAT OF DISEASE.	CHARACTER OF DISEASE.	CORRELATIVE PHYSICAL SIGNS AND SYMPTOMS.
Murmur most intense at the third left costal cartilage near the sternum, or even somewhat lower, or in the second intercostal space to the left of the sternum.	Pulmonary orifice.	With impulse, is <i>narrowing</i> ; taking the place of the second sound, <i>regurgitation</i> .	We have little knowledge, derived from clinical observation, of diseases of the pulmonary valves, of all the valves the ones most rarely affected. Nor does a murmur in the situation indicated, and hardly audible over the left apex or along the sternum, or in the course of the great vessels, having therefore the characteristics of a pulmonic murmur, warrant a diagnosis of disease of the valves: for it may be due to anæmia; be caused by deposits at the upper part of the left lung; or be observed immediately after or during the continuance of hemorrhage from the lungs. But these remarks scarcely hold good with reference to a diastolic murmur, and not at all as regards a double murmur. If this be present, and attended with thrill and with signs of dilated hypertrophy of the right heart, we are justified in concluding the disease to be a lesion of the pulmonary valves, or at the origin of the artery. But as regards the association with signs of hypertrophy especially, we must bear in mind that in rare instances of mitral disease, especially regurgitation, the murmur is loudest at the pulmonary area. Pulmonary narrowing is almost always congenital. Pulmonary insufficiency may be also congenital, or be due to malignant endocarditis.

In this manner are the symptoms and signs of valvular affections associated. It is not exactly the combination and precisely the way in which they happen in every instance. There are too many circumstances which modify them; disorders of several valves are too constantly conjoined; at the same orifice both narrowing and a state permitting of regurgitation are too often found to coexist,—to permit any tabular representation to express either all the symptoms or all the signs which may occur in individual cases. Apart from this difficulty, there is another: even where the affection of a second valve has been correctly fixed upon, the irregularity of the heart's action may be such that it is impossible to say whether the blowing sound heard be systolic or diastolic; whether, therefore, the orifice be narrowed or the valves insufficient. But this is not a matter of so much consequence; the

matter of consequence is, to determine that a disease of the valves is present.

Presuming that we have been enabled to fix, and to fix accurately, the state of each aperture, there is a point where all our skill invariably comes to a stand-still. We cannot tell how long it is possible for life to continue, or under what circumstances death will happen. It may take place suddenly and most unexpectedly in cases in which the amount of disease in the heart is not found to be great; and, on the other hand, life, and even a tolerable degree of health, may be maintained with valves so rigid and unyielding that the point of the knife can, at the autopsy, hardly be forced through them. In mitral disease the patient is liable to be worn out by the dropsy and by the increasing difficulty of breathing; and so, too, in that still more serious lesion, —tricuspid regurgitation. In affections of the aortic valves the patient suffers less, but he is more liable to sudden death.

Before dismissing these valvular affections, there are a few other matters which claim consideration, though the limits set to this work will prevent their full discussion. The blowing sound has been insisted upon as the diagnostic sign of a valvular lesion, and to insist upon this is to do no more than universal experience warrants. But there are undoubtedly instances in which no murmur reaches the ear to show that the valves are damaged.

I shall cite two examples. A man, thirty-five years of age, came under my care, complaining of palpitation of the heart, of occasional attacks of bronchitis, and of shortness of breath. His health was otherwise good. A physical examination of the chest showed the action of the heart to be extremely disturbed: the impulse was strong, and the extent of dulness in the præcordial region increased. A blowing sound was heard near the apex, but, owing to the great irregularity of the movements of the heart, it was impossible to say whether it corresponded in time to the contraction or to the relaxation of the organ. The pulse was small, frequent, and intermittent. The patient continued in this state for seven months, the beat of the heart becoming more and more tumultuous; but the murmur gradually disappeared. A peculiar clacking sound took its place, which was most distinct near the apex, and was faintly transmitted to other portions of the heart. It occurred with but one sound of the heart,—with which could



not be determined. For some time before his death he had considerable cough, with a frothy expectoration and great difficulty in breathing. His face and hands had begun to swell. The immediate cause of death was pulmonary apoplexy. The heart was found in a state of dilated hypertrophy, and the mitral valves had been converted into a calcareous mass, which had left but an extremely narrow chink for the blood to pass through.

The next case presents, in several respects, a striking similarity. A gentleman, about fifty years of age, who had led a gay and somewhat dissipated life, noticed that he experienced difficulty in breathing on the slightest exertion. He complained also much of loss of appetite and of distention of the stomach. I could not find any cause beyond flatulence to account for this. But to the dyspnoea, an inquiry into the state of the heart furnished a clue. The size of the organ was evidently augmented, and its rhythm very irregular. The impulse was strong; but the sounds were normal, except near the apex, where, taking the place of one, was heard a dull but very marked clack. When the hand was applied over this point, it felt a vibration of very much the same character as that which the ear could hear, and, like this, it was limited, or certainly only distinctly perceptible, at or near the apex of the organ. The diagnosis of disease of the mitral valves was made, and it proved to be correct. The dyspnoea became greater and greater; the feet, and subsequently the abdomen, were distended with fluid; and the patient died with all the symptoms of an unmistakable valvular lesion.

My note-book would furnish me with many more such cases; but these two present the main features of all. All the instances of valvular disease I have met with, unaccompanied by blowing sounds, have been instances of disease at the mitral orifice, and of extreme narrowing of that orifice. They were all attended with excessive irregularity of the action of the heart, and with hypertrophy. They all produced difficulty of breathing. They all presented the peculiar clacking sound most marked near the apex. In some, another sound, more like that heard in health, followed it; in others, not. In some, the blowing sound gradually disappeared; in others, none was perceived when first examined; and in others, again, it could be caught occasionally, as a very short whiff, along with the clacking sound. In all, the impulse was

strong and very variable in its rhythm, and a peculiar movement was felt near the seat of the apex,—not the purring tremor which so commonly accompanies the movements of a heart the valves of which are damaged, but a more localized vibration, similar, as far as such similarity can exist, to the sound the ear hears.

These cases are probably of the same nature as those that are every now and then reported as valvular lesions in which the sounds of the heart were normal. I cannot think that with a disease of the valves they ever are so. There may be no blowing sounds present, but the sounds of the valve affected must be different from what they are in health; and it may again, in all truth, be said that to hear the natural sounds of the heart well defined is to be able to exclude a valvular disease.

The other subject to which we may advert is the possibility of valves having been insufficient to perform their functions during life, and yet no signs of their incompetence being detected after death, at least none being indicated by any structural change in the valves. That such cases occur, is attested by more than one observer. In explaining them we must take into account those blowing sounds which are produced by mere abnormal action of the structures of the heart, the functional murmurs above described, and which may occur in hearts of healthy texture or in states of hypertrophy or dilatation.

Valvular disease may be at times suddenly developed, from *rupture of a valvulet or of a papillary muscle* by a severe strain. I have known such cases to happen where there was nothing in the history to lead to the belief of previous disease, though often there is some preceding disorganization, such as a granular or a fatty change. One of the most striking diagnostic features is the quickly-originating organic murmur attending the signs of disordered circulation and cardiac distress; another, the occurrence of pain in the region of the heart. Rupture may happen in the affected valve of an ulcerative endocarditis without any extraordinary strain. The previous history, the sudden aggravation of the cardiac symptoms, may furnish an explanation of the accident.

Let me also here briefly discuss another question,—whether the valvular affection shows any signs by which we can recognize it *before the development of a murmur*. We cannot do so with any certainty; although marked alteration—such as dulness of sound

confined to or most obvious at a particular valve, the signs of preceding or of growing hypertrophy, and, where the aortic valves are concerned, a distinct accentuation of the second sound, while the first has become dull and changed—might make us suspect what is about to happen. A doubling of either the first or the second sound, especially of the latter, is often, according to the observations of Sansom, an early sign of the development of mitral narrowing.

### Displacements of the Heart.

The heart is a very movable organ. Its apex is tilted upward by an enlarged liver, by an abdominal tumor, or by a pericardial effusion. It gravitates toward the median line when the walls of the heart have increased in weight and firmness. But these changes are hardly of a nature to attract as much attention as finding a heart beating on the right side of the sternum.

Now, it is not very uncommon to meet with it there; and the question immediately arises, What does this strange alteration in its situation signify, and how is it brought about? It is usually produced by pressure exercised on the heart by accumulations of fluid or of air in the left pleural cavity, and therefore denotes, as a rule, a pleuritic effusion or a pneumothorax of the left side, and is accompanied by distention of that side. In rarer instances, the heart is pushed across by a highly-distended emphysematous lung; in still rarer instances, it is drawn over to the right side by a shrinking of the lung, attended with dilatation of the bronchial tubes, the so-called pulmonary cirrhosis. It is sometimes found on the right side, because it had been forced there by a pleuritic effusion and had formed adhesions, so that when the fluid was absorbed it was unable to return to its natural place. In this case the left side will be markedly retracted, and not the right, as it is if cirrhosis of the right lung be the cause of the abnormal position of the heart.

The displacement may further have been brought about by a cancerous or an aneurismal tumor, or by any of the abdominal viscera having slipped into the chest through a hernial opening in the diaphragm; or it may be congenital. But these all are causes which seldom exist. Practically speaking, transpositions

of the heart are met with in connection with diseases of the lungs. We shall merely add that a congenital displacement cannot be diagnosticated unless all other causes capable of producing a displacement have been proved to be absent; and that a dislocated heart is able to perform all its functions. It may even be attacked by acute disease; the recognition of which,\* under such circumstances, belongs to the triumphs of physical diagnosis.



### SECTION III.

#### THORACIC ANEURISM.

An aneurism of the aorta, whether caused by a disease of the coats of the artery or not, whether true or false, may affect any part of the vessel. Yet it is chiefly at the ascending portion and at the arch that it is met with. When it occurs just after the artery has left the heart, it is prone to elude discovery. Higher up, nearer to or at the arch, it more rarely escapes detection. The tumor manifests itself by a local bulging, varying in extent and situation according to the extent and situation of the aneurism. A single rib alone may be raised, or nothing but a fulness may be observed. But some prominent spot is generally detected, and when this is percussed it is more resistant, and returns a duller sound, than normal. Yet neither the bulging nor the dulness on percussion is of as much significance as finding a distinct pulsation remote from the beat of the heart. Every time the latter is perceived, an impulse is communicated to the finger at the point in the chest-walls which appears to project; that is, usually on the right side of the sternum in the second intercostal space, or in the same interspace on the left side, or immediately under the top of the bone. Occasionally the beat is double, at times so violent as to shake the head of the listener, and almost always, unless the aneurism be filled with solid clots, stronger than the beat of the heart.

---

\* As by Stokes. See Diseases of the Heart, p. 463.



The impulse may be accompanied by a distinct thrill. But this is not always present, and, when present, is not always constant; since it may disappear and reappear. It is thus a serious mistake to regard the thrill as the requisite sign of an aneurismal enlargement; yet there is no mistake more common, except, perhaps, one,—to consider that the motion of the blood in the sac must necessarily engender a murmur. The ear, applied over the prominence, hears often nothing that in the least resembles a murmur, but sounds like those of the heart, sometimes two, the first weighty and prolonged; sometimes but one, and that one longer and more intense than the corresponding first sound over the ventricles.

Thus, then, neither thrill nor murmur is essential to the diagnosis of an aneurism. What is much more essential, is to find two points of pulsation in the chest,—two hearts, apparently, each with its own distinct beat, its own distinct sounds.

An aneurismal tumor in the chest gives rise to symptoms which vary somewhat according to its seat and size. Prominent among them stand those occasioned by pressure. The sac presses on the adjacent air-tubes, and shortness of breathing, or peculiar cough and signs counterfeiting those of a chronic laryngeal disease, are the result; or it presses on the œsophagus, and the patient suffers from difficulty in swallowing; or it presses on the subclavian artery, and the pulses at the two wrists are noticed to be strikingly different, both in volume and in time; or on the carotid, and pain in the head, dulness of mind, occasional giddiness, and flashes of light before the eyes, are complained of; or on the venous trunks, and the superficial veins of the neck and thorax are seen to be engorged, and the skin becomes very puffy and swollen; or on the trunk of the sympathetic nerve or on its ganglia and their communications, and marked contraction, or, in rarer instances, dilatation, of the pupil of the eye on the side of the aneurismal swelling, is perceived, or profuse sweating becomes an annoying complication. All these signs, then, denote pressure, and pressure connected with a pulsating tumor in the chest means an aneurism.

I say with a pulsating tumor, because a cancerous or other *intra-thoracic morbid growth* may produce exactly the same signs of compression as an aneurismal tumor,—the same stridor, the same cough, the same feebleness of respiration in one lung from partial

obliteration of its bronchial tube. But the solid tumor, large though it be, does not pulsate, or, if it do, pulsates but very feebly, and not with the heaving motion of a distending aneurismal sac. The tumor, which for the most part has its seat in the mediastinum, renders a large surface dull on percussion, and communicates a much greater feeling of resistance to the percussing finger. Yet the ear listens in vain over the prominence for the weighty sound with each beat of the heart, or for the hoarse murmur of the blood streaming through the sac. It is only where a solid growth presses on the artery that any murmur is perceived; and this is different from the superficial loud sounds or murmurs of an aneurism. Further, a tumor is not confined to the course of the aorta; it is more commonly connected with a distended state of the veins of the neck and thorax, and with œdema of the arm and chest; the pain it occasions is often more continued, and less neuralgic in its nature, and the dyspnœa is not infrequently paroxysmal. Moreover, as most thoracic tumors are cancerous, the violent constitutional disturbance, the formation of external swellings, the enlargement of the glands in the axilla and the neck, and the peculiar currant-jelly expectoration, aid us in arriving at a correct conclusion. Sarcoma, lymphomata, and lymphadenomata of the mediastinum come next in frequency to cancer.\* They all tend to grow inward rather than outward, and affect the anterior mediastinum far oftener than the other two spaces. Then the history is of some value in the diagnosis. In aneurism it points to gout, to aortitis, to atheroma, to alcoholism, to syphilis, to strain, to an embolic infarct, to infectious arteritis from mycotic invasion of the aorta.

The most difficult diagnosis—it is often an impossible one—is between an aneurism filled with solid clots and a tumor. The history of the case is here of the greatest importance; and there is generally less pain in these altered aneurisms than in tumor. The physical signs will not help us. As a rule, it may be said that we do not find in the latter the ringing second aortic sound, or the shock with this, which happens in aneurism,—happens mostly even when it is filled with clots.

As regards *abscess of the mediastinum*, we do not find the

---

\* Hare, Mediastinal Disease, 1889.

pressure signs generally so marked as in aneurism, and we may be able to detect fluctuation at the edge of the sternum or at the supra-sternal notch. The pain is usually very great; the elevation of temperature is significant. The sounds over the mass are not those of an aneurismal sac; there are certainly no distinctive murmurs, and we find no marked expansile pulsation. This absence of distinct pulsation was the main point of dissimilarity between an aneurism and an abscess of the mediastinum some time since under my care, which, after lasting a year, and simulating aneurism most closely in the pain, the dulness on percussion, the difficulty of breathing and of swallowing, and the altered voice,—having, therefore, pressure signs much more marked than usual,—got well by breaking internally and by the discharge, as expectoration, of large amounts of purulent matter. In inflammatory thickening of the mediastinum the impulse of the heart is weak and the sounds are faint.\*

The obvious inequality of the pupils, which is found in a certain number of cases among the signs of an aneurism, is of little aid in a differential diagnosis from intra-thoracic tumor, for a thoracic cancer has been noted to occasion the same.† The rarity of a non-aneurismal tumor in the chest is, however, very great; and, practically speaking, when the signs of intra-thoracic tumor are met with we shall be generally correct in thinking that it is an aneurism we have to treat, even should the pulsations not be very obvious.

The sphygmograph will at times aid us in the diagnosis of an aneurism, though its value is not on the whole great. Its most distinct significance is in showing clearly the difference between the two pulses. Of one radial the sphygmogram may be normal; the other tracing furnishes what is a characteristic record,—a sloping up-stroke, a rounded apex, an obliteration of the secondary curves.

Another sign of aneurism which has been of late much studied, and especially by MacDonnell,‡ is the so-called tracheal tugging. To obtain it the cricoid cartilage is firmly grasped and

---

\* Wilson Fox, *Treatise on Diseases of the Lungs*.

† MacDonnell, *Montreal Medical Chronicle*, June, 1858; Gairdner, *Clinical Medicine*, and Ogle, *Medico-Chirurgical Transactions*, vol. xli.

‡ *Lancet*, March, 1891.

the trachea put on the stretch by pressing upward. If an aneurism is adherent to it or near it, a significant tugging will be felt. In deep-seated aneurisms this sign is of special value, but it is not absolutely characteristic. It has been found by Grimsdale\* and by Ewart† in other conditions, and even in healthy subjects.

Let us suppose that we are satisfied, owing to a marked impulse, that we have not a solid growth or an abscess to deal with,—does a pulsation uniformly denote an aneurism? Can we say, on account of the impulse, that it is an aneurismal enlargement? If there be also swelling and signs of pressure, we can; should these not exist, we cannot be so sure. For a pulsation in the chest not immediately over the region of the heart, although it is nearly always indicative of an aneurism, may be owing to other causes. Where the aortic valves are insufficient, there may be a pulsation in the aorta; an empyema may pulsate; a dilated auricle may occasion an impulse separate from that of the ventricles; a pulmonary artery surrounded by consolidated lung may distinctly exhibit its beat. In all of these the signs of pressure on the surrounding parts are wanting; and, on the other hand, they show phenomena which an aneurism lacks.

*Insufficient aortic valves* are accompanied by hypertrophy of the left ventricle. So is at times a thoracic aneurism; but, instead of the throbbing at the upper anterior part of the chest being attended, as in aneurismal swelling, with a natural or with an unequal beat at the wrist, there, as well as in the larger trunks in the neck and arms, is perceived that strong and peculiar pulsation which is so characteristic of inadequate aortic valves. Then, again, a murmur is much more common in this affection of the valves than it is in aortic aneurism; and is usually a loud double murmur, most distinct at the right base of the heart, and associated with a double murmur in the femorals made evident by pressure with the stethoscope. This is very rare in aneurism of the aorta; moreover, the murmur heard over an aneurismal pulsation is better marked over its seat than over the heart, and is mostly single, systolic and short, hoarse and of low pitch. In truth, it differs in

---

\* Practitioner, London, Feb. 1892.

† Brit. Med. Journ., March, 1892.



distinctness as well as in quality from the murmur discerned at the base of the heart, which may be transmitted from the aneurism, or may depend upon coexisting cardiac disease. Then the sphygmographic tracings may also be of some value. Those of aortic regurgitation, as above described, are for the most part characteristic; while an oblique line of ascent, a loss of the summit wave, and a modification of the diastolic wave are usual when an aneurism is seated on a main trunk after its origin from the aorta. When the aorta is dilated, as well as its valves diseased, the diagnosis as regards aneurism is much more difficult. But even then we lack the distinct throbbing, the signs of pressure, and the unequal pulses.

While the diagnosis of aortic valve disease is under review, *coarctation or constriction of the aorta*, which in very rare cases is associated with the valvular affection, may be mentioned. It generally happens just at or below the insertion of the ductus arteriosus, and furnishes as its only special signs a dilatation of certain collateral vessels at the upper part of the thorax, and diminished size and feeble, retarded pulsation of the femorals. The arteries of the head and neck, as well as the epigastric and mammary arteries, throb, and there may be a marked thrill at the upper part of the chest near the sternum, and a murmur there louder than over the heart; pressure signs are absent, and the dilated vessels are often the seat of a purring noise.

A *pulsating empyema* is seldom met with; yet a collection of fluid in the cavity of the chest may vibrate with the motion of the heart, and throb with such distinctness as closely to simulate an aneurism. To determine the real nature of the pulsation in these cases, we must attach importance to the situation of the expanding mass, which is not often that of an aneurism, and to the signs which point out that liquid has accumulated within the pleural sac. We also note the circumstance that over the seat of impulse there are no peculiarly marked sounds, no murmurs, no thrill; moreover, the beat, which is wide-spread, is not apt to be so strong as that of the heart, which is displaced. The pulsation may happen both in acute and in chronic pleurisy, and be associated, as in Osler's \* case, with persistent tenderness of the thoracic walls.

---

\* Amer. Journ. Med. Sci., Jan. 1889.

There may be a number of these pulsating tumors.\* Pulsating pleurisies are nearly always left-sided and purulent; there is generally latent pneumothorax present.† In one of Wilson's cases‡ the pulsation disappeared immediately after aspiration.

A *dilated auricle*, the walls of which are at the same time hypertrophied, may give rise to a movement separate from that of the beat of the ventricle. Bouillaud cites an example of this nature, in which a double motion was perceptible in the second intercostal space of the left side, in a person whose heart was extensively hypertrophied and whose mitral valves were indurated. Such cases are extremely rare. The signs of an accompanying valvular affection and of enlargement of the ventricles, and the probable presence of dropsy, would serve to distinguish a dilated auricle from aneurism of the arch. And this is the only form of enlargement of the heart which is likely to be mistaken for an aneurism. In cases of hypertrophy or dilatation as we ordinarily meet with them, there is but one motion discernible,—that over the ventricles,—and not two beats at some distance from each other; the signs of pressure, too, are wanting. In dilatation of the right auricle, Sansom notes a vibration to the right of the sternum and a wedge-shaped line of dulness joining the dulness of the liver.

A *pulmonary artery* surrounded by consolidated lung-tissue may cause—especially if the vessel be somewhat widened—a distinct pulsation. But the seat of the dulness near the apex of the left lung; its non-extension over the median line; the limitation of the murmur to the site of the pulmonary artery, or, in some instances, to this vessel and the subclavian; the sharply-defined second sound of the pulmonary artery in the second interspace on the left side; the symptoms and physical signs of phthisis, the most common cause of the consolidation, and a morbid condition which of itself would appear to exclude an aneurism; the absence of pain and of the phenomena caused by pressure,—all these prove the murmur and the pulsation not to be due to an aortic aneurism. Absence of pain and of pressure signs, and accen-

---

\* Henry, *Proceed. Phila. Co. Med. Soc.*, vol. iii.

† Comby, *Arch. Gén. de Méd.*, April, 1889.

‡ *Transact. of Assoc. of Amer. Physicians*, 1893.

tuation of the second sound, are also the chief signs by which we distinguish those comparatively rare cases of murmur in the second interspace, close to the left of the sternum, which are due to retraction of the lung and uncovering of the heart and pulmonary artery. The murmur, which has been specially studied by Quincke\* and Balfour,† is systolic and loud, and mostly disappears on deep inspiration. The pulsation is marked, though not so strong as that of the heart; the singular murmur is supposed to be owing to compression of the pulmonary artery by the heart during the systole. In many respects it is like the murmur, which I have elsewhere investigated,‡ heard over the pulmonary artery in certain lung affections.

Another abnormal condition which may be mistaken for an aneurism is a *malformation of the chest*, particularly when produced by great prominence of the upper part of the sternum. This error is more especially apt to occur if there be coexisting disturbance of the heart, whether of functional or of organic origin. I have seen cases where the beating of the arteries of the neck, accompanied by an enlargement of the thyroid gland and by cardiac palpitation, was believed to be an aneurism, mainly because it was combined with very decided prominence of the upper portion of the sternum. But there were no distinctly localized tumefaction and pulsation, no altered sounds, no signs of pressure. I have also met with instances in which the active pulsation of the thyroid gland, both in exophthalmic and in ordinary goitre, gave rise to the idea of an aneurism, but in which no change of the chest-walls existed. In such cases the carotids and radials beat equally; a blowing murmur, attended by a continuous hum, is heard—certainly in instances of exophthalmic goitre—over the enlarged gland; there is nowhere a point of localized pulsation, and there are also no signs of pressure.

*Malposition of the aorta*, due to rickets, may simulate an aneurism closely. Balfour§ has pointed out how misleading may be the abnormal pulsation with the dulness on percussion, and the right-sided prominence of the chest. Moreover, thrill, murmurs

\* Berliner Klinische Wochenschrift, 1870.

† Lectures on Diseases of the Heart, London, 1876.

‡ Amer. Journ. Med. Sci., Jan. 1859.

§ Diseases of the Heart, London, 1876.

loudest over the pulsating mass, and cardiac hypertrophy, may coexist. We must be guided in our opinion by the history of the case; by the distortion of the spine; by the extended superficial dulness on percussion, out of proportion to the extent and strength of the pulsation of the tumor, which is less forcible than that of the heart; by the displaced position of the heart, which is tilted upward and thrown over more to the right; and especially by the absence of any signs of pressure.

The signs of pressure play, then, a very important part in the diagnosis of an aneurism. They are rarely absent, although they do not always manifest themselves in the same manner: sometimes it is bone, sometimes lung, sometimes œsophagus, sometimes nerve-fibre, which bears the brunt of the distending swelling. These signs of pressure are wanting if the sac be very small or be absent; or not prominent if the artery be simply dilated, in which case nothing but a constantly pulsating tumor can be detected. At times evidences of compression may be recognized by the attentive physician when no throbbing swelling can be discerned; and from them he infers the true nature of the case, although utterly unable to discover any of the ordinary physical signs of an aneurism. Whenever, indeed, obstinate and anomalous thoracic symptoms, which might be explained by the presence of an aneurismal sac, occur in a person whose lungs and heart appear to be in every respect sound and whose general health is not materially affected, we may suspect an aneurism to be the source of the disorder. So, too, if any laryngeal affection, or if any difficulty in swallowing, exhibit rather peculiar symptoms. It is, in truth, imperative in all cases of chronic disease of the larynx, or where there are indications of a stricture of the œsophagus, to examine the chest carefully, so as to avoid the grave error of overlooking what may be the only cause of the whole disturbance.

The symptoms of *chronic laryngitis* especially are at times astonishingly simulated, and it may happen that the patient, trusting to his feelings, refers obstinately to the chest as the seat of the disorder, while the physician as obstinately sees nothing but the presumed affection of the larynx. There is, as in chronic laryngeal disease, alteration of the voice, with stridor, and cough; but the voice is not so uniformly changed. Often it retains much of



its natural character; and the loss is not so progressive, and the aphonia not so permanent. Hoarse the voice may be, but, as the direction of the pressure varies, it alters rapidly both in pitch and in power. The cough is most commonly loud and paroxysmal, and has a ringing sound. Dyspnoea is a very constant symptom; it may be due to laryngeal disorder,—spasm or palsy of the muscles caused by pressure on the recurrent laryngeal nerve,—or to compression of the trachea and a large bronchus. It is often attended with wheezing or stridulous breathing, which is not persistent, and is sometimes produced only after a deep inspiration. The stridor, however, as Stokes points out, differs from that of an obstructive disease of the larynx by its seeming to issue from the notch at the sternum, and not from above, from the larynx itself. If, in addition, the respiration be found to be markedly unequal in the two lungs, the diagnosis of aneurism may be ventured upon; and it will be confirmed by finding no change in the larynx, when examined with the laryngoscope, sufficient to account for the laryngeal symptoms, or such a change—paralysis of only one cord, for instance, or paralysis of an abductor on one side—as could be readily explained by pressure on one recurrent nerve.\* Of course, the detection of dulness on percussion, of sounds stronger than or otherwise different from those in the cardiac region, or the occurrence of a hemorrhage, would place the diagnosis beyond doubt. A systolic sound or thud in the brachial artery is also a sign to which importance may be attached.†

In some cases of aneurism, pain is among the earliest symptoms, and the patient complains much of it before there is a single physical sign indicative of the presence of a tumor. The pain is dependent upon pressure on the nervous filaments: it may shoot toward the shoulder or the neck, along the arm, or deep into the centre of the chest. Dull, deep pain, boring and constant, occurs when the pressure of the sac is leading to absorption of the vertebræ. Over the seat of the swelling there is often pain, associated with great tenderness.

The severity of the pain may give rise to emaciation and ex-

---

\* In the chapter on Diseases of the Larynx, the forms of laryngeal palsy from an aneurism have been more specially examined into.

† Glasgow, New York Med. Journ., Sept. 1894.

haustion, and become a cause of death ; but death does not often take place from exhaustion. More usually the patient's life is cut short by the aneurism bursting, either externally or into internal parts,—into the trachea, bronchial tubes, œsophagus, pericardium, pleura, pulmonary artery, or spinal canal. Yet it is not always the first rent which leads to the fatal issue ; this, as we learn from the cases that Webb\* has analyzed, may, when the aneurism breaks externally, not happen for weeks after the accident.

Now, can we foretell the course of an aneurism, and the probable mode of death it is likely to occasion ? We cannot ; for in order to do so it would be requisite to determine accurately its seat, so as to know what tissues are likely to be encroached upon. And this is very difficult, nay, often impossible. It is true that, when the swelling gives rise to phenomena like those of angina pectoris, we may surmise it to be in the ascending portion of the aorta and near the cardiac plexus of nerves, and look for its breaking into the pericardium or the pulmonary artery ; when it is accompanied by laryngeal stridor or other laryngeal symptoms, it probably involves the posterior and lower portions of the arch, and will cause death by strangulation or by exhaustion ; when it produces much dyspnoea, it is apt to be seated in the descending part of the arch, and death may take place by the aneurism bursting into a bronchial tube, or by pneumonia. But in regard to all these matters we can usually do little else than conjecture ; because a tumor within the chest leads to such displacements that its relations to the surrounding structures cannot be clearly ascertained during life. The most valuable information we obtain is from a study of the physiological changes,—from the symptoms, therefore, of disturbed function.

An aneurism of the *descending aorta*, between the arch and the diaphragm, produces, if extensive, dulness on percussion and bulging posteriorly, and may exhibit the same physical signs and symptoms as an aneurism in the neighborhood of the arch. A gnawing sensation in the vertebræ has been especially noticed. Yet, in spite of the most careful scrutiny, an aneurism of the descending aorta often escapes detection, or its physical signs,

---

\* Amer. Journ. Med. Sci., Oct. 1874.

as a case recorded by Walshe\* proves, may exist to the right instead of to the left of the spinal column, because the vessel has been dragged across the median line by its enlargement.

An *aneurism of the heart* may in exceptional instances produce localized bulging in the cardiac region. But, whether it does so or not, it is beyond the reach of positive diagnosis. We may suspect it if the bulging have been preceded by signs of fibroid degeneration of the walls of the heart. Obstructed coronaries producing the myocardial changes are its most common cause. Pericarditis with adhesions near the aneurism has been also noticed. In a number of instances we have a syphilitic history.

In rare instances we find a varicose aneurism communicating with either the ascending or the descending vena cava. These aneurisms mostly present the ordinary signs of a thoracic aneurism; but, in addition, great venous enlargement above the diaphragm, with œdema of the face and hands and arms; a purple hue of the face and the upper part of the body, and spots of ecchymosis in the skin; a jerking pulse; a purring thrill; and a whirring systolic murmur,† diffused all over the front of the chest. The œdema and the symptoms of venous disturbance come on suddenly. In instances of occlusion of the vena cava the great venous distention is not accompanied by the physical signs of an aneurism, nor by the thrill, nor by the cyanosis and œdematous swelling.‡

Let us, in conclusion, glance at the other kinds of aneurism within the thorax,—that of the innominate and that of the pulmonary artery.

An aneurism of the *innominate artery* is strictly limited to the right side of the body. It differs from that of the arch by the higher situation of the pulsating swelling; by the displacement of the clavicle; by the comparative absence of signs of pressure on the larynx and œsophagus; and by the fact that compression of the right subclavian and carotid diminishes the beat of the tumor, while it exerts no effect on an aortic aneurism. Such are, at all events, the marks of distinction indicated by the observations in

---

\* Diseases of the Heart.

† As in Mayne's case, Dublin Quart. Journ. of Med. Sci., Nov. 1853; also in Glasgow's case, St. Louis Courier of Med., Jan. 1885.

‡ Arthur V. Meigs's case, Transact. Coll. of Phys. of Phila., 1886.

Holland's \* excellent memoir. An additional sign is mentioned by Wardrop.† It is that when the innominate is affected the difficulty will appear first on the tracheal side of the sterno-mastoid ; but on the cervical side if the aneurism be of the subclavian. In aneurism of the innominate, further, as the tumor is under the right sterno-articular articulation, percussion does not detect any distinct enlargement of the arch of the aorta.

An aneurism of the *pulmonary artery* is a rare disease. Its main phenomena are : a strongly pulsating swelling, perceptible to the left of the sternum, and limited to the second intercostal space near the costal cartilages ; a marked thrill with each expansion of the aneurism ; and in some instances a rough murmur, which is not discovered at the notch of the sternum or above the clavicles ; lividity of the face ; dropsy ; and great difficulty of breathing.‡ The most significant points of difference between an aneurism of the pulmonary artery and an aneurism of the aorta consist in the symptoms just mentioned, and in the absence of obvious evidences of pressure. The situation, too, of the physical phenomena is important ; yet we must bear in mind that an aneurism of the arch may occasion a pulsating tumor, mainly to the left of the sternum, and may even break into the pulmonary artery. A mere distinct beating of the pulmonary artery is discriminated from an aneurism of this vessel by the non-existence of a palpable swelling, of dropsy, of greatly-embarrassed breathing, of lividity of the face, and by the usually coexisting signs of some consolidation of the left lung.

Occasionally we meet under the outer half of the left clavicle with a pulsating tumor presenting thrill and murmur, and dilated veins above. These signs may be supposed to indicate a subclavian or axillary aneurism ; but they often suddenly disappear. These "mimic" or phantom aneurisms§ are apt to come back after excitement and after movement of the arms. They are thought to be due to temporary dilatation of the artery from vaso-motor paralysis, limited to a large vessel or to part of it.

---

\* Dublin Quarterly Journal, vol. xii.

† Holmes's Surgery, vol. iii. p. 562.

‡ In the case detailed by Skoda, Auscultation and Percussion, the dropsy was very great, and the face cyanotic ; there was a faint murmur over the base of the heart, but none over the pulmonary artery.

§ See paper by Samuel West, St Barthol. Hosp. Rep., 1880.



## CHAPTER V.

### DISEASES OF THE MOUTH, PHARYNX, AND ŒSOPHAGUS.

THE diseases of this part of the digestive system need not here be described at any length, because many of them have already been considered in treating of the affections of the larynx and of the heart and great vessels. But with the maladies of this part of the body may be considered the enlargement of glands at the angle of the jaw, as happens in mumps.

#### MOUTH.

Soreness of the mouth, pain in masticating, and a fetid breath are often complained of in diseases of the oral cavity. Let us suppose a patient to present himself with such symptoms. The interior of the mouth is exposed to a strong light, and its different parts are inspected.

*The gums are noticed to be swollen and injected, and the mucous membrane lining the cheeks reddened.*—This is a state of things observed in the different forms of stomatitis. In the *common diffused inflammation*, the result of direct irritation, such as of the swallowing of hot liquids or of corrosive substances, or an accompaniment and consequence of gastric disorder, the redness is marked; any attempt at chewing is painful; the taste is impaired; a flow of saliva takes place from the mouth, and superficial ulcerations occur at its various parts. In *mercurial stomatitis* there are much the same symptoms; but the more copious discharge of saliva, the pain in the jaws, the loosening of the teeth, the enlarged tongue, exhibiting their impress, the painful and swollen state of the salivary glands, and the peculiar nauseous breath, testify to the specific character of the inflammation. The sore mouth of scurvy may be distinguished from either of the preceding forms by the spongy, purplish, or livid gums, which bleed on the slightest

touch, by the eruption on the skin, and by the other signs which attend a scorbutic state.

*The gums and the inside of the cheeks and lips are covered with a whitish curd-like exudation.*—This constitutes the form of stomatitis known as *thrush*, so frequent in infants at the breast, and so constantly associated with intestinal disorder, with diarrhœa, with colicky pains, and with a feverish heat of skin and a hot, dry mouth. Very similar to it, regarded indeed by some as identical, is the aphthous ulceration, to which adults as well as children are liable. Here, too, a whitish deposit is perceived in various parts of the mouth; it is apt also to be combined with thirst and with gastric or intestinal disturbance, and the breath has a very disagreeable odor. The recognized difference consists in the presence of the small ulcers which may be detected when the white crusts that cover them are removed, and in the vesicular nature of the disease during its formative stage. Then more or less redness surrounds each spot, the ulcers are slightly raised at their borders, bleed easily on pressure, and may be irregular from several running together; their grayish covering is found to be soluble in ether, and to present many oil-globules under the microscope. On the other hand, the microscope shows us in thrush a special parasitic formation, the *Oidium*, or *Mycoderma*, *albicans*.

*Ulcerations are perceived on the gums, tongue, and various parts of the mouth.*—We meet with ulcers in the ordinary, in the mercurial, in the scorbutic, and in the aphthous inflammation of the mouth. But ulceration is apt to exhibit its most horrible features in the sore mouth of syphilis, and in that essentially ulcerative disease called *cancerum oris*, or gangrenous stomatitis. In the former the fauces as well as the mouth are, as a rule, involved, and the ulcers show peculiarities which we shall presently study. The latter is an affection which prevails especially in enfeebled constitutions. It is seen chiefly in hospitals, and not uncommonly in epidemics. It begins with pain in the gums, and these soon swell, redden, and bleed readily. They are covered with a soft, grayish exudation, which often extends to the soft palate. If the layer of exudation be scraped away, a bleeding, ulcerated mucous membrane comes into view, provided the swelling be not so great as to render a careful examination of the mouth impossible. The breath is most offensive; a profuse flow of saliva is noticeable;

perforation of the cheek quickly takes place ; the bones may be laid bare, the teeth loosened ; there is usually fever, often of hectic type ; yet the disease does not uniformly progress with activity ; it may last for weeks.

*The tongue is red and swollen.*—Changes in color and in appearance of the tongue occur in general diseases of the system, and more especially in those of the alimentary canal. The tongue is also more or less involved, at all events its mucous membrane is, in the different forms of stomatitis. An abnormal state of the covering of the tongue is, therefore, far from being a sign that the organ itself is primarily affected.

Occasionally, however, we do meet with affections of its deeper structures. Its nerves may be the seat of violent neuralgia ; its muscles may be paralyzed ; it may become hypertrophied or cancerous ; it may undergo progressive atrophy ; or it may be in a state of *acute inflammation*. The latter is, perhaps, the most frequent of its maladies, and is readily recognized by the red, swollen look of the organ, joined to a burning pain in it, and either to great dryness of the mouth or to constant dribbling. The swelling usually begins at the anterior portion, and may become so considerable as to threaten suffocation ; the inflamed tongue fills up the fauces and protrudes out of the mouth, and the unhappy patient can neither swallow nor utter a word. He has active fever, headache, great restlessness, and intense thirst,—symptoms which last for several days, and until the inflammation subsides. This may run on to suppuration or gangrene ; in some instances it leaves a permanent induration that may be mistaken for a cancerous nodule. Acute glossitis is a dangerous complaint ; fortunately, it is a rare one. Its most frequent cause, as now seen, is direct injury, either from wounds or the stings of venomous insects, or from the introduction of corrosive substances into the mouth. Its most frequent cause formerly was the abuse of mercury pushed to salivation. At times it is observed as a complication of scarlatina or of erysipelas.

Other affections of the tongue connected with diseases of its structure have been mentioned in the first part of this volume. *Cancer* of the tongue produces the greatest alteration in the form and texture of the organ. *Syphilis* of the tongue gives rise to deep fissures, ulcers, and gummous nodules which may be difficult

to distinguish from cancer, except by the history and the absence of pain. As a sign of recovery from syphilis, the tongue may present a quite peculiar indented appearance, similar to what is seen in the syphilitic liver.

### FAUCES.

The fauces—that is, the parts at the back of the mouth which are brought into view when the lips are widely opened, such as the half-arches, the uvula, the tonsils, the posterior wall of the pharynx—may be involved in the same diseases as the parts situated in front. The contiguity of these structures is in fact such that any morbid action is apt to spread to them, or to extend from them either forward or downward into the pharynx, and even into the larynx. Moreover, on this very account a disorder is rarely found limited to any one portion of the fauces, but transfers itself generally from one to the other, from the tonsils to the soft palate, from the soft palate to the tonsils. The most common affections of the fauces are inflammation and ulceration, both of which occasion a feeling of uneasiness in the throat, and also difficulty or pain in deglutition, and both of which are readily enough detected by looking into the mouth when the jaws are widely separated and the tongue depressed.

In the ordinary inflammation of the fauces, the *simple angina*, or sore throat, the parts are of a bright-red color, and the uvula is long and swollen, and by dropping on the tongue gives rise to a constant disposition to swallow, although the act of swallowing is attended with pain. Associated with the angina are coryza and febrile disturbance; and, owing to the inflammation travelling up the Eustachian tube, the sense of hearing is impaired.

**Tonsillitis.**—When the inflammation penetrates the substance of the tonsils, or in *quinsy*, much the same general symptoms occur as in ordinary angina. But the sense of constriction in the throat is greater; so is the difficulty in swallowing; and liquids are apt to return through the nose. The voice is thick, and has often a peculiar sound; it is painful to the patient to talk, and on looking into the throat the tonsils may be seen red, prominent, and covered with mucus which is not easily detached. Sometimes the swelling is so considerable that the tumid glands fill up the



space between the half-arches and leave but a small interval for the passage of food or drink. In some instances we cannot separate the jaws sufficiently to get a view of the throat, and have to trust to the introduction of the finger to tell us the condition of the affected parts. Occasionally the inflammation extends from the tonsils to the salivary glands; the submaxillary and parotid glands swell, and ptyalism takes place. It is necessary to be aware of this fact; for, if a mercurial cathartic has been administered, the profuse flow of saliva might be incorrectly attributed to it.

There is not much likelihood of confounding this, a form of *secondary parotitis*, with mumps, in which an outward swelling, visible beneath the ear, is found, but not a swelling within the throat, and in which no real difficulty in swallowing occurs, except, perhaps, when the tumefaction is at its height, and then only for a short time.

Tonsillitis terminates by resolution or by the formation of pus. There are no positive means of ascertaining that the inflammation is going to end in suppuration, although we may suspect that this will be the case when much pain is felt at the angles of the jaws and shooting to the ear, and when the symptoms have been severe and persistent for more than four or five days. Sometimes the pus may be seen through the covering of the tonsils; but often the vast sense of relief experienced by the patient, and the sudden improvement in deglutition, attended, perhaps, with an unpleasant taste, are the only signs that the collection of pus has been discharged. Attacks of tonsillitis are prone to be repeated, and may lead to permanent enlargement and induration of the tonsils. The enlarged tonsils, attended as they frequently are with cervical glandular swellings, may be mistaken for *cancer of the tonsils*. But in this affection sanious offensive discharges from the mouth occur, and, whether the disease be encephaloid or round-cell sarcoma, it extends rapidly; the neighboring lymphatic glands are early involved, the palate and the pharynx become implicated, and hemorrhage is frequent, as are also difficulty of deglutition and attacks of suffocation.\* Acute tonsillitis may be

---

\* Poland, Brit. and For. Med.-Chir. Rev., April, 1872; Newman, Amer. Journ. Med Sci., May, 1892.

seen in connection with malaria.\* At times the tonsils become gangrenous.†

*Acute Follicular Tonsillitis.*—There is a form of acute tonsillitis that is limited to the follicles and has well-marked clinical features. It begins with chilly sensations, to which a moderate fever, rarely exceeding 103°, succeeds. After a few days the fever disappears, a slight evening rise remaining, and in a week from the onset the patient is quite convalescent, though weak. At the height of the malady a swelling of the cervical lymphatic glands is often observed. But the characteristic feature of the disease is in the tonsils. These are red and slightly tumefied, and a thin yellowish or whitish punctiform exudation is seen in the crypts and around the follicular openings. This comes away gradually; in some parts much sooner than in others, and for days after convalescence from the general symptoms the appearance is found in some follicles. When cast off, superficial ulcerations on the glands may be noticed. This form of tonsillitis, called by some diphtheritic sore throat, is infectious, and various micro-organisms have been found in the exudate, particularly streptococci, staphylococci, and pyogenes aureus.

**Diphtheria.**—There is another kind of affection of the fauces which, in accordance with the clinical classification followed in this work, may be considered here, notwithstanding its specific character,—membranous angina, or diphtheria. Recent research leads us, indeed, to believe that the malady is primarily a local one, dependent upon the lodgement and multiplication of a specific bacterium. The constitutional symptoms are to be attributed to the absorption and action of the toxic products generated at the site of infection.

The bacillus of diphtheria was discovered by Klebs, and more fully studied by Loeffler. It is about as long as the tubercle-bacillus, but nearly twice as thick. It is, as a rule, curved, but its form is variable. It has rounded extremities, which are sometimes club-shaped. It is non-motile; it does not form spores. It may be stained in cover-glass preparations with Loeffler's alkaline methylene-blue solution, which consists of thirty cubic


---

\* Chassaignac, New Orleans Med. and Surg. Journ., Oct. 1888.

† Cragin, New York Med. Journ., Sept. 1, 1888.

centimetres of a concentrated alcoholic solution of methylene-blue and one hundred cubic centimetres of a 1 : 10,000 solution of

FIG. 52.



Klebs-Loeffler bacilli, from specimens prepared by Dr. Coplin and Dr. Bevan.

Fig. I. Fresh culture upon glycerin agar-agar. Eye-piece IV, Beck; Objective  $\frac{1}{2}$  ol. im. Leitz.

Fig. II. Fresh culture upon blood-serum. Eye-piece IV., Beck; Objective  $\frac{1}{2}$  ol. im. Leitz. This is also the appearance when obtained directly from the throat and subjected to the same power.

Fig. III. Old culture upon blood-serum. Eye-piece IV., Beck; Objective  $\frac{1}{2}$  ol. im. Leitz.

potassium hydroxide. The organism grows best upon a culture-medium consisting of three parts of blood-serum and one part of a mixture of meat-infusion with one per cent. each of peptone and grape-sugar and one-half per cent. of sodium chloride. This is sterilized and at the same time solidified in test-tubes supported at an acute angle in a steam or hot-air sterilizer at a temperature a little below  $100^{\circ}$  C. The inoculation is made by means of a pledget of cotton wrapped on the end of a steel rod, and the culture-tube is kept in a thermostat at a temperature of about  $37^{\circ}$  C. In the course of from twelve to twenty-four hours, dense, white, opaque colonies develop, and examination of cover-glass preparations will disclose the presence of the characteristic bacillus.

Bacilli resembling true diphtheria-bacilli in appearance and in culture, but wanting in virulence, have been described as pseudo-diphtheria-bacilli; but it is believed that these are merely a modified form of diphtheria-bacilli.

The disease begins usually as sore throat, with redness and swelling of the arches of the palate, and of the tonsils. There is slight stiffness of the neck, the cervical and submaxillary glands of the jaw are enlarged and tender, and the subcutaneous tissues may become involved in the swelling. Within a period varying from a few hours to a few days, an exudation takes place

on the tonsils, the uvula, and the soft palate. This exudation is more or less extensive, generally tough, and of a white or grayish hue. It may show but little tendency to spread; or it may extend to the gums, and along the walls of the pharynx into the windpipe, sometimes even into the bronchial tubes and the lung structure. In some cases it passes upward into the nares, yet it may begin there or in the larynx simultaneously with its appearance in the throat.

The false membrane in the throat, once formed, darkens, wastes from the circumference toward the centre, and gradually disappears. But sometimes the coat becomes for a time thicker and thicker by the constant addition of fresh layers. This happens particularly in the "croupous form" of diphtheria, in which the inflammation is more intense from the onset and the fibrinous exudations succeed one another rapidly until the dense thick coating of false membrane results. Under any circumstances, when artificially removed, the pseudo-membrane is soon re-developed. After the first week from its beginning, no further exudation is apt to take place, and the danger arising from the membrane may be generally looked upon as over, unless, as is not uncommon, a relapse of the malady occur. The specific bacterium may, however, be present in the fauces and nasal passages for many days, even for weeks, after the disease has apparently come to an end.\* It may be occasionally found in the absence of false membrane, and also, it is said, in the throats of healthy persons.† The Klebs-Loeffler bacillus has been also detected in instances presenting all the clinical features of follicular tonsillitis, and, however valuable it is as a sign, we are not justified, with our present knowledge, in abandoning well-attested clinical experiences and making the presence or absence of the bacillus the only test of diphtheria.

---

\* Tobieson, *Centralblatt für Bakteriologie und Parasitenkunde*, Bd. xii., No. 17, p. 587; Park and Beebe, *Medical Record*, vol. xlv., No. 1247, p. 1. Abel (*Deutsche Medicinische Wochenschrift*, 1894, No. 35, p. 692) has recorded a case in which virulent diphtheria-bacilli were found sixty-five days after the onset of the primary illness.

† Feer, *Correspondenz-Blatt für Schweizer Aerzte*, 1893, No. 8, p. 295; Welch, *American Journal of the Medical Sciences*, Oct. 1894, p. 427 *et seq.*; Park and Beebe, *Medical Record*, vol. xlv., No. 1247, p. 1.



The constitutional symptoms vary greatly. The pulse may be frequent, the skin hot, and there may be much pain in the head; in fact, the symptoms are those of fever, with a temperature of  $102^{\circ}$  to  $103^{\circ}$ . Yet the temperature is exceedingly variable; there is often, even in the worst cases, an almost normal temperature. A sense of weakness and prostration are prominent from the onset. Not rarely the urine contains albumin and casts, and there may be partial or complete suppression of the renal secretion. In some instances typhoid phenomena manifest themselves, especially when decomposition of the disintegrating exudation takes place, giving rise to the septic form of the malady; in this the temperature may be even below the normal. The nervous system becomes much affected, and the tendon reflexes are lost.\* In children exacerbations of pre-existing nervous symptoms may take place and give rise to a state resembling acute bulbar palsy.†

In diphtheria the danger is twofold: it arises partly from the depressing effect of the poison, increased as this effect must be by the absorption of toxic matter from the throat, partly from the extension of the disease to the larynx and lungs. Again, at the height or even at the decline of the malady there is risk of heart-palsy or heart-clot and of peripheral embolism.‡ Nor is the termination of the acute disorder always the termination of the complaint. A chronic irritation of the throat, lasting weeks or months, and readily relapsing, on exposure to infection, into a diphtheritic sore throat, remains; or albuminuria, which outlasts the acute manifestations; or pleurisy, or bronchitis and pneumonia, —both of which may be delayed until after the exudation has disappeared from the throat,—increase the list of the complications of the affection, and protract or imperil the convalescence. Occasionally, too, inflammation of the joints is observed in the course of diphtheria, or as a sequel, and sometimes trophic changes in the periarticular structures are met with.§

---

\* McDonnell, *Medical News*, Oct. 15, 1887.

† Guthrie, *Lancet*, April 18, 25, 1891.

‡ A case has been recorded in which embolic obstruction of the popliteal artery occurred during convalescence from an attack of diphtheria, and amputation of the affected member became necessary. Rooney, *Occidental Medical Times*, vol. vii., No. 4, p. 188.

§ Lyonnet, *Lyon Médical*, Jan. 4, 11, 1891.

Some morbid conditions there are which may be wholly looked upon as after-symptoms. A paralysis of the velum palati and of the pharyngeal arches, making itself apparent by a peculiar nasal intonation of the voice, and by proneness to regurgitation of fluids through the nostrils, is among the earliest of them; it manifests itself often, indeed, just at the termination of the acute malady. Later appear impairment of vision, gastrodynia, ulcers in various parts of the body, profound anæmia, and that gradual failing of muscular power with numbness which ordinarily does not take place until after complete convalescence, and which winds up in almost total loss of muscular force with anæsthesia, and absence of reflexes, constituting diphtheritic paralysis. In rare instances these symptoms occur early in the attack.\* They are attributable to the development of a peripheral neuritis dependent upon the action of the toxic products of the disease. Hemiplegia has been observed in some cases as a result of rupture of a cerebral blood-vessel or its occlusion by a clot.† Other symptoms of profound nervous derangement have also been recorded, such as peripheral neuritis, in which the sense of smell and the muscular sense were lost, and slight sensory irritation was not appreciated, while more profound impressions were referred to corresponding points on the opposite side of the body,‡ with temporary absolute deafness, unsteadiness of gait, and paralysis of the palate.§ Furthermore, I have known aphasia to follow the depressing complaint.

But to look at the differential diagnosis of the disorder. It varies widely from stomatitis, from tonsillitis, from pharyngitis,—in truth, from all the ordinary local inflammations of these structures,—by the presence of a membrane, by the striking constitutional symptoms, and by the sequelæ. The diagnosis becomes unequivocal if the characteristic bacilli are found on bacteriologic

---

\* As in two cases reported by Dabney, *Medical News*, Jan. 16, 1892, in which they appeared on the first and second days respectively.

† McPhedran, *Canadian Practitioner*, 1892, No. 19, p. 454; Allen A. Jones, *Medical News*, Oct. 22, 1892, p. 467; Edgren, *Deutsche Medicinische Wochenschrift*, 1893, No. 36, p. 864; C. W. Sharples, *Medical News*, Aug. 4, 1894, p. 124.

‡ Gay, *Brain*, part lxi., p. 431.

§ Tooth, *British Medical Journal*, 1893, No. 1680.

examination of some of the material taken from the throat or the nose.

Yet there are certain sources of error against which it is necessary to guard. In *simple pharyngitis*, a mass of mucus, in part derived from the nares, is apt to collect on the inflamed membrane, and looks at first sight like the coating from an exudation; but it may be easily removed, and a closer inspection proves its true nature. In *follicular tonsillitis*, liquid may ooze from the openings of the follicles on the surface of the swollen tonsils, or little yellowish or whitish points form there. But they are strictly confined to the gland, exhibit no tendency to spread or to coalesce, and are generally small white specks of roundish or oval shape. I desire particularly to call attention to the possibility of confounding these appearances with diphtheria, for I have known them to occasion more than one mistake. The error is most likely to happen in those mild cases of the disease in which the exudation is limited and the injection or superficial inflammation of the tonsils and back of the throat marked, which are sometimes described as the "catarrhal form" of diphtheria. What adds to the difficulty is that follicular tonsillitis is contagious. Should, in an individual instance, the facts mentioned be insufficient to solve the doubt, the microscope can do so; for it shows the white or yellowish masses to be largely composed of epithelium, and not, like the diphtheritic membrane, mainly of fibrillated fibrin, of granular corpuscles, and of leucocytes, besides epithelium in different degrees of development and retrograde change.\* Further, if the case is one of follicular tonsillitis, streptococci and staphylococci are likely to be found, though, as already stated, the Klebs-Loeffler bacillus has been occasionally observed in instances presenting all the clinical features of follicular tonsillitis.

*Ulcerative stomatitis*, the form of stomatitis most likely to be confounded with diphtheria, and especially with this malady when the exudation lines the gums, is discriminated by the ulceration or sloughing; whereas the mucous membrane in the pseudomembranous disease remains intact, save in the rarest instances. The same feature distinguishes diphtheria from *gangrene of the mouth*, for which, on account of the extreme fetor of the breath, it is some-

---

\* Senator, *Klinische Vorträge*, 1874.

times mistaken, and aids in distinguishing it also from other kinds of stomatitis, as from *thrush*. In the latter, too, the buccal mucous membrane, and not the throat, is chiefly affected, and the abdominal symptoms, and the other constitutional phenomena, are different. So are they in *aphthæ*, in which, moreover, the superficial ulcerations, which bleed when touched, the unbroken vesicles or pustules in other parts, and the seat of the disorder—usually on the edge of the tongue, on the internal surface of the lips, and on the gums and inside of the cheek—are points to be taken into account.

Besides these affections, there are others which must be distinguished from diphtheria. We occasionally find cases occurring in epidemics, and where the membrane is limited nearly altogether to the follicles, and chiefly to the tonsils. As the membrane passes away, ulcerations are obvious. Swelling of the glands of the neck, and fever, but not of acute type, attend this *ulceromembranous angina*, which, moreover, shows a strong disposition to relapses. But, though kindred to diphtheria, and in isolated instances perhaps difficult to discriminate, it differs from it in its seat and in its want of tendency to spread, in the formation of superficial ulcers, in its less marked constitutional depression, and in its invariably favorable termination.\* It is similar to herpes of the tonsils, described by Trousseau. Whether there be not also other kinds of membranous sore throat to be separated from true diphtheria, is a matter requiring further investigation. The pseudo-membranous inflammations of the throat attending scarlatina and measles and other of the exanthemata have been shown not to be diphtheritic, although they seem to predispose to invasion by the diphtheria-bacillus.†

There is an acute disease of the throat to which Todd especially has called attention,‡ and which presents also some strong points of similitude to diphtheria,—*erysipelas of the fauces*. Like diphtheria, it is a most dangerous ailment; as in diphtheria, the morbid process may extend to the larynx; as happens often in

---

\* See a paper in which I have described an epidemic of the kind, in the Amer. Journ. Med. Sci., July, 1870.

† Booker, Bulletin of the Johns Hopkins Hospital, vol. iii., No. 26, p. 129; Park and Beebe, Medical Record, vol. xlv., No. 1247, p. 1.

‡ Clinical Lectures on Acute Diseases.



diphtheria, the mucous membrane may exhibit a peculiar dusky-red color; as in diphtheria, the poison paralyzes the muscles of the palate and pharynx, and liquids are apt to be rejected through the nostrils and mouth. But the difficulty in deglutition differs from that of diphtheria in being present from the onset, and is not attended with enlargement of the glands of the neck, or with the formation of a false membrane. In some instances, too, we find vivid redness of the throat, which may be associated with much swelling. If the erysipelatous inflammation extend to the larynx, there is local pain, with urgent dyspnoea and hoarseness; and usually rapid exhaustion supervenes. In cases of this kind, the submucous tissues of the larynx are found extensively infiltrated with pus. The cases may happen without erysipelas showing itself on any external part of the body; on the other hand, erysipelas beginning in the fauces may spread to the face.\*

This erysipelas of the fauces is not a frequent disease; and it must be stated that there are cases of diphtheria which simulate it very closely. I have seen a number of instances of the malady in which the whole mucous membrane was of a vivid or dusky hue; in which there was much swelling, with an effusion of serum, especially in the submucous tissue of the uvula, causing it to look like a small transparent bag; in which immense difficulty or even impossibility in deglutition existed,—yet in which no membrane appeared for days after the violent inflammation of the throat had set in, and was, when it showed itself, very slight in extent, and out of all proportion to the inflammation. But the constitutional symptoms and the sequelæ were the same as those of ordinary diphtheria. In one of the cases of the kind referred to, suppuration of one of the tonsils took place in consequence of the inflammation; a layer of deposit had coated parts of the tonsils and of the half-arches and uvula.

How shall we separate diphtheria from *membranous croup*, a disease with which, indeed, it is by many regarded as identical? Yet this seems taking a narrow view of the facts. In the first place, ordinary membranous croup lacks the peculiar constitutional symptoms, the early depression, and the sequelæ of diphtheria. Secondly, an affection of the windpipe is not by any means an

---

\* Cases quoted in Schmidt's *Jahrbücher*, 1869, No. 1.

essential element of diphtheria, for in the majority of cases the disease does not spread to the larynx. Thirdly, when, from the paroxysms of hoarse, irritative cough, the labored breathing, the attacks of suffocation, the huskiness or extinction of voice, we may infer that the exudative process has reached the larynx,—when, in other words, the symptoms of croup arise,—we still recall that the first manifestations of the membranous affection were perceived in the throat, and not in the larynx. Indeed, save in the rarest cases, the disease does not begin in the windpipe; though the beginning above it may not attract attention, and may be readily overlooked. Thus, *laryngeal diphtheria* affects primarily the throat, and may extend to the windpipe; membranous croup affects primarily the windpipe, and may extend to the throat. Fourthly, croup is not contagious, as diphtheria is. And, even granting that as regards the membrane and the symptoms we may not be able to distinguish individual cases of membranous croup from laryngeal diphtheria, the origin of the diphtheritic complaint, and its spreading to other members of the household, if not in a membranous form yet in the form of sore throat with singular constitutional depression, show its peculiar and special traits. Of course the finding of the specific bacillus in the false membrane establishes the diagnosis, though we must always remember that this may be absent in diphtheria.

On one symptom we cannot lay as much stress as might be supposed. Albuminuria, the elaborate report of the committee of the Medico-Chirurgical Society has taught us,\* is not always present in laryngeal diphtheria, owing to the early fatality of the malady; again, in certain cases the mere dyspnoea of laryngitis may give rise to albumin in the urine. Yet when albuminuria is marked, and when it has happened where an affection of the fauces has preceded the laryngeal implication, it points to an infective cause,—to laryngeal diphtheria.

Lastly, diphtheria may be confounded with *scarlatina*. When, indeed, we reflect on the similar appearance of the throat, on the occurrence of albuminuria in both maladies, and on the frequency

---

\* Medico-Chirurgical Transactions, vol. lxii., 1879. Some of the anatomical points involved are also well discussed by Weigert in Virchow's Archiv, vols. lxx. and lxxi.

with which both are found to prevail at the same time as epidemics in a community, it is not astonishing that one should be looked upon as but a modified form of the other. Allied they certainly are, but not identical; for the poison of one leads to a thoroughly-defined rash, and leaves a protective influence against a second attack, and often also deafness, suppuration of the glands of the neck, and dropsy,—phenomena which are not encountered in the other. It is true that in very rare instances of diphtheria we encounter a slight erythema of the neck and breast, but it is not like the vivid, diffused rash of scarlet fever. Moreover, the exudation in the throat is not exactly similar in the two diseases. In scarlatina it is pultaceous, and not coherent, and has no tendency to spread to the respiratory passages. Bacteriologic examination, further, may disclose the presence of streptococci and staphylococci, but not the bacillus peculiar to diphtheria. Then the albuminuria happens at a different period. In scarlatina it is a sequel rather than a concomitant; in diphtheria it is a concomitant rather than a sequel. Further, the gravity of the symptom is not the same. In the latter malady it is an indication of danger; it has not so serious a meaning in the former.

Diphtheria may be intercurrent in various maladies: in typhoid fever, in the exanthemata, in pneumonia. Nor is the exudation always restricted to the throat. It may show itself in a wound or on excoriated skin, on the nasal mucous membrane, the conjunctiva, the nipple, the uvula, or around the anus; it may be found coating the stomach, the intestines, and the ramifications of the bronchial tubes.

*Nasal diphtheria* is a very grave form of the malady: it may either be present alone, or coexist with a deposit in the fauces and pharynx. It generally occurs with evidences of the septic form; the symptoms are of a low type, and we recognize the affection by carefully inspecting the posterior pharynx and seeing that the membrane extends upward; by noting the irritated, reddened look of the nostril, even when no membrane can be discerned in it; and by the coryza, the sense of obstruction in the nose, and the acrid sanious discharge which comes from it. In cases in which the nasal duct and the laryngeal canal are stopped up by the false membrane, tears are constantly rolling down the cheeks. Epistaxis is a not uncommon symptom; swelling of the

cervical glands may or may not be present. Recent bacteriologic investigation has shown that so-called membranous rhinitis is in reality often of diphtheritic origin.\* And in the enlarged glands in any form of diphtheria the characteristic bacilli are found in the opaque, yellowish masses, consisting principally of fibrin, which they contain.†

**Mumps.**—This, like diphtheria, is a general disease, and is only here described as a matter of clinical convenience. Parotitis is most commonly seen as an epidemic malady; but we occasionally encounter a secondary parotitis following typhus fever, scarlet fever, smallpox, measles, and dysentery. In this form suppuration is much more common than in ordinary mumps. The disease generally begins with pains at the angle of the jaw, which are soon followed by a marked swelling, first on one side, then on the other, which results in the head being kept rigid. The tumid glands are sore, and become more painful during attempts at swallowing and chewing, though there is really little, if any, difficulty in swallowing. If the patient be made to swallow slowly ten to thirty drops of undiluted vinegar, decided pain is produced in the affected glands,—an old and useful diagnostic test, to which Dr. Louis Starr called my attention. The mouth is generally filled with saliva, though it may be very dry; and the hearing may be impaired, or, for the time being, entirely lost, and ringing in the ears is very common. The temperature generally ranges between  $101^{\circ}$  and  $102^{\circ}$ , but in cases of orchitis following mumps, or of metastasis, I have seen it  $104^{\circ}$  to  $105^{\circ}$ . The nervous system may become decidedly affected. Acute mania has been known to become associated with mumps; so has peripheral neuritis.‡ Parotitis is easily recognized. There is no swelling of the tonsils, hence it cannot readily be mistaken for tonsillitis. Laveran and Catrin have found a diplococcus in mumps, in the secretions of the parotid and other glands, as well as in the blood.§

In cellulitis of the neck, *angina Ludovici*, the swelling may

---

\* Abbott, Medical News, May 13, 1893, p. 505.

† Bulloch und Schmorl, Beitr. zur Pathol. Anatomie, etc., von Ziegler, B. xvi., H. 2; Centralblatt f. Innere Medicin, 1895, No. 6, p. 156.

‡ Lancet, April 9, 1887.

§ Gaz. Méd., June, 1893.



mislead, but it is uniform and not confined to the region of the parotids; the constitutional symptoms are very severe, pointing to an infective malady. Ludwig's angina is met with as an idiopathic affection, or in certain fevers, such as scarlet fever or diphtheria.

**Chronic Sore Throat.**—Attacks of angina are prone to recur, and to lead to chronic inflammation of the structures. Now, an affection of this kind is liable, on any exposure, to be kindled into the acute complaint; besides, it yields at all times some manifestations of a disorder of the throat. A thickening of the folds of membrane forming the half-arches, a tumefaction of the follicles at the upper part of the pharynx, a lengthening of the uvula, are the visible signs of the chronic malady; a constant disposition to clear the throat, and a dry cough, are often the attending general symptoms. Owing to the habitual coughing, the patient may be suspected to be laboring under phthisis, and be treated accordingly, when the whole difficulty lies not in the lungs, but in the throat. Yet an error in the opposite direction is perhaps more frequently committed. Tonsils and uvulas are removed, with the view of curing a cough which is really kept up by a source of disturbance in the lungs, in forgetfulness of the fact that in scrofula and tuberculosis chronic enlargement of the tonsils and follicular pharyngitis are by no means unusual. A careful examination of the chest ought always to be made, even when inspection of the throat shows disease to be there present. On the other hand, we may find in the condition of the throat and of the nares the explanation of thoracic affections, for instance of asthma, a number of cases of which have their origin in irritation reflected from these parts.

*The follicular disease of the throat*, or "clergyman's sore throat," is the most frequent of all the morbid conditions which produce a chronic sore throat. The abnormal condition of the follicles of the pharynx and fauces often extends to the larynx. There are constant hawking and attempts at clearing the throat, and not unfrequently roughness of voice or decided hoarseness. On inspecting the throat, the enlarged mucous follicles can be readily discerned; those on the pharynx are very prominent. In cases of long standing, the follicles may ulcerate, and very commonly they pour out an acrid secretion. But, unless from coexisting

enlargement of the uvula, or an altered position of the epiglottis, or a laryngeal or bronchial complication, there is no decided cough. The follicular disease may occur in consequence of repeated attacks of sore throat, or be an attendant upon gastric disorder, or follow constant exercise and straining of the voice.

*Chronic rheumatic sore throat* gives rise to pain which is often referred to the hyoid bone, is increased by pressure, and is also felt in the tonsils. Ingals\* points out that the pain often entirely disappears while the patient is eating, but increases in cloudy and damp weather. There are signs of slight congestion in the throat, and generally in the larynx, yet mostly out of all proportion to the pain. The general health remains good, and we find no fever; there is apt to be a history of a rheumatic diathesis.

*Ulcers* are not often developed in the fauces during an attack of acute inflammation, except in the specific sore throat of scarlatina; in chronic inflammation, especially if occurring in scrofulous persons, they are more common. The most profound ulcerations are those of constitutional syphilis, implicating, as they do, not only the tissues of the fauces, but also the parts in front, and destroying both the fleshy covering of the bones and the bones themselves. With regard to treatment and to prognosis, it is of the utmost importance to distinguish these *syphilitic ulcers* from those produced by other causes. A cutaneous eruption of a syphilitic character, and enlarged lymphatic glands, or the history of antecedent syphilis, would lead us to a correct conclusion; but an accurate history of a syphilitic infection cannot always be obtained. The ulcers themselves furnish some information by which we may suspect their origin. They are not superficial and stationary, like those resulting from ordinary inflammation; on the contrary, they are deep and have a strong tendency to spread. They are rounded, or of a serpiginous form, with borders well defined and elevated, and surrounded by a distinct zone of redness; and the inflammation which precedes them is limited to spots, and is not so diffused, nor attended with so much swelling, as the inflammation which exists prior to simple ulceration. These ulcers must be distinguished from the deep ulceration with spreading destruction of tissue that occurs in *cancer of the tonsils*.†

---

\* Medical News, March, 1890.

† See Newman, Amer. Journ. of Med. Sci., May, 1892

## PHARYNX AND ŒSOPHAGUS.

In describing the affections of the fauces, those of that portion of the pharynx which is most usually the seat of disease have been at the same time described. Indeed, when we speak of acute or chronic pharyngitis, we generally mean acute or chronic inflammation of the fauces, to which the upper part of the pharynx belongs. Inflammation of the portion of the pharynx which is out of sight when the tongue is depressed, is rare. It may be presumed to exist if there be pain and an impediment in swallowing when the food arrives opposite the top of the larynx, while the respiration remains free and the voice unaffected. Abscesses sometimes form between the textures composing the pharynx, and between its posterior wall and the cervical vertebræ. These *retro-pharyngeal abscesses* mostly result from disease of the vertebræ. They occasion great difficulty in deglutition and in breathing; an altered voice; dull pain and stiffness in the neck; external swelling, which may or may not be œdematous; and commonly a tumefaction at the back of the throat, which can be seen, or which at least can be felt with the finger pressed against the posterior wall of the pharynx. On account of the obstructed respiration and the changed voice, the disease is liable to be mistaken for croup. Its differences have been already enumerated. Retro-pharyngeal abscess is often confounded with coryza and with tonsillitis. It differs chiefly from tuberculosis of the retro-pharyngeal glands by the presence of tuberculous lesions of the deep lymphatic glands on the same side of the neck.\* It may happen in infancy.†

There is a peculiar form of pharyngeal disease due to the accumulation on the mucous membrane of a micro-organism generally supposed to be the leptothrix, though Hemenway‡ in his elaborate article pronounces it to be the bacillus fasciculatus. The deposits in this *pharyngo-mycosis* take place largely in the follicles.

The œsophagus is not often the seat of disease. We meet with

---

\* Sokoloff, Vrach, May, 1891.

† See cases of Pollard, Lancet, Feb. 1892.

‡ Journ. of Laryng., Feb. 1892.

acute inflammation produced by swallowing boiling water or corrosive poisons, especially nitric or sulphuric acid, or ammonia. The symptoms of acute *œsophagitis* are usually mixed up with those of inflammation of the pharynx, or of the stomach. We may, however, infer its presence if difficulty and pain in deglutition exist for which nothing in the throat accounts, and if these phenomena be associated with hiccough and with a burning sensation between the shoulders, in the course of the tube. *Œsophagitis* is sometimes encountered in infancy.\*

Of the chronic diseases of the *œsophagus*, *stricture* is the most common. The narrowing may take place at any part of the tube, and results from preceding inflammation or ulceration, from cancerous degeneration of the walls, from polypoid growths projecting from the mucous membrane, or from the pressure of a tumor, of an abscess, or of an aneurism; sometimes it is congenital. The formidable malady manifests itself by impediment in swallowing: even liquid food cannot pass without great difficulty; and if the stricture go on increasing, the patient perishes miserably by starvation. In addition to the obstruction to the passage of food, we may find a peculiar pain occurring at a particular part of the tube, and that the patient raises, without cough or vomiting, clots of blood presenting the shape of the stricture.

The matter ejected in the attempts at deglutition consists simply of masticated food together with more or less mucus, and, unlike what comes from the stomach, has an alkaline reaction. If long retained, the albuminous materials are macerated; the starchy materials are in process of fermentation; fungi are formed in great quantities, although never *sarcinæ*.† Should there be doubt as to the seat of the obstruction, a bougie will clear up the doubt; and thus we possess in this instrument the most valuable diagnostic as well as therapeutic agent. But we must not immediately conclude, because the bougie meets with resistance, that an organic stricture is present. The narrowing may be only *spasmodic*, yet give rise to the symptoms of organic constriction. But they are not permanent: at times nourishment is readily swallowed, and a full-sized bougie passes with ease. Spasmodic stricture occa-

---

\* Brush, New York Med. Rec., 1883, xxiii.

† Ziemssen, "Diseases of the *Œsophagus*," in Ziemssen's *Cyclopædia*.



sionally accompanies ulceration of the larynx; but it is chiefly met with in hypochondriacs and in hysterical women. The latter, indeed, sometimes fancy that they are incapable of swallowing, and reject the food they take without there being even a temporary spasm to prevent its passage. Spasmodic stricture is also observed in hydrophobia and as an attendant on cerebral disease.

The distinction of the other causes of stricture is not always an easy matter. In the stenosis arising from *syphilis*, we lay great stress on the history. In the strictures due to *compression*, we discern the swelling that has occasioned them, and the œsophagus is apt to be pushed to one side. In strictures the result of *cicatrices*, we have the gradual development of the affection after an injury or the swallowing of some irritant poison, and the great resistance of the dense tissues to the sound is very significant. *Cancerous narrowing* occurs after forty years of age, progresses steadily, and, as Ziemssen has pointed out, is frequently associated with paralysis of the recurrent laryngeal nerves. It may affect the whole middle part of the œsophagus.\* Cancer of the œsophagus is most commonly epithelioma.

*Rupture of the œsophagus* may be met with as the result of protracted vomiting. The accident is apt to occasion great pain. It leads to a rapidly fatal result.†

*Dilatation* of the œsophagus above the seat of a stricture, or without a stricture existing, is, on the whole, a rare disease. Its chief symptoms, when extensive, are difficulty in swallowing, vomiting or regurgitation of food, a swelling in the neck coming on after eating and diminishing greatly after vomiting or by pressure, slowly-progressing inanition, and at times long spells of delusive improvement. The sound may penetrate through the neck of the sac with difficulty, or enter it readily, which largely depends upon whether the sac be empty or full; once in the sac, the end of the tube can be generally moved about with ease.

In all the diseases mentioned, the value of the *sound* as a means of diagnosis has been spoken of. A few more remarks about it may not be amiss. Great care should be always used in passing a sound. The patient's head should be well thrown back,

---

\* Moore, Lancet, London, 1883, i. 13.

† See for cases paper by Fitz, Amer. Journ. of Med. Sci. Jan. 1877.

and the instrument passed along the posterior wall of the pharynx with the utmost gentleness. There is a slight resistance as it goes past the cricoid cartilage. When an aneurism or an organic disease of the heart exists, it should not be employed at all. When the sound on reaching a particular spot always occasions pain, we may infer the existence of inflammation or ulceration at this point, and, in the case of ulceration, some pus or blood is likely to be brought up on the instrument. Should any doubt exist whether the sound have passed into the œsophagus or into the larynx, let the patient be directed to speak; he can make no noise if the tube be in the larynx. In cases remaining doubtful, a lighted candle may be placed before the end of the tube projecting from the mouth. If the instrument be in the windpipe, the flame will be wafted to and fro with the currents of air; if in the œsophagus, this is not observed, except when the tube is in the intrathoracic portion.

The diseases of the œsophagus may be studied by means of *auscultation*, listening while the patient swallows food or liquid; and we owe to Hamburger an elaborate description of the sounds.\* In health, the œsophageal sound is extremely distinct, but of very short duration; the pharyngeal swallowing sound is generally a loud gurgle. In a moderately advanced stage of stricture of the œsophagus, a noise similar to emptying a bottle, "clucking," "gurgling," is perceived; while in cases of dilatation we are apt to meet with a sound like that heard when rain driven by the wind impinges and is deflected. In cases of very marked stricture or of obstruction by an impacted foreign body, we find that the act of deglutition cannot be detected below a certain point, while it is distinct above. To auscult the œsophagus, the stethoscope should be placed in the vicinity of the hyoid bone, also to the left of the vertebral column from the upper dorsal vertebra downward. This method of exploration has not, however, proved itself of much value.

The disorders of the pharynx and œsophagus have as a common symptom difficulty in swallowing. But we must not forget that other causes may produce *dysphagia*, such as paralysis of the muscles of the throat, diseases of the larynx or trachea, particularly ulcerative diseases, and aneurismal tumors within the chest.

---

\* Jahrbücher der k. k. Gesellschaft der Aerzte in Wien, Bd. xviii.

## CHAPTER VI.

### DISEASES OF THE ABDOMEN.

THE abdominal cavity contains viscera of very varied functions: some form, others break down organic constituents; while others, again, excrete the broken-down material. They all, however, labor in one cause; they all work toward preserving a normal state of the blood, either by preparing fit matter for it or by removing such substances as would be hurtful if they were retained. Any serious derangement of any of these viscera, especially any serious chronic derangement of those which are not simply reservoirs, must therefore lead to a deterioration of the blood and to a defective nourishment of the body. But, independently of the change in the blood and the falling off in the general nutrition, there are no vital symptoms that characterize abdominal diseases as a group; and, as many causes may give rise to the same symptoms, they furnish but little information as to the particular organ at fault. This we learn to some extent by examining, where it can be done, the secretions or excretions; to some extent by noticing the peculiar appearances of the skin which are produced by alteration of the blood; and by the exploration of the organs through the parietes of the abdomen. It is, in truth, by means of the physical method of investigation that we often obtain the most valuable information not only as to the seat but even as to the nature of the morbid action; and, although physical exploration of the abdomen does not yield as perfect results as when this form of diagnosis is applied to the affections of the thorax, the senses of sight and touch still supply us with an amount of knowledge most valuable, and with which it would be difficult to dispense. I speak only of the senses of sight and touch, because the sense of hearing, save in so far as it enables us to judge of the sounds elicited by percussion, or of murmurs in the vessels, is not very applicable to the study of diseases below

the diaphragm. Let us pass in review the different methods of physical diagnosis with reference to abdominal disorders.

## Methods and General Results of Physical Examination of the Abdomen.

### INSPECTION.

By inspection we learn the size, shape, form, and movements of the abdomen. To inspect the abdomen satisfactorily, the patient should be placed in an easy attitude, either standing or sitting. The recumbent position is less eligible, though we are often obliged to examine sick persons in this posture. Whenever practicable, ocular inspection must be made not only from the front, but also from the sides and from the back. In appreciating the results thus obtained, it is necessary to bear in mind that the appearance of the abdominal walls is modified by certain physiological conditions. The abdomen is much larger, in comparison to the size of the chest, in childhood than in adult age. It is more voluminous in females, especially such as have given birth to children. It increases in size with advancing years, particularly when a tendency to obesity exists. Its shape is somewhat altered by the pernicious habit of wearing tight stays. Its upper portion is distended after a copious meal.

In disease we may observe either partial or general *abdominal enlargement*. The latter is caused by accumulations of air in the intestinal canal; by liquid in the peritoneum; by an œdematous condition of the abdominal walls; or by large tumors which fill up the whole cavity. A partial enlargement is mainly produced by an increase in size of particular organs. It may also be brought about by swelling of the mesenteric glands, or by tumor, —solid or hernial; and it is sometimes due to diseases above the diaphragm. A pleuritic or a pericardial effusion, or emphysema of the lungs, may give rise to a marked fulness below the margin of the ribs.

A *retraction of the abdominal parietes* is perceived in general emaciation, and is very obvious in that dependent upon a narrowing of the cardiac or the pyloric orifice of the stomach, or upon chronic diarrhœa or dysentery. It is also noticed in lead colic and in cephalic diseases, especially in tubercular meningitis.



There are further changes in the appearance of certain external parts which tend to elucidate the state of the parts within. Thus, we learn from the distention of the superficial veins that an obstruction to the flow of blood exists in the large veins of the abdomen, either in the portal system or in the vena cava. The lessening of the depression at the umbilicus, unless it be produced by pressure limited to the spot where the umbilicus lies, is a sign indicative of general abdominal enlargement.

While inspecting the abdomen, we may see distinct *movements*. The act of breathing gives rise to a motion which is very slight when a tumor or any other impediment interferes with the free action of the diaphragm, and which is much exaggerated by diseases within the thoracic cavity. The rolling of the intestines is sometimes visible on the exterior; so are at times those shiftings of accumulations of gas which give rise to a series of jerking elevations; so, too, are occasionally the spasmodic contractions and relaxations of the abdominal muscles. But none of these is as frequently encountered as a pulsation in the epigastric region, which is often mistaken for an aneurism. The inspection of internal organs, such as the stomach, will be considered in connection with those organs.

#### PALPATION.

We judge by the application of the hand of the size, position, and consistence of the viscera which are felt through the abdominal walls. We determine whether the parts are firmly attached or movable; whether they are smooth or nodulated; whether they possess a motion of their own; whether they are tender; and by tapping with the fingers of one hand, while those of the other are applied to another portion of the surface, we discover, by the peculiar feeling of fluctuation, the presence of fluid in the abdominal cavity. We satisfy ourselves further, by the sense of touch, of the state of the parietes, whether resistant or elastic, œdematous or not; and we may detect a friction fremitus.

In order to use palpation with most effect, the abdominal muscles must be relaxed; and to do this the patient should be placed on his back, and the thighs be flexed on the body. Occasionally it is essential to vary this position; to turn him from side to side, or to examine him when erect. The amount of pressure, too,

should not always be the same. When we wish to examine deep parts, the pressure is increased. The character and the intensity of the pain which pressure calls forth often throw considerable light on the disease we are investigating. Thus, if it take deep pressure to produce pain, we are usually right in concluding that the mischief is not superficially seated. The pain of inflammation of the serous membrane is commonly much augmented by pressure, and is of a very severe, cutting character. Pain due to inflammation of any part of the mucous membrane of the intestinal tract is duller. All neuralgic or nervous pain, such as that of colic, is, as a rule, relieved rather than augmented by pressure, and may be thus distinguished from inflammatory tenderness.

But we shall not enter into any fuller particulars as to what palpation teaches us in individual diseases of the abdomen; because, as there is hardly one of any importance in which it is not of some service, we should say here what it would be necessary to dwell on repeatedly hereafter. There is, however, one point connected with the subject which may be briefly mentioned,—the attempt to use palpation as a means of diagnosis by the introduction of the hand into the rectum. But the method is both disagreeable and not free from danger. Dilatation of the sphincter should be gradual, five minutes at least being allowed for its accomplishment. And, with all precautions, the information obtained may be indecisive. Strictures high up in the rectum or in the sigmoid flexure of the colon may be readily discerned, but a stricture below the descending colon may exist although the hand be unable to discover it.

We might with palpation consider the results obtained by the use of bougies and of tubes, such as the stomach-tube. But these will be more appropriately considered when discussing the diseases of individual organs.

#### PERCUSSION.

Percussion is, in the study of abdominal affections, even more valuable than palpation. By it we can circumscribe the different organs with accuracy; we can judge of the position of the stomach and intestines; we can limit the distended bladder, and fix the borders of the liver and spleen. By its aid, further, we can tell

whether a distention of the abdomen is produced by air, or by a solid tumor, or by liquid. But, without entering here into particulars as to its use in individual disorders, we shall examine the results it yields when applied to the healthy abdomen.

To render percussion a trustworthy interpreter of the state of the abdominal viscera, the patient should be placed in the same position as for palpation. The sounds are best elicited by mediate percussion. But to appreciate them fully we must be acquainted with the relations of the parts which the abdominal walls conceal from view, and take into account that during the digestive process the contents and position of these organs may vary sufficiently to modify the percussion sound.

To begin with the airless viscera. The *liver* is one of the easiest organs to limit. We determine its upper boundary by striking with moderate force in a line from somewhat above the right nipple toward the lower part of the thorax, until marked resistance and dulness tell us that a solid organ has been reached. At this point we make a mark; then we again percuss downward from near the median line, and above the dulness just obtained; then we percuss from the axilla downward; then posteriorly from beneath the lower angle of the scapula; and so on, until the line traced out reaches the vertebral column.

The dulness thus elicited marks the upper boundary of the liver; at least of the portion more directly in contact with the abdominal walls. Anteriorly it extends from the lower extremity of the sternum to between the fifth and sixth ribs; at the side, the dulness is generally in the seventh intercostal space; near the vertebral column, it is on a level with the tenth or the eleventh, more rarely with the ninth, interspace. The dulness of the left lobe reaches nearly two inches across the median line; but the heart lies here so near to the liver that we cannot with accuracy distinguish the flat sound of the one from the flat sound of the other; nor indeed is this, for practical purposes, of great consequence.

After the upper border has been fairly traced out anteriorly, laterally, and, if thought necessary, posteriorly, we determine the inferior margin of the organ. This is readily effected by percussing downward from the already-ascertained line of dulness, and noting where the large intestine sends forth its distinct tym-

panitic sound. To determine the lower border correctly, the pleximeter must be pressed firmly on the integuments, and the stroke of the finger be slight; for if it be strong, we obtain the sound of the surrounding hollow viscera through the thin layer of liver which covers them, and before we have arrived at its margin. This mode of procedure is different from the one pursued to determine the height to which the liver rises, because the position of the parts is different. Superiorly, the lung descends between the surface and that portion of the convex surface of the liver which fits into the diaphragm, and it requires strong percussion to bring out the dulness of the deep-seated solid organ. By forcible percussion, however, we detect a decided loss of the pulmonary resonance at about the fourth intercostal space.

The inferior border of the liver will, anteriorly, be generally found to lie immediately at, or to project below, the last rib; posteriorly, we cannot determine this border positively, for it becomes continuous with the dulness occasioned by the right kidney. The lower margin of the left lobe is commonly met with at the upper third of a line drawn from the ensiform cartilage to the umbilicus. A distended gall-bladder may cause a strictly-defined dulness lower than the dulness of the surrounding liver.

The *spleen* is not so easily circumscribed as the liver. Indeed, if the stomach or the intestines be distended, it is difficult to detect the dull sound of the spleen. To find its limits, we must place the patient on his right side, with his legs flexed; or let him stand erect, and then begin to strike with some force in a line from the axilla to the crest of the ilium. At the ninth, or sometimes at the tenth, rib, the sound becomes dull, and there is much greater resistance to the finger. Here is the upper boundary of the spleen. We mark the spot, and continue to percuss in the same line until, at about the twelfth rib, we arrive at the lower boundary of the organ, as indicated by the distinct tympanitic sound of the intestines.

After the vertical diameter has been thus ascertained, the horizontal is readily determined by percussing from the median line to a point between the lines which trace the superior and inferior margins, and by noticing where the sound of the stomach gives way to the dull sound of the solid viscus. When these three points have been decided upon, we have learned enough for practical purposes. We may then, if we choose, percuss posteriorly;



but we cannot circumscribe the spleen with any accuracy behind, because its dulness becomes continuous with that of the left kidney.

The average size of the spleen is four inches in length and three in width ; but it may in a diseased state increase to twice or three times that size. When the viscus eludes detection by percussion, we may infer it to be small ; provided the stomach and intestines be not much distended with gas.

The *kidneys* cannot be limited with anything like accuracy, except at their inferior and outer borders, where the dull sound they occasion is surrounded by the intestinal resonance. This dulness extends somewhat lower during a full inspiration.


To set limits to the *stomach* and *intestines*, by means of percussion, requires an ear accustomed to discriminate between shades of sound, since we have to judge more between sounds of different degree, but similar to one another, than between sounds of different character. Nor are the tones elicited always the same over the same spot ; they vary as the contents of the hollow viscera vary. We can make use of this circumstance for purposes of diagnosis. In percussion of the stomach and of the intestines we may often with great advantage resort to auscultatory percussion.

The stomach, when not unusually distended with gas or with food, renders a sound which is hollow, ringing, and tympanitic to a certain degree, yet which is not tympanitic as that of the intestine is. It is, in fact, a sound unlike any other, and experience soon enables us to distinguish it from that of the surrounding viscera. Sometimes the sound is distinctly amphoric. To determine the boundaries of the stomach, it is necessary to mark out first the lower margin of the liver, for it covers a portion of the stomach ; then the heart and the inner border of the spleen. The part which lies between these solid viscera yields the sound of the stomach, mixed at one point, namely, to the left of the apex of the heart, with the resonance of the lung. Near this spot, about opposite to the seventh rib, the cardiac extremity of the stomach is situated ; below it is the body of the organ ; the pylorus is on a level with the tip of the xiphoid cartilage, between the right edge of the sternum and a vertical line passing through the nipple.

To ascertain the lower border of the stomach, we percuss gently in a downward direction, until the alteration in sound

shows that we are striking over the colon. The difference is at times very obvious, at times very slight. It is readily detected if the stomach contain either solid or liquid ingesta. Availing ourselves of this fact, we may sometimes with advantage let the patient swallow a glass of water. By placing him in the erect position, the fluid gravitates to the greater curvature, and the line

FIG. 53.



Results of abdominal percussion, as set forth in the text. The dark shades indicate marked dullness; the light shading exhibits a lessening of the clear or of the tympanic character of the sound,—an approach to dullness.

of comparative dullness indicates the lower margin of the stomach, which is generally found near but above the umbilicus. In men the lower border of the stomach is a little higher than in women; in working-women it is higher than in other women; in children under fifteen years of age it very rarely extends to the umbilicus; in persons of fifty it is not unusual for it to do so. In strong,

healthy people the whole position of the stomach is more horizontal than in weak ones.\* Deep percussion is used to limit the superior line, and light percussion the inferior line, of the stomach. Direct insufflation of the stomach is also valuable to determine its size and its distensibility.

Another method to determine the limits of the organ, as well as whether the pylorus is still capable of self-closure in the direction of the duodenum, or is permanently patent, has been proposed by Epstein.† It consists in the distention of the stomach by means of carbonic acid, generated by first letting the patient swallow tartaric acid dissolved in lukewarm water and then rather more bicarbonate of sodium, about a full teaspoonful. The stomach becomes very much distended, and emits a deep tympanitic note on percussion, unlike that over the intestines; but if the pylorus be incapable of closure, the intestines too become swollen, and their tympanitic note is changed.

The *colon* yields, in all its parts, a sound of a purer tympanitic character than the stomach, the note of which is, indeed, in many respects more amphoric than tympanitic. When the tube contains faeces, the sound is modified; and as these are prone to accumulate on the left side in the descending colon, and especially where this passes into the iliac fossa, it is usually not so resonant as the ascending colon. The *small intestines*, unless they are filled with fluid or solids, or distended with gas, render a sound of higher pitch and of smaller volume than the surrounding large intestine, and by the less deep-toned sound their position may be accurately determined. Artificial distention of the colon, by generating carbonic acid in it by means the same as just mentioned passed into the lower bowel, has been advocated for diagnostic purposes by Ziemssen.‡ It enables us to distinguish with ease the outline of the large intestine, and shows whether there is communication with adjacent organs, such as the stomach, the bladder, or the small intestine. Anomalies of position and form of the bowel give rise to differences in the results of abdominal percussion, as has been well shown in a careful clinical study by Curschmann.§

---

\* Obrastzow, *Deutsches Arch. f. Klin. Med.*, Bd. xliii., 1888.

† *Klinische Vorträge*, No. 155, 1878.

‡ *Deutsches Arch. f. Klin. Med.*, Bd. xxxiii., June, 1883.

§ *Ibid.*, Bd. liii., June, 1894.

The position of the viscera in the pelvis cannot be ascertained by means of percussion. It is only when the bladder is much distended, or the uterus augmented in size, that the outline of either can be traced on the walls of the abdomen.

#### AUSCULTATION.

Auscultation is serviceable in aiding in the detection of an abdominal aneurism ; and sometimes an enlarged spleen gives rise to a distinct blowing murmur ; or the rubbing of a roughened peritoneum may occasion a friction sound ; but, on the whole, the application of the stethoscope to the abdominal walls is rarely of much aid. In health, no constant sound is heard save that of the aorta ; for the rush of blood through the other arteries, or through the veins, produces no appreciable murmur. When the stomach is distended with air and contains liquid, sounds possessing a metallic character are perceived, which an inexperienced observer is apt to consider as originating in the lungs ; over which, in truth, they are often audible. The passage of gas through the intestines gives rise to those peculiar noises termed “borborygmi.” In the pregnant state, auscultation is of value in detecting the pulsations of the foetal heart and the utero-placental murmur.



### SECTION I.

#### DISEASES OF THE STOMACH.

As the disorders of the stomach are so common, and as a patient hardly ever gives a history of his ailment without thinking it obligatory to enter into a minute account of the state of his digestion, it would be reasonable to suppose that no affections are so well understood and so susceptible of clear description as those of this viscus. But in point of fact it is only within the last few years that any attempts have been made to bring to bear upon them modern means of research. Most of these attempts have



had as their aim to ascertain the exact anatomical changes and the modifications in the secretions which give rise to the symptoms commonly referred to perverted function ; and to a decided degree they have been successful.

The stomach is examined partly by physical exploration by the methods just detailed, and partly by paying attention to the chemical changes which attend the digestive acts.

With reference to the physical examination, there are some special means that may be employed with advantage. To determine the relative sensitiveness over the epigastrium, Boas\* measures the pressure by an *algometer*. The normal tolerance is from eighteen to twenty pounds. In cases of gastric ulcer, pain is complained of at a pressure of from two to four pounds. The direct application of *electricity* to the coats of the stomach as a test of their motility has been also made use of ; but, valuable as this agent has proved itself therapeutically, it has not shown itself valuable diagnostically.

Ingenious instruments have been devised for illuminating and inspecting the interior of the stomach. By means of the *gastro-diaphane* of Einhorn,† which consists of a soft rubber tube through which pass wires connected with a source of electricity and provided with an incandescent lamp enclosed securely in glass, the size and outlines of the stomach, as well as the density of its wall, can be made out. The patient is, on an empty stomach, first made to swallow, or there are introduced through a tube, one or two pints of water ; the tube is passed into the stomach in the customary manner, and the appearance of the light is observed in a dark room. A reddish luminous zone upon the abdomen indicates the outline and the position of the stomach ; and dark spots may enable us to judge accurately of the size, shape, and position of tumors. The gastroscope of Mikulicz is a more complicated instrument, by means of which it is possible to inspect directly limited portions of the interior of the stomach.

A revolving sound, the *gyromele*, has been invented by F. B. Turek.‡ The revolutions can be felt upon the abdominal wall,

---

\* Münch. Med. Wochensch., Sept. 1893.

† New York Medical Journal, Dec. 1892.

‡ Journ. Amer. Med. Assoc., March, 1895.

and the various parts of the stomach, especially the greater curvature, accurately located ; we can also, by the use of different cables, measure the distensibility of the stomach. If the movements of the sound are distinctly felt on the parietes, tumors of the anterior wall and of the fundus can be excluded. The instrument is also valuable for obtaining materials from the wall of the stomach for microscopical and bacteriological investigation.

It is always important to study the activity of the movements of the stomach, and this is generally done partly by noting how long it will take a trial meal to digest completely, partly by chemical means to be presently detailed. But the object has been also sought to be attained by instrumental aid. With this view, F. B. Turck \* has introduced a *gastric motormeter*, which consists of a collapsed rubber bag with a fine rubber tube attached that is connected with a manometer ; thus both degree and force of movement are registered. The bag is inflated with air after being passed into the stomach. The instrument records not only the peristaltic, or active, movement of the stomach, but also the respiratory movement and the aortic movement.

Another way of determining the mechanical action of the stomach, as well as of recording its movements, is by the *gastro-graph*, the invention of Einhorn.† It is likely to prove of use in several pathological conditions of the organ, and will be valuable in obtaining tracings for comparison.

The accurate chemical study of the secretions and of the contents of the stomach is leading to great advances in the investigation of its affections, as has been proved especially by the labors of Leube,‡ of Ewald,§ of Boas,|| and of others. We get the contents of the stomach for examination from two to four hours after a full or "trial meal," given as a mid-day dinner, and consisting of four hundred grammes of soup, sixty grammes of scraped meat, and fifty grammes of white bread ; of this, if the act of digestion have been normally carried on and the chyme have passed on into the small intestine, nothing remains after the lapse of six or

---

\* Proceed. Amer. Med. Assoc., May, 1895.

† New York Medical Journal, Sept. 1894.

‡ Deutsches Arch. f. Klin. Med., Bd. xxxiii., 1883 ; also Schreiber, *ib.*

§ Klinik der Verdauungskrankheiten, Berlin, 1889.

|| Diagnostik und Therapie der Magenkrankheiten, 3d ed., 1894-95.

seven hours but a clear liquid. The material for examination is obtained by means of an elastic tube, preferably of soft rubber, about seventy-five centimetres long by six centimetres in diameter, and provided with an opening at its conical extremity and others at the side. The liquid is removed from the stomach by pressure over the epigastrium, or by aspiration by means of a hand-ball apparatus. Ewald has substituted a light breakfast trial meal, a small amount of dry bread or of toast, from thirty-five to seventy grammes, and a third of a litre, about eleven fluidounces, of warm water or weak tea, which, given on an empty stomach, allows the gastric contents to be tested in an hour, a matter often of great convenience. The results of these trial meals should be filtered for accurate examination.

The next points to determine are the composition of the gastric juice and its digestive power. We first have to ascertain if the liquid obtained be acid, how acid it is, and what its acid nature is owing to. In from ten to fifteen minutes after the trial breakfast acid salts and a free acid are found; the free acid is lactic acid. This disappears within the hour, hydrochloric acid gradually taking its place, and at the end of an hour only hydrochloric acid is found in the normal stomach contents.\* In the ordinary full mid-day trial meal, hydrochloric acid is not likely to be found for an hour or two after the ingestion. It is true that the acid phosphates in the ingesta determine somewhat the acid character of the gastric juice, but the hydrochloric acid is alone, for practical purposes, of importance.

To determine the presence of free acid in the gastric contents, the most delicate reagent is Congo red, which may be employed in solution or in the form of paper impregnated therewith. Free acid causes an azure-blue color; acid salts have no effect. A solution of methyl-violet may also be employed, which is turned into a deep blue; or tropæolin, which in a saturated watery solution is a dark yellowish-red liquid that on contact with any free acid becomes dark brown, while with acid salts it assumes a straw-colored tint. To ascertain the presence of hydrochloric acid the best test is Günzburg's phloroglucin-vanillin solution. It consists of two grammes of phloroglucin and one gramme of

---

\* Ewald, *op. cit.*, p. 21.

vanillin, with thirty grammes of absolute alcohol. A few drops of this solution, which is of a yellowish color, added to a similar quantity of a liquid containing hydrochloric acid, when gently heated, turn it at once a bright-red hue; while the reagent remains unchanged by organic acids, such as lactic or acetic acid. Boas \* recommends a solution containing resublimated resorcin five grammes, white sugar three grammes, dilute alcohol sufficient to make one hundred grammes. Of this, three or four drops are added to five or six drops of the gastric contents, and the whole is gently heated to dryness; a bright-red hue results if free hydrochloric acid is present.

To determine the presence of lactic acid, a matter often of very great value for diagnostic purposes, a solution is prepared of ten cubic centimetres of a four-per-cent. solution of carbolic acid, twenty cubic centimetres of water, and one or two drops of a solution of ferric chloride. This has an amethyst-blue color, which in the presence of lactic acid becomes lemon-yellow or canary-yellow.

The degree of acidity of the gastric juice is more difficult to determine than the presence of the acids. Ewald recommends, as a ready way, to titrate with a one-tenth normal sodium hydroxide solution, ascertaining the saturation point with litmus paper or with phenolphthalein. Wolff † puts one cubic centimetre of the filtered gastric juice into a graduated cylinder, and dilutes it repeatedly until there is no more reaction with the standard methyl-violet solution. The quantitative determination of the hydrochloric acid present is made by titrating with a decinormal sodium hydroxide solution until there is no further response with Günzburg's solution. We may test the solvent power of the gastric juice by taking a piece of hard-boiled egg albumen and adding the gastric juice in a test-tube. Heated in a culture oven, the egg albumen, if the gastric juice be normal, will be dissolved in about an hour. Propeptone and peptone are determined by the biuret reaction. The presence of the lab-ferment or rennet-ferment is shown by the coagulation of neutralized boiled milk by an equal quantity of neutralized gastric contents in ten or

---

\* Diagnostik u. Therapie der Magenkrankheiten, 3. Aufl., I. Theil, 1894, p. 149.

† Trans. Phila. Co. Med. Soc., Oct. 1889.



fifteen minutes in an incubator. After an hour from the time Ewald's trial-breakfast has been taken there should be no reaction for starch found by Lugol's solution in the filtered liquid of digestion.

The *absorptive activity* of the gastric mucous membrane is shown by the rapidity with which iodine appears in the saliva after the ingestion of one and one-half grains of potassium iodide carefully enclosed in a gelatin capsule. In health the characteristic blue coloration is, as a rule, obtained with starch-paper in the course of ten or fifteen minutes.

The *motor activity* of the stomach is determined by the development of a violet color on the addition of a drop or two of a neutral solution of ferric chloride to two or three drops of the urine placed upon bibulous paper, after the ingestion of fifteen grains of salol in gelatin capsules at the height of digestion. The violet color shows the presence of salicylic acid, which is in the majority of persons observed in the course of from sixty to seventy-five minutes. But there are still many clinicians who prefer the older method of examining the contents of the stomach after trial meals, with a view to determine the gastric motility. Leube's method consists in removing the contents of the organ six or seven hours after a trial dinner, or they may be examined an hour after Ewald's trial breakfast. In either case, the stomach should then contain nothing but the liquid of digestion; two hours after the trial breakfast it should be empty. If more than forty cubic centimetres are obtained an hour after Ewald's trial breakfast, it shows insufficient motor activity.

The symptoms which are constantly met with in derangements of the stomach, whether organic or functional, are loss of appetite, nausea and vomiting, acidity, flatulency, and pain.

**Loss of Appetite.**—This manifests itself in various ways. It may amount to absolute repugnance to taking any kind of food, or may be merely an inability to partake of certain articles. What the loss of appetite depends on, we do not know. That nervous influence has something to do with the anorexia, is shown by the sudden deprivation of all desire to eat when any strong impression is made on the nervous system,—such as that caused by the unexpected receipt of unwelcome news. The collection of epithelium on the mucous membrane is also connected with a

marked diminution of the appetite; for with a tongue much coated, absolute disgust at the mere thought of taking food often exists, which yields to relish for food as soon as the tongue begins to clear.

Attending lost appetite, we meet sometimes with great emaciation and with signs as if even the small quantity of food taken were not absorbed into, or utterly failed to nourish, the system. There is apt to be sensitiveness over the abdomen, and spots of particular sensitiveness exist which correspond to the situation of the mesenteric glands. We find, however, no evidence of organic disease, either in the abdomen or in the lungs; nor does this *pseudo tabes mesenterica*, if I may so call it, occur, like the disease it simulates, in scrofulous or tubercular patients. I have met with a number of cases, chiefly in young women with lowered vital force, fond of excitement, and living indolent lives. Some were hysterical, others not. But in all the complaint seemed to be due to deficient nerve-power, with impaired function of the stomach, and very possibly of the abdominal glands. This disorder is probably the same as that described by Gull as hysteric apepsia,\* and kindred to the affection delineated by Lasègue as hysteric anorexia.†

Instead of the appetite being lost, it may be capricious, or even ravenous. A craving after food is not often combined with a structural lesion. Yet we occasionally meet with it in persons affected with gastric ulcer. It is common to find it in those who suffer from neuralgia of the stomach. And sometimes in cases of mere nervous gastric disturbance, with or without pain, there is an extraordinary exaggeration of the appetite, a *bulimia*: the patient eats largely eight or even fifteen times a day, digests his food, yet is constantly hungry.

The feeling of *thirst* does not lessen when the desire for food does. On the contrary, it usually increases when the latter diminishes.

**Excessive Acidity of the Stomach.**—Excessive acidity occurs from various causes. The gastric juice may be secreted in great quantities, or it may contain an abnormal amount of acid. But excessive acidity is far more frequently due to the decom-

---

\* Transactions of the Clinical Society, vol. vii., 1874.

† Arch. Gén. de Méd., April, 1873.

position of food and to a process of fermentation dependent rather upon an insufficient amount or altered state of the gastric solvent. It then manifests itself only after meals. When the mucous membrane is covered with a tenacious mucus or with thick layers of epithelium, slow digestion and acidity from fermentation result; because, although the gastric juice is sufficient, it cannot mix as readily with the aliment.

The acids formed in the stomach are, besides the hydrochloric acid of the gastric juice, lactic acid, acetic acid, carbonic acid, butyric acid, and oxalic acid. Some articles of food produce these different acids in considerable quantities. Thus, sugar generates large amounts of lactic acid. The mode of detecting these acids, and of establishing whether the extreme acidity is due to excess of hydrochloric acid in the gastric juice or to other acids, as tested after a trial meal, has been above explained. In examining for acids, the two acids of greatest value to determine are hydrochloric acid and lactic acid.

The acids which are created in the stomach may give rise to various disorders. When much acid is present it occasions a sensation of heat which extends along the œsophagus. This "heart-burn" is apt to happen in paroxysms, and is attended with a feeling of constriction or with actual pain at the epigastrium. As a symptom it has no special diagnostic value, for it is met with both in functional and in organic diseases of the stomach. It simply denotes great acidity; it is common in gouty persons. It probably arises from the action of the acid contents of the organ on the sensitive nerves of the cardia and of the œsophagus, and the acid is mostly owing to fermentative changes. When the acidity is due to increase of hydrochloric acid, from excessive acidity or quantity of the gastric juice, it is the result of a gastric neurosis; there may be acid vomiting coming on irrespective of food, and happening in the night or during the early morning hours. What has been called *gastroxynsis* by Rossbach is a gastric neurosis appearing at intervals mostly after some psychical or mental disturbance, and marked by extremely acid vomiting and headache, like that of migraine.

**Flatulency.**—The gas in the intestinal canal may be merely air which is swallowed; or it may be generated from imperfectly-digested food; or it may be a secretion from the blood-vessels of

the part. In those who suffer from indigestion, it is produced in the last two ways, and the patient complains greatly of the annoyance it occasions. It causes a disgust for eating, a feeling of distention, and sometimes actual pain. By interfering with the downward movements of the diaphragm, it induces a sensation of constriction in the chest, shortened breathing, palpitation of the heart, and the sleep is broken by uneasy dreams.

An expulsion of the gaseous contents of the stomach by the mouth gives rise to *eructation*, or belching. The belching which follows the decomposition of food has sometimes the taste and the odor of rotten eggs, owing to the gas evolved consisting of sulphuretted hydrogen. At other times the eructation is odorless, because the gases formed are carbonic acid, or hydrogen or nitrogen, or some of their compounds. When the gas results from fermentation or decomposition of food, it frequently coexists with acidity occurring only after meals. When it is a secretion from the blood-vessels, it happens in an empty state of the stomach, and is often relieved by avoiding too long intervals between the meals. As a cause of flatulence and eructation which it is important not to overlook may be mentioned thoracic aneurism.\* Marked flatulency is often only a form of gastric neurosis. It is common in nervous dyspepsia and in hysteria.

**Nausea and Vomiting.**—These are often combined. But sometimes there is persistent nausea without vomiting; sometimes vomiting occurs without any or with but slight nausea. Yet they are both occasioned in much the same way: what gives rise to one will generally give rise to the other.

Vomiting is a complex act. But its causes, although various, may all be arranged under four heads. It either arises from an irritation of the peripheral extremities of the nerves which supply the parts more directly concerned in the act itself, such as the stomach, the diaphragm, and the œsophagus; or the irritation originates in the centres from which these nerves spring, and is referred to their peripheries; or there is a mechanical obstruction in the stomach or intestines; or the vomiting is purely sympathetic. Under the first head belongs the vomiting observed in acute or chronic inflammation of the stomach, in ulcer, or in can-

---

\* Walter F. Atlee, Amer. Journ. of Med. Sci., July, 1869.



cer; also that following a debauch, or the introduction of irritating substances into the viscus. Under the second head may be ranged the vomiting which occurs in diseases of the brain; perhaps, also, that which arises in morbid states of the blood, as in uræmia. Under the third head we may class the vomiting in narrowing of the œsophagus and of the pyloric or cardiac extremity of the stomach, and in obstructions of the intestine. The fourth group is exemplified by the vomiting in pregnancy, in wounds of the extremities, in inflammation of the peritoneum, of the intestines, and of the liver, in renal calculus, and in irritation of the fauces.

Connected thus with so many various conditions, the act of vomiting, taken by itself, is of little diagnostic value. It presupposes a certain amount of irritation existing in the stomach, or reflected to it; but nothing more. It is, of course, a frequent symptom in disorders of the stomach, especially in those which are organic; yet the error of considering it as having reference only to derangements of that viscus ought to be strenuously guarded against. As it is allied to morbid states too numerous to be here examined in detail, I shall content myself with making general statements regarding the indications to be drawn from it.

When vomiting is observed in a person previously in good health, we may suspect either the invasion of some acute malady, or that some poisonous substance has been swallowed. Again, it may come on suddenly from violent mental emotion. When everything that is taken is immediately expelled, the difficulty lies in the œsophagus, or at the cardiac orifice of the stomach, or in an extreme irritability of the viscus; and this irritability, attended as it often is with unceasing nausea, experience teaches to be more frequently due to sympathetic excitement of the organ than to structural gastric disease. But speedy vomiting, generally without preceding nausea, it must be remembered, is also among the symptoms of visceral hysteria, and is indeed, by some, regarded as the most frequent symptom.\* I have known it associated or alternating with extraordinary flatulency.

*Nervous vomiting* occurs where there is no lesion in the stomach or irritation of food as the cause. It is mostly due to reflected

---

\* Denian, *L'Hystérie gastrique*, Paris, 1883.

irritation of the nerve-centres controlling the act of vomiting, and is often found in disorders of the uterus ; or arises from direct irritation of the nerve centres in affections of the brain and cord. It is common in hysterical subjects. It is not associated with nausea, and may be of long duration. It is sometimes a primary gastric neurosis, and as such is seen, particularly in neurasthenics, in association with the condition described by Kussmaul as "peristaltic unrest." This is a very annoying symptom, in which there are loud borborygmi and gurgling, especially after eating.


As regards the vomiting which is brought about by gastric disorders, it is of much consequence to note the period at which it happens, whether before meals or after meals, and how long afterward. In some diseases, such as ulcer and cancer, it rarely occurs except when food has been taken. The act of vomiting then affords relief from the pain. In narrowing of the pylorus, it takes place some hours after digestion has begun. But, as vomiting will be described hereafter in its relations to the individual diseases of the stomach, we shall not dwell on what will be more fitly discussed elsewhere. Yet a few words on the characteristics of the ejected matter can hardly be omitted.

The nature and the quantity of the vomit are, of course, most various. The following are its most common kinds :

*Food* or *liquid*, mixed with saliva and some mucus, is expelled when the stomach is very irritable, or if an obstruction exist which renders the entrance into the organ difficult or impossible. Half digested food, in a state of acetous fermentation and with a strongly-acid reaction, is cast out when there is deficiency of hydrochloric acid, or when the food has been detained for a long time in the stomach. In the ejected matter the particles of food may be recognized ; but when the food has been kept for a prolonged period in the stomach, or when it has passed on into the duodenum and is returned, it is changed into an apparently homogeneous mass. Examined under the microscope, the structures of the animal or vegetable substances partaken of can even then be detected. Mixed with muscular fibre, elastic tissue, starch-corpuscles, and vegetable cells, is found usually a quantity of oil-drops and fat-crystals. The starch corpuscles are turned blue by a solution of iodine and iodide of potassium.

*Sarcinæ* and yeast fungi are sometimes discovered, by means of the microscope, in the vomit. These organisms, first described by John Goodsir, are the result of a process of fermentation, and are generally associated with copious vomiting. They are small square or slightly oblong bodies, divided into similar smaller portions by cross-lines, and each portion thus formed is again subdivided; but the markings of the smaller squares are not so distinct as those of the larger. The illustration shows a mass of *sarcinæ* found in the vomit of a patient who suffered from gastric ulcer.

FIG. 54.



Sarcinæ ventriculi.

Vomit containing *sarcinæ* is always indicative of structural change in the stomach. It is sometimes found in chronic gastritis of long standing; or in connection with ulcer, and yet oftener with cancer, and especially in those cases in which the narrowing at the pyloric extremity has led to distention of the organ; indeed, any form of dilatation or a condition preventing the stomach from completely emptying itself pre-eminently gives rise to it.

*Sarcina* vomit has an acid smell and reaction, and often a peculiar brownish appearance. After standing, it becomes covered with a dirty, frothy matter, like yeast. A solution of iodine and iodide of potassium turns the *sarcinæ* mahogany brown or a violet hue; but it is by the microscope that their presence can be recognized with greatest certainty. The process of fermentation attending the development of the *sarcinæ* occasions heart-burn and extreme flatulency, and the copious vomiting is a source of relief.

*Mucus* is occasionally ejected in large quantities, both mixed with food and pure. In chronic gastritis, and in the milder forms of acute gastritis, the mucous membrane is covered with a tenacious secretion, and a considerable amount of a glairy or stringy matter is expelled by the act of vomiting. As a general rule, indeed, it may be stated that, when much mucus is evacuated, a catarrhal state of the stomach is present.

A *thin, watery fluid*, looking much like saliva, is discharged in some cases of organic disease of the stomach, as well as in functional derangement of the organ brought on by eating coarse food. Now and then it is met with in pregnancy. This variety of vomiting is known as *pyrosis*; popularly, as “water-brash.”

It may be attended with a burning sensation extending to the fauces, and with pain running back to the spine. The fluid is commonly alkaline; sometimes, owing to its intimate admixture with the gastric contents, it is acid. Frerichs found that it possessed the power of converting starch into sugar. It is mostly regarded as being formed by the glands at the lower part of the œsophagus as well as of the stomach.

*Bile* may find its way into the stomach, and be expelled by the mouth, imparting to the vomit a greenish or yellowish color and a very bitter taste. The occurrence of bilious vomiting is commonly held to indicate a disease of the liver, or that the patient is extremely "bilious." It is not a proof of either. It is observed when there is much retching, and when the act of vomiting is protracted and frequently repeated, and is chiefly met with in the various forms of acute gastritis, and at the invasion of some acute malady which gives rise to sympathetic gastric disturbance.

*Fæcal* vomiting never depends upon a disease of the stomach. It may be possibly owing to a fistulous opening between the colon and the stomach; but such cases are extremely rare. Generally it is due to a mechanical obstruction to the passage of fæces. Occasionally it happens in fevers of a low type, or in peritonitis, and is then, perhaps, the result of paralysis of a portion of the intestinal tube, which acts, to some extent, as a mechanical obstruction. The matter that is ejected has the odor of fæces; but it is commonly of less firm consistence, and of lighter color, because it is the contents rather of the small than of the large intestine. Sometimes it is perfectly fluid. In fæcal vomit a considerable number of large comma-like bacilli have been noticed.\*

*Pus* in small amount is sometimes found mixed with the vomit in cases of large ulcers in the stomach, simple or cancerous. When in quantities, it is owing to an abscess in the neighborhood of the viscus having poured its contents into it. Still, pus is rarely met with in the matters expelled. And the same can be said of other substances that may find their way into the stomach, like echinococcus sacs and worms, and masses of false membrane.

---

\* Jaksch, Klinische Diagnostik.



*Blood*, on the other hand, is not infrequently vomited. Having described the appearance of the blood when it comes from the stomach, in treating of the diagnosis of hemorrhage from the lungs, I shall, before examining into the circumstances which cause a hæmatemesis, merely here recall the fact that it is preceded by nausea and followed by black stools, and that the fluid ejected is generally black, and presents an acid reaction.

The quantity of blood lost varies greatly; but the amount vomited is by no means a proof of the amount effused. The larger portion may pass off by the bowels, giving rise to peculiar tarry stools. Nay, the whole may be voided with the stools. Chocolate-colored material discharged by stool, and due to alkaline fluids acting on the blood after the effect of acids, is held to be a distinguishing trait between the blood passing by the intestines after a gastric hemorrhage and bleeding from the bowel.\*

*Hemorrhage from the stomach* is variously caused. It may spring from injury to the organ, or from disease of its coat; it may be vicarious; it may be the consequence of disorder elsewhere than in the stomach, as of a mechanical obstruction in the portal system; it may depend upon an altered state of the blood.

In the hemorrhage that follows blows or kicks on the stomach, an *active* hyperæmia of the mucous surface is occasioned, which leads to the extravasation of blood. An active arterial hyperæmia also precedes the hemorrhage that sometimes follows the swallowing of irritant poisons; and it is probably the cause of the hæmatemesis in several of the organic affections of the stomach. Of these, only cancer and ulcer are apt to present hemorrhage as a prominent symptom; and of these, again, it is much more frequent in the latter than in the former. The blood effused may be so slight in amount as to escape detection; and this is especially likely to happen when it is intimately admixed with food or with bile. Yet, by means of the microscope, the existence of blood-corpuscles in the ejected matter can be always demonstrated. The fulness of the vessels may be associated with degeneration of their coats, as, for instance, in amyloid degeneration of the stomach.

When blood has been detained for some time in the stomach, and has become intimately mingled with the acid contents of the

---

\* Bartholow, Practice of Medicine.

organ, it loses entirely its natural appearance. What is termed "coffee-ground vomit" is blood thoroughly intermixed with other substances. It is the result of a comparatively small or gradual hemorrhage; and, as this is the kind apt to happen in gastric cancer, it is common in this affection, though by no means limited to it.

Vicarious hemorrhage from the stomach is not infrequent, and especially frequent is that which takes the place of the menses. It is not dangerous. The blood escapes at the time of the normal discharge, and while the bleeding lasts the stomach is slightly tender, and the digestion impaired. But during the intervals there are no signs of disturbance of the functions of the organ, and no pain; both of which are points of importance in distinguishing between loss of blood caused by suppressed menstruation and loss of blood caused by disease of the stomach.

Gastric hemorrhage, dependent upon a state of *passive* congestion brought on by an obstruction to the flow of venous blood, is occasionally seen in organic affections of the heart. But it is much more common as the result of embarrassment of the portal circulation, from tumors, or from affections of the liver and spleen. It frequently attends, therefore, cirrhosis and enlargement of the spleen, and is often joined to intestinal hemorrhage.

In gastric hemorrhage resulting from changes in the blood the vessels themselves are toneless, and rupture easily or offer no resistance to their altered contents escaping. This kind of hemorrhage is met with in scurvy, in typhus fever, and in yellow fever.

We see thus that blood is vomited from various causes, and that merely from the occurrence of hæmatemesis we can determine nothing definite as to its origin. Yet the symptom—for a symptom it always is—is of serious import, and when taken in connection with others is of great service in diagnosis. We ought, in chronic cases, first to suspect the hemorrhage to be due to some organic disease of the stomach; when there is no other proof of a structural affection of this organ, we turn to the liver, spleen, or heart for its explanation, or examine carefully every part of the abdominal cavity, to see whether or not a tumor is the source of the disorder. If occasioned by none of these conditions, its cause lies probably in altered blood, or in suppressed discharges. The history of the case is indispensable to any

induction. Thus, in low fevers there is no doubt as to what has brought about the hemorrhage.

There is, however, one difficulty present in all instances; and that is, to tell whether the ejected blood has found its way into the stomach and has been subsequently expelled, or whether the hemorrhage is really gastric. The only method to avoid being deceived is to scrutinize closely the history and the attending phenomena. Blood may be introduced into the stomach by the bursting of an aneurism, or from an ulcerating pancreas; or it may have been swallowed during an attack of epistaxis or of hæmoptysis, or wilfully, to excite sympathy or to escape punishment. A strange result of gastric hemorrhage, first noticed by Graefe, is double-sided incurable amaurosis.

So much for vomiting of blood, and for the different characters presented by the vomit. In describing them we have been led away from the indications they furnish in diseases of the stomach. But it was more convenient here to consider vomiting connectedly and somewhat in detail, than to be obliged to treat of it in various chapters. To return now to the more special symptoms of a deranged stomach.

*Merycism, or Rumination.*—In this condition food that has been swallowed is brought up into the mouth, sometimes by an impulse of the will, but more commonly involuntarily, and remasticated and again swallowed. Rumination is recognized to be purely a neurosis, and may or may not be associated with other gastric disorder.

*Regurgitation* of fluid or partly digested food may take place in connection with a relaxed condition of the cardiac orifice of the stomach, and, if obstinate, may lead to pronounced derangement of nutrition. This phenomenon is to be distinguished from a similar occurrence that takes place when the œsophagus is the seat of a pouch or diverticulum that empties itself from time to time.

**Pain.**—Pain occurs in gastric disorders in every conceivable form. It is sometimes slight, at other times violent. It is often more a feeling of soreness than actual pain. It may or may not be increased by pressure, and may be either augmented or relieved by the taking of food. If persistent or severe, and accompanied by tenderness at the epigastrium, it is almost always linked to a morbid state of the tissues of the viscus. Mere uneasy sensa-

tions, on the other hand, also happen in functional derangement of the organ while the food is being digested, and may even be attended with slight tenderness at the epigastrium.

Now, as both pain and soreness to the touch may be present as well in functional disturbance as in organic change, how can we tell with which they are associated? As a rule, pain and soreness dependent on organic disease may be distinguished from pain and soreness that result from functional disorder by noticing the time at which they take place. If they are more severe soon after meals, or when the stomach is full, and worse after a heavy meal than after a light one, especially of a bland substance like milk, they point to a structural affection. If they occur only when the stomach is empty, and are relieved by food, they are indicative of a functional derangement.

Occasionally the stomach is the seat of violent paroxysms of pain. These are at times linked to a chronic organic affection; at others they are apparently connected with a perfectly sound state of the viscus, and coexist with a tendency to neuralgic pains all over the body, or with hysteria or neurasthenia; or they may appear as the gastric crises of locomotor ataxia; at others they are brought about by some article of food which the stomach does not tolerate or is unable to digest. The disorder is variously described under the name of *gastrodynia* or *gastralgia*. The pain is supposed to be associated with, or due to, a cramp of the stomach; but whether it is so or is a pure neuralgia is uncertain. When the predisposition to it exists, exposure to cold and damp, a draught of cold water drunk when heated, sudden and violent emotions, or a collection of wind in the alimentary canal, will bring it on. The predisposition is met with in gouty and rheumatic persons, and in those who are debilitated,—in women who are anæmic, and in men who have been exposed to exhausting influences. Then we also find the *gastralgia* interchanged with other neuralgic or spasmodic affections, giving way to asthma or to angina pectoris, or, on the other hand, occurring in their place.

The pain varies much in intensity: it is usually severe and agonizing; but it is not permanent; intervals of rest and comfort succeed to the paroxysms of distress. During a violent attack, the skin is cold, the pulse slow, there are frequently nausea, vomiting, sometimes fainting, and often sensations of utter prostration and



impending dissolution. The seat of the pain is in the epigastrium, immediately beneath the ensiform cartilage. The patient feels as if the coats of the stomach were being violently drawn together, or rent asunder, or rapidly pierced by a sharp instrument. Thence the pain extends toward the umbilicus and the hypochondria. It is sometimes relieved by the recumbent position and by external pressure. But relief depends much on the condition with which the pain is associated. If it be connected with a chronic gastritis or an ulceration, external pressure aggravates rather than alleviates it. This is certainly true as a rule; yet we cannot positively announce that pain with tenderness at the epigastrium is proof of an organic lesion. There is sometimes sensitiveness to the touch in purely nervous gastralgia; or slight pressure may augment the pain, but firmly compressing the pit of the stomach will diminish it.

In a practical point of view, it is very important to discriminate between the cases of gastralgia that may be viewed as pure neuroses and those in which the paroxysms of pain are combined with a chronic lesion. We infer that we have to deal with instances of the former, when the attacks occur in those whose impoverished blood or enfeebled health predisposes to neuralgia, and especially if they happen in women laboring under disorders of the uterus or of menstruation, or in persons who suffer from neuralgic pains in other parts of the body. But the broadest line of distinction is drawn by the state of the digestive apparatus during the intervals. The disordered digestion, the pain after eating, the tenderness at the epigastrium, the nausea and vomiting, and the other symptoms common in morbid alterations of the coats of the stomach, are not seen in pure neuralgic gastrodynia. Too much stress, it has been already stated, ought not to be laid on the influence of pressure on the paroxysmal pain during the paroxysm. A sign more trustworthy is the alleviation following the taking of food, for which, in truth, there may be a craving; and occasionally cases of gastralgia are met with in which the pain occurs only early in the mornings, and is very distressing, but is almost immediately eased by a hearty breakfast.

Gastralgia is common where there is an excess of hydrochloric acid in the gastric juice. The form of gastralgia which is produced by some article of food that disagrees with the individual

is readily distinguished from the other varieties by observing it to be transient, and by noting its cause. The indigestible substance undergoes fermentation in the stomach, and acidity, flatulent distention, and nausea attend the pain, which ceases when the offending matter is ejected and the gas expelled.

The remarks just made apply also, in the main, to other manifestations of perverted innervation of the stomach, such as hyperæsthesia, with or without persistent vomitings,—forms happening usually in weak or hysterical persons, or where menstruation is disordered,—but which in the present state of our knowledge are still conveniently classed with gastralgia as forms of gastric neuroses.

The nervous filaments, the irritation of which occasions pain in the stomach whether paroxysmal or not, belong to the vagus; sometimes, perhaps, the distress originates in the branches of the sympathetic that supply the organ. But we must be careful not to ascribe the seat of every pain which is felt between the umbilicus and sternum, or referred there, to the stomach. Diseases of the pleura, of the heart and its covering, affections of the intercostal nerves, abscess of the liver, intestinal disorders, rheumatism of the abdominal muscles, may give rise to pain in the epigastric region. And, again, spasmodic pain like that of gastralgia may be caused by colic, by disorganization of the tissue of the kidney and of the pancreas, and by the passage of gall-stones or of renal calculi. The great safeguard against error is to bear in mind that painful complaints of the stomach may be mistaken for those enumerated, and to ascertain carefully, in cases of epigastric distress, that there is no cause beyond the stomach to account for it. The nearer, in many instances, the pain is to the median line, or, should it occupy this, the more fixed and confined to a small spot, the greater is the probability of its being dependent upon gastric disease; and pain of the character alluded to is generally indicative of serious malady.

Pain is the last of the symptoms directly referable to the derangement of the viscus itself to which we shall advert. But when the great organ of assimilation is disordered, other organs suffer, either through sympathy, or because the irritation is transmitted to them, or because a similar state of their mucous surface is induced. The bowels are usually in a sluggish condition; it is

commonly only when the gastric acidity is extreme that they are relaxed. The viscera within the chest are frequently disturbed. The patient is annoyed by palpitation and shortness of breath after meals; and as he feels the agitation of his heart, and finds that always, after he has eaten, his face is flushed, the palms of his hands are hot, and his temporal arteries throbbing, he is apt to overlook the derangement of the stomach, and to fancy himself laboring under an incurable cardiac affection. A dry cough, also, is a not unusual concomitant; but a cough may be the result of coexisting catarrh of the bronchial mucous membrane, or of disease of the lung-structure; and sometimes the affection of the lungs precedes that of the stomach. Again, we may have an organic disease of the heart determining the gastric symptoms.

So, too, with the kidneys. They may be irritated by the crude material which has made its way into the blood, and which they are called upon to excrete. The urine often contains various abnormal constituents, especially urates; yet not seldom a morbid state of the urine is found previous to the derangement of the stomach, and the indigestion is the secondary rather than the primary ailment. Indeed, we must never be too hasty in concluding, when a disordered stomach is associated with diseases of other viscera, that it is their cause; it may exist as their consequence. Diseases of the liver and intestines are especially prone to induce a gastric affection.

One of the worst results of a disordered digestion is the state of mind it produces. It occasions listlessness and a disposition to look at all events in a gloomy light, and sometimes brings on inveterate hypochondriasis. Aretæus ascribed to the stomach as its primary power that it acted as the president of pleasure and of disgust, "being, from the sympathy of the soul, an important neighbor to the heart for imparting good or bad spirits." Now, although no one at present would agree with this physiology, who will deny that there is in the remark a germ of truth? But here, again, we must be careful not to confound cause with effect; for want of activity or a distressed state of mind may seriously impair the appetite and subvert the normal action of the viscus.

When the nervous symptoms are marked, the disorder is often called *nervous dyspepsia*. In this, while the gastric symptoms are slight, we may also have the gastric neurasthenia leading to

extreme acidity of the gastric juice, to eructations, to flatulency. There may be sensations of distress and uneasiness during the digestive act, but the gastric motility is not impaired. There is not always increased acidity of the gastric juice. The hydrochloric acid may be normal or diminished in amount. In all forms there are uneasy feelings after meals and great nervousness. Nervous dyspepsia is common in neurasthenics and in hysterics. The exact state of the stomach that coexists can be determined only by chemical investigation of the gastric secretion.

In the rough sketch just finished of the symptoms encountered in gastric disorders, no attempt has been made to separate strictly the signs which belong particularly to alteration of its coats from those which occur in mere derangement of its functions; in other words, I have not tried to dissociate the symptoms of so-called "dyspepsia" from those of actual lesions. And this for two reasons: in the first place, the most palpable indications of organic disease of the stomach are those of disordered function; and secondly, there are no symptoms which belong exclusively to functional indigestion.

Nor is it possible to present anything like a complete picture of merely functional, or, as it is still called by some, *atonic dyspepsia*; the combinations are too infinitely varied. The great difficulty is that the faulty performance of the act of digestion is too often regarded as the whole ailment. The time must come when the term dyspepsia will be banished. It is true that in an organ like the stomach it is particularly difficult to tell where disturbed function ceases and anatomical change begins. Yet that this can be done to a greater extent than is usually done, cannot be gainsaid. Moreover, there are many affections which have probably connected with them definite anatomical lesions and constant modifications of the gastric juice and of the secretions of the mucous follicles of the stomach, which we are as yet obliged to embrace under the name of dyspepsia, because we are unacquainted with their clinical expression. But we may fairly expect that, through those admirable clinical and pathological researches that have of late begun to illuminate the subject, the limits of purely functional indigestion will be much reduced; so that what the physician of the present day is still compelled to class under the term functional indigestion will be recognized



by the physician of the twentieth century as several distinct affections, each with its characteristic fine structural change or its recognized chemical or motor variations,—much in the same way that the physician of the eighteenth century was obliged to regard and to treat dyspnœa as an individual disease, while now we have learned to separate it into different varieties, in conformity with its prominent anatomical causes, and to treat it in accordance with its source.

### Diseases of the Stomach in which Pain and Soreness at the Epigastrium, and Vomiting, occur.

After what has been premised, it is obvious that the structural diseases of the stomach present but few symptoms that can be regarded as at all characteristic. Indeed, the only ones which can lay any claim to be so considered—and we have already seen that this claim is not always valid—are pain and soreness at the epigastrium, and vomiting. We may, then, take these symptoms as a starting-point in diagnosis, and describe the individual organic affections in which they chiefly occur, speaking first of the acute.

**Acute Gastritis.**—This malady is pronounced to be exceedingly rare, save as the result of irritant poisons. Undoubtedly, inflammation of an intense kind, involving more than the mucous membrane, originating spontaneously, and not from the introduction of any highly acrid or corrosive substance, is very seldom met with. But it is no less certain that inflammation of a less active character, limited to the most important part of the stomach, to the mucous membrane, and especially to its surface, the so-called gastric catarrh, is far from being a rare disease, and, whether as a concomitant of fevers or as a separate malady, is a disorder to which the physician's attention is constantly drawn.

Thus, then, acute inflammation of all the coats of the stomach, or even of the entire mucous membrane, is uncommon; acute inflammation of its surface is common. Yet it is the doctrine of the day not to regard any case as acute gastritis unless serious changes have been wrought by the inflammation in the tissues of the organ, so serious as almost to preclude recovery. But I think that, irrespective of the ordinary catarrhal cases, there are instances in which much more severe and deeper inflammation happens, and that a disease exists fully entitled to

be called acute idiopathic gastritis. I am sure that I have seen cases which differed in nothing from the typical and graphically-described cases of Andral,\* save in the fatal termination and in lacking the symptoms which immediately precede that termination.

I shall detail one which was striking. A robust woman, the mother of several children, whom she was obliged to support by hard labor, was suddenly seized with a pain in the epigastric region, and vomiting. There was no apparent cause for the attack : she had certainly not swallowed any irritating substance. Although at one time a sufferer from indigestion, her digestive organs had not been markedly disordered for weeks prior to the appearance of the pain and the irritability of the stomach. The former seemed to come on before the latter. It was of a dull character, increased by swallowing either solids or liquids, and associated with the greatest tenderness. Nausea was constant, and vomiting very frequent. Large quantities of a greenish fluid were ejected, as well as nearly everything she swallowed. The tongue was deeply coated ; its edges and tip were red. The bowels were constipated, but not painful on pressure. There was fever, —not, however, of an active type ; it rose toward evening ; the pulse was quick and small ; the breathing was hurried, and the patient exceedingly restless and prostrated. She complained most of the distress in her head, and of violent thirst. The treatment pursued consisted mainly in opening the bowels by enemata, and in administering ice and repeated doses of calomel, some of which she retained. After the symptoms had lasted for about ten days, they gradually disappeared, and she slowly recovered. The pain on swallowing and the soreness at the epigastrium were the last to leave. Indeed, when she passed from under my care they had not ceased entirely.

Now, here is a case which presented all the symptoms of a severe inflammation of the stomach, similar to that produced when an irritant poison has been received into the organ. In all such instances there are the same nausea and vomiting, and pain ; the same restlessness and headache ; the same form of fever and small or feeble pulse ; the same unquenchable thirst. Sometimes the

---

\* Clinique Médicale, tome ii.

pain is of a burning kind ; and in those cases which prove fatal—and many do prove fatal, as much perhaps from the destructive effect of the irritant on the tissues as in consequence of the inflammation—there is hiccough, the skin becomes cold, the features collapse, and the sufferer dies prostrated, yet frequently preserving his mental faculties to the last.

From these severe cases of acute gastritis, however caused, there exists every grade of inflammation, down to an active congestion of the mucous membrane, and to a mere reddening of its surface. Of course, there will not be in the milder forms the same intensity in the symptoms. But the outline is the same, although the filling in be in far less vivid hues. There is in all the same tendency to nausea and to vomiting, with more or less epigastric pain and uneasy sensations, and more or less tenderness at the pit of the stomach, and headache.

A mild gastritis is very commonly brought on by a debauch or by the introduction of irritating articles of diet into the stomach.

These cases are classed as *acute gastric catarrh*, and are popularly known as severe attacks of indigestion : that they are owing to an inflammatory state of the mucous membrane was proved by the ocular demonstration Beaumont had of the process in the person of Alexis St. Martin. The symptoms that inflammatory change, when it was marked, produced, were some tenderness at the epigastrium ; nausea ; vomiting ; constipation, or sometimes diarrhœa ; a coated tongue, and headache,—in fact, just the symptoms of which patients complain when they are suffering from an acute attack of indigestion.

Another common and kindred kind of mild inflammation of the stomach or acute gastric catarrh is that usually called a “bilious attack.” The French designate it expressively as *embarras gastrique*. It is a catarrhal affection, and often associated with catarrh of other mucous membranes. It may come on from indigestible food, or after cold and exposure ; it sometimes occurs in epidemics. The symptoms are those already detailed. There is nausea, and frequently bile is vomited. We do not usually observe much pain in the epigastrium ; but rather a feeling of uneasiness, and a slight soreness to the touch. The urine is dark, and deposits urates ; the tongue is much coated ; there is thirst, with generally a moderate or slight fever, which exacerbates at

night, and is of remittent type, and there may be a yellowish tinge of the conjunctivæ. In children acute gastric catarrh may become complicated with convulsions, or with symptoms simulating those of meningitis.

*Secondary* acute inflammation of the mucous membrane of the stomach is found in association with various disorders. It is met with in remittent fever, in typhus, in the exanthemata, in rheumatism, and oftener in gout, and partakes somewhat of the specific character of the malady with which it happens to be combined. Indeed, instead of being a secondary inflammation, it is oftener, to speak correctly, a local expression of a constitutional state.

Several writers describe a form of gastritis which occurs in very young children and leads to softening of the mucous lining of the stomach, a *gastromalacia*. This softening is most likely a post-mortem change due to the action of the gastric juice, and especially met with in the subjects of acute gastric catarrh. Kundrat has called attention to the occurrence of gastric softening with vomiting of blood in the brain affections of children, especially in tubercular meningitis.

Sometimes a *suppurative inflammatory process* takes place in the submucous and muscular coats of the stomach. This may be primary, arising through local infection, or secondary or metastatic. It is characterized by intense epigastric pain, which may set in suddenly or be preceded by vague symptoms of digestive derangement. In addition there are a sense of burning, intense thirst, dryness of the tongue, and complete anorexia. The general depression is profound, and the temperature is usually high. The vomited matter may contain pus. Death may be preceded by general restlessness, delirium, and coma.

*Chronic Diseases attended with Pain, Epigastric Tenderness,  
and Vomiting.*

The chronic diseases of the stomach may, like the acute, be considered in accordance with the pain, the soreness at the epigastrium, and the vomiting that attend them. At all events, these are the symptoms common to the chronic diseases which are susceptible of diagnosis. Besides these, there are some chronic dis-



orders with the morbid anatomy of which recent careful researches have made us familiar,—such as destruction of the tubular structures; hypertrophy of the solitary glands; interstitial growths leading to glandular wasting, and to a gradual fibroid thickening of the entire mucous or submucous coat; fatty degeneration of the atrophied masses,—but which we are as yet unable to distinguish with any certainty at the bedside, notwithstanding much recent valuable work.

Contrasting the chronic diseases with which we are clinically acquainted with the acute, vomiting is found to be a symptom of greater diagnostic value,—not the act itself, but the appearances of the ejected matter. And, further, the phenomena of dyspepsia stand forth much more conspicuously.

**Chronic Gastritis.**—In chronic inflammation of the mucous membrane, or *chronic gastric catarrh*, the symptoms of indigestion are persistent and manifold. They vary somewhat according to the extent of the mucous surface involved and the amount of mucus and epithelium which accumulates on it, and probably also according to the healthy or wasted state of the gastric glands. Generally there is a sensation of discomfort, of weight, and of soreness at the pit of the stomach, aggravated by food; the part is also tender to the touch. Sometimes, even when the stomach is empty, a burning at the epigastrium and an inward fever are complained of. The appetite is impaired or capricious. Fermentation, heart-burn, and flatulency frequently attend the slow digestion of the food; the tongue is usually heavily coated; it may, however, be clean. The bowels are constipated. The urine contains an excess of phosphates or of urates, or exhibits crystals of oxalate of lime. The patient's circulation is languid; he suffers from chilliness; his spirits are depressed. Not unfrequently he is annoyed by thirst, and by vomiting, after meals, the half-digested food mixed with strings of mucus. But the vomiting may also take place when the stomach is empty, and the ejected matter is then fluid and colorless. Drunkards who suffer from chronic gastritis often throw up a quantity of glairy fluid on rising in the morning. A colorless vomit, joined to symptoms of long-continued indigestion, is very characteristic of chronic gastritis.

The gastric contents removed after a trial meal show a diminution in the amount of hydrochloric acid present, usually in the

total acidity also, and in the activity of the digestive ferments; still, hydrochloric acid is generally present. When atrophy of the gastric tubules has taken place there is complete absence of hydrochloric acid and of the digestive ferments. The fasting stomach is usually found to be empty.

Thus, then, the results of chemical examination of the removed gastric contents, the character of the vomit occasionally, more frequently the coated tongue, the distress after eating, the soreness at the epigastrium, and the persistency of the symptoms, distinguish the dyspepsia of chronic inflammation of the stomach from that which is purely functional.

The causes of the malady are at times obscure. It certainly cannot be traced often to an antecedent acute attack, although those who suffer from the chronic disorder are particularly prone to acute exacerbations. It is more common in persons over than in those under forty years of age. It is especially common in gourmands and drunkards, and in those who live on coarse food or who eat irregularly. It is often found conjoined with chronic bronchitis, and sometimes with Bright's disease, with tubercular disease of the lungs, with gout, and with diabetes. Passive congestion undoubtedly acts as a predisposing element. The inflammation is seen to arise from this cause in the course of chronic affections of the heart or of the liver, and of obstructions to the portal circulation, whether complicated with a lesion of the liver or not.

Chronic gastritis is frequently associated with ulcers in the organ or with cancer, and many of the symptoms of these disorders are clearly attributable to it. Let us inquire whether there are any special symptoms to inform us that something more dangerous than chronic inflammation of the mucous membrane of the stomach exists.

**Gastric Ulcer.**—Ulcer of the stomach is a disease comparatively rare in this country; but it is not so in some parts of the Continent of Europe and in England. The affection is essentially dependent upon disturbance of the normal relation between the gastric secretion and the circulating blood, in that the one is unduly active or the other is deteriorated in quality. It is more common in females than in males, and between twenty and forty years of age than at any other period. It is generally associated

with anæmia, or follows chronic gastric catarrh, or embolic plugging of small arterial twigs, or other disturbances of the circulation in the gastric mucous membrane. Amyloid degeneration of the finer vessels, too, occasions these perforating ulcers. The acid gastric juice acts readily and destructively on the weakened tissues. Rarely, gastric ulceration is due to tuberculosis and to syphilis.

The ulcer or ulcers, for there are sometimes several present, are seated usually on the posterior wall of the stomach, in or near the lesser curvature and toward the pyloric extremity. The great danger arises from perforation of the coats and subsequent peritonitis. But the ulceration may prove fatal by opening a large blood-vessel. Again, the formation of a gastro colic or a gastro-pulmonary fistula may lead to death; or the protracted suffering and excessive vomiting may gradually exhaust the vital energies. On the other hand, the ulcers may heal by cicatrization; and this, William Brinton tells us, takes place in about half the instances. Perforation, Welch states, happens in about six and a half per cent. of all cases. Recurrence of the gastric ulcer is not uncommon.

In cases which may be regarded as typical, the malady is announced by symptoms exactly like those witnessed in chronic gastritis,—the same uneasiness and pain at the epigastrium, and occasional nausea and vomiting of food, or of a watery fluid. Perforation may at this early stage of the disease most unexpectedly cut short the patient's life. Should perforation not take place, hemorrhage from the stomach, with emaciation and anæmia, next appears. In this way the disease usually continues for months or years, the symptoms remitting from time to time, and showing singular variations in their severity. Welch\* states the average duration of gastric ulcer to be from three to five years. The majority of the cases recover.

Of the symptoms, pain and vomiting are the most characteristic. Pain is rarely absent; never, perhaps, except in cases which run a rapid course. It is generally a continuous dull feeling; sometimes a burning, at other times a gnawing sensation. As a

---

\* Pepper's System of Practical Medicine, article "Simple Ulcer of the Stomach."

rule, it is rendered more acute within a quarter of an hour after eating, and remains so as long as food occupies the stomach. Its situation is commonly in the middle of the epigastric region, and there it continues strictly limited. At that point, too, there is localized soreness, or even great tenderness to the touch. Sometimes the pain is seated behind the ensiform cartilage, or is referred to the right or to the left hypochondrium. It is often associated with a gnawing pain in the lower dorsal vertebræ, which may shoot between the scapulæ or down the spine; but the dorsal pain, like the epigastric, is, on the whole, very fixed, radiates but little, and is most severe when the ulcer is on the posterior surface. Besides this continued feeling of distress, there occur violent paroxysms of pain, which may last for several hours; nay, with trifling intermissions, for days. They sometimes come on suddenly when the viscus is empty, but are aggravated by pressure or by food; and, in fact, they are often thus induced. The patient refers the suffering chiefly to the pit of the stomach, or to the dorsal vertebræ. He is apt to seek the recumbent posture for its relief. Yet it is remarkable that there are at times long intervals during which all pain, whether paroxysmal or not, ceases, and during which food can be taken without inconvenience. The acidity of the urine is diminished; the reaction may even be alkaline; the chlorides are diminished or absent.

The peculiarities the pain exhibits form, on the whole, the most distinctive symptom of gastric ulceration. The paroxysms just spoken of might be mistaken for a purely nervous gastralgia. Indeed, when it is considered that both disorders are specially apt to occur in anæmic women, and in those whose menstrual functions are deranged, it becomes apparent how easily this mistake may be committed. The soreness at the epigastrium; the persistent symptoms of indigestion; the excess of hydrochloric acid in the gastric juice; the increase of pain after meals,—constitute, in a diagnostic point of view, the safeguard against error. To these might be added the vomiting of blood, were it not that vicarious hemorrhages are not at all unlikely to take place in young women who are troubled with amenorrhœa. This is, in truth, a matter having a close connection with the diagnosis of gastric ulceration. Persons who suffer from disturbance of the menstrual function are prone to be hysterical; and it may happen



that one of the most marked traits of the hysterical disorder is that it manifests itself by tenderness in the epigastric region, and by pain in the stomach.

We thus may have the most significant signs of gastric ulcer, occurring, as so many cases of amenorrhœa do, in chlorotic young women ; therefore happening in the class among whom ulceration of the stomach is most frequent. Nay, the very history may point to the probability of gastric ulcer.\* Yet generally, by close attention to all the phenomena of the case, we can arrive at a correct conclusion. The tenderness, as in all local hysterical affections, is great on the slightest touch ; and there is no severe pain posteriorly corresponding to the spot of soreness in the epigastric region. Pressure upon a spinous process may cause pain, but it is not the peculiar dorsal pain of gastric ulceration. Then, in the hysterical complaint there is often hyperæsthesia of the skin in various portions of the body, and the apparent gastric distress bears no relation to the taking of food or to the circumstance of its being of an irritating character or otherwise. The epigastric surface temperature is elevated in gastric ulcer, and may even exceed the temperature in the axilla.†

But to return to the vomiting of blood. When this is not traceable to a suppression of a natural discharge, and when it does not befall a person who suffers from disease of the heart, or liver, or spleen, or œsophagus, it acquires great significance. It is the only kind of vomit at all distinctive of a gastric ulcer ; for the substances ejected present otherwise appearances not different from what they do in chronic gastritis. The blood may be pure and red, but it is more frequently blackened by the gastric juice ; and large quantities are sometimes passed by stool. Now, hemorrhage does not take place in chronic inflammation of the mucous membrane of the stomach, except perhaps in drunkards. In those instances in which erosions exist on the surface, the vomited mucus may be a little streaked with blood ; yet anything like a profuse hemorrhage never happens. Hence its occurrence in a case with the symptoms of chronic gastritis, cancer being excluded, renders the presence of an ulcer probable. Yet there is a source

---

\* Case under my care, Philadelphia Hospital ; Med. and Surg. Rep., Feb. 1863.

† Hayem, *Revue des Sciences Médicales*, Oct. 15, 1888.

of fallacy, as I know by having met with such an instance, due to removal of the ovaries in an hysterical woman with marked gastric symptoms, in whose case subsequent hæmatemesis repeatedly occurred. It must also be borne in mind that we may have gastric ulceration without hæmatemesis.\*

The vomiting of the matters taken into the stomach may be immediate or not for some time after the food has been swallowed. Usually it happens speedily, and in some instances so speedily that there seems to be rather regurgitation than vomiting. But this is rare, and in the rarity is a safeguard against confounding gastric ulcer with the vomiting of cerebral disease, especially tumor; which I have known to happen in a young woman in whom, moreover, vomiting of blood had occurred. In the regurgitation, then, in the frequently absent nausea, in the clean tongue,—though coating may also be absent in ulcer,—in the want of oppression and weight at the epigastrium, and in the headache, altered vision, and other nervous phenomena, we have the distinguishing traits between *gastric* and *cerebral vomiting* on which to lay stress in the diagnosis between disease of the brain and gastric ulcer, or indeed any other serious stomach affection. The attacks of gastralgia that occur in the *gastric crises* of locomotor ataxia may be misleading. But the absence of knee-jerks and the eye phenomena explain their meaning. Perforating gastric ulcer may lead to abscess beneath the diaphragm, and be mistaken for pneumothorax. Indeed, this *pyopneumothorax subphrenicus*, as Leyden † called it, may show physical signs like those of pneumothorax. But they do not extend to the summit of the chest, and there is but little displacement of the heart. Moreover, the history points to long-existing gastric derangement. Pain in the front of the chest or in the abdomen, as the cases of Penrose and Dickinson ‡ prove, is an early symptom, and is soon followed by the physical signs of pneumothorax or of pneumonia.

In concluding this sketch of gastric ulceration, two questions arise which require solution: Does an ulcer always produce the peculiar train of symptoms mentioned? May not the same phenomena be met with in other disorders? The first question must

---

\* See a case reported by Walker, *Lancet*, No. 3691, p. 1301.

† *Zeitschr. f. Klin. Med.*, Bd i. p. 320.

‡ *Clinical Society's Transact.*, vol. xxvi., 1893.

be answered in the negative. Ulceration of the stomach may occasion nothing but the symptoms of chronic gastritis; and even these may not be marked. The second question is to be answered in the affirmative. There is a disorder with symptoms almost identical with those of gastric ulcer, the corrosive *ulcer of the duodenum*. Now, this affection, were it more frequent, would be a constant source of error. It may run an acute, or at least an apparently acute, or a chronic course. In either case it is scarcely distinguishable from gastric ulceration. Trier,\* from an analysis of twenty-six cases, mentions, among the most important grounds for a differential diagnosis, a sensitive tumor in the epigastrium, proceeding from adhesion with the pancreas, and jaundice or other hepatic phenomena. But these symptoms are far from constant; and, in accordance with his own showing, in acute cases, and in those chronic cases which run a latent course, the diagnosis is impossible. It may be added that the perforating ulcer of the duodenum is much more apt than ulcer of the stomach to remain latent and to lead rapidly to a fatal termination. The most certain signs of duodenal ulcer are the sudden and apparently causeless occurrence of intestinal hemorrhage, which may recur and be associated with gastric hemorrhage; violent attacks of pain referred to the epigastrium; and pain in the right hypochondriac region happening two or three hours after meals. Duodenal ulcer is thought by some to be almost invariably due to the action of a highly-acid gastric juice, and to furnish the best illustration of the so-called "peptic ulcer." It is most common between thirty and forty years of age, and, as Krauss proves, is ten times more common in men than in women.

---

\* Quoted in British and Foreign Medico-Chirurgical Review, Feb. 1864. See, also, the excellent monograph by Krauss, "Das perforirende Geschwür im Duodenum," 1865, and remarks on it in Niemeyer's work on Practical Medicine; Wadham and Barclay, London Lancet, Feb. and March, 1871; G. Ollive, Gaz. Méd. de Nantes, 1885-86, iv. 31; W. Osler, Canada Med. and Surg. Journ., Montreal, 1886-87, xv.; Buequoy, Arch. Gén. de Méd., Dec. 1887; W. H. Allchin, Transact. Pathol. Soc. Lond., 1887, xxxviii.; Schrötter, Aerztlicher Bericht des k. k. Allgemeinen Krankenhauses zu Wien (1886), 1888, 27; J. M. Emmert, Weekly Med. Rev., St. Louis, 1888, xviii.; W. W. Johnston, Amer. Journ. Med. Sci., 1888, N. S., xevi.; McPhedran, Canadian Practitioner, Toronto, Dec. 1890; Crozer Griffith, Transact. of College of Physicians of Phila., 1895.

There is yet another affection with symptoms like those of ulcer, an affection still more serious and destructive,—cancer.

**Gastric Cancer.**—Cancer is found more frequently in the stomach than in any other organ except the uterus. Of nine thousand one hundred and eighteen cases of cancer which occurred in Paris from 1837 to 1840, two thousand three hundred and three were in the stomach.\* The disease is generally primary. It is most often seated at the pylorus; next in frequency stands the cardiac orifice; most rarely does it involve the whole viscus. We find all the varieties of cancer affecting the stomach; but none is so common as scirrhus. Indeed, what is called cancer of the stomach means, in the large majority of cases, scirrhus; and, moreover, scirrhus at the pyloric extremity, deposited primarily in the textures which intervene between the mucous and the serous coat. It would be out of place to enter here into a minute description of the appearances of a gastric scirrhus. I shall only state that I have usually found it to present cell-growths less marked than those of scirrhus of any other part of the body. As found by an analysis of two thousand and thirty-eight cases of gastric cancer, three-fourths occur between forty and seventy years of age.†

The symptoms of cancer of the stomach are the same as those of chronic gastritis,—pain, tenderness in the epigastrium, disordered digestion, vomiting. In a more advanced state of the cancerous malady they may be those of gastric ulcer, hemorrhage being added to the list above given. There is only one symptom distinctive of cancer,—namely, the existence of a tumor; and this is so only when it is joined to digestive disorder and to increasing anorexia, debility, emaciation, and cachexia.

But let us see if there be anything in the pain and vomiting, or in the circumstances of the case, by which, even when a tumor cannot be discovered, the presence of a cancer may be suspected. Pain is a very constant symptom; quite as constant as in gastric ulcer. But the pain is, as a rule, more continued, much less influenced by the taking of food, and more radiating, being often referred to the right or the left hypochondrium. Its character is very varying. It may be dull, or gnawing, or it may be lanci-

---

\* Walshe on Cancer.

† Welch, Pepper's System of Practical Medicine.



nating. It may be slight, or it may amount to excruciating agony. It is often of the latter kind. But it is a mistake to suppose that a cancer of the stomach necessarily causes severe or lancinating pain. Again, it should be borne in mind that the part diseased may ulcerate, and then the pain is exactly like that of an ordinary gastric ulcer, and is affected in the same way by food. The most marked seat of the pain is sometimes under the shoulder-blade.

Vomiting is not an invariable result of cancer; yet it is a frequent one. The seat of the morbid growth determines, to a great extent, the occurrence of vomiting and the period at which it will happen. When the body of the stomach is attacked, and the orifices are not obstructed, it may not take place at all; or, if it take place, it is within a brief time after meals. When the disease has narrowed the cardiac extremity, vomiting supervenes almost immediately; the food has hardly been swallowed before it is brought up again. But when, as is much more common, the pylorus is constricted, the food is not thrown off until it attempts to pass through into the intestine; therefore not until a considerable time after meals.

With respect to the character of the substances ejected, this too depends on the seat of the cancer, and the time at which the vomiting arises. If it ensue several hours after meals, the cast-off matter consists of food partly digested, partly in a state of highly-acetous fermentation. An enormous quantity of acid material, the accumulation of several meals, is sometimes brought up during one act of emesis. The ejected matter may be intermingled with blood, and have a blackish or reddish-brown, "coffee-ground" appearance; or the mucus which is thrown up may be tinged with black flakes: in either case we find reduced hæmatin. Rarely is any considerable amount of unmixed blood vomited.

Free hydrochloric acid is, as discovered by von den Velden,\* often absent from the vomited contents of the stomach or from the "trial meal." But we must not forget that it is also absent in amyloid degeneration, in atrophy of the gastric tubules, in many fevers, and occasionally in chronic gastritis. Its presence,

---

\* Deutsches Arch. f. Klin. Med., Bd. xiii.

however, renders the existence of carcinoma improbable. The test is best made in the manner already described.

In many cases of carcinoma of the stomach lactic acid is to be found in the gastric contents after the administration of a special trial meal, free from lactic acid and lactates, and consisting of oatmeal gruel (a tablespoonful of oatmeal to a quart of water) with a little salt.\* This phenomenon is rare under other conditions, and, confirmed as it has been by the observations of Cohnheim † and of Stewart, ‡ promises to be a most valuable test, particularly of the early stages of gastric carcinoma. Microscopic examination may disclose the presence in the gastric contents of characteristic “concentrically arranged conglomerations of cells.” § .

A close study of the pain and vomiting may furnish evidence by which the existence of a gastric cancer may be suspected. There are a few other circumstances which would strengthen this suspicion: one of these is the acidity of the stomach, with the sour eructations; another, the extreme flatulency; another, the fetid breath, for, although fetor of the breath may result from putrefactive changes in the food in almost any form of gastric disorder, it is never perhaps so permanent as in cancer. A fourth is the obstinate constipation; a fifth, the progressive loss of flesh and the cachectic appearance of the patient, who is pale and tired-looking, or has a complexion slightly jaundiced, or whose face is of a color which seems to have arisen from a combination of the hue of chlorosis and that of jaundice. The supposed characteristic straw color of cancer is not often met with. The temperature is generally below the norm; but there are exceptional cases in which a moderate amount of irritative fever accompanies the gradual wasting,—gradual, because the duration of the malady averages more than a year. Edema of the ankles is a frequent symptom of the advancing disease. In some instances coma happens similar to diabetic coma, or tetany, as Kussmaul pointed out. There is an aseptic form, in which rapid enlargement of the liver, some fever, and erythematous eruptions occur.||

---

\* Boas, *Münchener Medicin. Wochenschr.*, 1893, No. 43, p. 805.

† *Deutsch. Med. Wochenschr.*, May, 1894.

‡ *Med. News*, Feb. 16, 1895.

§ Ewald, *Klinik der Verdauungskrankheiten*, 3. Aufl., II. Band, p. 342.

|| Hanot, *Archives Gén. de Méd.*, Sept. 1892.

Now, should all these symptoms be met with in a person who is steadily becoming feebler, whose age is above forty, in whose family cancer is hereditary; should cancerous tumors develop themselves in any other part of the body,—the suspicion entertained would be converted into almost a certainty. But it is not often that a perfectly typical case, presenting a combination of all the symptoms enumerated, is met with. And, I repeat, the most distinctive sign is a tumor: when this is not detected, considerable uncertainty hangs over any diagnosis of gastric cancer.

To contrast, then, cancer of the stomach with chronic gastritis and gastric ulcer:

CHRONIC GASTRITIS.	GASTRIC ULCER.	GASTRIC CANCER.
Pain at the epigastrium somewhat augmented by food; also soreness. Both constant, although comparatively slight.	Pain at the epigastrium much augmented by food; subsides when this is digested; paroxysms of pain, but not lancinating; a strictly-localized soreness to the touch in the epigastric region, sometimes a painful spot over the lower dorsal vertebræ. Intermissions in the pain of considerable length are frequent.	Pain frequently of a radiating kind, often paroxysmal, not unusually severe and lancinating, but not of necessity associated with soreness. Little or not at all affected by food. Pain rarely remits; never intermits for any considerable time.
Tongue usually heavily coated; may be clean.	Tongue dry, red, streaked in middle; or smooth and moist or slightly coated.	Tongue pale and thickly coated.
Acid eructations.	Eructations occur, are not acid.	Fetid eructations.
Symptoms of indigestion marked.	Symptoms of indigestion sometimes very slight.	Symptoms of indigestion marked. Anorexia; extremely sour stomach.
Sometimes vomiting; especially morning vomiting in alcoholic cases.	Vomiting usually immediately or soon after taking food; often an early symptom.	Vomiting a very frequent symptom; occurs sometimes on an empty stomach; usually preceded by other symptoms.
No hemorrhage, or but trifling hemorrhage; at most, bloodstreaks in vomited matter.	Abundant hemorrhage from the stomach common.	Hemorrhage not very abundant, but occasioning frequently coffee-ground-looking vomit.
Bowels constipated.	Bowels may or may not be constipated; usually are.	Bowels obstinately constipated.
No fever.	No fever.	Intercurrent attacks of slight fever may occur; but temperature often subnormal.
Not much emaciation; no cachectic appearance.	Frequently extreme pallor and debility.	Gradual and progressive loss of flesh, and debility; and at times with the cachexia hypertrophy of the peripheral lymphatic glands, especially above the clavicles.
Not confined to any age. More common in middle-aged or elderly people.	May occur in middle-aged persons; but is most frequently seen in young adults, especially in young women.	Most common in elderly people; rarely occurs in persons under forty years of age.

CHRONIC GASTRITIS.	GASTRIC ULCER.	GASTRIC CANCER.
Disease may be relieved or cured; is often of very long duration.	Duration uncertain; may get well, may run on rapidly to perforation; on the other hand, may last for years.	Average duration one year; may be shorter, but seldom longer; very rarely reaches two years.
No tumor.	No tumor.	Generally a tumor.
Contents of stomach contain almost always free hydrochloric acid.	Hydrochloric acid in excess in contents of stomach.	As a rule, no hydrochloric acid in contents of stomach; often lactic acid present.
No dropsy.	No dropsy.	Œdema of ankles often met with.

The differences laid down in the table are derived from an analysis of well-marked cases. In the early stages of the cancerous malady, a differential diagnosis is impossible. Subsequently, as already stated, the detection of a tumor plays an important part in any deduction. But this remark does not apply to cases of cancer of the cardiac orifice, which are rare, and in which a tumor, from its deep situation, almost always eludes discovery. Such cases are, however, discriminated by their presenting the same signs as a stricture of the œsophagus low down; indeed, they are very constantly combined with a narrowing of the tube, produced by the cancer spreading to it. Cancer at other parts of the organ occasions a perceptible tumor in about three-fourths of all the instances; its situation is, of course, not always the same. Where no tumor can be discerned, and particularly if, as may happen, portions of the stomach remain healthy and the digestive disturbances are slight, the existence of cancer may not reveal itself by any symptoms, and the case run a latent course.\*

A cancer of the anterior wall produces, as a rule, fulness, resistance, and percussion dulness in the epigastric region. A cancer involving the greater curvature gives rise to a swelling near the umbilicus, or to one extending toward either hypochondrium. The tumor formed by cancer of the pylorus is commonly felt plainly a little to the right of the median line, and one to two inches below the cartilages of the ribs. In women its position is apt to be even lower than this; and, indeed, in both sexes the situation of the indurated pylorus is very variable. It may be pushed down to near the umbilicus; nay, it has been discerned near the anterior superior spinous process of the ilium.† It is

\* See report of case under my care at the Pennsylvania Hospital, published in *Amer. Journ. of Med. Sci.*, vol. lii., 1866.

† See Lebert's cases in *Traité pratique des Maladies cancéreuses*.



rarely found in the left hypochondrium, but not unfrequently in the right. Then it may form adhesions to the liver, which viscus at times so completely covers the tumor as to render this impossible of detection.

The reason why the swelling, in not a few instances, shows itself much lower than the normal seat of the pylorus, is obvious. During meal after meal the organ seeks to overcome the resistance offered by the narrowed pyloric orifice, and does so with great and increasing difficulty. The constantly-repeated and long-continued struggle leads to hypertrophy of the muscular coat and to distention of the hollow viscus.

The tumor may or may not be movable,—generally is not; its surface may be either smooth or nodulated. It may be large and distinct, or small and requiring a careful examination to distinguish it from the surrounding and more yielding textures. Percussing over it elicits a dull sound, usually mixed with a tympanitic note. The tumor is much more perceptible on some days than it is on others. Its existence, as has been already insisted on, furnishes the most conclusive evidence in favor of a cancer.

But is a swelling in the region of the stomach strictly pathognomonic of gastric cancer? No; not even when the swelling has been ascertained to belong to that viscus. At times the cicatrix marking a previous ulcer, or even the indurated and thickened margins of an existing ulcer, may be palpable through the abdominal walls and raise the question of a new growth. A mere fibroid thickening of the pylorus will occasion a tumor, and, moreover, produce symptoms which resemble so closely those of malignant disease at the orifice that I much doubt the possibility of distinguishing during life, with any certainty, between the two affections. Let us take this case, which I saw with Dr. Moss,\* as an example.

A woman, aged forty, complained much of pain at the pit of the stomach, and of a heavy sensation throughout the abdomen. For some months she had been suffering from indigestion, and had been steadily losing flesh. Her countenance had a tired look, and she was very despondent. She had a slight cough; and on per-

---

\* Published in full in *Proceed. of Pathol. Soc. of Phila.*, vol. i.

cussing the lungs, impaired resonance was detected at the apices. The bowels were obstinately constipated, the tongue was smooth and red, the pulse feeble. She vomited shortly after meals, yet never anything but the ingesta. There was no pain on pressure over the pylorus; but a greater resistance to the finger than usual was detected. The further progress of the complaint was marked by incessant vomiting, only, however, after meals. Once, and once only, did it cease for several days; and then without apparent cause. As the case drew toward its fatal termination, the patient was much troubled with acid eructations, and had occasionally slight febrile attacks. The distress in the epigastrium increased in severity. About three weeks before her death she was seized with lancinating pains under both patellæ, which were neither relieved nor aggravated by pressure or motion. They were accompanied by pricking sensations and numbness in the legs, and an inability to walk. The pains gradually ceased, but the numbness and loss of motion increased from day to day. She died, utterly exhausted by the abdominal pains and the incessant vomiting, about three months after she began to reject her food. On post-mortem examination, tubercular deposits were found at the apices of the lungs. The abdominal viscera were healthy, except the stomach; and this, too, was healthy, save at its pyloric orifice, which was so narrowed that the tip of the little finger could hardly be forced into it. The mucous lining lay in folds, but on dissection was found to be perfectly normal. At the pylorus, but only there, the submucous and the muscular coat were uniformly thickened. Examined microscopically, they contained nothing but areolar tissue, spindle-shaped fibre-cells, and very distinct organic muscular fibres.

Now, here is a case which was not cancer; yet it had the symptoms of cancer. It is true that the absence of blood and of glairy mucus in the matter vomited, and the indistinctness of the swelling, in spite of the great emaciation, were against the supposition of cancer of the pylorus. Still, no inference based on these data alone could be strictly trusted, since every cancer is not associated with the vomit of coffee-ground material or of glairy mucus, or with a palpable tumor. The disease was combined with tubercular deposits in the lung. Nor is this the only example of the combination which has come under my notice. And when a

tubercular state of the lung has been fairly made out, and there exist at the same time signs of pyloric obstruction, I should make a diagnosis that this is not of a cancerous nature, but consists simply of an increased development of the submucous coat, with probably subsequent hypertrophy of the muscular tunic.

The *fibroid thickening* may extend throughout the whole stomach. Such cases differ from cancer by their long duration; the absence of hemorrhage, of vomiting, and of severe pain; and the more uniform gastric swelling. The affection is sometimes observed in spirit-drinkers. Its discrimination from cancer is never a certainty, but merely a matter of conjecture. I am not aware of any extended chemical examination of the contents of the stomach having been made. In a case reported by Cornell,\* which was complicated with tuberculous peritonitis, loss of digestive power was indicated by unbroken starch grains in the vomit. The absorptive activity of the stomach tested by iodine was normal. Boas† states that in these non-malignant cases with pyloric stenosis, though there are fermentative processes, lactic acid is absent.

There are other diseases than those of the stomach which may occasion a tumor in its region and are thus liable to be mistaken for gastric cancer. Prominent among these are enlargement of the liver projecting into the epigastrium, tumors of the omentum, and diseases of the pancreas and of the kidney. Of course, the stomach symptoms proper are not so marked in these affections, and in some they may be wholly wanting; examination of the gastric contents and of the urine, and due regard to the history of the case, will show us the truth about many; and, after all, the best way of preventing ourselves from falling into error is to seek in any case of supposed gastric cancer for these other diseases, and to see if their chief symptoms are present.

Resting with this general statement, I shall not take up the differential diagnosis of all the many affections mentioned; especially as some are referred to when treating of partial abdominal enlargements and of cancer of the liver. But there are two which may be here specially looked at: one is omental cancer, the other kidney affection attended with marked swelling, such as occurs

---

\* Montreal Med. Journ., Aug. 1892.

† Münch. Med. Wochenschr., Oct. 1893.

in hydronephrosis, pyonephrosis, abscess, hydatids, and morbid growths.

In *omental cancer* there is far less dyspepsia, hemorrhage and coffee-ground vomit are absent, the tumor appears to occupy chiefly the site of the greater curvature, the swelling is, or soon becomes, more generally diffuse, and hydrochloric acid and the digestive ferments are present in the gastric contents.

In the *kidney affections* referred to, the history is of great importance, and we include in this history the passage of renal calculi as bearing on some forms of kidney enlargement, especially abscess from impaction of stones; the limits of the mass, though this may project into the epigastrium, will scarcely be those of a gastric cancer. But the most certain safeguard against error is careful and repeated examination of the urine and of the gastric contents.

And as regards the urine, the observations of Rommelaere\* seem to show that its analysis may be of value even in the diagnosis of the different forms of gastric disease. Thus, a cancerous ulceration of the stomach is attended with decrease in the amount of urea, and the acidity and the chlorides are diminished. In simple gastric ulcer these are in normal quantity or in excess; so is the urea. In spreading gastric ulcer the chlorides in the urine are decreased, but there is normal or increased amount of urea and urates.

**Dilatation of the Stomach.**—This happens frequently in connection with obstruction of the pylorus, whether cancerous or fibroid, but it is also met with independently of this structural lesion. The latter form occurs from weakening of the muscular coats produced by malnutrition or impaired innervation, and has been noticed as an attendant upon anæmia or hysteria, or following fevers, or obstruction of the upper part of the bowel, or compression of the pylorus by an enormous gall-stone,† or, as Bamberger mentions, dislocation of the stomach by omental hernias. Elinger has proved that it may be associated with amyloid degeneration of the vessels or of the muscular coat of the stomach. The chief signs of a dilated stomach in either form are the rejection of food,

---

\* Journal de Médecine de Bruxelles, 1883; quoted in the Lancet, Sept. 1 and Oct. 27, 1883.

† Minkowski, quoted by Ewald.



mostly in large quantities and retained for days; fermented and vomited matter containing often *torulæ* and *sarcinæ*; extension of the tympanitic note of the gastric region, detected by percussion, to much below the umbilicus; a splashing sound when the patient moves, particularly after drinking, and gurgling on sudden pressure; the low line of dulness occasioned by fluids in the distended organ, and the change of the dulness with the position of the patient; and slowly-progressing emaciation. The character of the gastric secretion and that of the contents of the stomach after a trial meal vary with the nature of the causative condition. As a rule, there are increased acidity and diminished absorptive and motor activity. The general nutrition suffers as assimilation is more and more interfered with. In doubtful cases the organ may be examined and its limits traced by distending it with ordinary air, or with carbonic acid, generated by swallowing first bicarbonate of sodium and then tartaric acid. Displacement of the right kidney has been observed in a number of cases.

The sounds of the heart heard over the dilated stomach often have a metallic ring, but, irrespective of this, peculiar gurgling sounds, systolic in rhythm and evoked by the action of the heart, have been met with by Franck and other observers. Dilatation of the stomach may occasion nervous symptoms; even tetany has been noticed.\* The dilatation occasionally happens in an acute manner, and occurs in children † as well as in adults. As a rule, the muscular coat is not hypertrophied, but in the cases in which an obstruction at the pylorus exists, this is frequent at first, ultimately giving place to atrophy.

The stomach may be unduly large without giving rise to any symptoms. This condition of *megalogastria* is to be distinguished from *gastrectasia* by the absence of the symptoms of the latter, as well as of derangement of secretion, absorption, and propulsion.

To tell the atonic cases from those due to narrowing at the pylorus is generally not difficult: we can detect a hard swelling, or find the resistance with a stomach sound. In cancerous obstruction the gastric juice, as a rule, contains no hydrochloric acid, but we obtain lactic acid. In other forms of stomach dila-

---

\* *Bulletins et Mémoires des Hôpitaux de Paris*, t. xx., 1884.

† *Arch. Gén. de Méd.*, August, 1884.

tation, particularly in the atonic form,\* we find hydrochloric acid, as well as the acids of decomposition and fermentation, acetic acid, and butyric acid.

Enlargement of the stomach is to be distinguished from displacement of the organ,—*gastroptosis* or *Glénard's disease*. The latter is seen in persons with pendulous abdomen. The condition is due to relaxation of the ligamentous attachments of the stomach, and is often associated with a similar displacement of other abdominal viscera,—*splanchnoptosis* or *enteroptosis*. There are present, in addition to symptoms of digestive derangement and the obvious evidences of the dislocation of the viscera, manifestations of functional nervous disturbance. Among the first are impaired or perverted appetite, epigastric fulness and distention, eructation, acid taste and dryness of the mouth, burning or tearing pains at the pit of the stomach after eating, and constipation alternating with seeming diarrhœa. The nervous symptoms include a feeling of weakness, general irritability, mental depression, headache or a sense of fulness in the head, vertigo, heaviness of the lower extremities, coldness of hands and feet, palpitation of the heart, heavy sleep, and frequently sacral pains. Emaciation takes place, with impoverishment of the blood, pallor of the surface, changes in the skin, and falling out of the hair.

A certain number of cases are associated with dilatation of the stomach. This is also seen, as Reichmann has proved, in some persons who are much emaciated and in whom supersecretion of the gastric juice exists.

Dilatation of the stomach may be confounded with *dilatation of the large intestine*. But the gastric symptoms of the former malady are of great significance. Moreover, we may make use of the salol test in the discrimination. Salol is not acted upon by the acid gastric juice, but is changed into salicylic acid by the alkaline intestinal secretion. The salicylic acid manifests itself in the urine of healthy persons in from half an hour to an hour, as shown by the addition of a drop of tincture of chloride of iron to the urine giving it a violet color. In dilatation of the stomach salicylic acid does not appear for two or three hours after salol has been taken.

---

\* Germain Sée, Bull. de l'Acad. de Méd., May, 1888.

## SECTION II.

## DISEASES OF THE INTESTINES AND OF THE PERITONEUM.

In considering the diseases of the intestines, we meet with symptoms the import of which we have examined in connection with affections of the stomach. We encounter nausea, vomiting, and impaired digestion. These disturbances are sympathetic or dependent upon coexisting gastric disorder; they do not serve, therefore, as trustworthy guides in intestinal maladies. The signs upon which we rely more implicitly are pain and the faecal discharges. As regards the former, we draw the truest inferences from its kind rather than from its mere occurrence. The study of the faecal discharges tells us in a more direct manner what is going on.

**Alvine Discharges.**—To examine briefly their appearances:

*Watery* stools are observed whenever a large quantity of the serum of the blood finds its way through the intestinal coats. They are met with after the administration of saline purgatives, in serous diarrhoea, and in cholera. Their hue varies: they may be almost colorless, or tinged with yellow. Sometimes, although very thin and watery, they are decidedly yellow; again they are rendered turbid by the dissemination of whitish flocculi of cast-off epithelium, or by mucus. Whether they be yellow or colorless depends on the existence or non-existence in them of faecal matter and of bile. In a prognostic point of view, the most colorless evacuations are the most dangerous.

The presence of an excessive quantity of *mucus* renders the discharges less consistent than natural; yet, unless they contain more or less serum, they are not of necessity liquid. The appearance they present is similar to that of the white of an egg; or the whitish masses of mucus surround the lumps of faeces, or are intermingled with the fluid alvine discharges.

*Pus* in large amount and unmixed with faeces is discharged only when an abscess has ruptured into some part of the intestine. Stools composed of faeces and pus are encountered in chronic inflammation and in ulceration of the bowels; and whitish, creamy streaks indicate the presence of the foreign substance. Yet the

pus may be so intimately blended with the fæces, or with masses of mucus, as to require the microscope for its detection.

Stools consisting entirely of *bile* are rarely met with. More generally there are other elements joined to the voided secretion of the liver. An excess of bile in the alvine discharges gives rise to evacuations of a yellowish-brown or yellow hue, which darken on exposure to the air. When the alimentary tube is highly acid, the resulting color is green. Both these kinds of stools are commonly called "bilious;" but the latter is less absolutely so than the former. A deficiency of bile manifests itself by clayey, sometimes even by almost white, stools. The normal color of fæces is due to urobilin. It is the changed bilirubin from the bile. Bile-pigment is not found in healthy stools.

*Black* stools result from eating certain articles of food, such as blackberries; from the action of medicines, as iron, bismuth, manganese; from a vitiated condition of the bile and intestinal secretions; or from the effusion of blood into the alimentary canal. At all events, when the hemorrhage proceeds from the stomach or the upper part of the canal, the stools have a black, tarry appearance; when from the lower section of the tube, pure blood is passed, or, if it be small in quantity, a blood-streaked mucus. Should any doubt exist as to whether the dark discharges be dependent upon the presence of blood, let them be diluted with water; they will assume a reddish tinge if this be the cause of the abnormal color.

The *odor* of the evacuations is extremely offensive in fevers of a low type, and when the intestinal secretions are vitiated. So, too, at times in smallpox and in cholera. Acidity of the intestinal canal, as in the intestinal catarrh of children and of adults,\* or in rheumatism or gout, imparts to the stools a sour smell and an acid reaction. The reaction in health is mostly alkaline.

In cases of constipation it may be important to notice the *shape* of the passages, because this may show whether an impediment in the gut has flattened or otherwise altered them. In fevers, as well as in affections of the intestinal mucous membrane, whether inflammatory or not, we often derive information from studying the form of the voided matter. Figured stools succeeding to fluid

---

\* Jaksch, *op. cit.*



passages are always of favorable omen. We also note whether the stools contain masses of undigested matter, and its kind.

*Chemical and microscopical* examinations of the fæces are not often made; yet chemistry and the microscope may be frequently of great service. They enable us, for instance, to recognize with certainty that the yellowish lumps contained in the evacuation, or the greasy film which collects upon its surface, consist of fat. The microscope, too, detects masses of muscular fibre, of elastic tissue, of starch-corpuscles, of fat, coagulated albumen, crystals of cholesterin, red corpuscles, leucocytes, and various fungoid growths, micro-organisms, and parasites. Among the animal parasites, besides various infusoria and worms,—the main variety of which will be discussed farther on with the parasites,—we find the *amœba coli*, now known to be the chief cause of tropical dysentery. It is one of the rhizopods, varying in size from 0.012 to 0.035 millimetre, and when active has a characteristic movement. This will be best seen if the stage of the microscope be kept warm.

The microscope exhibits, in the faecal discharges of all diseases in which the stools readily decompose, masses of crystals of the triple phosphates; in acrid stools, yeast fungi; in typhoid fever, shreds of slough from the enteric ulcers, and bacilli; in tubercular ulceration of the bowel, tubercle-bacilli; and under many varying conditions both in the fæces and in different organs, as well as in peritoneal exudates and in appendicitis, the *bacillus coli communis*. This is, as a rule, a sluggishly-moving bacillus which grows readily on gelatin plates, the surface colonies being large and spherical and of a dull white. They are stained by aniline dyes, but are decolorized when treated by Gram's method. The main normal ingredient of faecal matter is mucin.\* Peptone occurs only in disease.† One drawback to the use of chemical research for clinical purposes is the uncertain composition of the fæces, owing to the number of elements derived from the food.

The study of the alvine discharges is of service not merely in intestinal complaints, but equally in the many maladies in which

---

\* Hoppe-Seyler, Handbuch.

† Jaksch, *op. cit* See also on this and other points Nothnagel's researches, Beiträge zur Physiologie und Pathologie des Darms, Berlin, 1884.

the alimentary tube sympathizes or becomes involved. But to return to the uncomplicated intestinal diseases, grouping them as they may be recognized by pain and peculiarity in the faecal discharges, and describing with them, for the sake of convenience, the affections of the peritoneum.

Diseases attended with Paroxysms of Pain referred chiefly to the Middle or Lower Part of the Abdomen, and not associated with marked Tenderness or with Fever.

The type of these is colic.

**Colic.**—This is an intestinal pain, paroxysmal in its character, and usually combined with constipation, but unattended with febrile symptoms. The pain is of a severe griping or twisting kind, and is commonly referred to the neighborhood of the umbilicus. It is generally relieved by pressure. Sometimes there is soreness with the pain, and, indeed, a slight soreness not unfrequently remains after the paroxysm has passed off. While the pain lasts, the countenance wears an anxious, frightened expression; the skin is cold; the pulse is depressed. Occasionally there is vomiting, and in severe cases the abdominal walls are tense or raised in hard knots by the spasmodic contraction of the muscles. An attack may last only a few minutes, or, with trifling remissions, for several hours.

Some persons are very liable to attacks of colic. Those who suffer from indigestion, or are enfeebled by exhausting maladies, are predisposed to them; so also are hysterical, gouty, and rheumatic individuals. As to the exciting causes, they are various; and somewhat according to its different causes, colic presents different forms. Let us indicate the more prominent.

*Colic, simple and unconnected with a disease of the bowel.*—Now, in these cases, which are generally called spasmodic colic, the paroxysmal pain may have a diverse origin. It may be the result of direct excitation of the peripheral intestinal nerves by the presence of irritating substances in the alimentary canal, such as indigestible food, cold or acid drinks, hardened faeces, gases, morbid secretions, ptomaines, worms, medicines, or poisons. It may proceed from an irritation of the central nervous system reflected to the

intestinal nerves. It may be sympathetic, and produced by a morbid state of the adjacent abdominal viscera.

1. Colic owing to food difficult of digestion is very common, especially at the time of year when fruit is beginning to ripen. It may be caused by food taken in quantities greater than the digestive organs can assimilate. Hence it is frequent in children at the breast who are overnourished, and in persons in delicate health with enfeebled digestive powers. The form of colic under discussion is often attended with vomiting and diarrhœa; it may be of only a few hours' duration, or it may last for several days.

Colic arising from distention of the intestines with flatus, or "flatulent colic," is the result of the decomposition of food in the alimentary canal; sometimes, however, the gases are extricated from morbid secretions, or are exhaled directly from the blood-vessels. The abdomen is very tympanitic and greatly distended, and the flatus is from time to time discharged, with evident relief. Hysterical persons are very subject to this form of colic.

Colic from accumulation of hardened feces is preceded by obstinate constipation, and is usually a tedious disorder. The accessions of pain are easily enough remedied by emptying the bowels; but they constantly recur.

Colic from the presence of morbid secretions in the intestinal canal is not so often encountered as that from indigestible food or retained fecal masses. Yet it is occasionally met with in cases of diarrhœa attended with a disordered state of the intestinal functions; even in the so-termed bilious colic the intestinal pain is not purely sympathetic, but is owing to the irritating character of the bile discharged into the intestine.

This "bilious colic" is often preceded by nausea, loss of appetite, and a coated tongue. The paroxysms of pain frequently go hand in hand with vomiting,—first of the contents of the stomach, then of bile. They are in general accompanied or soon followed by a yellowish tinge of the conjunctiva, by tenderness in the region of the liver, and by a desire to go to stool. The bowels are, however, apt to be obstinately constipated. Bilious colic is common in malarious districts; it occurs especially during the summer and autumnal months, and frequently follows exposure. It sometimes begins with a chill, and, unlike the other forms of

colic, it has as companions febrile excitement, and a full, frequent pulse. Malarial colic may occur in an epidemic form.\*

2. In the second class of cases to which allusion has been made, colic is dependent upon some abnormal condition affecting primarily the great centres of innervation. The colic arising from fright, from anger; that happening in nervous females and hypochondriac males; perhaps that proceeding from sudden exposure to cold; the form which is sometimes seen coexisting with neuralgic pains in other parts of the body,—in short, all those cases which are spoken of as nervous colic, might here be mentioned. The attack is sudden, and not commonly of long duration; but it is very apt to be repeated.

The “metallic colics” are further instances of colic produced through agents which act primarily on the nervous system. This is at any rate true of lead colic. *Copper colic* exhibits paroxysms of severe pain like those caused by the poisonous influence of lead; but it is attended with nausea, vomiting, diarrhoea, tenesmus, an abdomen distended and tender to the touch; in other words, it is rather an inflammation of the intestine with colicky pain, than uncomplicated colic. *Lead colic*, on the other hand, is a pure colic. The distinguishing marks of lead colic are the bluish-gray line along the gums; the contracted abdomen; the obstinate constipation; the great relief usually afforded to the pain by pressure; the duration of the pain; its marked and agonizing exacerbations; and the history of the case. The signs of the lead poisoning also manifest themselves in other parts of the body.

3. Affections of various organs may give rise to colic, by sympathy, and generally through the intervention of the nervous system, to which the irritation is first transferred, and from which it is then reflected. Thus, colic is a not uncommon attendant on morbid states of the kidneys, liver, bladder, testicles, uterus, or ovaries, and on disordered menstruation. Yet we must not forget that the pain, although spoken of as colic, is often not strictly intestinal, but is merely a pain radiating from the affected organs themselves. Again, how far it is due to neuritis is a matter to be taken into account.

---

\* American Journal of the Medical Sciences, April, 1872.



*Colic arising in consequence of some abnormal state of the bowel.*  
 —In the preceding illustrations of colic the disorder was viewed as occurring in a healthy bowel. But colic may have only the significance of a symptom, and be combined with an altered structure or a changed position of the intestine. We meet, indeed, with colicky pains, undistinguishable from those of pure colic, linked to an organic disease of the bowel, and under circumstances some of which forbid the idea of a spasm. They are encountered in dysentery; enteritis; hernia; ulceration; intussusception; strangulation; twisting; strictures; distention,—in fact, in the most various morbid states of the intestine. And colic as a symptom can be discriminated, as far as the pain is concerned, from colic as an idiopathic disorder, only by a careful study of the history and the concomitant phenomena of the case. In several of the maladies cited, however, the more transitory nature of the pain,—or gripings, as they are termed,—in others, the presence of fever and of tenderness, serve as guides in diagnosis. Fever and soreness to the touch are also met with in that form of inflammation of the bowel which happens after exposure or after the retrocession of rheumatism from some external part, and which is commonly known as rheumatic or inflammatory colic.

Having thus indicated the various forms of colic, and having mentioned the relation they bear to structural diseases of the intestines and to affections of adjacent viscera, I shall only here again insist on the necessity of tracing out in every case, as far as possible, the cause of the painful malady, so as to know if any serious mischief lie at the bottom of it; and shall add but a few words with reference to the disorders with which uncomplicated colic, or that which is held to be purely spasmodic, may be confounded. They are:

GASTRALGIA;

PERFORATION OF THE INTESTINE;

STRANGULATED HERNIA;

PASSAGE OF GALL-STONES;

NEPHRALGIA;

SPASM OF THE BLADDER;

UTERINE COLIC;

NEURALGIA OF THE DORSAL AND LUMBAR NERVES;

## ABDOMINAL ANEURISM AND TUMORS; DISEASES OF THE SPINE;

## ENTERITIS AND PERITONITIS.

*Gastralgia.*—In gastralgia the pain is seated in the epigastric region; whereas in colic, or enteralgia, as it is called by some, the pain is either in the neighborhood of the umbilicus, or rapidly shifts its position from that point to different parts of the abdomen, and is often connected with a spasmodic contraction of the abdominal muscles. Again, the history in cases of gastralgia; the fact that the attacks happen most frequently after meals; their association with signs of a disordered stomach,—indicate the organ in which the pain arises.

And much the same general signs, in addition to the marked constipation and the visible movements, enable us to distinguish those instances of peristaltic disturbance of the stomach to which Kussmaul\* has called attention, and in which the drawing pain is apt to be referred to the intestine; indeed, the peristaltic disorder may spread to it.

*Perforation of the Intestine.*—When paroxysms of pain have their origin in perforation of the intestine, the extreme prostration and collapse show that they are not produced by a harmless disorder like colic. Further, the abdominal distress is in such cases preceded by symptoms of a diseased state of the stomach or the intestines; and if the patient live sufficiently long after the accident, the pain is followed by great distention of the abdomen and extreme tenderness,—in fact, by the signs of peritonitis. However, the differential diagnosis is occasionally very difficult. Especially is it so in typhoid fever; for in this affection colic is readily induced, or perforation of the intestine may be brought on by very slight exciting causes; and, moreover, peritonitis may occur without perforation. A valuable sign of perforation and of air in the peritoneum is the obliteration of the dulness on percussion over the hepatic region, pointed out by Alonzo Clark.

*Strangulated Hernia.*—All mechanical obstructions of the intestine will lead to paroxysms of intestinal pain. They are met with in cases of intussusception and of ileus; they are equally frequent in cases of strangulated hernia. In all, the obstinate

---

\* Sammlung Klinischer Vorträge, No. 181, June, 1880.

constipation must arouse suspicion regarding the true nature of the complaint. To detect a hernia a local examination is required; and, therefore, a careful search at the usual seats of this affection ought to be made in every instance of severe or protracted colic. Lives have been lost in consequence of the neglect of this simple precaution against disastrous error.

*Passage of Gall-stones.*—The passage of a gall-stone is generally attended with paroxysms of intense pain which are readily mistaken for colic. There is, as a rule, the same absence of fever and of tenderness; yet fever of short duration does happen. Pressure is often resorted to in order to mitigate the suffering, and thus the resemblance to colic is heightened. The points of distinction from colic are, the position of the pain in the epigastric region; its sudden beginning and sudden termination; the severe nausea and vomiting attending the attack; the jaundice; and the voiding of gall-stones with the stools. The latter sign, though a positive one, assists less in the discrimination of the disorder than would appear at first sight; partly because it does not serve as a means of indicating the nature of the affection until its close, partly because the stone often escapes detection in the *faeces*.\* The other circumstances have, therefore, a more available diagnostic value. Yet even they do not enable us to distinguish positively between the transit of a biliary concretion from the gall-bladder to the intestine, and the bilious colic that is joined to derangement of the function of the liver. The repetition of the attack is always a strong reason for suspecting it to be owing to a discharge of calculi from the gall-bladder; and so are severe retching and vomiting, the sudden supervention of jaundice, and the localized epigastric pain. But these phenomena, too, are produced by *hepatic neuralgia*, which in rare cases is believed to happen without there being gall-stones or inspissated bile. There is nothing by which we can discriminate this malady—the very existence of which is, indeed, denied—except its recurrence after certain intervals, the alternations with other affections of the nervous system, and the slightest touching of the part inducing

---

\* The best way to find the stone is to pass the evacuation through a sieve: this is more certain than covering the discharge with water. The stone may not come from the bowels for some days after the attack of colic.

at times the acute pains.\* There is said to be an increase of temperature over the gall-bladder during an attack of gall-stones.†

Sometimes gall-stones are closely simulated by impacted fæces, which occasion colicky pains, and even jaundice, by pressure. The pain is at once removed by morphia given hypodermically, and a dose of oil brings away the hardened fæces. The attacks may recur, and are always relieved in the same manner. The swelling in the right side may be sometimes readily detected.

Among the rarer symptoms attending or following the passage of gall-stones, temporary dilatation of the heart and tricuspid regurgitation have been noticed,‡ just as temporary mitral insufficiency has been observed in jaundice.

Where the gall-stones are large and have become impacted in their course toward the intestine, they give rise to inflammation which may lead to ulceration and to the discharge of the concretion—generally then very large—into the intestine or stomach. Subsequently an obliteration of the duct may happen; or the inflammation and ulceration of the duct may result in perforation into the peritoneum. In some cases the gall-stones are voided through the abdominal walls, in consequence of their having caused inflammation of the gall-bladder and subsequent adhesions to the abdominal parietes. The fistulous passages discharge pus and bile, and occasionally fresh concretions: they may last for years; but in time they generally heal. As regards the other forms of fistulous communications alluded to, they very rarely present symptoms so peculiar as to warrant anything like a certain diagnosis.§

*Nephralgia.*—Paroxysms of pain with intervals of comparative ease and unassociated with fever occur in nephralgia, or pain of the kidney, and are, therefore, often mistaken for colic. Now, nephralgia is generally, although not invariably, caused by the passage of a calculus through the ureter. Its symptoms, besides the pain, are numbness of the thigh, nausea and vomiting, a constant desire to make water, and aching and drawing up of the

\* See the cases of Budd. on Diseases of the Liver; of Andral, *Clinique Médicale*, tome ii.; and of Frerichs, *Diseases of the Liver*.

† Jules Cyr, *Traité sur l'Affectio calculieuse*, Paris, 1884.

‡ Potain, quoted by Sée, *Maladies du Cœur*, Paris, 1883.

§ See a collection of cases by Murchison, *Edinb. Med. Journ.*, July, 1857.



testicle. The patient, as in colic, is restless, and seeks relief by frequently changing his position. The pain comes on suddenly, and is excruciating. It is felt in the loins, usually on one side, and shoots along the track of the ureter to the corresponding hip and thigh. It sometimes extends to the pelvis or toward the umbilicus, and is often attended with tenderness in the course of the ureter. Occasionally it is almost exclusively felt at the hip. When the stone reaches the bladder, the pain ceases as abruptly as it began; though sometimes there is still discomfort produced by the stone interfering with the act of micturition. During the attack the urine is passed in small quantities at a time. It is high-colored; sometimes it contains a little blood. If it be collected, and, after all pain has disappeared, be carefully examined, a small, hard body or a sandy deposit is generally detected, and reveals the cause of the past anguish. It is from the presence of the sandy deposit that the complaint has received popularly the name of a fit of "the gravel."

From the description given, it will be seen that in several respects the disorder is like intestinal colic. The seat of the pain is a point of distinction; yet in neither complaint is the seat entirely characteristic. It is not always strictly umbilical in colic; it is not always exactly in the region of the ureter or kidney in nephralgia. Of more importance is the state of the urinary functions, which are comparatively undisturbed in colic. Again, the numbness of the thigh and the retraction of the testicle are valuable diagnostic marks; they would be absolutely decisive, were they constantly present in nephralgia.

*Spasm of the Bladder.*—The bladder is sometimes the site of paroxysms of violent pain, supposed to attend upon a spasm of the viscus. There is an intense desire to urinate, which the passing of water does not allay. The pain is not steady; it is accompanied by a sense of constriction at or near the pelvis, and sometimes by tenesmus, and may extend to the kidneys, to the thighs, and to the sacrum; or the irritation may be communicated to the penis, and cause erections. If the sphincters be involved, the urine cannot be voided. The bladder distends; there is intense anxiety, with restlessness; the pulse is feeble; the skin is cold, and covered with clammy perspiration.

A spasm of the bladder may be caused by the presence of a stone

or of irritating urine. It is also encountered in gout and hysteria, and as the result of stimulating diuretics. Violent fright, too, may occasion it. It sometimes proceeds from a disorder of adjacent structures, such as of the rectum, or of the uterus. Now and then, as Sir Benjamin Brodie pointed out, it is associated with inflammation or suppuration of the kidney, and the vesical pain is so intense that it withdraws attention from the organ most affected. To distinguish it from colic is not difficult; the position of the pain and the disturbed condition of the urinary functions serve as guides. It resembles more closely nephralgia; as in nephralgia, too, after the fit is relieved, the important indication is to prevent its repetition by endeavoring to remove its source.

*Uterine Colic.*—The painful sensations experienced by some women at their menstrual periods may come on in paroxysms similar to those of colic. In truth, the pain is often spoken of as uterine colic, and at times continues for many days, persisting during the whole menstrual period, or even longer. In some of these cases the complaint is localized in the uterus; in others, more especially in the ovaries, which are then tender to the touch. Similar attacks of pain, also accompanied by congestion or even by inflammation of the ovaries, are occasionally met with as the result of falls or of blows on the hypogastric region.

Now, with reference to the disorder first alluded to, or ordinary dysmenorrhœa, it may be generally easily discriminated from colic by its occurrence with the setting in of the menstrual flow; by the pain remitting rather than intermitting; by the seat of the pain in the pelvis, or the lower part of the abdomen; by its not uncommon association with sickness, nausea, and vomiting; and by the fact that all the signs of disordered menstruation have happened over and over again at the menstrual periods.

Where the ovaries are very much congested or inflamed, whether or not the affection exist in connection with dysmenorrhœa, or occur in consequence of other causes, among which gonorrhœa may be one, the pain, tenderness, and swelling in the hypogastric region; the not unusual numbness and flexed position of one or both thighs; the febrile irritation, and the hysterical symptoms; the retention of the urine; the violence of the paroxysms of pain, and the duration of the malady,—form a group of phenomena very dissimilar to those of ordinary cases of colic.

*Neuralgia of the Dorsal and Lumbar Nerves; Abdominal Neuralgia.*—The dorsal and lumbar nerves are subject to neuralgic affections, which exhibit, like colic, paroxysms of pain unaccompanied by fever. But Valleix has taught us to look for spots painful to the touch in the course of the aching nerves, and has shown that the disturbance of the nerves supplying the abdominal parietes manifests itself on one side of the body only, whereas an irritation of the intestinal nerves obeys no such law.

In neuralgia of the lumbar nerves, or *lumbo-abdominal neuralgia*, the pain is commonly felt in the hypogastric region, a little to one side of the median line. In this situation, too, there is localized soreness on pressure; the other tender spots are, generally, one a little to the outside of the first or second lumbar vertebra, and one immediately above the middle of the crest of the ilium. In women, who are by far the greatest sufferers from the disease, there is sometimes also a painful place about the middle of the Fallopian tube, or on the neck of the uterus; in men, a point on the scrotum here and there is found sore to the touch. These spots of tenderness serve as characteristic signs; and they enable us to separate neuralgia not only from colic, but also from lumbago, and from rheumatism of the abdominal walls.

Besides these forms of neuralgia, we find other kinds of abdominal neuralgia, which may be mistaken for colic. They are attacks of pain affecting especially the mesenteric plexus or the solar plexus, happening in paroxysms of great severity, and attended with a sense of faintness and annihilation. The disorder is unconnected with lead poisoning or any of the causes that produce colic, is often excited by exertion, and is associated with debility and relieved by an antineuralgic treatment. In some cases it is clearly of malarial origin; and in every instance we must lay stress on the frequent recurrence of the pain and on the history to enable us to discriminate between the neuralgic complaint and colic. The distinction from gastralgia can be made only by the more marked gastric symptoms, and by the absence of, or the less decided, prostration and sense of fainting in this malady.\*

---

\* A number of cases of abdominal neuralgia are reported by Handfield Jones in his *Treatise on Functional Nervous Diseases*; and by Porcher in *Amer. Journ. of Med. Sci.*, July, 1869.

*Abdominal Aneurism and Tumors; Diseases of the Spine.*—In all of these we may find violent pain of a paroxysmal kind referred to various portions of the abdomen, and unaccompanied by fever. We judge that the pain is not colic, by its frequent repetition; by its want of association with intestinal or gastric disturbance; by its being, although liable to exacerbations, so steadily present at some part either of the spine or of the abdomen; and by the attending symptoms and signs occasioned by an abdominal tumor, or by a disease of the lower dorsal or of the lumbar vertebræ.

*Enteritis and Peritonitis.*—Inflammations of the intestines and of the peritoneum also give rise to severe abdominal pain. But it is more constant, and is linked to great tenderness, and, in acute cases, to symptoms of high febrile excitement. Thus enteritis and peritonitis belong to a different group of diseases,—a group of inflammatory affections, which I shall describe somewhat at length, before contrasting the symptoms of inflammation of the intestines or of the peritoneum with those of colic.

### Diseases attended with Pain and marked Tenderness in the Umbilical Region or diffused over the Abdomen.

**Acute Enteritis.**—Enteritis means, by common consent, inflammation of the small intestine, especially of the portion that lies between the duodenum and the colon. The morbid process may extend to the colon; if, however, it involves a large portion of the latter, it is colitis or dysentery. There are two forms of enteritis: one in which the mucous membrane of the bowel is alone affected; muco-enteritis or intestinal catarrh. In the second, more than the mucous tunic is implicated; there is also inflammation of the submucous and muscular coats, or even of the serous investment of the bowel. To this variety of the complaint the term enteritis is by several writers restricted; and it is to this rare form of the malady, a phlegmonous enteritis, occurring acutely, that the description about to be given more particularly applies.

The symptoms of an acute attack of enteritis are those of colic, attended with fever and tenderness. The disorder may begin with the symptoms of colic, or it may set in with chill and fever.



When the disease is fully established, the fever runs high; the pulse, tense and full at the onset, becomes small and wiry. There are nausea and vomiting, and sometimes distressing fits of retching. The tongue is covered with a white coat, or is red and dry. The bowels are constipated; sometimes there is diarrhœa, or constipation alternating with diarrhœa. The stools may contain a small quantity of blood, but they very rarely contain pus. The appetite is lost, the thirst great. The pain, as in colic, is paroxysmal. It begins near the umbilicus, and thence may shift to various parts of the abdomen, but not to the epigastrium; it does not cease as in colic, but rather exacerbates, and then changes to a dull feeling of distress. It is greatly increased by pressure, and the patient seeks relief, as in peritonitis, by lying on his back with his thighs flexed, so as to relax the abdominal muscles. Toward the right of the umbilicus it is not uncommon to find a marked pulsation, as if from throbbing of the abdominal aorta or of its large branches,—a sign to which Stokes\* directed attention. This pulsation may be very annoying. In looking over the notes of my cases on which the description of the symptoms of enteritis just given is based, I find one in which neither the thirst, nor the pain, nor the nausea and vomiting occasioned as much distress as the violent throbbing in the abdomen.

In those instances of the malady that advance to a fatal termination, the pulse becomes quick and irregular and loses its tense-ness; hiccough appears; the abdomen swells; the features are haggard; and the patient's strength becomes gradually exhausted. The worst and most hopeless cases of the disease are those dependent on mechanical obstruction of the bowel, whether it proceed from organized bands in which a loop of intestine is caught, or from invagination, or from accumulation of hardened feces, or from a hernial strangulation. The disease may lead to purulent infiltration of the submucous tissue and to abscess-formation.

Among the symptoms of enteritis mentioned, the pain is one of the most important for diagnosis. It is never absent, save in rare instances in which the inflammation is very intense at the onset.† Still more important is the great tenderness. This en-

---

\* Article "Enteritis," in *Cyclopædia of Practical Medicine*.

† Andral, *Pathologie interne*, tome i. p. 47.

ables us to say that the case, in spite of the colicky pains, is not colic. It tells us, when it succeeds to ordinary colic, that inflammation of the bowel has supervened. It admonishes us not to administer strong cathartics to overcome the constipation that appears in consequence of the severe inflammation.

The disease in its violent form just described bears a close resemblance to peritonitis: we shall presently see what are its distinguishing marks. But there is, as above stated, another variety of the disease, a mild variety, or *muco-enteritis*, in which the disturbance is limited to the mucous membrane. The main features of this intestinal catarrh are the same, but they stand out in less bold relief. There are griping pains, a slight soreness to the touch, general uneasiness, loss of appetite, thirst, nausea, and sometimes vomiting. But we find only slight fever; and the febrile excitement remits in the morning. Diarrhœa is present, and the stools are sometimes very offensive. This form of the disease may terminate, as the severer inflammation generally does, in less than a week; yet it may persist for several weeks, and thus gradually lapse into a chronic complaint. It is common in children, especially during dentition. It is also observed when irritating food or secretions occupy the alimentary canal for any length of time, or after exposure to cold and damp, particularly when the skin is perspiring freely, and as an attendant upon the exanthemata. It resembles typhoid fever. Indeed, it is sometimes difficult, especially in children, or in the intestinal catarrh of catarrhal fever, to know whether we are dealing with a case of simple intestinal catarrh, or with the intestinal symptoms of enteric fever. The state of the cerebral functions, the pain and gurgling in the iliac fossa, and the high temperature of the latter malady, may clear up the doubt; yet in some cases nothing but the eruption and the course of the symptoms will do so.

The symptoms just described belong to catarrh of the ileum, or of the ileum and the ascending colon. In catarrhal inflammation of the duodenum there is often constipation in place of diarrhœa. Pain between two and three hours after the taking of food, loss of appetite, coated tongue, fetid breath, marked digestive disorder, flatulency, and jaundice are prominent among the symptoms. The pain is apt to come on in paroxysms like gastralgia, although referred somewhat lower than the stomach; these seizures last several

hours, and slowly subside. We frequently find a certain amount of soreness developed by deep pressure in the right hypochondrium and the upper part of the umbilical region. There is weakness, with much despondency, and slight elevation of temperature. An acute attack lasts two or three weeks. In the chronic form the duration may be as many months.

Another affection which is liable to be mistaken both for enteritis and for typhoid fever has been described by Klob.\* The chief symptoms are violent pains in the hypogastric region, with vomiting, thready, frequent pulse, high temperature, and the rapid supervention of somnolence and coma. In some instances hemorrhages happen. Hemorrhagic erosions are found in the stomach, and bloody infiltrations in the jejunum; the parenchyma of the mesenteric glands, their lymphatics, and the thoracic duct are infiltrated with blood; the spleen is enlarged. The disorder shows then a striking hemorrhagic tendency, and is supposed to be a blood-affection similar to pseudo-leukæmia.

A *croupous* or *diphtheritic* enteritis is not seen except as a secondary process, if we except the instances in which it follows poisoning by mercury, by arsenic, or by lead. It is more generally encountered as a secondary affection in some infectious diseases, as in pneumonia, or in pyæmia, or in typhoid fever, or in cancer, Bright's disease, or cirrhosis of the liver. Its symptoms may be latent, but generally there are diarrhœa and pain without tenesmus.

**Acute Peritonitis.**—As in acute enteritis, so in acute peritonitis, pain and tenderness are the most significant symptoms. To these are joined fever, distention of the abdomen, and, frequently, cold sweats, nausea, vomiting, and obstinate constipation. The disease begins with chilly sensations or protracted rigor. To these succeed fever, and abdominal pain and distention. The fever runs high at the onset; it exhibits a dry, burning skin, an axillary temperature of  $103^{\circ}$  and upwards, a pulse frequent, but, as in acute inflammations of the mucous and serous membranes below the diaphragm, small and wiry. However, both the character of the pulse and that of the skin change as the malady progresses. The pulse will be less tense and more developed as

---

\* Wien. Med. Zeitung, quoted in Lond. Med. Record, Feb. 1875.

the inflammation subsides, or feeble and flickering if the disorder proceed toward a fatal termination. The skin is frequently covered with cold sweats. The temperature is irregular, and may sink below the normal. The features are sharpened and wear the look of death, even in cases which ultimately recover.

The pain is constant and severe. It may exacerbate, but it never intermits. At first the pain is confined to a particular point; but as the inflammation extends, so it extends over the whole abdomen. It is increased by the slightest pressure, be that pressure exerted by the hand or by movements of any kind. To obviate the pressure, the patient lies on his back with his thighs flexed, and, however tired of retaining the same position, he does not change it. The descent of the diaphragm augments the pain: instinctively, therefore, he refrains from drawing long breaths, and his respiration is short and frequent and purely thoracic.

The abdominal distention is in part owing to meteorism, in part to the liquid effused into the peritoneum. Percussion tells us in individual cases how far each factor acts as a cause of the enlargement, by the tympanitic or the dull sound elicited. Palpation, too, reveals the presence of liquid. Yet percussion or palpation ought to be employed only with the greatest care, on account of the pain they occasion. The fluid does not gravitate as invariably as in ascites to the lower portion of the belly. It is often caught in sacs formed by the membrane adhering in spots; and thus circumscribed dulness may be found at one or several parts of the abdomen. Sometimes the roughening of the membrane gives rise to a distinct friction sound.

Independently of the abdominal pain and swelling, we meet, in acute peritonitis, with constipation, nausea and vomiting, headache, a suppression of the urinary discharge, and in rare instances with priapism; of these symptoms, constipation is the most constant. The bowels are never relaxed, except in the puerperal form of the malady. The constipation is caused by the paralyzed state of the intestine, to portions of which the inflammation may spread; or by the lymph gluing together the coils of the bowels.

Death in acute peritonitis is commonly preceded by enormous tumefaction of the belly, cold sweats, a pinched countenance, and a rapid, flickering pulse. When recovery takes place—unfortunately a rarer issue of the malady than its fatal termination—it is



commonly very slow and gradual : the symptoms diminish one by one ; they do not cease suddenly ; and often morbid conditions remain which prolong greatly the patient's illness and may lead in themselves to a disastrous result. It is, therefore, impossible to foretell the duration either of the acute disease or of its consequences. Andral fixes the average length of an acute attack at between six and nine days, and of a subacute attack at from twenty to thirty days. But the nature of the malady is such that many cases last a longer, many a much shorter period.

The presence of gas in the peritoneal cavity is always a very valuable sign of perforative peritonitis. Tympanitic resonance over the hepatic region is thus occasioned. On the other hand, when the hepatic dulness is found, the inference is a fair one against perforation of stomach or intestine as a cause of a peritonitis that is detected.

Acute peritonitis arises only very occasionally idiopathically from exposure to cold and wet ; much oftener in consequence of injuries to the abdomen, such as blows, stabs, or kicks ; or from perforation or laceration of some of the abdominal organs, such as perforative ulcer of the stomach, intestine, or gall-bladder, and discharge of their contents into the peritoneal cavity, or from a ruptured tubal pregnancy. Uterine injections passing into the peritoneal cavity may cause peritonitis. It also results from rheumatism,\* or from a poisoned state of the blood, as, for example, the peritonitis of childbed fever, or from acute tuberculosis, or from Bright's disease. It sometimes originates from an inflammation of the abdominal viscera, especially of the spleen, intestines, or uterus and its appendages, spreading to their serous covering, and thence extending more or less rapidly. Again, other morbid states of the abdominal organs, such as cysts of the ovaries, intestinal intussusception, or strangulated hernia, may compress or irritate the membrane, and lead to inflammatory action. Owing to these diverse sources, peritonitis presents varieties which exhibit points of difference that require special notice.

The inflammation produced by *extravasation* into the peritoneal sac is characterized by its sudden development. The matters ex-

---

\* Schmidt's Jahrbücher, No. 9, 1873.

travasated may be blood, or bile, or urine, or the contents of the stomach. Most frequently perforation of the stomach or intestine lies at the bottom of the mischief. Whatever its cause, the perforation is immediately followed by collapse; and tenderness and distention of the abdomen soon make their appearance. Yet peritonitis may set in rapidly in cases in which there has been no rupture; and, on the other hand, in rare, very rare, instances, the contents of the alimentary canal may be discharged into the sac without giving rise to inflammation.\*

The peritonitis of childbed fever, or *puerperal peritonitis*, is principally distinguished by its occurring during the puerperal state. Its symptoms are, so far as the peritoneal inflammation is concerned, those of any other kind of peritonitis, except that diarrhœa, instead of constipation, is often present. The disease is generally ushered in by chills. The temperature rises speedily to a considerable height, to  $104^{\circ}$  or  $105^{\circ}$ , and continues high with irregular remissions. The uterus or the uterine appendages are generally first attacked; and it is in these regions that pain and tenderness are first felt. The inflammation spreads to their serous investment, or it may be primarily seated in that investment: in either case it soon involves the entire membrane. But, independently of the symptoms of the local disorder, there are phenomena which clearly belong to the diseased state of which the inflammation of the peritoneum is but a local expression; there are evidences of a poisoned state of the blood, of a septicæmia. We find delirium, black vomit, exudation into the pericardium and pleura, features of disease not met with in the purely local malady, and showing that, while childbed fever occasions peritonitis, peritonitis does not constitute childbed fever. Fortunately, the diagnosis is one we are now less and less often called upon to consider, for antisepsis has almost put a stop to the disease.

*Partial or local peritonitis* is almost invariably owing to a pre-existing morbid condition of some abdominal viscus. Sometimes the circumscribed inflammation is protective rather than calculated to work mischief. It arrests a destructive perforation of the mem-

---

\* Cases reported by Bardeleben and Siebert, quoted in Henoch's *Clinic of Abdominal Diseases*. Instances of rapid peritonitis without perforation are given by Thirial, *L'Union Médicale*, 1853.

brane, or it limits the matter discharged to a certain spot ; it may at least do so for a time, for general peritonitis is very apt ultimately to follow.

Partial peritonitis often pursues a subacute rather than an acute course. It may end in adhesions or lapse into a chronic state. Its symptoms are much the same as those of a more general inflammation,—the same fever and constipation, the same pain and tenderness. The fever does not, however, run so high, and the pain and the great tenderness are much more localized. The abdomen, also, is not so swollen or so tympanitic. But perhaps even more frequently than in general peritonitis are found accurately-limited spots of dulness on percussion corresponding to circumscribed collections of pus in the peritoneal cavity.

Partial peritonitis is more liable than the general disease to be confounded with other disorders. Yet error can hardly arise, or, should it arise, it is not of much consequence, provided we bear in mind that it is precisely with the morbid states of the viscera which lie below the peritoneum that the circumscribed inflammation of the serous membrane is usually connected, and that local peritonitis, therefore, frequently attends the very disorders from which it is sought to be distinguished. Let us, however, examine into some of the complaints with which peritonitis, whether local or general, may be confounded. They are—leaving for consideration elsewhere obstruction of the bowel, appendicitis, and perityphlitis—

- ACUTE GASTRITIS ;
- ACUTE ENTERITIS ;
- ACUTE PANCREATITIS ;
- METRITIS ;
- CYSTITIS AND DISTENTION OF THE BLADDER ;
- RHEUMATISM OF THE ABDOMINAL WALLS ;
- ABDOMINAL HYSTERIA ;
- COLIC.

*Acute Gastritis.*—Acute inflammation of the stomach can scarcely be mistaken for inflammation of the peritoneum, provided attention be paid to the history of the case and to the seat of the pain. The former disorder begins with vomiting, and this continues a prominent symptom throughout ; whereas vomiting is not so constant, nor does it occur so early, in peritonitis. The

pain and tenderness are limited to the region of the stomach in gastritis; they are diffused and accompanied by general abdominal enlargement in peritonitis. They may, it is true, be localized when the peritonitis is partial. But acute inflammation of the gastric peritoneum is hardly encountered, save as an attendant on severe inflammation of the stomach, or on a destruction of its coats,—the form of gastritis which results from irritant poisons.

*Acute Enteritis.*—Enteritis differs from general peritonitis by the less extended tenderness; by the seat of the pain near the umbilicus, and its more paroxysmal character; by the comparative absence of tympanites and abdominal tumefaction; and by the greater prominence of nausea and vomiting. It is, moreover, a disease far less violent and dangerous than acute peritonitis; yet it cannot be distinguished with certainty from the partial form of this disorder, to which, in truth, some of its symptoms are clearly owing.

*Acute Pancreatitis.*—This is a cause of peritonitis which may be easily overlooked. The pancreatic inflammation mostly arises in consequence of the extension of a gastro-duodenal inflammation along the pancreatic duct; or it may follow hemorrhage into the pancreas. In the former case we find sudden pain, deep-seated, constant, or paroxysmal, tenderness, and tympany in the epigastrium in the region of the pancreas, with nausea and vomiting. This is gradually followed by peritonitis at the same place, and by a low fever. Constipation is frequent, and, with the other symptoms, has led to the diagnosis of acute intestinal obstruction, and to laparotomy. In *hemorrhagic pancreatitis* the symptoms run a rapid course. The disease occurs in persons over thirty years of age. The attack begins with violent pain in the abdomen; nausea, vomiting, and tympanitic distention soon follow, and signs of collapse appear. There is usually constipation. The temperature, as we know from Fitz's\* comprehensive study, may remain normal. The disease is most likely to be confounded with acute perforative peritonitis. It usually proves fatal in from two to four days. The hemorrhage may lead to gangrene; in either case the signs of peritonitis are marked. Hemorrhage may occa-

---

\* Middleton-Goldsmith Lecture for 1889.



sion sudden death.\* *Abscess of the pancreas* has much the same symptoms ; but it does not run so acute a course,—is, indeed, often chronic ; the pain is much less ; there is apt to be irregular fever.

*Metritis*.—Inflammation of the womb is not likely to be mistaken for general peritonitis ; the pain on pressure, which they have in common, is confined in the former disease to the uterus and its annexes, and there is little or no tympanites. It is thus only that the acute metritis of childbed fever may be distinguished from the acute general peritonitis of the same malady. When the puerperal malady attacks the uterus as well as the whole peritoneal surface, the signs of inflammation of the serous membrane mask those of inflammation of the womb.

A local inflammation of the peritoneum occurs still more constantly as an attendant on inflammation of the womb and its appendages, whether the disorder of the sexual organs be or be not puerperal. It frequently leads to collections of pus, which can be readily felt through the parietes of the abdomen or through the rectum and the vagina, and which sometimes discharge into the bowel or vagina after a lingering sickness. The proofs that the uterus is involved in these cases of partial peritonitis, are the signs of its disordered functions and the excessive pain occasioned by pressing on the cervix during an examination per vaginam.

*Cystitis and Distention of the Bladder*.—Both inflammation and distention of the bladder are occasionally mistaken for general acute peritonitis. An acute inflammation of the bladder gives rise to frequent calls to pass urine : yet the act is performed with great difficulty, and in severe cases may become impossible ; the bladder distends ; a sense of uneasiness is felt in the perineum ; the region above the pubes becomes tender, and sounds dull on percussion ; the unhappy sufferer is restless and distressed ; he has the excited pulse and the hot skin of fever ; at times vomiting and hiccough supervene ; and death is preceded by gradually-deepening coma. Such cases resemble those of peritonitis with suppression of the urinary discharge and with strangury. But the urine voided in peritonitis is simply high-colored, like that of any febrile state. In cystitis it contains large quantities of mucus and pus, and often blood and crystals of phosphates.

---

\* Draper, Transact. Assoc. Amer. Physicians, 1886.

Again, the abdominal tenderness is localized, and is frequently accompanied by a smarting in the course of the urethra. Neither of these signs is encountered in peritoneal inflammation, and, as a rule, the temperature in this is higher. The disturbance of the urinary organs which not unfrequently takes place in the latter disorder is attributed to inflammation of the part of the peritoneum covering the bladder or its immediate neighborhood.

An overdistention of the bladder, not the result of inflammation of its coats, may produce a local tenderness spread over a considerable portion of the lower part of the abdomen. But the outline of the dulness, which is coextensive with that of the tenderness, the fact that the patient has generally not passed urine in any quantity for a considerable time, the almost normal temperature, and the sudden cessation of the supposed peritonitis on passing a catheter, show the true nature of the malady.\*

*Inflammation and Abscess in the Abdominal Muscles.*—When the abdominal walls become inflamed, symptoms are occasioned that are not always easily distinguished from those of acute peritonitis. The disease is attended with some fever, with pain increased by movement, by the act of coughing, and by pressure, and sometimes with excessive tenderness. The seat of the inflammation is generally the rectus muscle and the surrounding cellular tissue. The parts on one side of the umbilicus are most commonly attacked, and it is there that a hard swelling is perceived, over which the skin is rather hot and sometimes red. The tumefaction gradually disappears by resolution, or else fluctuation becomes from day to day more distinct, showing that suppuration is taking place; and the pus being discharged, immediate relief follows, and the pain and febrile symptoms instantly cease.

Now, the disease rarely runs a very acute course; it lasts at least a week or two, and often much longer. Where much of the muscle is involved, the complaint simulates peritonitis,—more, however, the partial than the general kind. Where the inflammation of the muscle is not extended, the resemblance to inflammatory affections of the organs lying underneath the point of tenderness is even greater than to inflammation of the peritoneum.

---

\* A case of this kind, occurring after delivery, is given by Lever, *Guy's Hospital Reports*, 2d Series, vol. viii. p. 41.

Hepatitis, splenitis, and gastritis have been mistaken for the affection of the abdominal parietes. These errors can be avoided only by taking into account the absence of disturbed function of the suspected viscus; often, too, the peculiar swelling furnishes a clue to the real nature of the case. But as regards signs of disturbed function, we must bear in mind that these are produced occasionally by disorder of adjoining viscera. Thus, we have jaundice in abscesses seated in the walls in the right hypochondrium.\* Abscesses in the abdominal walls are sometimes symptomatic of a more distant lesion, as of caries of a rib.†

Can we distinguish, with anything like certainty, between abscesses in the abdominal walls and instances of partial peritonitis leading to *collections of pus in the peritoneal cavity*? I believe not; for in both there is a tumefaction; in both the general symptoms are much the same; and, as happens sometimes in peritoneal abscesses, the pus presses its way through the parietes of the abdomen. How, then, are we to know where was the seat of its formation? Whenever we find a swelling which has come on gradually, or has followed a blow or a kick on the abdomen, or a swelling which is very hard before fluctuation appears; whenever the softening of the tumor is immediately preceded by distinct chills, and the skin covering it is tense, and heated, or reddish; wherever there is nothing pointing to the occurrence of partial peritonitis, as an attendant on visceral disease, or as a consequence of an attack of general peritonitis,—we may infer, from the history and the signs, that the affection lies in the abdominal walls. But the skin is not always discolored or hot, and the beginning of the swelling is sometimes veiled in obscurity. In some instances I have seen, in which there was great doubt, the aspirator drew off a very offensive pus and broken-down material; and I looked upon this—as the sequence proved, correctly—as indicating abscess in the abdominal walls. Abscesses within the abdomen seated at the upper part, if not caused by abscess of the liver, are, as Bristowe accurately points out,‡ largely due to perforation of one of the hollow viscera with circumscribed peritoneal suppuration.

---

\* As mentioned by Habershon, *Diseases of the Abdomen*, 1878.

† Oppolzer, *Wien. Med. Wochens.*, 1862.

‡ *Lancet*, Sept. 1883.



But it is not every case of abscess in the walls which is attended with symptoms that render it likely to be mistaken for the results of inflammation. Sometimes the preceding tumefaction is so hard, or it is so long before the process of suppuration sets in, that the affection is more liable to be confounded with abdominal tumors. The most trustworthy points of difference are furnished by a study of the history of the case, and of the mode of invasion; by the slow growth of the tumor on the one hand, and its far more rapid growth on the other; and by the absence, or at all events the comparative absence, of signs denoting serious disturbance in one or several of the abdominal viscera. Then, in doubtful cases, the aspirator or the exploring needle will be of use. The fluid thus obtained shows, under the microscope, shreds of broken-down muscle and of areolar tissue, mixed, if suppuration have commenced, with pus. Again, stress may be laid on the occurrence of chills preceding the softening of the mass. In some patients the inflammation is unaccompanied by any appreciable signs; it leads to gradual changes in the muscular fibres, which do not reveal themselves until the disorganized muscle gives way. The fibres undergo softening or a true fatty metamorphosis, and the slightest force suffices to produce a rupture. Not a few cases have been reported in which one of the recti muscles has been torn asunder during a fit of coughing. The seat of laceration is generally about midway between the umbilicus and the pubes, a little to one side of the median line; the rent fills with blood, occasioning a circumscribed swelling and rigidity of the abdomen. There is sometimes pain, with nausea, vomiting, and obstinate constipation. Nay, the symptoms have mimicked so closely a strangulated ventral hernia as to have led to the performance of an operation.\*

*Rheumatism of the Abdominal Walls.*—Occasionally rheumatism

---

\* Richardson's case, *American Journal of the Medical Sciences*, Jan. 1857. Further instances of this accident are given by Virchow, in the *Würzburg. Verhandl.*, Band vii. The description of abscesses in the abdominal parietes I have drawn from cases coming under my own notice, from manuscript notes taken by Dr. J. K. Kane at the Philadelphia Hospital, and from the cases collected in the *Dictionnaire des Dictionnaires de Médecine*, art. "Abdomen." See also Paul Deriencourt, *Thèse de Paris*, 1886, No. 153; Marsigny, *Arch. Méd. Belges*, Bruxelles, 1886, 3e sér., xxix.; Weisz, *British Med. Journ.*, April, 1891; Moran, *Physician and Surgeon*, Ann Arbor, Mich., Aug. 1893.



attacks the abdominal muscles, and gives rise to local symptoms similar to those of peritonitis. But the pain is not so constant, nor is it spontaneous, as in this disorder. It is also less affected by movements or by pressure. Not that these diminish it; on the contrary, they aggravate it. Yet deep pressure causes little or no more pain than slight pressure; and it is only during certain motions—when the muscles are placed on the stretch—that the pain is severe, or sometimes, indeed, at all produced.

The pain is often one-sided, or, at any rate, much more marked on one side, and we find no meteorism, and but slightly elevated temperature, and not the anxious countenance of peritonitis. Moreover, the attack is apt to happen in those of rheumatic tendencies, and there is concentrated, highly-acid, scalding urine. So strong a degree of similarity may, however, exist between the two diseases as to keep judgment in suspense. In such cases it is better to treat the disorder as if it were inflammation of the peritoneum. In point of fact, it may happen that such inflammation does succeed to the rheumatic affection of the abdominal muscles, and this occurs chiefly when the disturbance in the muscles forms part of an attack of acute rheumatism having a decided tendency to shift its seat.

*Abdominal Hysteria.*—No disease simulates peritonitis more closely than hysteria. The abdomen may be extremely painful to the touch, swollen and distended with gas, fever may set in temporarily, and yet the whole disorder be purely hysterical. To illustrate:

An unmarried woman, twenty years of age, placed herself under my care, on account of extreme tenderness of the abdomen and febrile irritation, both of which had become developed in a few days. The abdomen was swollen and tympanitic, and so sensitive that it would not bear the pressure of her clothes; the pulse was frequent; the skin dry and warm; the tongue slightly coated; the bowels constipated; the countenance expressive of distress. Here was certainly a group of symptoms like those of acute peritonitis. But the absence of the wiry pulse, the comparatively slight fever,—slighter, certainly, than was to be expected from such general and great tenderness,—and the expression of countenance, arrested my attention. I found that the patient had had similar attacks previously; that they had come on sometimes

shortly before, sometimes shortly after, her menstrual period ; but that for several months her menses had ceased to flow. The abdominal tenderness was in reality, as she represented it to be, very great ; yet strong pressure produced no more pain than the lightest touch. Nor was the pain increased by deep inspiration, or by coughing, or by extending the thighs. Taking all these circumstances into account, as well as her age and sex, and her nervous temperament, instead of treating her for acute peritonitis, cold-water injections, mild purgatives, and a mixture of assafetida and valerian were employed. Under these remedies, all the symptoms of the apparent peritonitis speedily vanished.

Yet all cases of abdominal hysteria do not pass off so quickly ; sometimes they are much more persistent, or recur frequently, as I know from personal observation. They are from the onset unattended with fever, or, as the thermometer shows, the fever is fitful and soon ceases. The absence of febrile excitement, too, especially if taken in connection with the several localized and more or less distinctly circumscribed spots of tenderness, enables us to distinguish between peritonitis and those instances of neuralgia of nerves supplying the abdominal parietes, to which women who are laboring under disorders of the uterus are so liable. It is in these cases, as well as in all instances of abdominal hysteria, that the thermometer proves a most useful aid in the diagnosis.

*Colic.*—As already stated, the pain of colic is paroxysmal, and not attended with fever, or with much, if any, tenderness ; while the pain of an inflamed peritoneum is constant, and associated with the greatest tenderness and with fever. Cases of colic do indeed occur in which we find fever and some tenderness ; but these signs are then still out of proportion to the amount of pain. The pulse is not wiry, nor the tenderness so exquisite or so diffused. Further, it is not unlikely that in such cases the peritoneum is really in parts injected or slightly inflamed.

The same remarks are applicable to those severe paroxysmal pains which accompany the passage of gall-stones or of urinary concretions, or which occur at the menstrual periods. They are frequently spoken of as varieties of colic, and, as far as their discrimination from peritonitis goes, there is no difference,—it rests on the same grounds precisely ; for when there is fever or tenderness on pressure, it is likely that inflammation has been set up in

those parts in which, or in the neighborhood of which, the pain is felt. In the so-called uterine colic, an injection of the peritoneum has positively been demonstrated.

**Chronic Peritonitis.**—An acute attack of peritonitis may imperceptibly assume a chronic form. The fever gradually disappears, or at all events lessens; but the exudations into the peritoneal cavity, whether organized or not, remain, and so do some abdominal pain and tenderness. In this condition the patient may continue for many months, now and then a fresh inflammation starting up in the peritoneum and giving rise to acute symptoms, or an intercurrent severe diarrhœa leading to rapid loss of strength. Again, the disease may develop slowly, be latent from the onset, and may not attract attention until the abdomen swells. In all cases, no matter what their origin, if they last for any length of time, debility and emaciation become marked symptoms; then hectic fever is observed; decided effusion into the peritoneum is generally noticed; the legs become œdematous; and the patient may die worn out and presenting the symptoms of septic poisoning. Where recovery takes place, the exudation into the peritoneal cavity is either discharged through adjacent viscera; or may be gradually absorbed; or may be transformed into tissue. When the disease terminates in this way, it is apt to leave its traces in a chronic thickening and roughening of the peritoneum. A friction may be often felt. Chronic peritonitis of latent origin and leading to much thickening is sometimes found to attend cirrhosis of the liver or of the stomach, or contracted kidney. Under no circumstances is chronic peritonitis likely to be an independent affection.

Chronic peritonitis may be confounded with affections of the liver attended by impediment in the portal circle; and what adds to the difficulty in diagnosis is, that the liver is apt to atrophy in chronic diffuse peritoneal inflammation. The most trustworthy signs of distinction are that, in the latter affection, tenderness exists, and is under any circumstances much greater and more diffuse; that there are evening exacerbations of temperature, a quickened pulse, dark stools; and that, if the veins of the abdomen are dilated, their dilatation is slight and uniform.

Chronic peritonitis more usually comes on and ends in a particular fashion. It is insidious in its approach, and its fatal ter-

mination is preceded by evident signs of *tubercular* or *cancerous deposits* in the abdominal cavity or in the lungs. The disease is not then simply chronic peritonitis, but chronic peritonitis in connection with a cachexia. Cases of the kind are commonly of long duration. They are attended with ascites, and often with very considerable abdominal distention. I shall, therefore, postpone most of what I have to say about their diagnosis until I come to abdominal enlargements, and shall then consider what differences there are between these various forms of chronic peritoneal affections and other disorders leading to ascites and to consequent abdominal distention.

### Diseases attended with Pain and Tenderness in the Right Iliac Fossa.

**Appendicitis.**—Inflammation of the appendix is pre-eminently the disease attended with pain and tenderness in the right iliac fossa. The appendix has an average length of four inches, and the diameter of a goose-quill. It lies in the right iliac fossa, but is variable in position. It points for the most part downward, or downward and inward. A. T. Bristow and Fowler\* locate for it a central point by drawing a line from the anterior superior spinous process of the ilium to the median line, and placing the central point from two to two and a half inches within the anterior superior spinous process. From this central point the appendix will radiate in different directions. The usual location of the appendix is at the edge of the right rectus muscle below a line drawn from the centre of the umbilicus to the anterior superior spinous process of the ilium. Appendicitis is essentially a disease of adolescence and of young adults. It presents itself clinically in these forms: acute catarrhal appendicitis; ulcerative and suppurative appendicitis; perforative appendicitis; chronic recurring appendicitis.

Acute catarrhal appendicitis may come on from exposure to cold and wet. Fowler cites two such cases. Much more generally it is an infectious process due to hardened faecal masses leading by the irritation they produce to exudations in which extraordinary development of bacteria, as of the bacterium coli commune, takes

---

\* Appendicitis, Philadelphia, 1894.



place. It may also be caused by other infecting processes or micro-organisms. The far greater prevalence of appendicitis since the recent wide-spread epidemics of influenza suggests that this subtle poison, too, may act as an exciting cause. The disease may also result from vascular disturbances or torsion of the part. It is at the bedside always extremely difficult to say which cause has given rise to the attack.

Whatever the immediate cause, whether it is a quickly-acting one, or, as is more common, has been silently working, the attack itself is generally sudden, and announces itself by acute abdominal pain, by tenderness in the right iliac fossa, by nausea and vomiting. The pain and the tenderness are very significant. The pain may be referred to the lower part of the abdomen, but it is very often referred to the umbilicus or to the epigastrium. It has, especially at first, the character of colic. It is soon noted to be associated with tenderness, which is especially manifest at or near McBurney's point. This corresponds to the outer edge of the right rectus muscle, and is most readily located by fixing a spot midway between the anterior superior spine of the right ilium and the umbilicus. The patient lies on his back, because to do otherwise means to him increase of pain, and very often the right rectus muscle is somewhat tense, a fulness or a slight tumefaction can be perceived in the right iliac fossa, and there is some impairment of tympanitic resonance on percussion. Tenderness and swelling, as well as the shape of the appendix, may at times be recognized by deep pressure, and palpation of the appendix, as recommended by Edebohls,\* may thus become of value. In some instances, and I have met with a number of them, the sensitiveness is not in the right but in the left iliac fossa, and pressing with one finger readily develops it. Sensitiveness is always a very important sign, and when it lessens both in degree and in extent it denotes decreasing inflammation. The nausea and vomiting disappear in the progress of the case, though vomiting may return should there be perforation. If the peritonitis become general, abdominal distention will be marked. Other symptoms met with in appendicitis are constipation, urine diminished and frequently containing albumin and indican.

---

\* Amer. Journ. Med. Sci., May, 1894.

Some cases do not begin so acutely, but are rather subacute. The complaint presents the following history and symptoms. The patient has been suffering for some time from constipation, or alternately from diarrhoea and constipation. He has a dull pain referred principally to the iliac fossa, and radiating to the hips. When the iliac region is examined, it is tender to the touch, full and hard, and sounds dull on percussion, while around the dulness there is a very tympanitic sound, if the intestine, as is often the case, is much distended with gas. Colicky pains occur from time to time, but are mainly confined to the lower portion of the abdomen.

No matter what its beginning, the case in its further progress exhibits varied features: it may end in resolution, and hardened faecal matter is often passed; or the tenderness in the iliac fossa may become greater, and vomiting, decided fever, and the marked signs of an extending peritonitis appear; or ulceration of the appendix may allow a discharge of extraneous matter into the peritoneal cavity, which produces violent general peritonitis, or an abscess forms that ruptures and perhaps leads to the same results; or, again, the bowel may become so paralyzed or so constricted that it can no longer propel its contents, and the patient dies with all the distressing signs of intestinal obstruction.

There are other terminations with which experience makes us familiar. The attack may end in a chronic appendicitis, indicated by persistent tenderness and some swelling, pain on walking, and often dyspeptic symptoms; or the chronic inflammation may lead to a series of recurring acute attacks, of which in one instance, under my observation, there were forty-seven before the case was operated on. Then as complications in appendicitis we may have thrombosis of the iliac vein, iliac phlebitis, post-cæcal abscess, fistula into the bladder or rectum, hepatic abscess.

There are two most important questions that always arise in appendicitis: Is there pus present? Has perforation occurred?

It is always very difficult to determine the presence of *pus*, and there are no certain signs. Chills are generally absent; the temperature is of little value. The most trustworthy signs are very decided tenderness, a local swelling, and waves of pain in the affected region.

*Perforation of the appendix* is most often seen among healthy

young men. It is found chiefly in the form of appendicitis that has been caused by seeds and concretions of various kinds, cherry-stones, and foreign bodies. In a certain proportion of cases the symptoms have been latent until the perforation happened. Its most constant and the first decided symptom is sudden, severe abdominal pain. It occurred in eighty-four per cent. of the cases which Fitz in his admirable essay has analyzed.\* The pain is mostly at first in the right iliac fossa, and is followed by tenderness which gradually extends. It may be accompanied by a chill, but I have known it absent where a chill was very decided. Fever, with a temperature of between  $100^{\circ}$  and  $102^{\circ}$ , is next observed; but it is not constant, for I have met with a temperature nearly normal in a case in which a gangrenous perforation of the appendix was found.† A circumscribed resisting swelling in the right iliac fossa, which forms in from two to five days, with impaired resonance on percussion and with a sense of fluctuation from the abscess that develops, and disturbed micturition, establish the diagnosis. A rectal examination may aid us in detecting the tumor, but, as I know from experience, is not absolutely to be depended on as a means of recognizing the swelling or the pus that has formed. In the majority of cases general peritonitis begins from the second to the fourth day after the perforation. The cases that die from shock die before the second day; but, as a rule, the collapse comes on more slowly than in other forms of perforative peritonitis. Leucocytosis, Richardson tells us, is invariable in perforative appendicitis.

A question that arises is whether we can distinguish inflammation of the appendix from an *inflammation of the cæcum*, both of which were formerly included under the name *typhlitis*. There is no certainty in the diagnosis. But these facts will often aid us greatly. Most of the cases of inflammation of the cæcum are due to impacted feces, and the history of preceding long-continued constipation, a resisting elongated mass in the right groin, slight pain, and absence of fever, are very significant. Then, perforating inflammation of the cæcum is very rare, while perforation of the appendix is of frequent occurrence.

---

\* Transact Assoc. Amer. Physicians, 1886.

† Seen with Dr. Morton.

Much used to be said about inflammation of the loose areolar tissue around the cæcum, *perityphlitis*, and consequent abscess. But we now know that the abscess nearly always has its origin in disease of the appendix, and is often the persisting result of an acute or a subacute attack. The collection of pus may find its way into neighboring viscera, or be discharged externally, or become encysted, or the sac rupture and fatal peritonitis ultimately ensue. The tumefaction which the abscess occasions is generally very evident. When, however, the pus burrows under the iliac fascia, the swelling may be slight. But under such circumstances there appears a characteristic sign: the pain, on moving the right foot, is intense, because the iliac muscles become involved in the disorder. If the swelling be great, there may be œdema of the foot and numbness of the thigh, from the pressure on the vein and nerves.

Chiefly on account of the pain and tenderness, acute appendicitis may be confounded with a number of diseases, prominent among which are colic; bilious colic; renal colic; typhoid fever; ulceration of the lower part of the ileum; obstruction of the bowel; tumors of the kidney and abscesses in or around it; floating kidney; inflammation of the right ovary; extra-uterine pregnancy; pelvic hæmatocele; rupture of the gall-bladder; dropsy of the gall-bladder; abscess in the abdominal walls; psoas abscess; hip-joint disease; abscess of the liver; distention of the cæcum; cancer of the cæcum; pneumonia.

The sudden pain, the acute indigestion, the nausea and vomiting may cause appendicitis at its beginning to be mistaken for *colic*, especially for bilious colic, but the localization of the pain and particularly the tenderness in the right iliac fossa are very different. On the other hand, the jaundice that attends or follows *bilious colic* is not a symptom of appendicitis, and the pain of this does not radiate to the shoulder and the scapula. The same localization of the tenderness is of value in distinguishing *renal colic*, where the tenderness, if it exists at all, is most marked over Poupart's ligament. Moreover, rectal and vesical tenesmus and retraction of the testicle, common in renal colic, are very rare in appendicitis.

Pain and tenderness in the right iliac fossa may be the cause of *typhoid fever* being confounded with appendicitis. But neither pain nor tenderness is great in typhoid fever; then the character-



istic temperature record, the nervous symptoms, the diarrhœa, the eruption, furnish striking points of difference.

Ulceration of the lower part of the *ileum* produces pain and tenderness in the iliac fossa. But, combined as the ulceration generally is with tubercular disease, the history of the case gives a clue to the nature of the malady. Moreover, diarrhœa occurs, and there is not present a tumefaction which sounds dull on percussion. Should, however, perforation of the bowel take place before the patient is seen, and general peritonitis come on, the diagnosis is not so readily made, because we are deprived of the decisive proof furnished by the swelling.

Another very difficult diagnosis is at times as regards *obstruction* of the bowel; the more difficult because appendicitis may become a cause of intestinal obstruction. In both there is pain; in both constipation; in both vomiting. But the pain in obstruction is not localized, or attended with such a significant point of tenderness as McBurney's point; the constipation in appendicitis is not so absolute, and flatus passes; the vomiting in this disease occurs early, then generally stops; late vomiting is the rule in obstruction, and it becomes fecal. Though fever is not a marked symptom of appendicitis, there is generally some. An *invagination of the bowel* has a different history, and makes its appearance suddenly with such peculiar signs that, although it may be likewise the occasion of a tumor in the right iliac region, it can be generally distinguished from appendicitis. Yet, where the latter leads to intestinal obstruction, the diagnosis is not always obvious; and tenesmus and discharge of bloody mucus from the rectum may happen in appendicitis.

As regards *tumors of the kidney and abscesses in it or around it*, the situation of the swelling is not exactly in the ileo-cæcal region, or at all events it is not confined to this region. The mass of the tumor lies in the loin, or above the anterior termination of the crest of the ilium; and the urine contains ingredients, such as pus, or blood, or heavy deposits of urates or phosphates, which show that the secretion of the kidney is abnormal. Moreover, there is no intestinal disturbance or marked local tenderness, such as we find in appendicitis. In *floating kidney* the mobility of the displaced organ, and the slight tenderness, are very significant.

An *inflammation of the right ovary* gives rise to pain and ten-

derness in the right iliac region, and to fever. But it is attended with disturbance of the uterine functions, and occasions no very perceptible swelling. A tumor of the ovary or of the uterus may produce a visible tumefaction; but, springing as it does out of the pelvis, its exact seat, its bulk, its shape, the absence of marked intestinal symptoms, and a per vaginam examination, will permit its cause to be discovered.

*Extra-uterine pregnancy* may be mistaken for acute appendicitis in consequence of the sudden rupture of a sac. But the previous history, the great prostration, the excessive thirst, and a pelvic examination will explain the true meaning of the symptoms. In *pelvic hæmatocele* the pain and the suddenness of the attack make us think of acute appendicitis. But the tumor that forms is generally larger, doughy; there are no localized spots of tenderness, no marked intestinal symptoms; and the history of irregular menstruation and a per vaginam examination will remove all doubt.

*Rupture of the gall-bladder* may give rise to right-sided tenderness, to rigidity of the rectus muscle, and to severe pain, and may be unavoidably mistaken, as the case of Fowler\* proves, for acute appendicitis. Several instructive cases reported by the same surgeon prove that unless the history help us, and point conclusively to the occurrence of gall-stone, it is extremely difficult to distinguish *dropsy of the gall-bladder* in which a swelling may be felt low down, from appendicitis.

An *abscess in the abdominal walls* furnishes very many of the signs of abscess around the appendix. The most trustworthy point of distinction is that the former moves with the abdominal walls and is unassociated with intestinal irritation, while the latter is commonly so combined. Then the peculiar spots of tenderness, the outline of the swelling, its want of prominence, are unlike what is found in abscess of the abdominal walls.

In *psoas abscess* we have the association with caries of the vertebræ: an excurvation of the spine, dorsal pain and tenderness, testify to this connection. It occurs in scrofulous persons, and, although gradual in its formation, is often sudden in its manifestation; for not unusually a fluctuating, painless tumor appears

---

\* Op. cit.

below Poupart's ligament as the first positive sign of this formidable affection. This is very different from the history of an appendicitis which has led to post-cæcal abscess. Moreover, preceding the pointing of the psoas abscess at the spot mentioned, there are often indications of irritation in those muscles in the sheath of which the pus travels; there is difficulty in extending the leg, with inability to stand upright.

In *hip-joint disease* the inclination of the pelvis, and the inability to move the joint normally, furnish trustworthy points of distinction.

It is sometimes difficult to distinguish between appendicitis, especially in its chronic forms, and *abscess of the liver*; the more difficult because, as I know by experience, they may coexist, the hepatic abscess being consequent to the appendicitis. Another point that makes the diagnosis difficult is that the pain and tenderness in appendicitis do not always exist in the right iliac fossa, but may be found at various parts of the abdomen; the abscess following appendicitis may extend high up towards the liver. In these difficult instances the history of the case, as well as the study of the sequence in which the phenomena appeared, becomes of the greatest value.

A *distention of the cæcum* may be mistaken for chronic appendicitis. It gives rise to fulness in the right iliac fossa, and to pain, but, unless associated with inflammation, not to tenderness or to fever; purgatives, too, clear out the fæces which accumulate from want of power of the bowel to propel them, and the dulness on percussion vanishes after the free evacuations. Another element of distinction is furnished by the circumstance that those who suffer from atony of this portion of the alimentary tube labor under it for a long time; they are generally highly nervous persons, with impaired digestion, whose bowels are habitually constipated, and who complain of attacks of spasmodic pain and fulness in the iliac region. Yet, although there is fulness, there is no dulness on percussion, and no hard swelling is detected, unless the cæcum be loaded with fæces. On the contrary, the cæcum and ascending colon generally show, by the excessive tympanitic resonance when they are percussed, that they are distended with flatus.

In that rare disease, *cancer of the cæcum*, there is a fixed, firm

swelling; but it is of very gradual growth, and the disorder generally produces a stricture of the bowel and is associated with malignant disease in other parts of the body.

Other affections than those of the bowels may give rise to signs supposed to indicate appendicitis. It does not at first sight seem likely that this would be the case with *pneumonia*. Yet the mistake has been committed. Pain is sometimes referred to the right groin in pneumonia, and there is soreness there, connected probably with the efforts at coughing and the disordered breathing. Nay, I have known poultices applied to the right iliac fossa to relieve the inflammation which really was in the chest. An examination of this part of the body will of course at once explain the true character of the symptoms.

### Disorders attended with Constipation, and of which it is a Prominent Symptom.

An inactive state of the bowels is often but a concomitant of some disorder which presents phenomena much more striking than the imperfect voidance or the prolonged retention of the fæces. But there are cases in which the constipation is the most important symptom, and in which it furnishes by far the most decisive proof of a serious morbid condition of the intestine. Now, these cases are either those in which the constipation arises suddenly, or at any rate becomes suddenly aggravated, is attended with severe symptoms, and is often insuperable; or those in which it is an habitual state and not associated with any signs of urgent distress.

**Intestinal Obstruction.**—Intestinal obstruction, when coming on suddenly, manifests itself generally in the following manner. A person, previously in good health, or perhaps of costive habit, notices that his bowels have not been moved for several days, and that he has an uneasy feeling in the abdomen in consequence. He takes the purgative he is wont to employ, but without the usual effect. Something more active is tried, and still the bowels remain obstinately bound. Colicky pains have in the mean time made their appearance. He becomes alarmed, and sends for his physician, who sees that there is indeed cause for alarm. The abdomen is found to be distended, but not painful, or only slightly painful, on pressure. But through its parietes



may be noticed the violent, rolling motion of the irritated intestine. Vomiting sets in,—first, of the substances contained in the stomach or of a bilious fluid, and, as the case progresses, of stercoraceous matter. In this way, unless nature or art comes to the rescue, the disease continues; and signs of inflammation of the bowels, and with them fever, appear as preludes to the fatal termination. Sometimes, however, the patient becomes gradually exhausted; there are no tenderness and fever, but a cool skin, a quick, small pulse, a countenance ghastly and panic-stricken. Severe paroxysms of pain, alternating with intervals of ease, may occur to the last moment. But, in spite of the utter prostration, the mind generally retains its clearness. Should recovery take place, large quantities of faecal matter are discharged, and the symptoms of the impediment speedily disappear.

These phenomena are too striking to permit of errors in diagnosis. Yet errors are of frequent occurrence, because the history of the attack and the sequence of the symptoms are not taken into account. Many a person laboring under *peritonitis* has been violently purged to remove the stubborn constipation believed to be due to a mechanical hinderance in the bowels; and, on the other hand, many a case of intestinal obstruction has been treated solely with reference to the inflammation that may attend it, and without regard to the source of the inflammation. Yet it is not ordinarily difficult to distinguish which is cause and which effect. A case that begins with colicky pains and obstinate constipation, in which at first, in spite of the pain, there is little or no tenderness; in which the thermometer does not indicate materially-raised temperature; in which vomiting and tympany soon occur; in which fulness on palpation and dulness on percussion may be detected at or above the point of stoppage; and in which faecal matter is ejected by the mouth after a stoppage of the bowels of a few days' duration,—is not primarily, whatever may be the ultimate complications, enteritis or peritonitis. A case presenting almost from the onset fever and great and extended tenderness; in which vomiting of faecal matter, if it happen at all, does not happen until late; in which diarrhoea is sometimes found to supersede the enduring constipation,—is inflammation of the peritoneum, but not a mechanical obstruction. Only in very rare instances, and especially when the bowel is invaginated, is the malady so

quickly succeeded by inflammation as seemingly to make its appearance with the signs of peritonitis. On the other hand, perforative peritonitis, with its signs of collapse, shows a much stronger likeness to acute obstruction of the bowel than ordinary peritonitis does.

The symptoms dwelt upon as pointing to an intestinal obstruction bear a close resemblance to those of *external strangulated hernia*. In truth, they not only resemble but are identical with those of this affection. Hence in every case of obstinate constipation each point which may be the seat of a hernia must be explored by the eye and the hand. No motives of false delicacy, no reluctance, should prevent the physician from insisting on a search, the neglect of which may cost a life.

It would be foreign to the object of this work to discuss the external signs by which a strangulation of the intestine at a hernial opening manifests itself. It need only be mentioned that it is at the groin, at the umbilicus, at the side of the anus, or through the ischiatic notch that the gut descends and forms a tumor, and that these are, therefore, the regions to be scrutinized. But there is one part of the subject, of importance alike to the physician and to the surgeon, which cannot be passed by without a few words, since it may be a cause of much perplexity,—namely, the possibility of intestinal obstruction taking place in a person laboring under an irreducible hernia and simulating strangulation without any strangulation having occurred. Of this the following case furnishes an example.

A number of years since I was requested by a physician to see with him a woman, the mother of thirteen children, who had been for days laboring under obstinate constipation. Large doses of mercurials, croton oil, and turpentine enemata had failed to procure a passage, and the patient was becoming much frightened. Nor was her situation free from danger. She had considerable pain in the abdomen; she had been vomiting stercoraceous matter profusely; the rolling of the intestines could be plainly perceived. On her right side was a small irreducible femoral hernia, which, on inquiry, was found to have existed for many years. It was not painful on pressure, nor was the skin covering it discolored; neither did the mass itself communicate an impulse during the act of coughing. Here were signs of a serious impediment to the

onward passage of the intestinal contents, as the fæcal vomiting and the rolling of the intestines showed plainly. But was it due to strangulation at the hernial opening? Was it an internal intestinal obstruction?

An accurate examination of the abdomen did not throw much light on these questions. The belly was moderately tympanitic, and not painful to the touch, except when the pressure was considerable. The rolling of the intestines was perhaps more obvious on the left side; but nowhere could a tumor be felt. Taking all the circumstances of the case into account,—the fact that the patient was of costive habit; that she was subject to attacks of colic and of obstinate constipation; that there was nothing to prove that the hernia had recently increased, or was in any way inflamed,—the conclusion arrived at was that the case was not one of hernial strangulation, but of internal intestinal obstruction. Copious warm-water injections were thrown into the colon through a flexible tube; her abdomen was rubbed with mercurial ointment. But all in vain: she continued vomiting fæcal matter.

Her situation now appeared desperate. She had not had a passage for six days; she was steadily sinking. Knowing that sometimes the gut may be strangulated at a hernial opening without much pain or tenderness, the counsel of an eminent surgeon was sought, to aid in determining whether this was not the cause of the impediment. He thought it probable that it was, and proposed an operation, to which consent was reluctantly given. The patient was etherized, and the hernial section performed; but no constriction was found. The wound was closed, and large doses of opium were administered, so as to mitigate, as far as practicable, the torture of the only termination to the case which seemed possible. On the day after the operation, the intestines had ceased to roll; there was no vomiting. But stercoraceous vomiting reappeared two days afterward, and the rolling of the intestines was occasionally, although faintly, perceptible.

The patient's exhaustion was now extreme; her pulse was very quick and small; her skin cold, of a dirty look; the odor of the breath and of the whole body offensive; and the eyes sunken and surrounded by a broad leaden ring. There was slight pain on pressure between the umbilicus and the sigmoid flexure. The vomiting had ceased, or occurred only very occasionally. Al-

though there was little hope, we had, as soon as admissible after the operation, recommenced rubbing mercurial ointment over the abdomen, and giving injections in the manner before described. This was continued until, to our great gratification, one morning, after a tube had been passed a distance of several feet into the colon, the patient had a copious discharge of tarry faecal matter from her bowels,—seventeen days after the symptoms of complete intestinal obstruction had declared themselves by the occurrence of stercoraceous vomiting.

This case is instructive in more than one respect. It teaches that recovery may take place most unexpectedly after many days; and, in a diagnostic point of view, it illustrates a difficulty which any physician may have to encounter in attending a patient the subject of a long-standing hernia.

Supposing that the symptoms are altogether owing to an obstacle at some portion of the intestine within the abdomen: can we determine the exact position of the impediment, and its nature? We know how varied are the conditions which lead to sudden and invincible constipation. We know that strangulation from bands and adhesions, or gaps in the omentum, or the pedicle of an ovarian tumor; that intussusception; that twists and knots; that strictures and tumors; that abnormal contents, such as foreign bodies, impacted faeces, gall-stones, worms, concretions of drugs, as of bismuth, may all occasion intestinal obstruction. We also know that in certain cases the obstruction is from spasmodic contraction of the intestine,\* or paralysis of the bowel; in others is congenital.† Can we distinguish these different lesions at the bedside? In certain cases we can,—we can determine exactly both the position and the character of the lesion; in others there is no clue to an accurate discernment of either. From the method of the introduction of the whole hand into the rectum much has been expected; but experience has not confirmed these expectations. Obstruction of the bowel may clinically present itself as an *acute* or as a *chronic* malady. The same symptoms occur in both. It is the mode of origin that is different. Nay, the same lesion may occasion in some instances an acute, in others

---

\* Archives Générales, Aug. 1868; Flint, Practice of Medicine.

† Gould, Transact. Clin. Soc. Lond., 1882.



a chronic, affection. Invagination, internal strangulation, volvulus, impaction of a large gall-stone, are generally acute ; strictures, tumors, contractions, and, for the most part, fecal accumulation, lead to chronic obstruction.

Among the forms of intestinal obstruction, *intussusception* or *invagination* is very frequent and at the same time the least difficult of recognition during life. Part of the bowel becomes inverted, slipping into the cavity of the adjoining upper or lower portion. Inflammation is soon set up, produces infiltration of the tissues, and often leads to adhesions between the opposed serous surfaces, and to effusions of blood and mucus into the canal. The inflammation may spread rapidly over the serous membrane, and the patient may die from general peritonitis. But sometimes in this inflammation that is lighted up at the seat of the ileus lies safety. It may give rise to adhesive inflammation of the opposed serous coats of intestine, and ultimately to a sloughing off of the invaginated part and its discharge into the bowel, while the annular mass of adhesive lymph surrounding the seat of ulceration maintains the continuity of the intestinal canal, and thus the inflammation may pave the way to a favorable issue by restoring the calibre of the tube,—sufficiently, at any rate, to permit of the transit of its contents.

These pathological peculiarities develop special symptoms which frequently enable us to determine the nature of the obstruction. When the intussusception takes place rapidly, a sudden local pain is produced, recurring in paroxysms, and likely to be referred to the seat of the disturbance. The pain is quickly followed by vomiting, by constipation, by tympany, and by peritonitis. But the constipation is not so absolute as in other cases of intestinal impediment. Not unusually, in fact, owing to the invaginated bowel remaining open, the liquid contents of the intestine pass through the intussuscepted part and produce a deceptive diarrhœa; yet oftener will occur tenesmus, and discharges of the bloody mucus and serum which have accumulated in the intestine. Both of the latter signs are eminently diagnostic of the lesion. Still more so is feeling the end of the invaginated bowel by an exploration of the rectum, or finding the loosened segment in the stools. But it is only in cases in which the lower portion of the canal is affected, or which have been sufficiently protracted to allow of the curative

efforts of nature being accomplished, that signs so pathognomonic are met with.

The casting off of the sloughed portion of the intestine is attended with hemorrhage. Whether this be the only cause of the hemorrhage or not, it is undoubted that purging, or sometimes vomiting, of blood, is among the differential signs of intussusception. A sign more valuable, because so much more common, is the presence of a tumor, frequently of cylindrical shape. Its seat varies with the seat of the lesion; and as the most common invaginations are those of the ileum and cæcum into the colon, or those at the inferior portion of the ileum, it is at the lower part of the belly, and generally passing in direction from left to right, and in the right iliac fossa, that the swelling is detected, and often it may be felt through the rectum. The malady is generally due to irregular peristalsis. It is often preceded by diarrhœa; there is rarely much tympany. Sometimes it is caused by tumors of the intestine, particularly by lipoma.\*

The majority of cases of invagination happen in children under ten years of age. The course the affection pursues is rapid; the patient dies generally in less than a week after the occurrence of the accident, utterly prostrated. Yet the records of medicine furnish us with instances in which life has been prolonged for months. The cases which get well recover either gradually after the invaginated bowel has been discharged, or, in very rare instances, more quickly by the inverted bowel righting itself.

Acute obstruction from *internal strangulation* is almost invariably seated in the small intestine. Its most characteristic feature is furnished by the history of an old peritonitis, an operation on the abdomen, or an appendicitis. There is rarely fever; the obstruction soon becomes complete; nausea and vomiting set in early; fecal vomiting usually begins from the third to the fifth day. It is the exception to find a tumor; tympany is often marked, but no flatus escapes by the bowel. Of further significance in the diagnosis of internal constriction are the signs dwelt on by Fagge:† the sudden onset of the illness; the occurrence of collapse at its beginning; the frequency with which it is found in

---

\* Clos, De l'Invagination intestinale, etc., Paris, 1883.

† Guy's Hosp. Rep., 3d Series, vol. xiv.

young adults; the severity of the pain, which is generally referred to the umbilicus; the absence of external or of discoverable obturator hernia; the absence of precursory symptoms and of visible peristole,—such as happen in stricture and contractions,—of tumor, of hemorrhage, and of dysenteric symptoms,—as seen in intussusception,—and of that extreme intensity and rapidity of development which characterize the more acute forms of volvulus. Obstruction by a band connected with a diverticulum scarcely ever occurs except in males under twenty years of age.\*

Acute obstruction from *volvulus* or twist begins with severe abdominal pain, soon becomes associated with nausea and vomiting and extreme distention of the abdomen, and rarely presents a tumor or visible intestinal coils or elevation of temperature.† It most frequently affects the sigmoid flexure. But it cannot with any accuracy be separated from internal strangulation. It is most commonly met with in men between thirty and forty years of age.

As regards *other forms of intestinal obstruction*, they are, with our present knowledge, undistinguishable from one another. However desirable it might be on therapeutic grounds to be able to diagnose with certainty a blocking up of the intestine by hardened feces, or gall-stones, or enteroliths, or its strangulation by bands or by rents in the mesentery; however desirable to know whether, if medical means do not bring relief, laparotomy may be attempted with hope of success, or whether the impediment is not even to be removed by such a mode of succor,—it must be confessed that there are few signs which enable us positively to decide on the nature of the obstacle.

Yet there are sometimes circumstances in the case which help to a correct decision. For example, if the complaint occur in one who has suffered from the passage of *gall-stones*, especially in a fat, elderly woman,‡ it is likely that a large concretion of this kind has been arrested in its passage through the small intestine. Should the disorder be encountered in a person over forty years of

---

\* Fagge, *Practice of Medicine*, vol. ii.

† Fitz, *Acute Intestinal Obstruction*, *Transact. of Congress of American Physicians and Surgeons*, vol. i., 1889

‡ Fagge, *Practice of Medicine*, vol. ii. p. 210.

age, who at all times has difficulty in voiding the contents of the tube ; whose fæces present peculiarities in shape and size, and are sometimes mixed with blood ; whose health has been gradually breaking ; whose abdomen is much distended and yields a ringing tympanitic resonance on percussion,—should such a person have an attack of constipation unusually protracted, attended with enormous distention of the bowel, and in which the remedies that hitherto barely procured a passage now fail utterly, it would not require much sagacity to discern that a *stricture of the intestine*, probably of a cancerous kind, is the source of the cruel and irremediable suffering. If, in addition to the symptoms enumerated, a bougie passed into the rectum meet in its course with a decided obstacle, an error in diagnosis is hardly possible. When, however, the stricture is not accessible to instrumental examination, although we can commonly recognize its presence, we cannot fix its site. The distention above the narrowed part is often so extreme as to lead to displacement of the colon and to an almost uniform swelling of the whole abdomen. For instance, in a case reported by Albert H. Smith, the enormously-dilated colon had broken loose from its attachments and concealed the rest of the viscera. It was in several places eighteen, in none less than fifteen, inches in circumference ; and fully two gallons of liquid fæces were found in the bowels.\*

In the majority of cases the stricture is either in the rectum or in the sigmoid flexure. Other causes of stricture besides cancer, though less common ones, are cicatrization of extensive syphilitic, tuberculous, or dysenteric ulcers. A *contraction* in the small intestine similar to the true stricture of the large bowel is seen chiefly as the result of chronic peritonitis binding down the bowel, and may lead, like a stricture, to chronic obstruction.† Faecal accumulations also produce chronic obstruction. We distinguish this form chiefly by its occurrence in women, especially hysterical women, or in hypochondriacs, by the tenderness over the faecal tumors, the gradually-increasing constipation, the late occurrence of pain and of vomiting, and the extreme foulness of the breath.‡

---

\* Proc. of Pathol. Soc. of Phila., Dec. 1858, vol. i.

† Fagge, Guy's Hosp. Rep., 3d Series, vol. xiv.

‡ Treves, Lancet, Oct. 29, 1887.



With reference to the *frequency* of the different forms of intestinal obstruction, the elaborate studies of Fitz \* give us valuable information. Strangulation is the most frequent cause of acute obstruction, occurring in fully one-third of the cases ; a number of cases are noted to follow operations upon the pelvic organs in women, though the disease is very much more common in men than in women. Intussusception comes next in frequency, and is especially seen among children and young adults. Volvulus or twist is mostly encountered in men, and in half the cases is in the sigmoid flexure. Strictures and tumors, that are such common causes of chronic obstruction, very rarely lead to acute obstruction.

In any kind of obstruction the *location of the lesion* is difficult to determine. There are, however, a few circumstances which may aid us in arriving at such a determination : one is the fact pointed out by Barlow,† that the higher up the obstruction is in the canal, the nearer therefore to the stomach, the smaller is the quantity of urine passed ; another is the early occurrence of the vomiting and the want of stercoraceous character of the matters ejected,—both of which render it likely that the impediment is in the small intestine and remote from the cæcum. Still another is the speedy presence and the greater severity of hiccough when the mischief is in the small intestine. Sometimes the patient is himself aware of the exact seat of the cause of his suffering ; he notices that the injecting tube or the enemata seem to reach a certain point and go no farther ; so, also, with the rumbling of the wind. Again, these borborygmi are especially apt to occur in obstructions of the large intestine, and, if joined to tenesmus, are signs of some importance. Indican is found in the urine in greatly-increased quantities in stoppages of the small intestine. We may also be able to come to some conclusion about the seat of the lesion by inflating the large intestine, or by finding out how many quarts of warm water we can inject into it.

The position of the pain, too, may furnish a clue to the position of the impediment. If this be in the small intestine, the pain is

---

\* Transact. of Congress of Amer. Phys. and Surg., vol. i., 1889.

† Guy's Hosp. Rep., 2d Series, vol. ii. Brinton accepts this statement only in so far as the amount of vomiting, which is apt to be greatest when the obstruction is high up, influences the amount of urine passed.

apt to be chiefly, if not entirely, in the neighborhood of the umbilicus. Another circumstance on which some stress may be laid is the distention of the intestine above the point of intussusception. Indeed, this distention may occasion a visible fulness, sounding extremely tympanitic on percussion; at times, too, a slight dulness is found, attended with some resistance at or immediately above the seat of the obstruction. But with reference to the swelling and the tympanitic dilatation of the bowel there are—as William Brinton\* sets forth—several reasons which render these signs uncertain guides. The distended intestine may not be capable of being traced by the eye or by percussion, owing to its occupying a large portion of the abdominal cavity. Moreover, a stoppage at the descending part of the large intestine, for instance at the sigmoid flexure, may lead to most palpable distention of the cæcum, and to pain in that region; while pain and swelling are also observed in the same locality in obstructions which affect the small intestine. Thus there are several modifying circumstances which prevent too much importance being attached to any of the signs mentioned as proofs of the seat of the obstacle; for, with the exception of a tumor dull on percussion and resistant to the touch, there is nothing absolutely indicative of the lesion being at a particular spot. It is hardly necessary to say that a swelling of this kind cannot always be found.

In referring to the usual seat of pain and swelling in the right iliac fossa, and to the difficulties which on this account beset the recognition of the precise site of the hinderance, one source of error deserving of special notice was not mentioned. The pain and the fulness in this region may be caused by an *appendicitis*. Moreover, affections of this part of the alimentary tract, like intestinal occlusion, give rise to constipation which is most obstinate and in some instances incurable. Therefore they in reality enter at times into the category of intestinal obstructions, from the other varieties of which they are undistinguishable save by the history of the case, the localization of the symptoms, and the different sequence of the phenomena. The tumor and the other local signs do not follow the insuperable constipation, but precede it. Stress may be laid upon the occurrence of

---

\* Croonian Lectures, and work on Intestinal Obstruction.

chills, of waves of pain, and of marked local tenderness, as indicative of the presence of pus, or upon the signs of collapse in perforative appendicitis. In *acute hemorrhagic pancreatitis* there may be also the signs of intestinal obstruction, not to be distinguished except perhaps by the history, the extremely rapid course of the disease, and the marked peritonitis.

**Habitual Constipation.**—This is a chronic state, unattended with urgent symptoms of any kind. Still, it is an annoying and very prevalent disorder. The symptoms encountered, independently of the rare and difficult faecal evacuations, are headache, giddiness, sluggishness of the mind, a want of the natural appetite, anæmia, cutaneous eruptions, and, joined as the complaint not unfrequently is to derangement of the stomach and of the biliary secretion, digestive disturbances and a sallow complexion: an altered state of the blood from the absorption of ptomaines may exist. In women there are also often added to the list of evils to which costiveness gives rise, neuralgic pains, palpitation of the heart, cold feet and hands. Infrequent evacuation of the bowels does not always produce such unpleasant consequences. It may, indeed, in individual cases be compatible with perfect health; for what is costiveness in one person may be a natural state in another.\* But when the bowels are acting less frequently than is their wont, the disagreeable symptoms mentioned are apt to arise.

Habitual constipation is produced by various causes. It may be brought about by the peculiar nature of the diet. It may depend upon a deficiency or a faulty composition of the intestinal secretions, or upon disorders of those neighboring glands which pour their secretions into the intestines. It may result from impaired power of the bowel to propel its contents, the consequence either of some mechanical interference with its action, or of nervous influences, or of exposure to the poisonous effects of certain substances, as of lead. To particularize the numerous conditions which furnish illustrations of each of these different causes would serve no useful purpose. A few only need be specially noticed.

We have often to treat constipation in those who are dyspeptic and suffer from piles. In them there is, in all probability, some

---

\* In the American Journal of the Medical Sciences, Oct. 1874, a case is reported in which the constipation lasted eight months and sixteen days.

congestion of the portal system, and not unfrequently a constant derangement of the flow of blood through the liver. The normal secretion of intestinal juices is interfered with, healthy bile is not supplied, and thus costiveness results. A similar congestion of the intestinal mucous membrane has its share in producing the constipation which is encountered in diseases of the heart. Sometimes, however, enough healthy fluid is poured out within the intestine; yet there is a deficiency, because the inclination to go to stool is resisted, and the liquid that has been mixed with the matter to be voided is reabsorbed. In women who neglect the calls of nature because circumstances prevent their being obeyed at the proper time, this is a common cause of constipation.

The influence of the nervous system on the alimentary tube is shown by the confined state of the bowels which attends excessive intellectual exertion and violent emotions. And when these states are protracted, they lead to a permanent and annoying debility of the intestine. The colon especially becomes torpid in its action, and all the evil results of constipation show themselves in their most marked degree. Not that an atony of the bowel is always due to psychical agencies. Any disorder which induces loss of power in the muscular fibres may give rise to it. We find it where the blood is watery and deficient in red corpuscles, and in those who lead, as far as bodily exertion is concerned, a sluggish life. In some cases—fortunately rare—the weak intestine distends greatly, and becoming, as above explained, unable to propel the accumulated fæces, insuperable constipation occurs. The same complete paralysis of the tube, attended with the same unfortunate consequences, may be brought about by chronic lesions of the brain or spinal cord. Yet the inveterate constipation which is so constant an accompaniment of these states is partly owing to the powerless condition of the abdominal muscles.

Among the different organic changes in the intestine which, by interfering *mechanically* with the peristaltic wave, set up constipation, we find distention of the tube, with atrophy of the muscular fibres; various infiltrations into the walls, producing a narrowing of the calibre, as in carcinoma; and adhesions between the serous coats of the intestines, or between these viscera and the parietes. Of the first, it need only be said that the symptoms are due to the same paralyzed condition of the intestine, whether complete or in-



complete, which has been already considered. The second group embraces those infiltrations which result from inflammations, and new growths of different kinds which lead to strictures.

The former of these are recognized, as far as they can be with certainty, by the history of the case. The latter present peculiarities in the form and size of the fæces, distention of the bowels above the seat of the narrowing, vomiting, attacks of colic, gradual wasting and exhaustion; besides which, extreme costiveness, deepening gradually into invincible constipation, furnishes a key to the grievous nature of the affection.

When the constipation arises as the result of peritoneal adhesions, there are sometimes signs in the case—such as tenderness at a particular spot from still existing inflammation, or partial distention or retraction of the abdomen—which point out its nature. In the absence of these, the history is our only guide, except in those instances in which, as Bright\* first informed us, a peculiar sensation is communicated to the touch, varying between the crepitation produced by emphysema and the feel derived from bending new leather in the hand.

From long-standing constipation *stercoral ulcers* may arise. The sacculi of the colon are filled with little hard fæcal balls, which irritate the mucous membrane and produce ulceration. Mucus, or muco pus, with stains of blood, is occasionally discharged with the small scybala, and at times there is diarrhœa.

### Disorders in which Morbid Discharges from the Bowels occur.

Matters very unlike the healthy alvine evacuations are often voided from the intestinal canal: loose watery stools, large quantities of mucus, pus, or blood, may be discharged. The disorders which occasion these discharges may be here described.

**Diarrhœa.**—Like constipation, diarrhœa will be merely treated of as we meet with it constituting the entire ailment, or at all events its most prominent symptom. There are several varieties of diarrhœa. Difference in time gives rise to marked varieties,

---

\* Cases illustrative of the Diagnosis of Adhesions and other Morbid Changes of the Peritoneum, Med.-Chir. Transact., vol. xix.

—to an acute and to a chronic form ; and of both it has been pointed out already how often the lesion is an intestinal catarrh.

*Acute Diarrhœa.*—Acute diarrhœa proceeds from more than one cause : it may be excited by the irritating character of the food taken, or by impure water ; it may be brought about by the morbid nature of the secretions poured into the intestines ; it may be owing to atmospheric influences,—to heat, to moisture, to contaminated air ; it may be caused by chilling of the surface of the body, or by irritant poisons, retained fæces, or worms. It may be occasioned by pyæmia and septicæmia, by reflex irritation, as in dentition, or by mental emotions, and especially by fear. Sometimes it occurs in an epidemic form due to some unknown miasm. Its symptoms are thirst ; abdominal uneasiness ; griping pain in the bowel ; pallor ; slight debility ; and frequent fluid alvine evacuations, which may finally become almost colorless.

In the diarrhœa caused by a debauch or by indigestible food, nausea and a furred tongue are added to the list of symptoms mentioned. This kind of diarrhœa is generally of short duration. It is an effort of nature to get rid of obnoxious matter ; and when this is effected, the looseness of the bowels ceases.

The variety of diarrhœa under consideration sometimes goes hand in hand with a disturbance of the biliary functions, and the stools discharged are fetid, and present the appearance generally described as bilious. This “bilious diarrhœa,” too, is not uncommon in persons whose livers are habitually sluggish. It is also frequently encountered during the hot months of summer and early in the autumn, and has a tendency to run on.

There are cases of diarrhœa attended with pain, considerable soreness to the touch, and, what is not ordinarily met with in diarrhœa, some febrile disturbance. These kinds of acute diarrhœa, or rather of acute intestinal catarrh or of muco-enteritis with diarrhœa as a symptom, are often the result of irritant poisoning, or are common as the result of the influence of cold, or of acrid drinks and unripe fruit. They are also observed as secondary disorders in typhoid fever and in the exanthemata.

*Chronic Diarrhœa.*—In chronic diarrhœa the lesions encountered are much more marked than they ever are in the acute form. The mucous membrane is tumid and discolored ; its follicles are not unfrequently ulcerated. Chronic looseness of the bowels

originates in a diarrhœa which is permitted to continue, either from neglect or because the patient remains for a long time exposed to the original cause. But the disorder, no matter under what circumstances it originated, is apt to prove rebellious, and to end by breaking down the constitution. When of long standing, the patient becomes gradually weaker and weaker, and more and more emaciated. The abdomen is sunken; the complexion is pale; the eyes are surrounded by a dark ring. The character of the discharges is various. They are often dark-colored and very offensive. Sometimes the looseness of the bowels alternates with an opposite condition; but the irritability of the intestines never intermits.

This morbid excitability of the intestinal tube is especially brought about in persons of nervous temperament and of dissipated habits. The abuse of purgatives, too, induces it, and in consequence chronic diarrhœa is not an uncommon result of the cathartic pills which patrons of quack medicines habitually swallow.

But perhaps the most persistent irritability of the intestines is found in the diarrhœa to which *soldiers* are so liable, and which is apt to pass, no matter what its beginning, into the chronic form of the disease. This complaint, which follows impure water, defective diet, exposure, malaria, and scurvy,\* which is generally associated with a morbid state of the large intestine as well as of the small, and which combines therefore some of the features of chronic dysentery with those of chronic diarrhœa, is one that often clings to its victim through life: many a soldier, in truth, escapes the bullet and the sword, only to die of the intestinal affection long after his return to his home.

Then, the affection is very often witnessed as a complication of other disorders. Two-thirds of the fever patients received in the hospitals of Constantinople during a long period of the Crimean War were affected with diarrhœa or with dysentery. Diarrhœa was so very general that nearly all disorders were preceded by acute diarrhœa and terminated in chronic diarrhœa.† It was

---

\* Woodward, *Outlines of the Chief Camp Diseases*, p. 253; see also the elaborate analysis of the alvine fluxes in vol. ii. of the splendid "*Medical and Surgical History of the War of the Rebellion*," Washington, 1879.

† Baudens, *La Guerre de Crimée*.

much the same in this country during General McClellan's peninsular campaign.

But chronic diarrhœa, as the practitioner of medicine commonly sees it, is often attendant on general constitutional affections, or on abdominal diseases that have led to a secondary disorder of the secretions, or even of the coats of the intestine. Thus, we find chronic looseness of the bowels in scurvy, in pyæmia, in Bright's disease, in scrofula of the mesenteric glands, and in tuberculosis. In the last of these complaints the diarrhœa may be occasioned by changes in the secretions of the intestinal glands; but it is not seldom dependent upon a true *tubercular disease of the intestines*, which, like the disease of the lung, leads to softening and ulceration. The discharges are generally copious and very offensive, and show traces of blood. The diarrhœa is continuous and intractable; the abdomen is retracted, and presents spots very tender to the touch. There are marked fever and emaciation, and there may be severe intestinal hemorrhage. Yet, after all, only the signs of tubercle elsewhere furnish any positive indications by which the true nature of the wasting malady can be discerned. Indeed, it may happen that the reverse of diarrhœa occurs, for acute primary miliary tuberculosis may simulate an acute intestinal obstruction.\* In all cases of suspected tubercular diarrhœa the stools should be examined for tubercle-bacilli, and these will be found very generally.

Tubercular ulceration is the most prominent type of ulcerative enteritis. But *ulceration* of the bowel is also met with under other circumstances. We find it in the diarrhœas of children; it occurs then as *follicular ulceration*. Ulceration is also occasionally observed from *cancer*, or as a *solitary ulcer* leading to perforation. The seat of the latter is generally the cæcum or colon. *Albuminuric ulceration*, the careful analysis of Dickinson † shows, is almost invariably associated with contracted kidney. *Simple ulcerative colitis* is usually met with in middle-aged persons. It lasts generally about two months,‡ and is ushered in with abdominal pain, which remains a symptom. There is diarrhœa with very fluid movements, but there are no dysenteric

---

\* Thoman, Allg. Wien. Med. Zeit., 1887.

† Med -Chirurg. Trans., vol. lxxvii., 1894.

‡ Hale White, Guy's Hosp. Reports, 3d Series, vol. xxx.



stools; blood in the discharges is common. The diarrhœa may alternate with attacks of constipation; often there is vomiting. The disease may lead to perforation. *Unhealed typhoid ulcers* form another variety of ulceration of the bowels.

In the diagnosis of all forms of intestinal ulceration, we must lay stress on the diarrhœa, on the character of the discharges, on the pain, and on the occurrence of hemorrhage from the bowels. In the discharges, mucus and pus and shreds of tissue are valuable signs. In follicular ulceration little sago-like masses of mucus are met with, but they are not pathognomonic. The stools may be very frequent; this is especially the case if the ulcer is in the lower part of the colon. Abdominal pain may or may not be associated with tenderness. Pain, as in other forms of colitis, is often referred to the præcordial region. With reference to the frequency of this, Potain \* tells us that, of one hundred persons complaining of heart-disease, about seventy have an affection of the colon.

In the *chronic diarrhœa of strumous children* there is sometimes a scrofulous infiltration into the intestinal walls, sometimes marked scrofulous enlargement of the mesenteric glands, sometimes both, but in some cases neither. Improper nourishment, however, may be here, as in any other form of the diarrhœa of childhood, the exciting cause of the continued purging.

At times chronic diarrhœa assumes an *intermittent type*, and its malarial nature is clearly proved by the readiness with which the disorder yields to quinine.† In this respect malarial diarrhœa differs from cases of diarrhœa we sometimes encounter, in which the pain and discharges come on at an early hour of the day and cease toward evening and during the night.

Another form of looseness of the bowels is the *membranous*. Here the discharges show shreds of membrane, either in connection with the loose stools, or sometimes in such quantities that the whole mass voided seems to consist of them. Gripping pains and tenderness usually precede this kind of diarrhœa, which may happen in attacks of a subacute form, or as a persistent and very

---

\* L'Union Médicale, Nov. 1894.

† See contribution by Sanford B. Hunt on *Diarrhœa*, in Medical Memoirs of U. S. Sanitary Commission, p. 306.

obstinate disorder : the former variety is the more common. The faecal discharges are loose, but occasionally there is constipation. The disease is often associated with peculiar hysterical symptoms or occurs in neurasthenics. The so-called membranes, in this membranous enteritis, contain a large amount of mucus, as I have elsewhere described.\*

**Dysentery.**—Frequent and painful passages of mucus mixed with blood, accompanied by straining and bearing down, are the characteristic symptoms of dysentery. In the acute form we find thirst, restlessness, and heat of skin superadded ; and sometimes, especially when the disease prevails epidemically, those symptoms of prostration which are commonly designated as typhoid.

*Acute Dysentery.*—The acute disorder is at times ushered in by a chill ; at times it is preceded by diarrhoea. The fever which attends it is not generally intense. It is the exception to find it exceed  $103^{\circ}$ , and in light cases the temperature is only slightly raised ; the pulse is not tense. More or less pain is always present. It has its seat mostly at some part of the colon, and this is tender on pressure. It is intermitting and shifting, and is often accompanied by a feeling of weight near the anus, which causes a continual desire to go to stool. Yet no relief follows the frequent attempts ; the violent straining only adds to the discomfort.

The matters voided are small in quantity. They consist of blood mixed with mucus ; but they are composed not simply of mucus, but also of leucocytes, granules, and large quantities of cast-off epithelium, with many swollen round or ovoid epithelial cells. The stools are in some cases highly offensive, and resemble the washings of meat ; in others they are like jelly, or greenish in color. They do not contain faeces, or only here and there small, firm lumps of faecal matter. When the dysenteric inflammation subsides, the bowels are unloaded of their contents ; in consequence, the passage of quantities of small, hard masses of faeces is generally a sign that the acute malady is inclining to a favorable termination. Sometimes the stools are very dark and slimy and have a putrid odor, and here and there pieces of sloughed-off tissue can be detected. This kind of stool marks

---

\* American Journal of the Medical Sciences, Oct. 1871.

the diphtheritic or gangrenous variety of the malady, though it is not constant even in this.

How long it will take for the disorder to run its course, or whether the acute disease will pass into chronic dysentery, cannot be foretold. Generally this is not its termination; it very often ends, within a week from its beginning, in recovery. But severe cases occur which are of much shorter duration, in which the symptoms hasten on to complete prostration and death takes place early in the malady. In these frightful cases—most frequently encountered in epidemics—collapse may happen with almost the same rapidity as it does in malignant cholera.

Dysentery is essentially a disease of hot climates. It is very common in this country in summer and in autumn. Eating green fruits, exposure to a chilly night after a hot day, and sleeping on damp ground, are prolific exciting causes. It is occasionally found in combination with malarial fevers, adding greatly to their danger, or with scurvy. It also occurs from drinking water full of impure substances or micro-organisms. It may be seen in a sporadic or in an epidemic form. It is very common in armies and in jails. The immediate cause of most of the symptoms is inflammation of the large intestine, and especially of the descending colon. Yet in many cases of dysentery we see phenomena manifested which are clearly not to be accounted for solely by the local morbid appearances detected after death, and which show that dysentery mostly belongs to the infectious maladies. In truth, inflammation of the colon may give rise to the symptoms of acute diarrhœa; for it is a great mistake to suppose that the cause of diarrhœa is to be sought only in some abnormal change in the small intestines. Thus, colitis is not always dysentery; and dysentery is often more than mere colitis.

But, whatever be the ultimate cause or the form of dysentery, we find that it presents peculiarities which render it easy of recognition at the bedside. Yet we must take good care to ascertain that the supposed characteristic tenesmus and bloody discharges are not really owing to *piles*, or to morbid, especially cancerous, growths in the rectum, or to its ordinary limited inflammation. In the latter case, or *proctitis*, there is much pain when the hardened fæces are discharged, the rectum is forced down during the efforts, the sphincter contracts spasmodically.

Strangury and hemorrhoids are not uncommon symptoms ; and, as the consequence of the inflammation extending to the parts around the anus, an abscess may follow. Rectal pain often extends to the thighs.

Dysentery is not apt to be confounded with *diarrhœa*, for symptoms exist in the former which do not belong to the latter. Diarrhœa differs essentially from dysentery by the liquid fecal evacuations, and by the fact that neither tenesmus, nor bloody stools, nor discharges of mucus occur. Yet in practice we meet with cases which begin with diarrhœa and terminate in dysentery, or begin with dysenteric symptoms and terminate in diarrhœa, and in which it becomes, therefore, puzzling to say which disorder we are dealing with.

There are some clinical varieties of dysentery which it is important to separate. The ordinary form seen in temperate climates to follow errors in diet or exposure is the catarrhal form. In tropical climates, where dysentery is very common and is met with frequently as an epidemic, we find mostly a kind that is characterized by the presence of the *amœba coli*, or *amœba dysentericæ*, as Councilman and Laffeur,\* to whom most of our knowledge on the subject is due, call the micro-organism. *Amœbic dysentery* does not, as a rule, run so rapid a course as ordinary catarrhal dysentery, and local tissue degenerations in the liver, or single or, it may be, multiple abscesses of the liver, are common attendants. The abscesses, like the discharges from the bowels, contain amœbæ. The evacuations, as the disease progresses, lose their dysenteric characteristics, except the mucus, and become very liquid ; the tenesmus disappears. The amœbæ are most active in alkaline stools. The diarrhœa has marked exacerbations and remissions, and is attended by striking anæmia. The fever is very moderate. In some instances hemorrhage from the bowels, in others peritonitis, happens. It is not unusual in protracted cases for the urine to become albuminous and to contain casts.

In tropical climates, too, though also seen elsewhere in persons who have low forms of pneumonia, or who have become cachectic from scurvy, from Bright's disease, or from long-standing disease

---

\* Johns Hopkins Hospital Reports, vol. ii.



of the heart, a form of dysentery attended by extensive exudation and sloughing of the membranes is met with. This *diphtheritic dysentery*, as it is called, has generally high fever, much abdominal pain, great prostration, and delirium. The discharges are very frequent; the blood gradually disappears from them. Vomiting, especially at the onset, is common. In the progress of the case, which is generally to a fatal issue, the temperature becomes irregular, and hiccough is not uncommon.

*Chronic Dysentery.*—In chronic dysentery this mingling of the two complaints is especially apt to happen. We rarely see chronic dysentery without chronic diarrhœa. At all events, we seldom find instances of the former in which the tenesmus and the discharge of blood and mucus mixed with pus are not accompanied by frequent loose alvine evacuations, by griping, by the same gradual wasting and the same irritability of the bowels as are encountered in chronic diarrhœa; nay, the symptoms of the latter, and the difficulty of determining the presence of pus when mixed with fluid fæces, may so obscure the true nature of the malady that what has been regarded as chronic diarrhœa turns out, at the autopsy, to be chronic dysentery. The mucous membrane of the colon is found to be extensively inflamed; its texture altered and irregularly thickened; its surface riddled with ulcers. In such cases the patient goes on steadily losing flesh, has some elevation of temperature; but no pain on pressure or localized distress exists to denote the ravages the disease is making in the alimentary tube. Many die from exhaustion; others, in consequence of abscess of the liver, which chronic as well as acute dysentery may induce.

**Intestinal Hemorrhage, or Melæna.**—This is commonly the result of a mechanical hinderance to the flow of blood through the liver, as in cirrhosis, or of disease of the heart, or of a depraved state of the blood,—such as exists in typhus fever, in yellow fever, in scurvy, or in purpura. Occasionally the bleeding proceeds from a fungoid growth in the intestine, or from an ulcer in the duodenum or ileum, or from an invagination, or from fecal impaction, or from amyloid degeneration of the mucous membrane of the bowel, or is due to a disease of the spleen, or to the bursting of an aneurism, or follows extensive burns of the abdominal parietes. In very young infants a dis-

charge of blood, both by the mouth and by the rectum, is not unusual.

The blood passed by stool is generally of dark color, like tar. When it is not, we may fairly infer that it flows from the lower part of the intestine and has not had much chance to become admixed with other matters. In all such cases, however, we must make sure, before arriving at any conclusion as to the source of the bleeding, that it does not proceed from *hemorrhoids*. The exact seat of the hemorrhage cannot be determined; nay, blood may be evacuated by the bowel and not be poured out at all from the intestine, but from the stomach. In some instances the blood accumulates in the bowel, and, before the clots moulded to its shape are discharged, death results.\* When the bleeding proceeds from hemorrhoids it is very seldom vicarious.†

In point of diagnosis the first thing to determine is, that what is supposed to be blood is really blood. Very dark bilious stools, or stools blackened by iron, may mislead. If doubt exist, water should be poured on the stool, and, when blood is present, a reddish tinge is imparted to the water; still more accurate is it to examine with the microscope or the spectroscope.

We next have to ascertain the disease with which the intestinal hemorrhage is associated; and this is often a very difficult matter. We must lay the greatest stress on the history of the case, look for the complaints—of which most have been above mentioned—that are apt to give rise to the bleeding, especially investigating for cirrhosis of the liver; searching for intestinal ulcers in connection with typhoid fever, or tuberculosis, or a duodenal affection; or examining for the evidence of scurvy in the gums and skin; or for purpura with its characteristic spots; or for splenic enlargement, the result of chronic malaria, or perhaps combined with bone disease or syphilis and joined to amyloid degeneration of liver, kidneys, and intestinal walls. Embolism of the superior mesenteric artery may also occasion intestinal hemorrhage. But unless we have with the bloody stools marked abdominal pains, peritoneal exudation, and obvious causing ele-

---

\* See observations of Cheyne, Dublin Hospital Reports, vol. i, and of Belcombe, Medical Gazette, vol. iv.

† Lee on the Rectum.

ments of embolism or signs of it elsewhere, the diagnosis is most uncertain.

**Fatty Diarrhœa.**—The occurrence of cases in which large quantities of fat, mixed or pure, are voided by the rectum, is well attested. In some of these cases oil was at the same time passed with the urine; in others the urinary secretion was healthy; some cases ended fatally, others in recovery; some were found to be connected with a disease of the pancreas, others were not; in some the disorder was not of long continuance, while in others it lasted, with intervals, for years.

As a rule, the occurrence of fatty stools is a matter of serious concern. The recognition of the malady is easy. The white, fatty masses, or the oily matter which collects on the discharges, are soluble in ether, and are readily proved to be fat by the microscope; they burn, too, like fat, with a flame. In some instances the bowels are very constipated, and lumps of hard feces are discharged along with the fatty substance. This happened in a marked example of the disorder that came under my observation. The patient, a man twenty-six years of age, passed a considerable amount of fat both by the rectum and with the urine. He suffered much from digestive disturbance, from constipation, and from weakness. He had a good appetite, but a dislike to fats of any kind. In his case there was, as far as the other symptoms and the physical signs indicated, no tumor in the region of the pancreas. The man's condition was much improved by the administration of cinchona and rhubarb; but whether permanently or not I cannot say, as I lost sight of him.

I have also met with instances of fatty diarrhœa associated with saccharine diabetes and with disease of the pancreas. In examining into the subject of fatty stools it must be borne in mind that the clay-colored stools of jaundice, owing to the absence of the emulsifying properties of the bile, contain considerable fat, which may be found in oil-drops or as fine needle-shaped fat-crystals.

### Diseases attended with Vomiting and Purging.

There is a group of diseases in which vomiting and purging are very prominent symptoms. It embraces those disorders in

which the intestine and the stomach are equally involved. To this group belong some affections which have already been considered, which begin in one viscus and then spread to the other. But those in which both are primarily affected still remain to be described. The most important of them are the various forms of cholera. Now, there are several very different complaints classed together under the head of cholera.

**Cholera Infantum.**—And first, of the so-called cholera of infants. It is an endemic in the larger cities of the United States during the hot months, and one fraught with danger to all young children. It begins generally with diarrhœa. Vomiting soon follows ; and for a time the two go hand in hand ; but, unless the case be of short duration, the spontaneous vomiting ceases, or at all events gives way to occasional exacerbations of irritability of the stomach, while the looseness of the bowels remains, or even augments. The discharges are colorless, or yellowish, or greenish. There is thirst ; sometimes fever. The abdomen may be sunken or swollen ; and it may be tender. Sometimes the disease runs its course within three or four days ; at the end of which time the child dies, worn out by the constant vomiting and purging. More generally the disorder is of longer duration ; for weeks or for months it continues, the diarrhœa improving and then returning with redoubled severity, and kept up or increased by the irritation of teething. The irritability of the intestinal canal, and the utter impossibility of retaining enough food to nourish the wasting body, gradually wear out the system. The child before death is wan and distressingly emaciated ; sometimes hypostatic congestion of the lungs, broncho-pneumonia, boils, suppression of urine, plaintive cries, rolling of the head, strabismus, and coma precede the fatal termination.

Such is a sketch of grave and intractable cases. Yet very many cases are far from being desperate. Under judicious treatment a large number are annually saved. Recoveries would bear a still higher proportion to the deaths, were it not that the greatest sufferers from the disease, the children of the poor, are unable to obtain the means most certain to restore them to health,—change of air. Cooped up in crowded neighborhoods, surrounded on all sides by filth rapidly decomposing under the burning rays of the sun, they are compelled to breathe the hot, noxious atmos-



phre which has been the chief agent in producing the liability to the complaint.

The disease is an entero-colitis from a milk infection leading to bacterial fermentation in the intestines, with enlargement of the solitary glands, and even at times of Peyer's patches. The researches of Vaughan have demonstrated that a ptomaine appearing in milk, tyrotoxin, is its most frequent source. Temporary diarrhoeas in children occurring in hot weather could alone be mistaken for the disorder. But the fact that they are temporary, not followed by vomiting, and not associated with the grave symptoms of approaching collapse, shows us the difference.

**Cholera Morbus.**—This, or *cholera nostras*, is, like cholera infantum, a disease of the hot season; yet it is also observed at other times of the year. But, although the chief predisposing cause is undoubtedly heat, there is generally an exciting cause which develops the disorder,—such as exposure, checked perspiration, drinking large quantities of ice-water, or imprudence in eating. The attack is characterized by spasmodic pains in the abdomen, by cramps in the legs, by rapid loss of strength, and by repeated vomiting and purging. The matter ejected both from the stomach and from the intestines is liquid, and contains a large quantity of bile. In truth, the affection is in reality a cholera, a flow of bile, which its more formidable namesake, Asiatic cholera, is not. Finkler and Prior have found in the stools a comma-bacillus, *vibrio proteus*, which is larger and thicker than the bacillus of Asiatic cholera, but with shorter spirilla, and cultures of which, unlike the latter, rapidly liquefy in gelatin, and grow on potato even at ordinary temperatures.

Cholera morbus may be preceded by colicky pains, nausea, and rumbling in the intestines. More generally it comes on suddenly. When at its height, the cramps in the calves of the legs cause the muscles to rise up in hard, knotty masses; the stools are fetid; the vomiting is constant; the thirst is great, and the skin is cool or cold. But the patient does not remain long in this condition. In the course of a few hours, or at the utmost of a day, the symptoms mitigate, or yield entirely to treatment; and, pale and visibly emaciated though he be, he speedily regains his health. Only in some cases the disease proves intractable, and, after

running on for several days, passes into a state of hopeless collapse.

There are not many morbid states with which cholera morbus is likely to be confounded. It may be mistaken, as we shall presently see, for epidemic cholera. We find many points of similarity between it and irritant poisoning. But there are also strong points of difference. The vomiting and purging produced by an irritant poison do not come on at the same time: the vomiting precedes the purging. The pain is first in the epigastrium, thence it may spread. Moreover, we often detect signs in the mouth or fauces which prove the irritating character of the substance swallowed. The vomiting and the subsequent acute gastritis are accompanied by fever and a small, tense pulse; whereas the skin of cholera morbus patients is commonly cool, and the pulse very compressible and feeble. In acute gastric catarrh, as we meet with it mostly from indigestion, there is vomiting, but not purging.

**Cholera.**—The formidable complaint known as epidemic cholera, Asiatic cholera, malignant cholera, or by the simple name of cholera, has some striking features of resemblance to the disorder just considered. It shares with cholera morbus the vomiting and purging, the cramps, the sudden depression; but it is an affection of different origin and of much more serious import, and presents symptoms not encountered in the cholera that occurs yearly during the hot weather. And although, on account of the gastric and intestinal disturbances which form so prominent a part of its manifestations, it is here described among the disorders of the alimentary tube, I am doing so for the sake of clinical convenience, and contrary to sound pathology; for cholera is not an affection either of the stomach or of the intestines; it is an epidemic constitutional disorder of the most formidable character, generated by a poison transmitted to us from the East. The poison leads to a casting off of the epithelium of the mucous membrane of the alimentary tube; perhaps to changes in the membrane. But the engorged veins all over the body; the ready exosmose of the watery parts of the blood; the frightfully-rapid prostration; the sudden blight which befalls the nervous powers,—are elements which are even more characteristic, and which throw more light on the nature of the fearful malady, than the

comparatively uncertain and far from uniform appearances of irritation in the intestinal canal.

The access of cholera is at times sudden and most unexpected; the patient, previously in good health, is stricken down without warning by the force of the poison. More generally there is a premonitory stage: a stage of languor, low spirits, uneasiness, headache, and diarrhœa. The effects of the morbid matter are indeed visible in hundreds of individuals who, during the prevalence of cholera, suffer from these premonitory symptoms without any of greater danger arising. Nay, the same influences which give rise to a *choleraic diarrhœa* in healthy persons have the effect of rendering the bowels of those habitually constipated regular, and sometimes even loose.

When the malignant disease is fairly developed, there is vomiting as well as purging. The contents of the stomach and intestines are first voided, and then large quantities of a rather turbid fluid resembling rice-water, with whitish particles like rice floating in it. They are the epithelial cells of the alimentary tube, which have been thrown off from the mucous membrane; and in the dejecta we find the comma-bacillus discovered by Koch. This may be seen by examining microscopically the bacilli obtained from a small amount of cholera dejection that has been mixed with an equal amount of alkaline meat broth at a temperature of 30° to 40° C. and allowed to stand for twelve hours in an open glass. The cholera-bacilli develop on the surface. They are readily stained, in about ten minutes, with a diluted alcoholic solution of fuchsine or methyl-violet. They are decolorized by Gram's process already described. After the staining, which must take place with the infected side downward, the cover-glasses are taken from the solution with the forceps, washed in water, dried with the prepared side uppermost, and mounted in Canada balsam. Prior to the staining a drop of the infected broth or a particle from a stool is dried in air, after having been rubbed between two cover-glasses and passed three times through the flame of a Bunsen burner. The bacilli of cholera may be recognized even without the microscope by a rose-violet color, the *cholera reaction*, that becomes apparent in a few minutes if a ten per cent. hydrochloric acid solution is added to cholera cultures. The cholera-bacillus is confined to the intes-

tine. In the extensive observations made by Shakespeare in India and elsewhere\* it was not found in the blood or in the tissues or organs outside of the intestinal canal.

FIG. 55.



The comma-bacillus of Koch, from culture in blood-serum. Zeiss  $\frac{1}{2}$ , homo. im., Oc. 4.

Simultaneously with the vomiting and purging, or very shortly after, come on severe spasmodic pains in the abdomen and cramps of the muscles of the belly and of the extremities. With all this there are a burning sensation in the epigastric region; an unquenchable desire for cold drinks; a cool skin; a pulse slightly more frequent than normal; a temperature which may be normal or may fall to about  $95^{\circ}$ , and which often shows many degrees of difference between the rectum and the axilla; an oppressed breathing; and a rapidly-progressing exhaustion. The case now stands on the verge of collapse. Should this follow, a state of things is witnessed which, once seen, remains indelibly engraved on the memory. The pulse is quick, but hardly perceptible. The discharges cease, and so do often the cramps. The skin is cold,

---

\* Report on Cholera in Europe and Asia, Washington, 1890.



covered with a clammy sweat, and has a bluish look. The nails and the lips have the same unnatural appearance. The whole body shrinks, and seems at times almost to wither visibly even while under inspection. The countenance assumes the aspect of death; the eyes are sunken and have a glassy look. The temperature is low; it may fall to  $79^{\circ}$ . The intellect is commonly clear; but, when the patient talks, the words fall strangely on the ear. It seems as if a corpse had spoken, and the voice is husky and faint. The tongue and the expired air are cold. No symptom, indeed, has struck me more forcibly than the icy breath.

But the symptoms do not always take place in the order described, nor are they all uniformly present. The vomiting and purging may be wanting from the onset, and so too may the cramps. Only one symptom is never absent,—the tendency to early sinking. Sometimes a stage of perfect collapse is reached with frightful rapidity: instead, as is commonly the case, of several hours elapsing before complete prostration comes on, the vital powers are at once laid low by the assault of the dreadful malady. When cholera prevailed in Philadelphia some years since, I attended a woman who at six o'clock in the morning was in perfect health and who in a little more than half an hour afterward was lifeless. There was neither vomiting nor purging; nothing but cramps, stupor, and speedy collapse. Such cases are not uncommon in the home of cholera,—India. Post-mortem inspection shows the thin rice-water fluid locked up in the alimentary canal. Nature makes an effort to eliminate the poison; but before she completes her task, life is palsied.

In those cases that recover, or in those of light character, *cholérine*, the vomiting and purging gradually subside, the skin becomes warm, the pulse fuller, the urine—which, while the disease was at its height, was not passed, perhaps not secreted—is again voided, the patient falls into a refreshing sleep, and, the symptom most favorable of all, bile reappears in the stools. Even in apparently hopeless cases of collapse we may be fortunate enough to witness these favorable changes. But where the prostration has been great, the reaction is apt to be violent. A decided fever of low type, with rapid pulse and heat of skin, and attended very often by alarming cerebral symptoms, succeeds; and the urinary secretion, even if it had been restored, becomes again very scanty.

Thus the period of reaction brings with it new dangers, and of a kind which are sometimes insurmountable. And this low form of fever, very similar to typhoid, though readily enough distinguished by the preceding symptoms, may last for upwards of a week before death takes place or the signs of danger gradually yield. Now, this cholera typhoid may be preceded by scanty urine and marked uræmia, but it may also exist independently of this morbid state, though probably equally due to the blood being loaded with broken-down material. In cases in which uræmia sets in, whether it be followed or not by a fever of low type, there is at first but little, if any, heat of skin, and a slow pulse; the patient is wild, restless, or drowsy; the kidneys act very imperfectly, the urine is greatly deficient in urea, and usually contains albumen. These are very dangerous cases, and if the secretion be seriously retarded for more than twenty-four hours they are likely to perish.

In any case of cholera, convalescence is apt to be slow. For weeks or months irritability of the intestinal canal remains; and I have met with instances in which it has never disappeared.

It would be needless to go into any minute description of the differences between cholera and other affections: its features are not to be mistaken. *Cholera morbus* is the only disorder which really resembles it. The dividing-line is drawn by the absence of bile in the discharges, the rice-water evacuations, the greater severity and more rapid progress of the symptoms, the bluish color of the surface in the stage of collapse, and the epidemic character of the more fatal disease. In the truly epidemic nature of the distemper, in the presence of the cholera-bacillus in the evacuations, and in the speedy collapse, which shows but too plainly that some highly deleterious matter has poisoned the system, lie, even in doubtful cases, the proofs that we are dealing with malignant cholera; for sometimes rice-water discharges occur in bad cases of cholera morbus; occasionally, too, this disorder appears to be epidemic; but it is only on a very small scale. To speak more accurately, it is an endemic on a large scale. We find no proofs of a virulent poison directly conveyed by human intercourse and traffic, and so noxious as to smite animals as well as man. Certain rare cases of *irritant poisoning*, especially from arsenic, bear some resemblance to cholera, although generally

more to cholera morbus. The severe vomiting in advance of the purging, the usual absence of rice-water stools, the presence of bloody evacuations, and the traces left by the poison in the mouth, furnish significant features of distinction.

The mortality of cholera is very various. In many epidemics one-half, or more than one-half, die. In some the havoc is far less. The first cases that occur almost invariably perish; and, taken altogether, the malady ranks among the most destructive to life. Its epidemic visitations are what the plague was to the Europeans of the seventeenth century, and what yellow fever still is to the inhabitants of this continent. It is at least as dangerous; its true nature is as hidden; its management is as unsatisfactory.



### SECTION III.

#### DISEASES OF THE LIVER.

We have already inquired into the clinical methods of examining the liver with a view to forming a judgment of its physical characteristics. Let us now look at some of the symptoms.

*Pain* is one of these. It is generally dull, and radiates from the seat of the liver to the upper portion of the thorax, to the scapula, to the shoulder, and to the umbilicus. Commonly it is persistent and increased by strong pressure; yet the exceptional cases are numerous. As happens with other symptoms of disease of the liver, with vomiting, with jaundice, it may be noticed that the pain is sometimes strangely periodical, suggesting malaria, but uninfluenced by quinine.\*

*Digestive troubles* are usual accompaniments of hepatic affections. They are of all grades, from mere indigestion to the signs announcing chronic gastritis.

*Disturbance of the portal circulation* is another frequent consequence of disease of the liver. The flow of blood is interfered

---

\* See on this subject a paper by Cyr, Arch. Gén. de Méd., May, 1883.

with, and the result is seen in the occurrence of dropsy, of piles, of partial peritoneal inflammation, of hemorrhages from the engorged stomach and intestines, and of enlargement of the spleen and of the veins on the surface of the abdomen.

**Jaundice.**—The most significant manifestation of hepatic disorder is jaundice. This marked sign shows itself by the yellow tinge imparted to the skin and to the conjunctiva. Yet the yellowness is not confined to these structures: it may be found in internal organs. Besides the peculiar aspect of the surface, icterus is usually attended with depression of spirits; with slow pulse; with itching of the skin; with high-colored urine, in which the main ingredients of bile can be detected, and sometimes small quantities of albumin, or hyaline and epithelial casts without albumin; with constipation, the *feces* passed being hard and knotty, and often of bad odor, and almost devoid of color, or of a leaden hue.

Jaundice is due to the presence of biliary constituents in the blood; they get there from the bile, in consequence of some impediment to its outward passage, being reabsorbed and conveyed into the circulation; or it happens because the liver-cells cannot perform their functions; or because some poison changes the proper relation between blood destruction and liver secretion, and the elements of bile accumulate in the blood. In the jaundice from reabsorption, the most common form of jaundice, it has been demonstrated that the bile may get into the blood in consequence of altered pressure in the vessels. Lower pressure in the vessels causes the bile to flow into them.

The *diagnosis* of jaundice is easy. The only morbid signs with which it is liable to be confounded are the slightly yellowish hue of chlorosis, or of some cachectic conditions combined with organic visceral disease, and the yellow appearance of the conjunctiva which is natural to some persons. The changed color of the countenance due to chlorosis is told by its association with a bluish-white or pearly-tinted eye, and with pale lips and tongue and transparent ear. The absence of a yellow tint from the conjunctiva is of equal importance in discriminating from jaundice the yellowish hue of cancer, of malaria, of lead poisoning, and of granular kidneys. The history of the case also aids us. The yellow look of the eye sometimes found in health, and at times



dependent on subconjunctival fat, is known by the unequal distribution of the color and by the absence of a yellow hue of the complexion. But in negroes—and it is in them especially that we meet with the discolored conjunctiva—we have to judge by the character of the coloration alone. In any doubtful case, the chemical tests for bile-pigment in the urine will solve the doubt.

When once jaundice has been recognized, the difficulty in diagnosis may be said to begin. Of the many distinct sources of icterus, which one is before us? Now, clinically speaking, the causes may be thus grouped: 1. Diseases of the liver. 2. Diseases of the bile-ducts. 3. Diseases of parts remote from the liver, or general diseases leading to a disorder of the viscus. 4. Certain poisons acting upon the blood. In the first two of these causes there is, as it were, a mechanical difficulty impeding or arresting the excretion of bile; in the third and fourth no impediment exists. Let us look at some of the peculiarities of these groups.

1. The jaundice connected with diseases of the liver is, as a rule, recognized by its association with changed dimensions of the organ, and with pain or other palpable signs referred to the hepatic region. It is met with in all disorders of the liver, but does not exist in all in the same degree of intensity. It reaches a high development and is combined with cephalic symptoms in acute yellow atrophy. In fatty liver, in waxy liver, in cancer, in cirrhosis, and in acute hepatitis, it is not marked, and may be, indeed, absent: in truth, it can hardly be looked upon as belonging to the first-mentioned morbid states. The jaundice of this class of cases is due to interference with the secreting function of the liver-cells.

2. Jaundice arising from disease of the larger biliary ducts, such as their catarrhal swelling; or in consequence of their obstruction by pressure exercised by a morbid enlargement of adjacent parts, as of the pyloric extremity of the stomach or the pancreas; or by tumors, aneurismal, cancerous, or fæcal, closing the orifice of the duct; or by the stoppage of the ducts by inspissated bile or a biliary calculus, or by hydatids or foreign bodies from the intestines,—is a form of the malady in which the icterus is commonly intense. The obstructive jaundice occasions no head symptoms; and when these are absent in a case of very deep jaundice,

when, further, the stools are completely discolored, we are generally correct in attributing the morbid phenomena to an impediment to the flow of bile through the common bile-duct or the hepatic duct. If this impediment be due to the impaction of a gall-stone, a constant sense of weight and recurring severe colicky pains are encountered in addition to the signs just mentioned.

In the jaundice due to reabsorption—precisely the form of jaundice, therefore, that happens if any serious obstacle in the biliary passages exist—the biliary acids pass into the blood, and thence into the urine. But this is not a certain sign of obstructive jaundice. Traces of the bile-acids have been found in healthy urine.

3. Illustrations of jaundice following some local lesion of other parts of the body, or appearing in the course of a general constitutional affection, are furnished by the jaundice which happens in some cases of pneumonia, or which is encountered in remittent, in typhus, in relapsing, or in yellow fever. In these fevers the yellow hue is generally found to be connected with an acute enlargement and with structural changes in the organ; and in the latter malady, with disordered hepatic circulation and a fatty degeneration of the secreting-cells. But, besides the interference with the secreting power of the cells, the blood alterations in these manifestations of jaundice must be considered.

To recognize the form of jaundice under discussion, we must examine all the viscera of the body with care, laying stress upon the history of the case and the phenomena attending the jaundice. Otherwise, too much importance will be attached to this symptom, and the disturbance of the liver will be regarded as forming the whole complaint, when it is but a small part of it.

4. Poisons acting upon the blood sometimes give rise to jaundice very rapidly; for instance, the jaundice from snake-bites or from pyæmic affection is apt to be suddenly developed. As a rule, the tint is light. In the history of the accident and the signs of alteration of the blood we possess the means of distinguishing this form of jaundice. Certain mineral poisons, such as phosphorus, copper, antimony, come into the same category. Chloroform and ether, too, may lead to abnormal blood changes producing jaundice. The urine enables us to a certain extent to tell blood jaundice from jaundice caused by liver disorder. We find hæmoglobin in the urine, or get from its hæmatin the hæmin crystals

of Teichmann. These are obtained by drying urine on a slide, adding a little salt, and then glacial acetic acid under the cover-glass. The slide is heated until bubbles rise, and on cooling the characteristic blood-crystals form. If these or hæmoglobin be found, there is a strong probability of the jaundice being of blood origin.

Thus, then, we can bring, clinically speaking, most of the varieties of jaundice under one or the other of the four heads mentioned; and, roughly speaking, they come really under two,—obstructive jaundice, where the disorder results from obstruction of the common duct, and jaundice without such obstruction. But there are a few kinds of jaundice which it is not easy to classify with precision: one of these is the jaundice from mental emotion; the other, the jaundice of newly-born children.

As regards the former, no satisfactory explanation has been given. All we know is, that violent anger or fright may lead within a very brief space of time to the development of jaundice, and that the quickly-occurring discoloration is not dangerous or of long duration. The perverted innervation caused by concussion of the brain leads to a similar kind of jaundice.

The jaundice of newly-born children—*icterus neonatorum*—is ordinarily a mild complaint, which appears soon after birth and rarely lasts over two weeks. The yellow hue of the skin is often very deep; yet the child does not suffer, and has no febrile excitement. The bowels are constipated, but the stools are not necessarily altered in their color; nor do they usually present the clayey look which might be expected from the aspect of the skin and of the conjunctiva. West states that the disorder is most frequently observed in children prematurely born.

The prognosis of jaundice depends upon its cause. In general terms, we may say that if the icterus last upwards of two months it is always a matter of some danger, as showing, in all likelihood, an organic lesion of the liver or of the biliary passages, or unyielding pressure on them. If the discoloration of the skin be attended with cerebral symptoms, the patient's state is precarious. Icterus accompanying affections of the blood, peritonitis, or pneumonia is an unfavorable sign; so are wide-spread ecchymoses, or a very dark color of the skin. Indeed, cases of "green" or "black" jaundice generally prove fatal.

Before examining the hepatic maladies according to their clinical features, let us look at their pathological classification :

## DISEASES OF THE LIVER.

Diseases of hepatic parenchyma.	{	Hyperæmia .....	{ Acute congestion. Chronic congestion.
			{ Acute hepatitis. Chronic hepatitis. Interstitial inflammation; cirrhosis, atrophic and hypertrophic.
	{	Inflammation and its consequences .....	{ Abscess. Softening.
			{ Syphilitic hepatitis.
	{	Atrophy .....	{ Acute yellow atrophy. Simple chronic atrophy. Red atrophy.
		Hypertrophy .....	{ Partial. General. Fatty liver. Waxy liver. Pigment liver.
	{	Degeneration and new formations.....	{ Cancer. Sarcoma. Lymphatic growths. Gummata. Tubercle. Hydatids. Simple cysts.
Diseases of biliary passages.	{	Inflammation of gall-bladder and gall-ducts .....	{ Catarrhal. Exudative. Suppurative.
		Occlusion of biliary passages.	
	{	Dilatation of gall-bladder.	
		Morbid growths.	
		Foreign bodies; concretions, such as gall-stones.	
Diseases of blood-vessels.	{	Biliary fistulæ.	
	{	Of hepatic artery .....	{ Inflammation. Sclerosis. Aneurism.
		Of hepatic vein.	
	{	Of portal vein.....	{ Suppurative inflammation. Thrombosis.



Acute Diseases of the Liver attended generally with Slight Enlargement of the Organ, and with more or less, though rarely very much, Jaundice.

**Acute Congestion.**—This arises from organic disease of the heart, from obstructed portal circulation, from irritating food and drink and disturbed digestion, or from malarial poison ; sometimes it is caused by a high temperature, by a blow on the hepatic region, by arrest of the menstrual flow, by a protracted chill, by violent exercise, or, as Frerichs points out, by injury to the semi-lunar ganglia. The acute congestion is characterized by pain in the right shoulder and loin, by an unpleasant sensation of weight and of tension in the right hypochondrium, increased after meals, and by nausea and vomiting. At the same time the action of the bowels is deranged, being generally too frequent ; the tongue is coated ; there is flatulency, as well as depression of spirits, with loss of appetite and of strength ; and the liver is enlarged. But we find ordinarily only slight jaundice, and no fever. Gradually these signs disappear ; the increased hepatic dulness, however, remaining for some time after the gastric and intestinal disturbances have abated. Not unfrequently the acute disorder passes by imperceptible degrees into a chronic state.

**Acute Hepatitis.**—The symptoms of this affection are much the same as those of acute congestion, except that we observe rise of temperature, and in some cases enlargement of the spleen, and albumin in the urine. The pain is dull, and is increased on pressure, yet not much so, unless the peritoneal covering of the liver be involved. Jaundice is not generally marked ; indeed, at the beginning of the disease it is often absent.

Acute hepatitis is common in hot countries, and many of the cases are connected with dysentery, particularly with amœbic dysentery. It may end in resolution ; but, especially in persons of indolent or intemperate habits, it often terminates in suppuration, and pus collects in the substance of the liver. The occurrence of this, the *tropical abscess*, as Murchison \* called it, is indicated

---

\* Diseases of the Liver, 2d edit., 1877.

by recurring rigors, by fever of remittent type, by clammy perspirations, by prostration and loss of flesh. Not unfrequently, too, a decided fulness of the side may be noticed, and occasionally careful palpation detects deep-seated fluctuation. After an abscess has formed, the danger is great; secondary abscess may follow, and the patient is apt to perish from peritonitis, or from blood-poisoning. Yet recovery may take place. The matter may be discharged through the abdominal walls, or burst into the intestine, or find its way through the diaphragm into the pleural cavity, to be discharged through the lung. But, as the phenomena of abscess of the liver following acute inflammation are in the main the same as when the collection of pus is consequent upon other morbid states, we shall not here consider what we shall presently fully examine.

Let us now examine the maladies with which acute inflammation of the liver may be confounded, premising that, making allowance for the febrile phenomena and the other slight signs of difference just indicated between hepatic inflammation and hepatic congestion, the same remarks will apply to the distinction between this morbid condition and the affections about to be mentioned. The complaints resembling acute hepatitis are:

PERIHEPATITIS;

INFLAMMATION OF THE PORTAL VEINS;

PIGMENT LIVER;

CHRONIC HEPATIC DISEASES WITH ACUTE SYMPTOMS;

ACUTE NON-HEPATIC DISEASES WITH JAUNDICE;

DIAPHRAGMATIC PLEURISY;

ACUTE INFECTIOUS JAUNDICE;

INFLAMMATION OF THE BILIARY PASSAGES;

ACUTE YELLOW ATROPHY.

*Perihepatitis*.—Inflammation limited to the serous covering of the liver is not a frequent disease. Unless it be of syphilitic origin, it is scarcely ever observed as a primary affection; it is generally caused by the extension of inflammation from organs adjacent to the liver,—as from the stomach, intestines, diaphragm, or pleura,—and may therefore be looked upon as a local peritonitis; or it is an attendant upon disease of the liver itself. In the latter case it presents no peculiar symptoms, except that it adds tenderness to the signs of the hepatic malady it complicates.

Under other circumstances it is more likely to be confounded with acute inflammation of the liver-texture. Yet the far greater tenderness, the severe pain upon motion or deep inspiration and its marked increase when the patient lies on either side, an occasional grating friction sound, the normal size of the gland, the history of the case or the evidences of a disease in the neighborhood of the liver that is likely to cause the malady, the absence of jaundice, and the slight fever, distinguish the perihepatic inflammation.

*Inflammation of the Portal Veins.*—An inflammation of the portal veins, terminating in suppuration or in their infection by a general pyæmia, or through local processes in the portal circle, is very liable to be mistaken for acute hepatitis. Nor are there, in truth, any positive symptoms by which we can discriminate between the two maladies. Still, we may suspect that the veins, rather than the structure of the liver, are the seat of inflammation, if, with the signs of acute and painful enlargement of the organ, we find jaundice, thin and copious stools, recurring chills and profuse sweats, emaciation, increase in size of the spleen, without any apparent fluctuation or other signs of an hepatic abscess; if there exist pains between the ensiform cartilage and the umbilicus, or in the epigastrium or right hypochondrium, or shooting to the lumbar and sacral regions; if following these symptoms appear swelling of the veins of the abdominal walls and striking evidences of hectic fever or of peritonitis; and if these phenomena be encountered in a person who, on account of a previous affection of the intestines or the appendix or the spleen, or of any other organ having a connection with the portal circulation, is liable to disease of the portal system. Marked enlargement of the spleen is a constant feature of impediment in the portal vein, whether from inflammation or from thrombosis.

*Pigment Liver.*—In accumulation of pigment in the liver, which is most common as the result of a deep malarial poisoning, the liver is not the only organ implicated in the morbid process: the spleen is commonly affected; the blood becomes watery, its corpuscles are broken down, and it contains the malarial corpuscles and large quantities of pigment; and pigment accumulates in the kidneys or in the brain. Now, the effect of all this is to occasion marked symptoms, besides those referable to the derangement of

the liver; for it is not unusual to find grave cerebral disturbance, albuminuria, hemorrhage from the intestines, profuse diarrhœa, and enlargement of the spleen. Irrespective of these manifestations, we may note the singular ash or grayish-yellow color of the skin, and the evident anæmia. The fever that accompanies the morbid condition is of an intermittent type; the pulse is not, as a rule, much accelerated, and the jaundice is generally slight. In India, pigmentary degeneration of the liver tends to suppurative hepatitis.\* When we contrast the phenomena described with those of acute hepatitis, we see at once the difference. The fever, the aspect of the patient, the blood full of dark pigment and malarial corpuscles, and the frequency of cerebral symptoms, are entirely unlike.

*Chronic Hepatic Diseases with Acute Symptoms.*—We occasionally meet with patients who seem to be laboring under an acute affection of the liver, either some form of inflammation of the liver-structure or of the biliary passages, or congestion of the liver, but in whom the acute symptoms have merely supervened upon a chronic complaint. Such cases are puzzling; we may have to wait for their solution until the acute symptoms subside. In hepatic cancer the sudden and rapid development of the malady amid the signs of acute congestion is not very uncommon. Occasionally the peculiar physical phenomena of individual hepatic diseases, such as the nodular tumors of a malignant growth, or the fluctuation of a hydatid cyst, will assist materially in the diagnosis.

*Acute Non-Hepatic Diseases with Jaundice.*—There are many acute affections, such as pneumonia, pyæmia, puerperal fever, and some forms of poisoning, in which jaundice may coincide with febrile symptoms and excite suspicions of acute hepatitis. But the yellowness of the skin which may attend the non-hepatic disorders mentioned is accompanied by symptoms so different that a mistake is not likely to arise if the history of the case be taken into account and other viscera besides the liver be explored.

*Diaphragmatic Pleurisy.*—The manifestations of inflammation of the pleural covering of the diaphragm are in several respects similar to those of inflammation of the liver. We find in this

---

\* Aitken's Practice of Medicine, vol. ii.



diaphragmatic pleurisy pain in the right hypochondrium, nausea and vomiting, dry cough and embarrassed respiration, occasionally jaundice,—much the same symptoms which we observe in hepatitis, especially if the serous envelope of the liver be at the same time implicated. But the pain in diaphragmatic pleurisy is far greater, more suddenly developed, is much more aggravated by movements and by full inspiration, and is always evoked by pressure. The diaphragm on one side is immovable; the hypochondriac region is retracted; the breathing is purely costal and short; the difficulty in breathing amounts to orthopnoea; the body is bent forward. We often encounter hiccough, great anxiety, and delirium, sometimes a sardonic grin on the features, and the cough comes on in frequent paroxysms; and although, as a case recorded by Andral\* proves, there may be jaundice, yet this is in reality so generally wanting as scarcely to belong to the symptoms of diaphragmatic pleurisy. Then in this complaint we may find friction sounds,—though the physical signs will not always aid us, being often uncertain,† and consisting simply in enfeebled breathing, with perhaps a few fine moist rales at the lower portion of one side of the chest. The fever with these imperfect physical signs may be slight or be very marked; it is generally ushered in by a chill. There is generally, in addition to the pain along the cartilages of the false ribs, a tender spot in the epigastrium, on a level with the tenth rib, one or two finger-breadths from the linea alba. There are shooting pains along the clavicle and in the tract of the superficial cervical plexus, and the phrenic nerve of the affected side, pressed on in the neck, is very sensitive. The pain on pressure is generally most intense along the costal insertions of the diaphragm, especially of the tenth rib; it is stated that upward pressure affords a means of diagnosis, as it relieves the pleuritic pain.‡ The difficulty in expectorating, owing to the pain, may be so great as to hasten death.§

*Acute Infectious Jaundice.*—This malady,|| also known as *Weil's disease*, presents many symptoms of acute hepatitis. It is very

---

\* Clinique Médicale, tome ii.

† Cases by Habershon, *Guy's Hospital Reports*, 1869.

‡ *British Medical Journal*, Aug. 1871.

§ Frank Donaldson, Jr., *Amer. Journ. Med. Sci.*, April, 1885.

|| Described by Weil, *Deutsches Archiv für Klin. Med.*, Bd. xxxix.

doubtful, however, whether it is an affection of the liver, but is not rather an infectious fever due to the invasion of a specific micro-organism through the gastro-intestinal tract. Jaeger\* has, in cases of Weil's disease, isolated from the urine during life and from the tissues after death a short curved rod, provided with cilia, which he designates "*bacillus proteus flavescens*." Weil's disease is marked by jaundice, swelling of the spleen, nephritis, and blood-alteration. It mostly affects vigorous young men in hot weather; butchers and soldiers are especially liable to it. It has been also observed in persons who have bathed in water contaminated by fowls suffering from an analogous disorder.† It begins abruptly with headache, dizziness, and decided elevation of temperature. The jaundice is moderate, the liver slightly swollen and painful; there is great weakness, with delirium and somnolency, increased thirst, and general malaise, with loss of appetite. Besides albumin and tube-casts, the urine may contain blood; both bile-pigment and bile acids are found in it. There are pains in the limbs, especially in the calves; the bowels are usually loose. The symptoms abate quickly; from the seventh to the eighth day the temperature falls gradually to normal. A return of fever after a period of its absence from one to seven days may happen, but this return does not last more than three to six days. The convalescence is extremely slow. Fatal cases have presented fatty degeneration of the liver, acute parenchymatous nephritis, and enlargement of the spleen.‡ The disease resembles *relapsing fever*, but the spirilla have not been found in the blood. Nor is defervescence attended with a critical discharge followed by sub-normal temperature. Further, the ascent of the temperature of the secondary fever is gradual, while that of the paroxysm of relapsing fever is sudden. The return of the fever makes it unlike *abortive typhoid* with bilious symptoms. Then, it shows no eruption, except herpes and an erythema.§ Besides, jaundice and nephritis are rare in typhoid fever.

Between *acute yellow atrophy* of the liver and Weil's disease

---

\* Zeitschrift für Hygiene und Infektionskrankheiten, Dec. 9, 1892.

† Jaeger, *loc. cit.*

‡ Ibid.

§ Fiedler, Deutsches Archiv f. Klin. Med., Feb. 1888.

there is a close resemblance. But the former has a prodromal period, while the onset of the latter is abrupt. The second is attended with elevation of temperature of peculiar range; in the first the temperature is, as a rule, not elevated, and may be sub-normal, and the bowels are constipated. In acute yellow atrophy the jaundice is gradually progressive and the liver is at first enlarged and subsequently reduced in size; the jaundice of Weil's disease is slight and soon subsides, and the liver remains enlarged throughout the attack. In acute yellow atrophy the urine may contain albumin and tube-casts, but there are not the pronounced symptoms of nephritis that Weil's disease presents. The tendency to hemorrhages is far greater in acute atrophy of the liver than in infectious jaundice. The one condition is almost invariably fatal; the other is, as a rule, followed by rapid improvement and recovery. Weil's disease in some respects resembles yellow fever, but it is an affection of several paroxysms.

*Inflammation of the Biliary Passages; Acute Yellow Atrophy.*—Both of these maladies may be confounded with hepatitis. But the former, although presenting more jaundice than the other maladies of the group now under discussion, is otherwise so similar that it will be described as one of the main affections of this group; and, in truth, in temperate climates acute affections of the liver are mostly catarrhal jaundice. Acute yellow atrophy belongs clinically to diseases characterized by decrease in size of the liver; and it is while discussing these that we shall point out its characteristic differences.

**Inflammation of the Gall-Bladder and Gall-Ducts.**—Inflammation, when it attacks the biliary passages, is most apt to affect the gall-bladder and the ductus choledochus. Frequently the morbid process is propagated from the stomach or intestines, and nausea, furred tongue, a feeling of weight in the epigastrium, feverishness, and diarrhoea, occur previous to the discoloration of the faeces, to the jaundice, to the increased hepatic dulness, and to the slight tenderness on pressure in the right hypochondrium; in other words, the symptoms of gastric or gastro-intestinal catarrh precede those of “*icterus catarrhalis*,”—by far the most common form of inflammation of the gall-bladder, for suppurative inflammation is very rare.

Catarrhal icterus does not cause any great enlargement of the

liver, and the swollen organ remains smooth on palpation. Nor is the tenderness decided, except over the tumid and projecting gall-bladder. The jaundice, at first slight, becomes after a few days, as the bile-ducts are obstructed, intense, and the stools are white and devoid of bile. There is now no fever, or this is but very slight; the pulse is usually slow. The affection is the most common cause of marked jaundice in young persons; when found in the middle-aged or in the old it is apt to be associated with a gouty diathesis or to have followed syphilis; and at any age it may be secondary to other diseases of the liver, and is then apt to be lasting.

Generally catarrhal icterus is a tractable disorder; and after continuing for two or three weeks, it usually subsides. But it may persist for as many months; and in rare instances the inflammation leads to an occlusion of the bile-ducts, and to a fatal issue. I had such a case in 1863 under my charge at the Philadelphia Hospital. The patient, a man upward of sixty years of age, died deeply jaundiced and comatose. He had presented, during life, the signs of enlargement of the liver; little or no tenderness in the hepatic region; no fever; but much gastric irritability and obstinate constipation, both of which had existed for three weeks prior to a noticeable discoloration of the skin. The whole disease was, as far as could be ascertained, of only two months' duration; and the jaundice steadily deepened from the time of its first appearance. At the autopsy, the gall-bladder was found enormously distended, its coats thin, yet otherwise scarcely abnormal; but the common duct was obliterated by inflammation. The stomach and the upper bowel were congested, while the coats of the stomach toward the pylorus were thickened. A similar case has been described by Tyson.\*

Now, in point of diagnosis, it is not generally difficult to distinguish the catarrhal inflammation of the gall-bladder, except in those rare instances in which the common duct or the hepatic duct is obliterated. It differs from *congestion of the liver* by the different etiological elements in the history of the case,—the one disorder happening commonly in connection with disease of the heart, or an obstruction of the portal circulation, or a miasmatic

---

\* Transactions of the Pathological Society of Philadelphia, vol. iv.



poison, the other following usually exposure to cold and damp, or the eating of quantities of indigestible food. Then, inflammation of the gall-ducts gives rise to decided jaundice.

Catarrhal jaundice may occur as an accompaniment of some general morbid condition, or in an epidemic form. These cases are distinguished by the history, by the tendency to acute disease of other organs, such as the lungs and kidneys, and by enlargement of the spleen.\*

From the jaundice of chronic hepatic maladies—such as *cancer* or *cirrhosis*—we separate catarrhal icterus by the non-existence of the physical signs of these maladies, by its acute course, and by the dissimilar progress of the symptoms. Still, as regards cancer we must bear in mind that we encounter in elderly gouty persons cases of long-persisting catarrhal icterus attended with frequent vomiting and marked emaciation which strongly resemble cancer, yet slowly yield to treatment. Inflammation of the biliary passages with the jaundice arising in consequence of *biliary calculi* is distinguished by the severe pain, the sudden appearance of the icterus subsequent to the paroxysms of pain, its increase after them, and its often rapid fading after the gall-stone is voided. The symptoms of the early stages of *acute atrophy of the liver*, as well as those of some cases of acute inflammation, may be so like the symptoms of inflammation of the gall-bladder and gall-ducts that their discrimination is for a time impossible; but the phenomena which soon follow clear up the obscurity.

In some cases of inflammation of the biliary ducts, especially where an occlusion of the ducts takes place, a peculiar paroxysmal fever is developed, with temperature ranging from 103° to 105°, which might readily be mistaken for a malarial outbreak. This *hepatic fever* is generally ushered in by a violent chill, and the paroxysms, which are repeated at irregular times, are apt to be followed by increased jaundice. Their irregularity,—to which, however, there are exceptions in the earlier part of the case,—their resistance to quinine, the frequent occurrence of vomiting and of pain in the region of the liver, and the history of the case, distinguish them from malarial fever. From abscess of the liver the affection is more difficult to discriminate, and we must lay

---

\* Heitter, Wien. Med. Wochenschr., 1887.

stress on the deep jaundice, which mostly happens after the fever outbreaks, and on the different physical phenomena. Sweats occur in both, but they occur only at the end of the marked paroxysm in the so-called hepatic fever. The febrile attacks are explained by Pepper \* as mostly due to a purulent lesion at some point of the biliary canals, or to the development of ptomaines, owing to the ptomaine-destroying function of the liver being interfered with when the bile is pent up. We also find similar attacks of rigors and intermittent pyrexia associated with hepatic pain in obstruction of the common bile-duct from gall-stones. Charcot looks upon them as septic, as does Osler; † Ord ‡ holds the fever-outbreak to be due to local irritation of the mucous membrane, and this is the view I hold. Hepatic fever bears a close relation to malaria. Those who have malarial poison in their systems are more liable to it, and it is likely to be in them connected with a biliary catarrh, and with inspissated bile rather than with an impacted gall-stone.

Now, considering the question of operative interference that may arise, it is of the utmost importance to distinguish the cases in which the obstruction is purely catarrhal and not connected with gall-stones from those in which it is. The most certain test undoubtedly would be having found gall-stones on previous occasions, or finding them after the fever paroxysms. The cases with gall-stones are very much more frequent than the cases of hepatic fever without them; the jaundice is more distinctly connected with the attacks, and generally passes off more completely between them; the pain is greater and ceases more abruptly; and the febrile paroxysms are not brought on by cold, exposure, and fatigue, as they are often in hepatic fever without gall-stones.

### Acute Diseases characterized by a Decrease in the Size of the Liver and by Deep Jaundice.

**Acute Yellow Atrophy.**—This dangerous affection consists in a rapid diminution of the liver, with disintegration in the secreting-cells. Its functions are almost wholly suspended, and

---

\* Medical News, March 29, 1890.

† Johns Hopkins Hospital Reports, vol. ii. No. 1, 1890.

‡ Boston Med. Journ., 1887.

the evil effects of the accumulation of the elements of the bile in the blood show themselves in the deep jaundice and in the profound disturbance of the nervous system. To this disease belong most of those cases of malignant jaundice which terminate rapidly in death after violent cerebral symptoms. The malady scarcely ever lasts a week ; generally a few days only elapse before the patient becomes comatose and dies.

The complaint is sometimes ushered in by nausea, a coated tongue, irregular action of the bowels, and a frequent pulse ; at other times it begins abruptly with pain in the head, and with vomiting, at first of the contents of the stomach, but soon of coffee-ground material, which is evidently altered blood. The skin is yellow, and becomes from hour to hour more discolored. Jaundice is, indeed, never absent : it may not make its appearance before the other urgent symptoms, but sometimes it precedes the signs of serious difficulty for several days, or even for longer,—perhaps for upwards of two weeks.\* That the jaundice is not due to obstruction is proved by the stools containing bile. There are not uncommonly pain at the epigastrium and in the hepatic region, muscular and arthritic pains, dyspnoea, meteorism, enlargement of the spleen, epistaxis, and hemorrhage from the bowels. The pulse exhibits extraordinary changes : it is generally very rapid, but sinks at times, without any assignable reason, to a normal frequency ; during the deep coma of the last stages of the malady the beat of the artery is apt to become slow and full, but it may be very quick and very small. There is fever, not, however, active ; the temperature may be, indeed, after the early stages of the disease, below the norm until towards the end, when it has been known to be  $104^{\circ}$  or  $105^{\circ}$ . The surface may be covered with petechiæ. But, if we except perhaps the deep jaundice and the lessening hepatic dulness, the most significant symptoms are those referable to the nervous system. Severe headache, delirium, involuntary discharges, tremors, spasms, convulsions, or a constantly-increasing stupor and sluggish pupils, show clearly what disturbance the poisoned blood is creating in the nervous centres.

Acute atrophy of the liver scarcely happens in children or after

---

\* As in Observation No. XVII. of Frerichs on Diseases of the Liver.

forty years of age ; it is much more common in women than in men. We find it not unusually following violent mental emotions or drunkenness and venereal excesses ; or it occurs during pregnancy, and is then accompanied by renal disorder.

Now, how does this fatal malady differ from *acute inflammation of the liver* ? By the marked jaundice, the cerebral symptoms, the rapid diminution in the volume of the liver, the dry, brown tongue, the frequent, changeable pulse, and the occurrence of hemorrhages. Then the circumstances under which acute atrophy makes its appearance are very dissimilar. Indeed, the diagnosis is not generally a difficult one,—not nearly so difficult as between acute atrophy and typhoid fever, or between the former affection and yellow fever or certain local diseases, such as peritonitis, pneumonia, and meningitis, when accompanied by jaundice and delirium. The character of the eruption, the presence of diarrhœa instead of constipation, the milder nature of the mental wandering, the significant temperature record, and the slower progress of the disease are of much value in enabling us to distinguish between *typhoid fever* and the typhoid symptoms of acute yellow atrophy of the liver. From *yellow fever*, acute atrophy differs by the epidemic character of the former, by the injected eye, by the intense pain in the back, limbs, and forehead, by the stages the febrile malady presents, by the decided fever temperature, by the comparative absence of cerebral symptoms, and by the enlargement rather than the lessened size of the liver.

From the other affections named, the hepatic disorder may be discriminated by a thorough examination of the various organs of the body, and by a careful weighing of all the symptoms. In truth, it is thus only that we can avoid error ; since, unless we can establish the most positive sign of acute atrophy, the diminution of the area of percussion dulness of the liver,—and there are cases in which we cannot establish this, particularly if there have been enlargement from previous disease,\*—there is no manifestation of the hepatic malady that may not occur in the diseases mentioned, when they are complicated by jaundice. It is true that vomiting of blood is scarcely among their symptoms ; but this does not invariably happen in acute atrophy. In many cases

---

\* As in a case in my ward at the Pennsylvania Hospital.



of doubt we may seek in the urinary secretion for the sediments of tyrosine or for leucine; and the test for urea, which is greatly deficient or absent. So may be the uric acid, the chlorides, the sulphates, and the earthy phosphates. We may in this connection remark that leucine and tyrosine have been also found in the blood and in many tissues. This was observed in a case which I saw with Dr. H. C. Wood, and which he has carefully reported.\*

Acute yellow atrophy may happen occasionally in children.† An affection like it occurs from phosphorus-poisoning; and indeed there are those who believe that acute yellow atrophy is really due to phosphorus accidentally introduced into the system.‡ The occurrence of the fatal malady in pregnant women has already been referred to. Jaundice from mental emotion, or produced by the pressure of the gravid womb, is in them not unusual; and we may be called upon to distinguish this harmless form of icterus from that of yellow atrophy. In the serious derangement of the nervous system, and the graver character of all the symptoms, lie the marks of separation.

### Chronic Diseases attended with Enlargement of the Liver, and with slight or no Jaundice.

**Chronic Congestion.**—This morbid condition is observed chiefly in persons of sedentary habits, or in those who indulge too freely in the pleasures of the table, or use large quantities of alcoholic drinks or fermented liquors. It is frequently met with in hot climates and in malarial districts. It may also occur in scurvy, and in connection with abdominal affections which interfere with the portal circulation, or it may happen in consequence of a disturbance of the flow of blood through the liver, dependent upon disease of the heart.

Whatever the source of the hyperæmia, the symptoms are similar. They are impaired appetite, bitter taste in the mouth, a coated tongue, flatulency, a feeling of tension and weight in the

---

\* Amer. Journ. Med. Sci., April, 1867.

† Duckworth, St. Barthol. Hosp. Rep., vol. vi.; Tuckwell, *ib.*, vol. x., 1874.

‡ Perls, Handb. d. Allg. Pathol., i., points out an anatomical distinction: in acute atrophy there is fatty degeneration; in phosphorus-poisoning the liver-cells are only infiltrated with fat.

right hypochondrium, depression of spirits, loss of strength, impoverishment of blood, deposits of lithates from the highly-colored urine, headache, dry cough, and occasional nausea and diarrhœa, or looseness of the bowels alternating with constipation, and, in protracted cases, hemorrhoids. The conjunctiva has constantly a more or less jaundiced tinge; the dulness on percussion in the hepatic region is increased in extent. In some cases the habitual congestion leads to an altered condition of the bile-ducts and of the secreting-cells of the liver; but ordinarily, unless the hyperæmia be kept up by some exciting cause which it is impossible to remedy,—such as an abdominal tumor, or an organic affection of the heart,—it can be removed. A troublesome feature of the malady is its disposition to return.

By attention to the signs mentioned, there is usually little difficulty in recognizing chronic hepatic congestion. How it may be discriminated from other forms of enlargement of the liver, we shall presently inquire. It is sometimes confounded with, or rather there is sometimes mistaken for it, a liver which has been pushed downward by the habit of *tight lacing*. But the absence of any signs of hepatic derangement, and the lowered outline of the upper border of the displaced right lobe, will generally enable us to distinguish this state from chronic congestion of the liver.

Chronic hepatic congestion, as indeed any disease of the liver which leads to its enlargement, may be confounded with *chronic gastritis*. But the outline of the dulness when the liver is increased in size, the jaundiced hue of the conjunctiva, the altered character of the stools, and the less marked gastric symptoms will enable us to arrive at a correct diagnosis. Yet we must not forget that the two morbid states may be conjoined.

*Hypertrophy of the liver* may present the manifestations of congestion. The little we know of an increased formation of the liver-cells teaches us that this may happen as a partial hypertrophy, to compensate for loss of substance, in instances in which a portion of the gland has been destroyed; or as a more general increased growth in diabetes, in leukæmia, and as a consequence of malaria. Perhaps the history of the case may enable us to arrive at the discrimination of the rare disease. Yet there is never any certainty in the diagnosis.

So-called *torpor of the liver*, in which there is supposed to be a

deficient excretion of bile, has much the same symptoms as congestion. Indeed, it is a question whether this is not often present as at least a secondary result. In persons of middle life who eat freely and take too little exercise in the open air, or those of sedentary habits in whom anxiety and worry have lowered the nervous tone, the well-known symptoms of headache, languor, depression of spirits, loss of appetite, drowsiness after meals, sallow hue of skin, dingy conjunctiva, urine depositing lithates, stools black and offensive, or more often pale or whitish, bespeak this "bilious" state, and we can only distinguish the functional disorder from the ordinary forms of chronic congestion by the history, the concurrent symptoms, the tension in the region of the liver, and the enlargement of the organ, which these present.

The symptoms of chronic congestion of the liver, as indeed of other hepatic derangements, show themselves at times more particularly in the *nervous system*. Headache, vertigo, dimness of sight, and noises in the ears are common; and I have often known the same to happen that Murchison states to be not infrequent,—I have known tingling and pricking sensations and a feeling of creeping in the extremities cause needless alarm that paralysis was imminent, and disappear under blue pill and a few saline purgatives. On the other hand, we must be careful not to regard as evidence of an hepatic disorder signs of stomach and liver derangement which are really due to an affection of the nervous system. Twice it has come under my observation that altered character of the stools, bitter taste in the mouth, vomiting, and slight discoloration of the conjunctiva, existing in connection with tumors at the base of the brain, were considered as purely of hepatic origin. Clifford Allbutt\* cites a case of Ménière's disease, in the person of a physician, where the vomiting and giddiness received this false explanation. In such instances, of course, attention to the occurrence of disordered gait, and of the persistent noises in one or both ears, and to the loss of power of hearing of one ear, shown when a tuning-fork is placed in contact with the skull on the affected side, tells the true meaning of the other symptoms.

**Chronic Hepatitis.**—It is difficult to say what are the symptoms of the malady, because most of the chronic affections of the

---

\* St. George's Hosp. Rep., vol. viii.

organ, especially the congested, the fatty, the albuminoid liver, and hypertrophic cirrhosis, are included in its description. The liver is enlarged in size. The inflammation may be chronic almost from its onset, and be developed under much the same circumstances as chronic congestion; or it may succeed an attack of acute hepatitis. But chronic hepatitis is not a common disease, except in hot climates, and is scarcely to be distinguished from persistent hyperæmia of the organ, unless when the inflammation leads to the formation of abscesses.

**Abscess of the Liver.**—Hepatic abscesses may form as the result of inflammation of the liver. In the tropics this is not unusual; in temperate climates we seldom encounter the affection, save as the consequence of embolic or pyæmic inflammation of the liver, or in connection with some disease of the intestines, or of abscesses around the rectum, or as a sequel of gastric ulcer, or of gall-stones which have produced ulceration of the gall-bladder and gall-ducts and secondary abscesses of the liver, or of traumatism, or of suppurative disease of bones.

The symptoms of hepatic abscess are obscure. Sometimes the only symptoms are debility, great irritability of the nervous system, and irregular slight febrile attacks. More usually the formation of pus gives rise to rigors, leads to night-sweats, and not unfrequently to the development of a fever simulating that of a quotidian or tertian intermittent or remittent, and attended during certain hours of the day with considerable elevation of temperature. Jaundice occurs, but is generally slight, and is often absent. There is no enlargement of the abdominal veins, nor is there, save exceptionally, ascites or œdema of the lower extremities. Dry cough, quickened breathing, and gastric disorder, especially loss of appetite, are frequent, and obstinate vomiting, singultus, and meteorism are not unusual. In the advanced stages of the malady typhoid symptoms are apt to develop. But the disease may be latent. The local signs, too, are far from being always obvious, or indeed uniform. In some instances the hepatic region is more prominent than natural, and we can detect fluctuation over portions of the enlarged gland; but neither sign is constant, and the latter depends greatly upon whether or not the abscess is deeply seated. Tenderness, either general or limited, is found only in a certain proportion of cases, especially when



the abscess is near the surface. It is frequently associated with a throbbing or a dull pain, which may be transmitted to the right shoulder. According to Annesley,\* this sympathetic pain in the right shoulder indicates that the convex part of the right lobe of the viscus is affected. Conjoined to the feeling of weight, and to the throbbing in the hepatic region, is at times a tension occasioned by palpation of the abdominal muscles, especially of the rectus. Twining† regards this as very significant of deep-seated abscess. Cyr‡ tells us, with reference to the exact position of the abscess, that when it is in the front convex part of the liver there is pain radiating to the chest and shoulder, dyspnœa, but rarely jaundice; when in the central part of the organ, there are few signs of local affection of the liver itself or adjacent organs, except decided jaundice if the abscess be large. In abscess limited to the under surface, thoracic symptoms are absent, but gastric symptoms, especially uncontrollable vomiting, occur; the pain is apt to radiate towards the groin.

A positive diagnosis of abscess of the liver is often a very difficult matter; for there are a number of affections with which it may be readily confounded. Prominent among these are hydatids, cancer of the liver, actinomycosis of the liver, affections of the gall-bladder, and a pleuritic effusion on the right side.

From *hydatids* of the liver, the febrile symptoms, the disturbed nutrition, and the pain distinguish an hepatic abscess, except in those cases in which the cyst becomes the seat of suppuration. Under these circumstances error can scarcely be avoided, unless we are fully cognizant of the previous history.

*Cancer* of the liver differs from an abscess by its dissimilar history, by the hard nodular masses, and by the absence of fluctuation. It is only in rapidly-growing medullary cancer that we can discern a sense of fluctuation; but even here we can generally distinguish some nodules which do not fluctuate; and should the soft cancerous matter impart a feeling of fluctuation, it is rarely as distinct as that of an abscess. Further, the marked fever and the other constitutional symptoms are not like what occur in hepatic cancer; for in this affection, as in all cancers, the tem-

---

\* Researches into the Diseases of India.

† Diseases of Bengal.

‡ *Traité des Maladies du Foie*, 1887.

perature, except in instances of large rapidly-spreading growths, is but little affected,—may, indeed, be subnormal.

*Actinomycosis* of the liver may give rise to a collection of pus, and the abscess may discharge through the loins or through the lungs, as in hepatic abscess. The hepatic swelling is painful on pressure, but is unlike that of hepatic abscess in arising suddenly from the parts beneath, and in being surrounded by a firm base in the liver. These characters distinguish it from an ordinary abscess as well as from hydatid of the liver.\* Yet it is by the history, and by finding the ray fungus in pus from other diseased parts of the body, that the diagnosis is mostly established, for actinomycosis of the liver is almost never primary.

Of the *affections of the gall-bladder*, the one most liable to be confounded with hepatic abscess is distention. This occurs either from a closure of the cystic or of the common duct, especially the former, or from an inflammation of the gall-bladder itself, and perhaps a subsequent closure of the ducts. In such a case the gall-bladder may become enormously distended with decomposing bile and puriform matter, and thus may be occasioned a fluctuating tumor, tender on pressure, and readily mistaken for an abscess. Now, we are sometimes able to distinguish the soft swelling caused by a diseased gall-bladder by its situation, its pear-shaped form, its mobility and the absence of adhesions to the abdominal walls, its distinct and persistent fluctuations; by its never having been hard; by the normal appearance of the parietes of the abdomen; by the absence of tenderness over the liver, merely tenderness over the tumor being found; and by the fact that affections of the gall-bladder are frequently preceded by repeated attacks of violent pain due to the passage of biliary calculi, or by bilious fever. Then we find little jaundice, or none at all; and no hectic fever. But to neither of these circumstances can we trust implicitly. For there is apt to be intense jaundice in an affection of the gall-bladder, if the common duct also be implicated; and jaundice is, in abscess of the liver, a symptom more frequently absent than present. And with reference to hectic fever, the continued suppuration in the distending sac may produce it, and lead, indeed, to great constitutional dis-

---

\* Harley, Med.-Chir. Transact., vol. lxi., 1886.

turbance.\* Further, these *biliary abscesses* may, like hepatic abscesses, open externally, or burst into the chest. At times the communication is with the bronchial tubes, and gives rise to very anomalous symptoms. Thus, Simmons† details a case in which there was a circumscribed tumor in the epigastrium, fluctuating with a sense of intervening air or gas, and resonant on percussion; a blowing sound was distinctly discerned synchronous with the respiratory act, and occasionally accompanied by a gurgling noise; there were profuse sweats and extreme oppression, but no signs of pneumothorax.\* At the autopsy a biliary abscess was found communicating with the right bronchus.

As regards the shape of the swelling due to an enlarged gall-bladder being diagnostic, we must bear in mind that it may be changed by contraction of the muscular coat.

A *pleuritic* effusion on the right side is distinguished from an hepatic abscess by the same phenomena that we found, in discussing pleurisy, to separate this affection from all forms of enlargement of the liver. But abscesses of the liver may open into the right pleural cavity. Then we observe the physical signs of a pleuritic effusion subsequent to those of hepatic abscess. Generally, too, the pus which has made its way through the diaphragm destroys the lung-texture, until it reaches the bronchial tubes, when large quantities of purulent sputa are expectorated; in rarer instances it is discharged through the walls of the chest. In the former case, the accumulation of pus in the pleura may be very limited; the inflammation of the pleural membrane may be circumscribed, while the signs of an inflammation at the lower portion of the right lung, dulness on percussion, tubular breathing, and rusty-colored sputa, are evident. These phenomena may subside, and the respiration in parts become inaudible, when a discharge of a large quantity of a reddish or whitish pus takes place, in which the elements of bile and the microscopical appearances of the hepatic tissue may be detected. Gradually this expectoration ceases, and the affected textures heal. But in some instances the discharge never stops, and the patient dies worn out by the constant drain.

---

\* As in a case reported by Pepper the elder, Amer. Journ. Med. Sci., Jan. 1857.

† Amer. Journ. Med. Sci., Oct. 1877.



In *subphrenic pyopneumothorax*, cavities full of air form beneath the diaphragm and extend into the thorax. When situated on the right side they may be mistaken for the breaking of an hepatic abscess into the chest. The history of the affection is generally significant; the subphrenic abscesses are the result of perforating ulcers of the stomach or of the duodenum, and their development is preceded by the symptoms of general peritonitis or by the discharge of pus by the bowels. The signs of pneumothorax, as Leyden\* has found, subsequently show themselves, with distinct metallic tinkling and succussion sound; yet, while all breath-sound is sharply cut off below the fourth or fifth rib, up to this point the normal vesicular murmur is heard on deep respiration, and there are no signs of pressure in the pleural cavity or of distention of the chest, and the marked alteration, by change of position, of the dulness on percussion, from the exudation at the lower part of the chest, is strictly limited to this part. The liver reaches to the umbilicus or lower, and when a canula is passed into the cavity beneath the diaphragm and a manometer is attached, inspiration shows increased pressure, expiration the reverse,—exactly opposite, therefore, to what happens if the canula be in the pleura.

When an hepatic abscess forces its way *externally*, it may, prior to its discharge through the thoracic or abdominal walls, occasion difficulty in diagnosis from abscesses originating in these walls. Nothing but a careful consideration of the attending symptoms and of the history of the case will lead to a differential distinction. Nor does the difficulty wholly cease when the slowly-developed tumor, which an hepatic abscess forms, has opened, since it is far from always that we find in the pus the evidences of the broken-down liver-tissue, and it is only occasionally that the fluid is of yellow or greenish color and yields the reactions of bile. The means of discrimination most to be relied upon is a probe; for by the depth to which it can be passed, the direction it takes, and the feel of the structures it encounters, we are placed in possession of many important facts bearing on the diagnosis. In doubtful cases, also, we may employ the aspirator, and a chemical and microscopical examination of the pus, other than that oozing

---

\* Zeitschrift für Klin. Med., Bd. i.



out of the opening, may tell the nature of the abscess. Indeed, the aspirator may be made a means of diagnosis of abscess of the liver under some of the circumstances above mentioned, where abscess is closely simulated by other hepatic affections. If no abscess be found, no particular harm results from the exploration; nay, it has even been affirmed that the local depletion does good.\*

Occasionally a *hernia through one of the recti muscles* is mistaken for a projecting abscess of the liver. I was called some years since to see such a case, in which the opinion that it was an abscess of the liver had been long entertained. The sound of the mass on percussion; the clearly-defined limits of the liver; the absence of hepatic and gastric symptoms,—taught the true nature of the malady.

Much has been said of the distinction between the abscesses which are developed in the course of embolism or of pyæmia—"the pyæmic abscess"—and the abscess, common in tropical climates, which forms as the result of hepatitis, "the tropical abscess." This kind of abscess is often met with following dysentery. One of its forms occurs in connection with the *amœba coli*, though we may have abscess of the liver due to the *amœba* without dysenteric symptoms. The points of distinction between pyæmic and tropical abscess may be thus tabulated:

## PYÆMIC ABSCESS.

Many in number; small in size.  
Uniform enlargement of liver; only exceptionally bulging of ribs.  
No fluctuation; always pain and tenderness.  
Jaundice present in the majority of cases.  
Enlargement of spleen usual.  
Rigors and night-sweats marked; great tendency to symptoms of blood-poisoning.

## TROPICAL ABSCESS.

Usually a single large abscess, seated in right lobe, towards the convexity of the liver.  
Enlargement not uniform; bulging of ribs, or in epigastrium, or in right hypochondrium.  
Fluctuation usual; pain and tenderness always absent.  
Jaundice exceptional.  
Enlargement of spleen unusual.  
Rigors and night-sweats less marked; obstinate vomiting often present.

---

\* Maclean, *Lancet*, July, 1873.

## PYÆMIC ABSCESS.

Course rapid; three weeks to three months.

Arises after external injuries and operations, or internal suppurating cavities or ulcerations, such as ulcers of the stomach or gall-bladder.

## TROPICAL ABSCESS.

Course less rapid; often extends to three or six months, or longer.

Arises in tropical climates, chiefly in free livers; dysentery frequently coexists.

**Fatty Liver.**—A fatty liver occurs in drunkards; in persons who lead indolent lives and are large eaters; in wasting diseases, especially in phthisis; in the course of protracted diarrhœa; and sometimes in children after exanthematous fevers. But of all these causes, pulmonary consumption is the most common.

A knowledge of the sources of fatty liver is the most important element in the diagnosis; for neither the physical signs nor the symptoms present anything which is characteristic. The physical signs are simply those of an enlarged liver; the enlargement is generally moderate and uniform, and the lower margin rounded. In thin persons it may be possible to discern the doughy consistence of the organ. The symptoms are much the same as those of hepatic congestion, except that there is perhaps greater tendency to diarrhœa, and that we find in some instances a pale, greasy-feeling skin. There is neither pain nor ascites. The amount of jaundice is always very slight; in truth, jaundice is most frequently wanting.

**Waxy Liver.**—A peculiar infiltration into the structure of the liver, or its degeneration into a substance rendering it firmer and more glistening, gives rise to that appearance of the liver which is variously designated as waxy, lardaceous, or amyloid liver.

The symptoms of a waxy liver are those of an hepatic derangement which manifests itself rather by the signs of disturbance of other organs than by the direct proof of altered function of the viscus really affected. Thus, disordered digestion, nausea, vomiting, tympanites, discolored stools, and diarrhœa are much more frequent than jaundice, which, indeed, is very much oftener absent than present. There is a feeling of fulness in the hepatic region, but little or no pain; while physical exploration exhibits an increased percussion dulness, and shows the dense organ to have a well-defined though somewhat rounded margin. The enlargement is uniform, but considerable; at times so great that the liver oc-

cupies a large part of the abdomen, producing a visible bulging. The smoothness and the regularity of outline are lost if waxy liver coexist with diseases of the liver which may harden the organ in nodules, such as cancer, fibroid changes, or cirrhosis.

Enlargement of the spleen is commonly associated with the enlargement of the liver, and in many cases the urine is albuminous from waxy disease of the kidneys. Dropsy, as a rule, is not encountered; but in this respect much depends upon the state of the kidneys and of the blood, or upon the existence of secondary peritonitis.

Waxy liver is much more common in males than in females. It is usually caused by constitutional syphilis or coexists with scrofulous diseases of the bones, with unhealed ulcers, especially rectal ulcers, with discharges from or collections of pus in various parts of the body, with repeated attacks of intermittent fever. In some instances it is associated with cancer or with phthisis, or it results seemingly from the abuse of mercury. We cannot trace always the pathological process to any known cause; but we always find it attended with signs of impaired nutrition and occurring in persons evidently cachectic.

The disease is one lasting for years. In advanced cases, besides the spleen and the kidneys, the stomach and the intestines are apt to be implicated; looseness of the bowels, with dysenteric symptoms, arises, and the skin and breath have a musty, disagreeable odor.

Now, when we contrast a waxy liver with other hepatic complaints in which the liver is enlarged, we find it resembling most closely the *fatty* and the *syphilitic* affections. But in the former, although there is enlargement, there is not often so much increase in volume as in the waxy liver. Besides, the organ feels softer on palpation, and the disorder is not associated with a diseased spleen or kidney, and is much less likely than a waxy liver to give rise to dropsy. Then the history of the case is very significant. A syphilitic hepatitis, with which indeed the waxy liver is at times combined, is mainly distinguished by the prominent nodules felt on the surface of the liver. From *congestion of the liver*, waxy liver is readily discriminated. A comparatively slight affection in which jaundice is frequent is very different from a malady in which the hepatic disease forms but part of a

general cachexia and in which jaundice is very infrequent. In *leukaemic liver* we may have considerable and smooth enlargement and marked cachexia. But the history of the case and an examination of the blood tell its true nature.

**Cancer of the Liver.**—In cancer of the liver the organ is almost invariably large, and sometimes it reaches an enormous volume. The form of the gland, too, is altered. It is irregular and uneven, nodules of various size being developed in its substance and projecting from its border and surfaces. These prominences are harder than the surrounding hepatic tissue; but there are exceptions to this rule, for sometimes, especially in the encephaloid variety, the elastic tumors impart, when pressed, a very deceptive sense of fluctuation. The cancerous masses increase, and in some cases with great rapidity.

The malignant disease is rarely confined to the liver; it frequently supervenes upon cancer of the mammary gland or of the uterus or of the stomach. It is an affection pre-eminently of middle life or of old age; yet it occasionally occurs in young persons. I have met with two cases of primary cancer of the liver in women not twenty-five years of age, and two in children. In primary cancer of the liver we generally find a history of cancer in the family; and protracted grief or anxiety, Murchison tells us,\* precedes the development of the malady, whether a family taint can be traced or not. The disease rarely lasts beyond a year, and it may run a rapid course.

Now, many of the pathological facts just mentioned have a strong bearing on the diagnosis of hepatic cancer. They especially throw light on the most important signs of the malady,—to wit, the increased percussion dulness in the hepatic region, and the uneven surface detected on palpation. The enlarged liver is found extending across the epigastrium far into the left hypochondrium; it reaches at times lower than the umbilicus, and presses the diaphragm upward. The nodules can often be felt distinctly through the abdominal walls. The diseased organ is painful, and tender to the touch. In cases in which the peritoneal covering is affected, the tenderness is greatest. And, although any of these three phenomena—the enlargement, the uneven

---

\* Lectures on Diseases of the Liver, 2d edit.



surface, and the tenderness—may be absent, they are tolerably constant attendants on cancer of the liver. The tenderness is, I think, the sign least frequently wanting.

Among the symptoms of hepatic cancer, we find gastric and intestinal disturbances; pain in the right shoulder; rigidity of the abdominal muscles; a disordered nutrition of the whole body; a cachectic look; occasional febrile attacks, yet, on the whole, normal or subnormal temperature; and, in the later stages, sometimes hemorrhages from the stomach or bowels, and diarrhœa. Ascites, too, is observed among the symptoms of the malignant malady, and is generally dependent either upon chronic peritonitis attending the development of the cancer, or upon the pressure this exerts upon the larger branches of the portal vein. Jaundice may or may not be present; it is most frequently wanting. I have seen it, however, intense when the cancerous growth presses on the bile-ducts: in any instance in which it occurs it persists until death. There are cases in which all these symptoms are perceived; while in others only some occur, and in others, again, even these few may not be well defined. Indeed, when we consider the amount of deposit which is generally present; when we regard its character; when we take into account the necessarily impaired function of one of the most important glands in the body; when we reflect upon the pressure which the enlarged organ must occasion,—it is truly astonishing that often so little dropsy, so little jaundice, so little pain, so little constitutional disturbance, are produced by the disease.

Yet in point of diagnosis we can generally discern the malady by the combination of the symptoms and signs indicated. It is only at an early stage of the disease, or when the liver is not enlarged, that we are apt to be in doubt. Under the former circumstance, a swelling in the hepatic region, pain upon pressure, associated with retching, with nausea and vomiting, and with failing health and strength, occurring in a person above forty years of age, may well excite our suspicion. But, unless there be a history of cancer in the family or a cancer in some other part of the body, we cannot be certain that the beginning swelling in the right hypochondrium is malignant. When the liver is the seat of cancer, but is not increased in size, the recognition of the malady is next to impossible. In these obscure cases, the persistent tenderness in

the hepatic region, accompanying the evidences of disturbed function of the liver, ascites, anæmia, and a cachectic appearance, are the signs most trustworthy and most likely to lead to a correct conclusion. In any instance, jaundice coming on in a person over forty years of age, lasting for months, and associated with gastric disease and failing health, must, in the absence of a history of gout or of syphilis, be looked upon as pointing to hepatic cancer. Again, we must remember that loss of flesh and of strength not unfrequently precedes jaundice and pain,—in fact, all signs of disorder of the affected organ.

Let us pass in review the complaints with which well-marked cancer of the liver may be confounded. Omitting, because elsewhere discussed, hydatids, abscess of the liver, and cirrhosis, they are :

WAXY LIVER ; FATTY LIVER ; CHRONIC CONGESTION.

ACUTE CONGESTION ; ACUTE HEPATITIS ; CATARRHAL JAUNDICE.

SYPHILITIC LIVER ;

AFFECTIONS OF THE GALL-BLADDER ;

CANCER OF THE STOMACH ;

CANCER OF THE OMENTUM ;

ENLARGEMENT OF THE RIGHT KIDNEY.

*Waxy Liver ; Fatty Liver ; Chronic Congestion.*—A waxy liver presents often as much increase in size as cancer ; moreover, like cancer, it is associated with evident signs of cachexia. The main points of distinction are the smooth feel and uniform increase of the liver in waxy disease, its painlessness and slow progress, its combination with enlargement of the spleen and albuminous urine, and the history of the case pointing to constitutional syphilis, or to diseases of the bones, or to long-continued suppuration,—in fact, to the causes which generally lie at the root of a waxy or lardaceous state of organs. In the differentiation of cases of infiltrated cancer without distinct nodules, the physical exploration does not aid us, and we have to lay stress on the other points.

A fatty liver is easier to discriminate from hepatic cancer. The occurrence of the non-malignant malady in consumptives or in drunkards, and the total absence of pain,—in truth, of any decided indications of hepatic disease, except increased size of the organ,—

enable us to distinguish between the two affections with certainty. The slighter signs of disturbance, both constitutional and local, the dissimilar history, and the uniform enlargement of the liver separate chronic congestion from cancer. As a mark of distinction, too, of the cancerous from all of these non-malignant disorders, Virchow lays stress on the existence of swollen jugular glands; and a small cancerous induration in the abdominal walls, around the umbilicus, also not unfrequently aids the diagnosis.

*Acute Congestion; Acute Hepatitis; Catarrhal Jaundice.*—It is rarely indeed that these ailments are confounded with cancer of the liver, because the history and the course the latter malady takes are so dissimilar to those of an acute hepatic disorder. Yet there are cases in which the malignant disease is either developed with great rapidity, thus simulating an ordinary acute affection, or has lain dormant and passed unnoticed until it begins suddenly to increase. Under such circumstances, even, we may be able to recognize the malignant complaint, if its physical phenomena be well defined; but if these be not clearly marked, the diagnosis becomes one of great difficulty.

To cite a case in illustration:

A married woman, twenty-five years of age, was admitted into the Philadelphia Hospital on January 14, 1862, with jaundice and slight fever. She stated that she had been in excellent health until about two weeks before, when she caught cold by sleeping in a damp apartment. Her appetite and digestion had been good previous to her present illness, and she had been fully able to perform her household work. Since she was taken ill she had noticed a feeling of weight in the region of the stomach and liver. Rales indicative of bronchitis were found in the chest, and the impulse of the heart was feeble. The hepatic percussion dulness was increased in extent, especially that of the left lobe; but the outline of the organ appeared regular and even. Tenderness of the abdomen, more particularly in the epigastrium and right hypochondrium, was also noted. There was nausea, but no vomiting; the tongue was clean; the evacuations were discolored.

Now, here was certainly a patient presenting none of the signs of hepatic cancer, except, perhaps, the tenderness over the enlarged gland. Yet at the autopsy, which was made within a week

after her reception into the hospital, and therefore not three weeks from the apparent beginning of the complaint, whitish nodular cancerous spots, many of them soft, were found in the substance of the liver, but not at its edges, nor forming anywhere distinct protuberances which could have been detected during life.

To the similarity of certain cases of protracted catarrhal jaundice in elderly persons, presenting emaciation, with nausea, retching, and vomiting, we have above alluded. The physical signs of the enlargement of the liver may or may not assist us, according to their character.

*Syphilitic Liver.*—As a consequence of constitutional syphilis, the liver may at times exhibit cicatrices on its surface, and scattered nodules, consisting of connective tissue, and extending into the parenchyma. This condition is styled syphilitic inflammation of the liver, or the syphilitic liver. The organ becomes uneven from the contraction of the cicatrized parts, and is apt to be somewhat increased in size, from coexisting waxy degeneration or interstitial hepatitis. The patient has a pale, cachectic look, but is not jaundiced,\* except from a temporary catarrh of the bile-ducts, produced by the syphilitic poison; nor is dropsy present, unless there be at the same time an affection of the kidneys or enlargement of the spleen. But the most important elements in the diagnosis are the age of the patient, the history of the case, and the detection of syphilitic cicatrices in the throat. When contrasted with cancer, we find, besides these points, the chief distinctive marks to be: the much more usual absence of jaundice and of dropsy, the not uncommon increase in size of the spleen, the want of local hepatic tenderness,—unless this be due to passing attacks of perihepatitis,—and the smaller size and softer feel of the nodules. Syphilis of the liver may be hereditary.†

*Affections of the Gall-bladder.*—Dilatation and cancer of the gall-bladder are both very liable to be mistaken for cancer of the liver. The former affection may result from occlusion of the he-

---

\* No jaundice is mentioned in the cases of Dittrich, Prag. Vierteljahrsschr., Bd. vi. and vii.; of Gubler, Mémoires de la Société de Biologie, tome iv.; of Bamberger, Krankheiten der Leber, in Virchow, Pathologie, etc.; or of Moxon, in Guy's Hospital Reports, 1867. In the cases of Murchison, Diseases of the Liver, 2d edit., 1877, it was a passing or an absent symptom.

† Arch. Gén. de Méd., June, 1884.



patic and common bile-ducts, produced by pressure of surrounding tumors or by an impaction of gall-stones; or it may be owing to the distention of the bladder with an albuminous fluid,—the so-called dropsy of the gall-bladder. In either instance the bladder may attain an enormous volume, and give rise to a marked tumor at the lower margin of the liver. The prominence is apt to be rounded or pear-shaped, and, except in those cases in which the occlusion is in the cystic duct or at the neck of the gall-bladder, the impediment to the flow of bile is accompanied by intense jaundice and by decided hepatic swelling. Hence, in the deep hue of the skin, the uniform enlargement of the liver, the peculiar contour of the prominence, the absence of ascites, the paroxysms of pain preceding, not following, as in cancer of the liver, the other marked symptoms, and the history of the case, which not unfrequently points to repeated attacks of colic from the passage of gall-stones, we find the clue which permits us to determine that we are not dealing with hepatic cancer.

*Cancer of the gall-bladder* is scarcely ever met with in young persons, and is, as a rule, associated with cancerous formations in the liver or in other organs. It is difficult to make out a certain diagnosis of the affection, for it presents a strong likeness both to cancer of the pyloric extremity of the stomach and to cancer of the liver. From the latter it is undistinguishable, unless the situation and form of the tumor be such that we can clearly recognize it as belonging to the gall-bladder. Sometimes it is preceded by a history of gall-stones.\* Jaundice, as in cancer of the liver, may be absent or present: in five cases reported by Bamberger† it was found in all, and was even intense. Frerichs, on the other hand, states that in most instances it is wanting. Musser‡ finds it reported in sixty-nine out of a hundred cases. In sixty-eight out of one hundred cases analyzed by him a tumor was discovered, the position of which is most frequently in the right hypochondrium and the umbilical region, and which is painful on pressure. There is also generally gradually-increasing pain and a sense of weight in the right hypochondrium. The disease is more common in women than in men. The signs of the cancerous cachexia

---

\* Murchison, *op. cit.*

† Krankheiten des Digestions-Apparates.

‡ Transact. Assoc. Amer. Phys., vol. iv., 1889.

are always strongly marked; as a rule, more strongly than in hepatic cancer.

*Gall-stones* occasionally accumulate in the gall-bladder in such numbers as to give rise to a hard, even nodulated swelling, which may be mistaken for cancer. But the tumor is generally movable, is not painful on pressure, and does not alter in size, or does so but slowly. Sometimes the patient complains of the feeling of a weight rolling from side to side when he turns in bed, and on palpation a crackling sound is produced, which is readily discerned with the stethoscope. Generally we obtain a history of bilious colic. There may or may not be jaundice; there is an absence of the cachectic symptoms of cancer.

*Cancer of the Stomach.*—This is discriminated from cancer of the liver by the far more constant vomiting, by the dark appearance of the ejected matter, by the more obvious symptoms of indigestion, the persistent pain in the stomach, or the pain radiating from there to either hypochondrium. Moreover, the seat of the tumor is different; it is epigastric, or extending downward, but not often passing into the right hypochondrium, and it shows on percussion a very different contour from an enlarged liver. Yet there are cases in which we are kept in doubt; especially those in which the left lobe of the liver chiefly is affected with cancer and presses upon the stomach, inducing perhaps—and thus making the likeness still closer—obstinate vomiting. The only traits of distinction are then found in the presence or absence of marked derangement of the functions of the liver, and in the chemical examination of the contents of the stomach.

*Cancer of the Omentum.*—The absence of jaundice, and the unaltered appearance of the stools, are here, too, of great value in indicating that a tumor near or joining the left lobe of the liver is not due to cancer of that viscus. Moreover, the boundaries of the morbid mass are different from those of a diseased liver. But we cannot always trust to this. Cancerous tumors of the lesser omentum may so surround the liver, and correspond so closely to the regular form produced by hepatic cancer, that the two maladies cannot be distinguished; at least not by the local signs. Again, a loop of intestine may be thrust across the enlarged liver at a point corresponding to the usual limit of the percussion dullness of its left lobe, thus dividing the most prominent nodules

from the greater portion of the viscus, and making it appear as if the tumor were to the left of, and below, the stomach, and belonged, therefore, probably to the omentum.\* In such cases we have to depend entirely upon the signs of disturbed liver function.

*Enlargement of the Right Kidney.*—A tumor formed by an enlargement of the kidney does not present the same outline of percussion dulness as a cancerous liver. The dulness is, moreover, bounded by the tympanitic sound of the intestine, and is not lowered by a deep inspiration; and the signs of disturbed function of the kidney, and an examination of the urine, will generally materially assist the diagnosis. Still, cases may occasionally happen in which, owing to a peculiar shape of the diseased kidney and to the obscurity of the symptoms, an error in diagnosis can scarcely be avoided.† The difficulty in discrimination is heightened by the circumstance that many cases of morbid growth of the kidney, at least of one-sided growth sufficient to give rise to a palpable tumor, are cancerous, and are therefore, so far as the manifestations of a cachexia go, similar to cancer of the liver.

Finally, in reviewing the diagnosis of cancer of the liver, we must inquire whether other than cancerous growths, such as sarcoma, melano-sarcoma, myxoma, epithelioma, cysto-sarcoma, angioma, lymphadenoma, can be distinguished from true cancer. They may produce identical physical signs and symptoms; indeed, a distinction is impossible, unless the history of the case enable us to make it. Much the same may be said of that rare disease, tubercular formations in the liver. Leukæmic livers may attain enormous size, and be mistaken for cancer; and the cachexia that attends them makes the error more likely. But the swelling of the spleen and of the lymphatic glands and the microscopical examination of the blood furnish the points in diagnosis.

**Hydatids of the Liver.**—The development of one or of several cysts in the liver, containing within them echinococci, is

---

\* See case, Proceedings Pathological Society of Phila., vol. i. p. 275.

† Vidal (Bulletin de la Société Médicale des Hôpitaux, 1874) cites errors in diagnosis between tumors of the kidneys, especially hydronephrosis, and diseases of the liver attended with enlargement, like abscess or cancer, made by such masters in our art as Velpeau, Nélaton, Gosselin.

not, as a rule, a disorder which occasions serious disturbance of the general health. Nor do the hydatids usually give rise to either jaundice, dropsy, or any marked signs of gastric or of intestinal irritation, or to fever, or to local pain. Their most constant manifestations are a decided increase of the size of the liver, and the presence of elastic tumors discernible in the hepatic region. In some instances xanthelasma has been noticed.\* This disorder of the skin, however, is not peculiar to hydatids, but has been observed in connection with other forms of hepatic enlargement associated with chronic jaundice.

The growth of the hydatid is generally very slow, and usually in one direction only,—upward, downward, laterally. Very commonly the hydatid tumor grows from the right lobe. In most cases it attains considerable dimensions, and the liver may be found to encroach upon the lung as far as the second intercostal space, or to extend far down into the abdominal cavity. On percussion, the line of dulness either of the upper or of the lower boundary of the viscus, or of both, is perceived to be very irregular, and occasionally on striking a series of abrupt blows on the pleximeter, or on the fingers of the left hand used as such, we discern a peculiar vibration, similar to the sensation perceived on striking a mass of jelly, and very significant of the existence of the cyst. Owing to the pressure the increasing tumor may exert on adjacent structures, we observe in some cases dry cough; palpitation and displacement of the heart; vomiting; possibly jaundice and ascites.

Hydatids ordinarily last for years. The echinococci may die, the sac become much reduced in size, or obliterated, and recovery take place; or the cyst may discharge its contents through the stomach and intestines, through the bronchial tubes, or through the walls of the abdomen, and the patient then gets well. But so favorable a termination cannot be counted upon. A fatal issue may at any time ensue by the hydatid tumor bursting into the pleura, or the pericardium, or the peritoneum, and leading to violent inflammation; or by suppuration occurring in the sac, when the symptoms become those of pyæmia. Even when the hydatids are discharged through the stomach, intestines, bronchial tubes,

---

\* Duckworth, St. Bartholomew's Hospital Reports, vol. x., 1874.



or abdominal parietes, recovery is apt to be slow; nor is it, indeed, unusual to find the patient's strength giving way before the contents of the sac have been entirely voided and it has closed.

In some countries hydatids are frequent; it is not so in this country. In Iceland these growths developed from the eggs of a tape-worm are so common that they cause one-seventh of the human mortality. In point of diagnosis, it is not generally difficult to detect the presence of hydatids. It is true that when these are small or deep-seated it may be impossible to discern them. But a large and superficially-seated hydatid tumor can usually be distinguished, and can be separated from the maladies to which it bears a resemblance. It differs from an *abscess of the liver* by the want of febrile action, pain, and great constitutional disturbance; indeed, the latent character of the hydatid tumor becomes of much importance. Its slow growth, too, is very significant, much more so than the physical characteristics, which are here not to be trusted to. When, as sometimes happens, a hydatid tumor inflames and suppurates, we have nothing to guide us in the differential diagnosis but the history of the case previous to the development of the urgent symptoms. From *cancer of the liver* we distinguish hydatids by the long duration of the case, by the absence of evident cachexia, of local tenderness, and of the unevenness of the surface which the small, hard, cancerous tumors projecting from it occasion. On the other hand, we have in hydatid tumor the sensation on palpation of elasticity or fluctuation. Under rare circumstances this may happen in medullary cancer, and the rapid growth of the latter and the cachectic symptoms would determine the diagnosis. A *distended gall-bladder* may, like hydatid tumor, be free from pain on pressure, but, unlike this, it is movable, is preceded by attacks of colic, is generally accompanied by deep jaundice, and its situation corresponds to that of the normal gall-bladder.

An *aneurism of the aorta* differs from hydatids in the severe—for the most part neuralgic—pain the patient suffers, so utterly dissimilar to the absence of pain or to the mere feeling of tension and weight of a hydatid swelling. Then the pulsation and the other physical signs aid us. In *aneurism of the hepatic artery*, which may also present a smooth, throbbing tumor, we are apt to have deep jaundice from compression of the biliary ducts.

*Pleuritic effusions* have many features in common with those cases of hydatids of the liver in which the growing tumor extends upward into the chest. All the physical signs of a large effusion may be present, even the dilatation of the thorax and a sense of fluctuation in the intercostal spaces. But the absence of constitutional symptoms, the irregular outline of the dulness on percussion of the hydatid cyst, the great displacement of the heart, and the decided lowering of the upper margin of dulness upon deep inspiration, enable us commonly to detect the real nature of the disease. When the cyst has opened into the lung and the hydatids are being expectorated through the air-passages, the harassing cough, the copious sputum, and the inflammation of the pulmonary tissue which is apt to be occasioned, may cause the affection to be mistaken for pulmonary abscess or phthisis. The surest marks of distinction are furnished by the changed form of the lower part of the thorax, and by finding bile and the hooks of the echinococci in the sputum.

*Renal enlargements*, such as cysts, hydronephrosis, cancer, are discriminated from hydatids of the liver by the same physical signs by which we found them to be distinguished from hepatic cancer,—chiefly by the renal tumor having the tympanitic sound of the colon in front of it, by not being affected in position by deep inspiration, and by the direction of its growth. Moreover, the history and an examination of the urine will greatly assist.

*Ovarian cysts*, unlike hydatids, grow from below upward, are not influenced by deep inspiration, and produce enlargements greatest below and not above the umbilicus; then they have a different outline on percussion from hydatid liver.

But, though we may thus generally distinguish hydatids of the liver from the maladies which have similar symptoms, there are unquestionably cases in which it is extremely difficult to arrive at a satisfactory conclusion. Under these circumstances, an exploratory examination with an aspirator would be proper. We may at times detect shreds of striated hydatid membrane, and portions of echinococci. Besides, the character of the fluid drawn off will assist us materially in diagnosis. It is as clear and colorless as water, has a specific gravity of 1007 to 1011, and contains not a trace of albumin or of urea, but large quantities of chloride of

sodium. No other fluid in the human body, whether in health or in disease, presents these peculiarities.\*

Occasionally portions of the liver are transformed into a mass consisting of connective-tissue stroma and numerous large and small cells filled with a gelatinous substance. The disorder looks like alveolar carcinoma, but it is really *multilocular hydatids*, or echinococcus tumors. The centre of the mass suppurates, but even this does not diminish the great resistance of the hepatic tumor; nor is fluctuation, save in the rarest instances, perceptible. The liver may retain its normal shape, or elevations may be perceptible, such as we observe in carcinoma and syphiloma of the organ: indeed, the affection is not to be distinguished with any certainty from either, except it be by the history and the attending constitutional symptoms. No jaundice usually accompanies the hard hepatic swelling; but in cases in which the bile-ducts are obstructed we meet with jaundice without dyspeptic symptoms or previous paroxysms of pain, and usually without enlargement of the gall-bladder. In cases with icterus, unlike what we find in syphilis or in cancer, there is complete discoloration of the feces.†

Let us now, in concluding the review of the hepatic maladies which are attended with decided increase of the size of the organ, briefly contrast their most important manifestations. We have found that, as regards the enlargement, they differ materially. Simple congestion, chronic inflammation, fatty liver, do not attain nearly the volume of cancer, of hydatids, of abscess, of waxy disease of the liver. The three affections first mentioned differ, moreover, from all the others, except the waxy liver, by presenting a uniform and not an irregularly-shaped swelling or an uneven outline of the percussion dulness.

Concerning the symptoms, we observe that, although these hepatic disorders all agree in not being in any way characterized by *jaundice*, yet this sign is more commonly present and more distinct in some than in others. In hydatids, and in the syphilitic liver, there is no yellow hue of the skin or of the conjunctiva; so,

---

\* Murchison, *Lectures on Diseases of the Liver*, 2d edit., p. 61.

† See the cases of Friedreich and of Niemeyer, referred to in Niemeyer's *Practice of Medicine*.

too, as a rule, in waxy liver. In fatty liver and in abscess it is, on the whole, most frequently wanting. The same may perhaps be said of cancer, though sometimes there is decided icterus in this malady. In chronic congestion and in chronic inflammation we ordinarily find jaundice, though it may be but a slight yellow tinge of the skin and the eye. With reference to *dropsy*, we are not apt to encounter it in any of the hepatic affections under consideration except cancer, and waxy disease, when more than the liver is implicated. It is in these two complaints, also, that the most obvious signs of a cachexia are met with; while in abscess we find fever, and perhaps the greatest constitutional disturbance.

As regards *pain*, the fatty liver, hydatids, simple hypertrophy, and the waxy liver are painless; while, generally speaking, congestion, catarrhal inflammation or obstruction of the bile-ducts, chronic hepatitis, intestinal hepatitis, hepatic abscess, and cancer, are more or less painful affections.

### Chronic Diseases attended with Decreased Size of the Liver, and with Abdominal Dropsy.

**Cirrhosis.**—A liver reduced in bulk, very dense and hard, exhibiting granulations of various size separated by bands of fibrous tissue, and surrounded by a thickened serous envelope, presents the morbid state known as cirrhosis, or hob-nail liver. The inflammation which leads to these alterations in the fibrous tissue is generally consequent upon the abuse of spirituous liquors. But this cause does not explain all cases: in some, the malady is connected with disease of the heart; in others, with constitutional syphilis; in others, with anthracosis; in others, again, it cannot be attributed to any known agency. Sometimes it is combined with fatty or waxy degeneration. Again, there may be granular livers in which the fibroid matter preponderates and which never contract,—an interstitial hepatitis, or hypertrophic cirrhosis. The disease is essentially a disease of middle-aged men; it is far less common in women, and rare in children.\*

In the first stage of cirrhosis, the organ is somewhat increased

---

\* See, however, cases by Howard, Transact. Assoc. Amer. Phys., 1887.



in size; then, as Glisson's capsule thickens more and more, the bulk becomes lessened. It is, however, doubtful whether the stage of enlargement invariably precedes that of shrinking: the process of reduction constitutes not unfrequently the first change. But, without entering into this question, we may state that there are no symptoms by which we can recognize the disease at an early period, for the symptoms at first are the same as those of chronic congestion,—dull pain, perhaps tenderness at the hypochondrium and pain referred to the shoulder, disordered digestion, and a sallow or a jaundiced hue of the skin. Nor can we say, even after the stage of contraction is fairly developed, that the diagnosis of the affection is always possible. It may rest on no stronger grounds than finding in a person who is known to be a spirit-drinker, “a tippler,” an intractable ascites, without any obvious cause to account for the dropsy. The dropsy, due to the obstruction of the portal circulation, consists throughout strikingly of ascites; as it increases, œdema of the legs may be developed, and passing albuminuria, from pressure on the renal veins.

Besides the dropsy, the other clinical features of the malady are not very marked. The most significant signs consist in the diminution of the percussion dulness in the hepatic region, and the detection, by the touch, of firm, irregular granulations on the margin and under surface of the liver. But both these signs are very difficult to discern, on account of the distention of the abdomen with fluid, and the displacement of the liver this may occasion. In fact, it is often only after the performance of paracentesis that the abdominal walls will permit us to judge with any accuracy of the shrinking and altered state of the organ. This is especially true with reference to palpation; as regards percussion, it may be possible, even when the abdomen is still full of dropsical effusion, to detect the lessened extent of the hepatic dulness. In rare cases cirrhosis happens without abdominal dropsy.\*

Irrespective of these phenomena, we find at times other manifestations of disease which assist us in the diagnosis of cirrhosis. They are enlargement of the spleen; dilatation of the veins of the abdomen; gastric and intestinal derangements; hemorrhoids; marked loss of flesh and strength; jaundice; a decidedly

---

\* Arch. Gén. de Méd, Nov. 1886.

cachectic appearance, with sunken features; and hemorrhages from the nose and mouth, or from the stomach, or into internal cavities. Hæmatemesis in an alcoholic must always arouse suspicion. The increase in size of the spleen is far from constant, and rarely reaches a considerable extent. The dilatation of the abdominal veins is not perceived until an advanced stage of the disease, and is sometimes connected with a peculiar vascular net-work, stretching from the umbilicus upward and downward, and, as Sappey \* was the first to describe, with a decided enlargement of the epigastric and mammary veins, the blood flowing through the former in a reversed direction from what it does in health,—namely, not toward the liver, but from it to the veins of the abdominal wall, and thence to the vena cava. Other external veins share in the enlargement; the veins of the legs may be varicose, and the venous twigs on the cheeks become developed. In some cases an irregular but moderate fever not exceeding  $102.5^{\circ}$  is also noticed; generally there is none.

Another symptom to which I have had my attention strongly directed is the presence of small amounts of sugar in the urine. Thus, in two cases which I saw with Dr. Simpson, Trommer's test readily detected sugar in the urine. In the one case the secretion was scanty; in the other it was abundant. One had lasted for several years, and was slowly developing; the other had existed about sixteen months, and was rapidly progressing.

Cerebral symptoms due to a toxic cause sometimes appear. They show themselves frequently in a wandering delirium of mild type, attended with confusion of persons and places. The delirium is often like that of uræmia, but there is nothing in the urine to account for it. It may not show itself until toward the end of the disease; on the other hand, it may be of long duration. In a case I saw with Dr. Lloyd, it lasted four months. Coma and convulsions also occur occasionally.

The gastric and intestinal derangements, the result of a congested or inflamed mucous membrane, are rarely wanting: they manifest themselves by failing appetite, impaired digestion, both gastric and intestinal, morning sickness, flatulency and constipation, or the frequent voiding of pale-colored stools. The jaundice

---

\* Bulletin de l'Académie de Médecine, tome xxiv.

does not often attain a high degree; when it does, it has a bad meaning. It shows itself usually in a yellowish tinge of the skin and conjunctiva; but in some cases even this hue is absent, and we find the pale skin and pearly eye of anæmia.

Yet not one of these symptoms is really characteristic; they become so only when viewed in connection with the dropsy, with the local signs in the hepatic region, with the history of the case, and with the absence of any organic disease of the stomach or the intestine, which might explain them. Then the age of the patient, generally above thirty-five years, and his habits, must be taken into account. The cirrhosis of young children is generally due to inherited syphilis. Gout seems to predispose to the disease. Murchison tells us that the condition of the liver which develops gout renders it liable to suffer from alcohol. Cirrhosis of the liver often becomes associated with acute tuberculosis. At times cirrhosis runs a rapid course.\*

There is a form of cirrhosis due to infection. It has been described as *subacute infectious hepatitis*.† It is attended with irregular fever of remittent or intermittent type, with decided enlargement of the spleen and splenic pain, with urobilin in the urine, with greatly-lessened renal excretion of urea, but ureic sweating, with slight jaundice and cirrhotic diminution of the size of the liver. The infection occurs, probably, through the intestine, and from the liver spreads along the hepatic veins to the vena cava, and may ultimately infect the arterial system, giving rise to infectious nephritis and purulent meningitis.

A very similar disease is met with in children, a cirrhosis with jaundice after infectious maladies, such as scarlet fever or measles.

Another form of cirrhosis, if it be a form and not a separate disease, by comparison rare, has been mentioned,—*hypertrophic cirrhosis*, or “interstitial hepatitis,” or cirrhotic enlargement. It has the same symptoms as atrophic cirrhosis, and is undistinguishable, except by the increased percussion dulness it presents, and by the signs of enlarged liver being usually attended with more jaundice and greater tendency to slight febrile attacks, and

---

\* Hanot, “Cirrhose atrophique à marche rapide,” Arch. Gén. de Méd., June, 1882.

† Levi, Arch. Gén. de Méd., April, 1894.

to peritonitis.\* A peculiar mawkish odor of the breath has been spoken of as present.†

But, with reference to these symptoms, there are forms of hypertrophic cirrhosis with but slight jaundice, without ascites or marked development of the abdominal subcutaneous veins, terminating in a slow cachexia. Generally, however, the disease begins with the signs of congestion, acute or chronic, with jaundice, and with some pain in the right hypochondrium, and lasts for years; at the end there is marked jaundice, and the patient sinks into a typhoid state. Ascites may, as already indicated, be wanting throughout, or, as is more usual, it comes on late in the malady. The disease is, in my experience, not unfrequently complicated with a fatty liver, forming "a fibro-fatty liver." As regards the cirrhotic state in the markedly-enlarged liver, it is asserted that besides the increase of fibrous tissue, both within and without the lobules, the smallest biliary ducts are much developed.‡ In some instances of hypertrophic cirrhosis there is organic disease of the heart. The infectious nature of hypertrophic cirrhosis has been of late often affirmed.

Cirrhosis of the liver due to *malarial infection* is also associated with enlargement, at times very great. It presents, moreover, a persistent chronic jaundice, which may last for years, and is combined with marked enlargement of the spleen and manifestations of the malarial poisoning. Bleeding from the nose, gums, and intestines is frequent; dropsy and distention of the abdominal veins are absent.§

Let us now look at the distinction between ordinary cirrhosis and some of the maladies which resemble it; and first let us compare its traits with those of other *hepatic affections*. From diseases of the liver attended with enlargement, such as waxy liver, fatty liver, and chronic congestion, fully-developed cirrhosis is discriminated by the presence of ascites and the other signs of seriously-obstructed portal circulation, by the diminished, or certainly not augmented, size of the organ, and by the different history of the disorder. From *hydatids* of the liver we diagnosticate cirrhosis

\* Hayem, Archives de Physiologie, Jan. 1874.

† Duckworth, St. Bartholomew's Hospital Reports, 1874.

‡ See an excellent review by Hanot, Arch. Gén. de Méd., Oct. 1877.

§ Lancereaux, quoted in Sajous's Annual, 1888, p. 335.



by the irregularity of outline of the enlarged liver in the former complaint, by the sense of fluctuation, and by the comparatively unimpaired general nutrition of the body. *Cancer* of the liver is unlike cirrhosis in the distinctness and size of the protuberances, in the obvious hepatic enlargement, in the less marked or absent ascites, and in the normal size of the spleen. But when a cirrhotic liver is associated with syphilitic nodules, or when its volume is augmented by waxy infiltration, the discrimination from cancer becomes a matter of extreme difficulty; indeed, it may be impossible to avoid erroneous conclusions. Hypertrophic cirrhosis may also be very difficult to distinguish from cancer, except by the history of alcoholic dyspepsia and the enlargement of the veins, and, though large and nodulated, the liver is rarely so tender.

We shall now consider and compare the clinical traits of some diseases of the liver producing, like cirrhosis, atrophy of the organ.

As the result of repeated attacks of perihepatitis, we find great thickening of the capsule, with fibrous bands passing into the interior of the organ, and some atrophy. This condition, described as *simple induration of the liver*, is met with chiefly in connection with constitutional syphilis, though it is also seen following a right-sided pleurisy and diseases of parts contiguous to the liver, producing inflammation which spreads to it. The affection is not to be distinguished from true cirrhosis, except by the causing elements, particularly by the syphilitic history, and by the absence of the habit of spirit-drinking; the greater and more persistent pain and tenderness in the hepatic region are of significance; sometimes there is coexisting heart disease.

In *red atrophy*, too, we have greatly-diminished hepatic dulness with the symptoms of portal obstruction; it, too, is therefore undistinguishable from cirrhosis by the symptoms alone, unless the difference may be thought to consist in the doubtful points of far less frequent or decided jaundice and in outbreaks of diarrhoea. But, in reality, the only traits of importance on which to base a diagnosis are that the dense, reddish, homogeneous liver occurs not preceded by alcoholic dyspepsia or valve disease, but generally in those with a most marked history of malaria or of dysentery or of ulceration of the intestine.

*An inflammation of the portal vein, with coagula forming in it,* may occasion the same manifestations of deranged abdominal circulation, the same or greater tumefaction of the spleen and decrease of the liver, as cirrhosis. And what complicates the diagnosis very much is, that cirrhosis is the chief disease which leads to thrombosis of the portal vein. Indeed, we cannot, under any circumstances, positively discriminate this affection from cirrhosis. Still, we are sometimes enabled to distinguish the venous disorder by laying stress on the sudden development of the symptoms, especially of the engorgement of the portal system, and by noting the rapidity with which the ascites returns after paracentesis, the copious gastric or intestinal hemorrhage, the severe vomiting and diarrhœa, the great enlargement of the abdominal veins, and, when not too soon fatal, the marked emaciation. Other causes, of course, than inflammation of the coats of the vein produce coagulation. We may have thrombosis from mere weakness of the circulation, or as the result of disease of the liver structure or of compression by enlarged cancerous or tubercular glands, or in consequence of the perforation of the vein by gall-stones. Compression of the portal vein and of the biliary ducts in the fissures of the liver, from inflammation of the surrounding areolar tissues, may be separated from cirrhosis chiefly by the intense icterus and the complete decoloration of the stools.

Of non-hepatic affections, cirrhosis is most liable to be confounded with *chronic peritonitis*; a mistake rendered the more likely because chronic congestion or even chronic inflammation of the peritoneum may exist as a complication of cirrhosis. But, even when no such complication is present, the diagnosis may be difficult. It rests chiefly upon the greater and more extended tenderness of the abdomen in peritonitis, the febrile signs, the absence of splenic enlargement and of dilated veins, the usually unchanged, or certainly not jaundiced, hue of the skin, the association with signs of disease in other viscera, especially of the lungs,—for chronic peritonitis is generally tubercular.

Under rare circumstances, *cancer of the stomach* may simulate cirrhosis. I had some years since a case under my charge at the Pennsylvania Hospital, in which, with very slight digestive symptoms, and without discernible epigastric tumor, considerable ascites and effusion into the left pleural cavity existed. Owing

to this effusion, the state of the spleen could not be accurately ascertained. There was some fulness of the abdominal veins, and the hepatic percussion dulness did not extend entirely to the margin of the ribs. Bile-pigment was present in the urine, the bowels were loose, and progressive emaciation ensued. The man had been very intemperate, and his case might certainly have been selected as an illustration of cirrhosis; yet at the autopsy the liver, though small, rather hard, and deeply congested, was not cirrhotic, and a cancer involving the whole stomach, except the pylorus, was found.\*

**Chronic Atrophy of the Liver.**—Although cirrhosis is the most frequent it is not the sole cause of dwindling of the liver. We have just spoken of its diminution in consequence of obstruction of the trunk of the portal vein, as well as of other causes; but besides these causes we find some, such as a decrease of the organ from long-continued closure of the common duct, or its atrophy in old age, or in connection with grave disease of the heart or lungs obstructing the circulation and causing long-standing hyperæmia of the liver, or as an accompaniment of chronic disease of the intestine. The first of these morbid states is mainly discriminated by the deep jaundice; the second, by the absence of any important symptoms referable to the liver and associated with the diminished hepatic dulness; the third, by the history of the case and the physical signs of cardiac or pulmonary difficulty, the more general dropsy, or at least by the œdema of the legs preceding the ascites. The fourth form, partly already mentioned under red atrophy, which it may become, presents the phenomena of cirrhosis, and cannot be distinguished from this unless the surface of the liver can be distinctly felt through the abdominal walls and ascertained not to be irregular. We may sometimes suspect the cause of the shrinking of the organ from the persistent and intractable diarrhœa and disturbance of the stomach. But, on the whole, this decrease in size of the liver following gastro-enteric inflammation is not frequent: in truth, there is no cause of simple atrophy of the liver so common as thrombosis of the portal vein.

---

\* For a fuller report of this case, see Proceedings of the Pathological Society, *Amer. Journ. Med. Sci.*, vol. lii., 1866.

## SECTION IV.

## ABDOMINAL ENLARGEMENT.

In describing the causes of abdominal enlargement, I shall view them as they occasion a general and uniform or a more circumscribed and partial swelling.

## General Abdominal Enlargement.

**Ascites.**—The collection of serous fluid in the peritoneal sac gives rise to dropsy of the belly, or ascites. This may form part of a general dropsy, and be dependent upon an organic disease of the kidneys or of the thoracic viscera, or the accumulation of liquid may be confined to, or occupy principally, the abdomen. In either case the local signs are much the same. They are: enlargement of the belly; a dull sound on percussion, due to the presence of liquid; and the sense of fluctuation imparted to the hand on one side of the abdomen by a wave of fluid put into motion by a tap on the other side.

As regards the former of these signs, it is uniform and progressive, and is generally very evident; although, of course, when the quantity of liquid is small, enlargement of the abdomen may escape detection. The percussion dulness is most readily perceived at the lower portion of the abdomen, where the fluid gravitates, unless when prevented from so doing by being circumscribed by peritoneal adhesions. The bowels float usually to the upper part of the liquid, and at this spot their tympanitic resonance may be distinctly discerned. When the patient is in the erect position, the intestinal percussion note is commonly discoverable in the epigastric and umbilical regions. If he be placed upon his back, the tympanitic sound is, for the most part, found to extend lower than the umbilical region, while dulness will be elicited in the hypogastric region and the flanks. If the person affected with ascites be placed upon his side, the flank which is uppermost becomes resonant. This alteration of the level of the fluid with the change of position is thus a significant sign, and always happens except when the effusion is encysted; it is detected without



difficulty, save where great flatulent distention of the bowels or impaction of fæces accompanies the accumulation of liquid.

Ordinarily, the fluctuation wave felt by the hand is easily discerned. It is obscured by thickening of the abdominal walls from œdema, or from the accumulation of fat in the subcutaneous tissues; it is, moreover, indistinct if adhesions circumscribe the fluid in the peritoneum. The amount of albumin in the fluid rises with the ascites and its duration. For all practical applications the specific gravity determines the proportion of albumin, and the urinometer may be employed for the purpose.\*

There are no means of distinguishing the character of the fluid except by direct observation. It must be inferred from the attending symptoms. Chyliform ascites has been not unfrequently found associated with tubercle.†

The other symptoms often found in ascites, such as a pushing upward of the liver, spleen, and stomach, embarrassed breathing, compression of the lungs, and digestive disturbances, need not be specially described, as they present nothing characteristic. Nor is it necessary to insist upon the self-evident fact that a diagnosis of ascites is only half a diagnosis, and that we should in every instance endeavor to ascertain the cause of the collection of fluid in the peritoneal sac; and we may at once proceed to consider the morbid states with which dropsy in the peritoneum is liable to be confounded. They are chiefly:

OVARIAN DROPSY;

CHRONIC PERITONITIS;

DISTENTION OF THE BLADDER;

GRAVID UTERUS;

CHRONIC TYMPANITES.

*Ovarian Dropsy.*—It is not until an ovarian cyst rises above the brim of the pelvis that it occasions a swelling marked enough to be mistaken for abdominal dropsy. Supposing that it has led to considerable enlargement of the belly, we are yet able to discriminate between the two disorders by attention to the physical signs of the history of the case.

---

\* Runeberg, "Eiweissgehalt der Ascitesflüssigkeiten," Deutsches Archiv f. Klin. Med., September, 1883.

† Busey, Amer. Journ. Med. Sci., Dec. 1889.

As regards the former, we perceive these differences : the sound on percussion over an ovarian cyst is dull in the umbilical and hypogastric regions, while at the sides the tympanitic resonance of the intestines may be obtained. Moreover, when the patient assumes different postures the dulness in ovarian dropsy does not change its position ; and, like all ovarian tumors, the ovarian dropsy causes a projection in the centre of the abdomen, not a flattening there and a bulging of the flanks, as is common in ascites. Bacelli\* states that in ascites there is a deep tympanitic sound during percussion in the region of the intestines, while an ovarian cyst presents dulness on the side in which the cyst has its origin, and a tympanitic sound on percussion on the other. In ascites, vaginal and rectal touch detect fluctuation at once, and the uterus is normal in size and in mobility, sometimes it is prolapsed ; in ovarian dropsy, fluctuation is less distinct, and may not be reached at all, or may not exist in case of polycyst, and the uterus is generally displaced behind the cyst.

The fluctuation from an ovarian cyst is apt to be very unequal at different parts of the distended abdomen. When the effused fluid is free in the peritoneal cavity, fluctuation may be perceived beyond the line of dulness as the fluid is thrown in waves among the intestines ; but when it is confined within a cyst, fluctuation cannot be perceived beyond the cyst-walls : hence the outline of the cyst as obtained by percussion, and that of the area within which fluctuation is perceived, must be the same. It should be remembered, however, that fluctuation in an ovarian cyst may entirely escape detection on account of the great thickness of the cyst-walls, or of the unusual tenseness of the cyst, even though it be large, or of the great density of the fluid, or of the small amount of fluid in each cyst. In ovarian cyst there is, for the most part, impairment of the general health, and the color of the face is that of cachexia. Lastly, the pulsations of the aorta are transmitted by an ovarian tumor to the anterior surface of the abdomen, and can be there felt by the hand.

When, however, there is ascites complicating an ovarian tumor, the diagnosis is very difficult. Finding the fluctuation unequal, and an irregular outline of the ovarian growth, may aid us ; but

---

\* Wien. Med. Wochensch., April, 1890.

a preliminary tapping, though now mostly condemned by gynaecologists, may be necessary to arrive at an opinion. Entire reliance cannot be placed on the chemical character of the fluid, since the rule that paralbumin is significant of ovarian fluids and fibrin of serous fluids has many exceptions. Spencer Wells\* accepts the presence of the "granular cell," as shown by Drysdale and W. L. Atlee,† to be characteristic of ovarian fluid. This granular cell, as described by Drysdale,‡ is generally round, sometimes oval, varies in diameter from one five-thousandth to one two-thousandth of an inch, is very elevated and transparent, is much smaller and far less opaque than the compound granular cell of inflammation, and contains a number of fine well-defined granules which become more distinct on the addition of acetic acid, and nearly transparent under ether, while the appearance of the cell is not changed. There is no nucleus. In several very doubtful cases of abdominal tumor the diagnostic import of the cell was well attested.§ The cell differs, Drysdale teaches us, from any other granular cell found in the abdominal cavity.

In uncomplicated cases, the history assists us greatly in reaching a correct diagnosis. In ovarian dropsy, we can, as a rule, make out that the distention of the abdomen has begun at its lower portion, and has gradually spread upward, one side being very much more prominent than the other, until the abdominal enlargement has become considerable and the relative position of the umbilicus is altered. Again, we do not find those signs of disease of the liver, heart, or kidneys which are so apt to co-exist with ascites, or that the swelling is temporarily reduced by the use of hydragogue cathartics and diuretics, as in the latter disease.

Attention to the history and progress of the complaint is especially valuable in the class of cases in which the physical signs are modified by the intestines not being able to float to the surface of the fluid in the peritoneal cavity, in consequence of adhesions to one another, or of a diseased omentum, or in which the fluid has

---

\* Brit. Med. Journ., June, 1878.

† Ovarian Tumors.

‡ Transactions of the American Medical Association, 1873.

§ See Transactions of the Pathological Society of Philadelphia, vol. vii., 1877; American Journal of Obstetrics, vol. xii., 1879; also Gynecological Transactions, 1883.

been limited in sacs by inflammatory adhesions. These are cases in which a peritoneal inflammation has led to the effusion of liquid ; and the history of antecedent peritonitis, or of peritonitis in connection with tubercular disease, will go far toward elucidating the diagnosis. On the other hand, an ovarian cyst may contain air, either from a communication with the intestine, or after tapping and decomposition of the contained fluid, and percussion would then give a clear note in front and a dull note below. Under either of these circumstances physical signs alone could not enable us to make a diagnosis, and we should have to seek further light from the history and the general condition of the patient. This is especially true in the diagnosis between encysted dropsy of the peritoneum and an ovarian cyst. If we obtain by tapping a spring-water fluid, it points to cyst of the broad ligament.

*Chronic Peritonitis.*—Excluding the kind of chronic inflammation which is due to an attack of acute peritonitis passing into a chronic state, let us inquire how cases of chronic peritonitis, in which the disease was gradual in its development, can be distinguished from pure dropsical effusion.

Now, these cases of chronic peritonitis are, with the exception of those unfrequent instances of chronic diffused peritonitis of latent origin which we have already discussed, almost invariably associated with tubercle or with cancer, and only under rare conditions with chronic dysentery and dilatation of the colon. In *tubercular peritonitis* the malady generally occurs in those who have at the same time tubercles in the lungs or enlarged caseous glands ; and when we find such patients complaining of abdominal pain and uneasiness, of soreness to the touch, of nausea and vomiting, of diarrhoea alternating with constipation, of having more or less fever, and of losing flesh and strength ; when we discover the tender abdomen to be tense and much distended, in part with liquid, but especially with wind, and sometimes very resistant to the touch, and exhibiting on its exterior the tracings of the convolutions of the intestines ; when in addition there is œdema of the lower limbs, and we find the fever to be irregular, at times high, at times almost ceasing, and a growing cachexia ; when we are able to exclude as the cause of the dropsy disease of the heart, disease of the kidneys, and cirrhosis of the liver, —we can hardly



be wrong in presuming the signs of chronic peritoneal inflammation to be owing to the presence of tubercle. Even when the signs of disease of the lungs are wanting, or are not well defined, we shall generally be correct, if the abdominal symptoms mentioned exist, in determining the peritoneal affection to be tubercular. In some instances the tubercular abdominal disorder develops with rapidity, and the disease has not so much the aspect of a chronic as of an acute complaint. The tumefaction and tension of the belly may be so great as to simulate an abdominal tumor.\*

A *cancer of the peritoneum* gives rise to many of the same phenomena as tuberculous disease. But the affection is far less common, and there is this difference: the malady usually happens consecutively to an external or an internal cancer, and scarcely ever save in persons advanced in years; there is little or no fever, or, indeed, a subnormal temperature; no diarrhoea, or but little diarrhoea, and no profuse sweats, occur. Pain, on the other hand, or at least attacks of spontaneous pain, are more frequent; the lymphatic glands enlarge; and, as the omentum is the most common seat of the cancerous growth, we can generally detect a tumor stretching across the upper portion of the abdomen, and extending perhaps from the epigastrium nearly to the pelvis. The morbid mass is unequal, and usually detected readily, except where separated by fluid from the abdominal parietes. Hemorrhage into the abdominal cavity or the effusion of bloody serum occurs here as it does in tubercular peritonitis. In cancerous peritonitis the ascitic fluid has a turbid gray look. In the sediment that forms there is a rich cell-growth with many red blood-corpuscles. The cells are for the most part peculiar, large swollen nucleated cells, in size like those of the white corpuscles of the blood.† In primary cancer of the peritoneum, or that following *cancer of the retro-peritoneal glands*, the diagnosis is very obscure, unless the tumors are marked. The cancerous malady is apt to pursue a slowly progressive course, lasting months; but it may develop as an acute miliary disease. Retro-peritoneal tumors

---

\* See case in Liverpool Hospital Reports, 1868.

† Runeberg, *Deutsches Archiv f. Klin. Med.*, Sept. 1883; also Coe, *New York Med. Journ.*, July, 1888.

may be readily mistaken for diseases of the liver. They may occasion jaundice from pressure on the common duct. The fact that they do not move with the acts of breathing, as well as that there is often a line of resonance between the dulness they occasion and the liver dulness, is a point of value in diagnosis.\*

*Distention of the Bladder.*—This may give rise to a sense of fluctuation and to very marked abdominal enlargement; so marked, indeed, that patients have been tapped, under the supposition that they were laboring under dropsy of the abdomen. But when the bladder is so much distended as to simulate ascites, there is usually more or less tenderness on pressure over the seat of the obvious swelling; which, moreover, presents a rounded outline of dulness on percussion. Again, we have the history either of retention or of apparent incontinence of urine.† But, to avoid all possible chance of error, in any case of doubt a catheter should be introduced into the bladder. This mode of procedure, it may here be mentioned, is the one which leads most speedily and decisively to a true appreciation of the abnormal phenomena in those rare cases of anasarca which are produced by distention of the bladder, and of which Trousseau has recorded several.

*The Gravid Uterus.*—A gravid womb is readily distinguished from abdominal dropsy by the peculiar form of the dulness on percussion, its steady and uniform increase corresponding to the enlargement of the womb, the absence of fluctuation, the detection of the sounds of the foetal heart, the alteration in the color and appearance of the mammary areola, and the production of movements in the womb on making an examination per vaginam.

*Chronic Tympanites.*—Great prominence of the abdomen, due to flatulent distention of the bowels, is, if at all persistent, very apt to be mistaken for ascites. But the large abdomen yields not a dull, but everywhere a tympanitic sound, and there is no fluctuation. Then the history of the case and the attending symptoms throw light upon the nature of the ailment.

Besides the complaints just reviewed, which are those most commonly confounded with ascites, there are a few very rare disorders

---

\* Vander Veer, Amer. Journ. Med. Sci., Jan. 1892.

† In a case recorded by Watson, in his Lectures on the Practice of Physic, although the bladder was enormously distended, large quantities of urine were constantly passing from the patient.

which might be mistaken for collections of fluid in the peritoneal sac. They are: dropsy of the womb; dropsy of the Fallopian tubes; dropsy of the omentum; very large serous cysts in the kidney; hydatids of the liver, of size so great as to lead to general abdominal distention; and a dilatation of the stomach so extensive that the viscus occupies almost the whole abdomen. With reference to the latter affection, we may distinguish it from ascites by the history of the case and the vomiting and other marked gastric symptoms, by the extended tympanitic percussion note, by the indistinct fluctuation, which is not noticed except over the most dependent part of the organ, by the splashing or the metallic or amphoric sounds which are perceived when its contents are agitated, by the length to which the stomach-tube can be introduced, and by the chemical examination of the gastric contents. The other maladies mentioned can be separated only by taking into account their history and progress, and by laying stress upon the absence of those morbid states which generally cause ascites, and upon the occurrence of special phenomena which point to the structures implicated.

**Chronic Tympanites.**—A collection of gas in the cavity of the peritoneum is of rare occurrence, but is frequent in the intestinal tube, and the accumulation becomes sometimes a chronic condition, and leads to very great and uniform enlargement of the abdomen. We find this form of tympanites in some cases of hysteria; in instances of constriction of portions of the intestinal canal, in consequence either of cicatrization, or of cancer of the bowels, or of their compression by a morbid growth, or of great dilatation of the colon; as a sequel of enteritis or peritonitis, or of a spinal lesion; and we also observe it in persons whose digestive powers are weak and who partake much of food—such as cabbages, beans, and peas—which is apt to occasion flatulency.

Among soldiers this chronic tympanites—owing, perhaps, in many cases to the character of their diet and consequent digestive disturbances—is far from being an uncommon disorder, and may be a very obstinate one. It gives rise to abdominal enlargement, which is constantly mistaken for dropsy, but which does not yield a sense of fluctuation, or return on percussion any other than a well-marked tympanitic sound. The distention produces, moreover, an inability to take active exercise, sensations of cutting pain

under the ribs, and palpitation of the heart ; pressure on the abdomen occasions much discomfort ; the soldiers, therefore, walk with their clothes unbuttoned, and find it very irksome to wear their belts. They are sometimes troubled by indigestion, and feel particularly uncomfortable after meals ; or the symptoms of indigestion, although they may have been present at the beginning of the complaint, disappear, but the swelling of the abdomen persists for many months. According to my experience, the ailment is always gradual in its development.

### Partial Abdominal Enlargement.

**Abdominal Tumors.**—Even at the risk of some repetition, it is for clinical purposes a matter of convenience to point out connectedly the relations an abdominal swelling is likely to bear to the normal structures of the abdominal cavity, and to consider, moreover, the swelling as constituting the starting-point of our diagnosis.

Let us first examine the meaning of an abdominal tumefaction occupying solely or principally one region of the abdomen.

*Right Hypochondrium.*—The most usual cause of a tumor in this region is an enlargement of the liver. Sometimes a tumor which is in the lower part of the right hypochondrium, or proceeds from the termination of this region, is simply a displaced liver, or an affection of the gall-bladder. In the first instance, the recognition of the disorder—such as a pleuritic effusion—which has given rise to the displacement ; in the second, the history of the case, the shape of the swelling, and the symptoms attending it,—will give us an insight into its cause. Again, a tumor in the parts mentioned may be due to an enlarged kidney, cancerous or cystic, or especially hydronephrosis. Careful examinations of the urine and the history of the case furnish the most certain means of discrimination. Then we must also bear in mind that all enlarged kidneys displace the bowel in a particular manner ; they press it forward, and the dulness over the tumor is largely mixed with a tympanitic sound, or the dulness is, indeed, not very appreciable.

*Left Hypochondrium.*—The most usual tumors in this region are those produced by enlargement of the spleen. An increase



in size of this viscus, if acute, is generally owing to altered blood conditions and infectious maladies, as pyæmia, puerperal fever, acute tuberculosis, typhoid fever, relapsing fever, or the malarial fevers. The cause of the swelling is disclosed by the history of the case and by the accompanying symptoms.

*Inflammation of the spleen* is an affection very difficult to recognize. The most trustworthy symptoms are: pain in the left hypochondrium, radiating thence in various directions, as far as the left shoulder, and augmented by pressure, especially if the serous envelope be implicated, by coughing, and by a deep inspiration; nausea and vomiting; fever having irregular fits of exacerbation; sometimes delirium, dry cough, and a sense of suffocation. The extent of the splenic percussion dulness is decidedly increased, and, when we are sure that the spleen is not displaced, the suddenly-widened area of dulness forms a most important element in the diagnosis. Splenitis is very rarely primary, generally metastatic. It is often observed to be connected with emboli from endocarditis, and, these being wafted also to the kidneys, albumin and blood are found in the urine. When suppuration in the spleen ensues, the fever may assume a hectic character and the patient lose flesh rapidly, while the spleen increases in size. But there is no certainty in these signs, nor, indeed, in any of the signs of splenic abscess; this may be latent and suddenly rupture into the abdominal cavity or the stomach. Then there may be abscesses around the spleen with manifestations similar to those in its substance or to pyo-pneumothorax.\* An acute enlargement of the spleen may also be owing to hemorrhage from injury.

*Chronic enlargement of the spleen* may be caused by hypertrophy, by waxy disease, by leukaemia and lymphadenoma, by malignant growth, by hydatids, by syphilitic tumor, by congenital syphilis, and by structural changes from malaria. There are scarcely any symptoms characteristic of these states, except the alteration the blood undergoes, as evinced by a diminution of the red globules and an increase of the white; and even this may not be marked. Waxy hue of the face, dropsy, bleeding from the nose, from the stomach, or from the intestinal canal, and digestive disturbances, though far from infrequent, are less constant. In truth, all the

---

\* Zuber, Revue de Médecine, Nov. 1882.

phenomena mentioned, except perhaps the microscopical evidences of deteriorated blood, are, in the recognition of a splenic tumor, of secondary importance as compared with the extended percussion dulness in the splenic region. In some cases the symptoms are very ill defined, and death may result from rupture of varices of the enlarged viscus, without any other signs of a lesion than those of increased size of the organ.\* When enlargement of the spleen has reached a certain point, the organ curves into the hypogastric and right iliac regions, and a notch or notches may be felt on its anterior and inner surfaces.† This sign may be very valuable in distinguishing the enlarged organ from cancer of the kidney, for which it has been mistaken.‡ In some instances enlargement of the spleen is hereditary.§

Having determined the persistent swelling to be due to the abnormal size of the spleen, we must next endeavor to ascertain the cause of it. The history of the case, such as the proof of leukæmia, of protracted suppuration, of malaria, forms, with the coexisting phenomena in other organs, the main element in diagnosis.

A fulness projecting from the left hypochondrium toward the umbilical or lumbar region may be owing to *fæcal accumulations* in the colon. Although these fæcal accumulations do not occur so often in or near either hypochondrium as they do in the iliac regions, yet they are not very uncommon, and we should be on our guard against confounding them with organic disease, whether of the stomach, spleen, liver, kidneys, peritoneum, or ovary. Their irregular outline, a doughy consistence and painlessness, and close attention to the history of the case and to the accompanying disorder of the digestive functions, will generally enable us to detect the true nature of the swelling. But we must not lay too much stress on the non-existence of constipation, for sometimes great irritability of the bowels or persistent diarrhœa is kept up by a large collection of fæcal matter in the colon, and an irritative fever superadded gives a strong resemblance to typhoid.|| Repeated attacks of colicky pains and some soreness to the touch

---

\* Traube, Virchow's Archiv, 1869.

† Fagge, Guy's Hosp. Rep., 1868.

‡ Lancet, July, 1873.

§ Wilson and Stanley, Clin. Soc. Trans., 1893.

|| As in a case seen with Dr. Arthur V. Meigs.

are not unusual in cases of extensive fæcal accumulation, and jaundice and anæmia have been also noticed. Besides looseness and mucus, the stools are apt to show small hard fæcal masses, which may be of leaden hue. In cases of doubt, laxatives, especially castor oil, should be employed before any opinion is given, and with the voiding of large masses of fæces the tumor and the attending symptoms may disappear.

As regards swellings of any kind situated in either hypochondrium, or in fact at any portion of the upper third of the abdomen, it is always to be inquired into whether they are affected by the act of respiration. This, as Kennedy\* has pointed out, is a valuable sign, for if the morbid mass move in consequence of the depression of the diaphragm, it is because structures are involved, such as the stomach and transverse colon, the liver or spleen, which admit of some mobility; whereas a tumor that is uninfluenced must appertain to a fixed part,—for instance, to the aorta.

*Epigastrium.*—The most common cause of an epigastric tumor is cancer of the stomach. The swelling is then associated with the symptoms above described.

But a tumor in this region may be also produced by a *disease of the pancreas*. A swelling occasioned by *fatty degeneration*, or by *uniform simple hardening of the gland*, cannot, as a rule, be discerned at the bedside. In *pancreatic fat necrosis*, the areas of white necrotic tissue are usually also found in the mesentery and in other seats of abdominal fatty tissue. There are no diagnostic signs. In *chronic pancreatitis*, deep-seated epigastric pain with colicky attacks, a large quantity of matter like saliva passed by stool, profuse salivation, sugar in the urine, fatty stools, and jaundice have been observed to attend the appreciable swelling extending across the epigastrium. The association of chronic pancreatitis with diabetes has been frequently noticed. *Suppurative pancreatitis*, as we know from Fitz's analysis, is much more common in women than in men. Though chronic, it begins with sharp epigastric pain and vomiting, and is not unfrequently attended with irregular fever. It may last weeks or months. As regards *cancer*, which can be recognized with more certainty, the most trustworthy symptoms are: a tumor in the epigastric region; pain

---

\* Dublin Quarterly Journal, August, 1864.



there or in the back, not increased by the taking of food, but usually augmented by the erect posture; progressive emaciation and debility; an appetite capricious rather than diminished, and in some instances, indeed, a ravenous desire for food; constipation, and at times, but far from invariably, fatty stools, or fat-crystals in abundance in the grayish stools.\* Besides these indications, we commonly find, as the disease advances, obstinate jaundice and occasional vomiting. Many of these phenomena belong also to cancer of the stomach; in truth, we never can be certain of the existence of the pancreatic malady until we have excluded the gastric affection. In a differential diagnosis of this kind, the early presence and habitual occurrence of vomiting after meals, the sour eructations, the hæmatemesis, the absence of free hydrochloric acid in the stomach-contents, and the absence of jaundice, assist us in locating the seat of the disease in the stomach. *Calculous disease* of the pancreas is a very rare affection. There are, in addition to the dull sense of weight at the epigastrium and other symptoms of pancreatic disease,—such as sugar in the urine, vomiting, fatty stools,—sharp, irregular attacks of paroxysmal pain, due to the passage of calculi. In cases of large concretions these attacks of colic may become associated with jaundice.† Pancreatic calculi may lead to atrophy of the gland and become associated with diabetes.‡

An epigastric tumor is sometimes simulated by a *contraction of the upper portion of the rectus muscle* on palpation; but the swelling soon subsides, especially if rubbed. Occasionally, however, a tumefaction due to contraction of an abdominal muscle may be of some duration.§ I have known a contraction of the rectus muscle in a case of gastric cancer occasion so obvious a resistance and swelling that it was looked upon as due to malignant disease of the intestine or of the peritoneum. Moreover, the rigid muscle gave rise to dulness on percussion. But,

---

\* But collections of fat-crystals, Gerhardt has found, are also detected in the pale stools of icterus without pancreatic disease: when the bile reappears in the stools the crystals are no longer seen.

† Pepper, Medical News, Dec. 25, 1882; and Johnston, Amer. Journ. Med. Sci., Oct. 1883; see also eighteen collected cases in Sajous's Annual, 1889, C-44.

‡ Lichtheim, Berlin. Klin. Wochenschr., 1894, No. 8.

§ Greenhow's cases, Lancet, 1857.



though the phenomena were for a long period a marked feature of the case, it was observable that the muscle was raised and rigid to a decided degree only in certain positions; at all events, that certain positions gave a distinct outline to the swelling, and that the latter then, like the line of dulness, was regular and straight, evidently corresponding to the contour of the muscle. And this occurs in all instances of contraction of the rectus, no matter with what associated.

The muscular contractions are not always confined to one muscle, or to the whole of one muscle, and when irregular, and particularly when associated with tympanitic distention of the intestine, give rise to most of the so-called "phantom tumors" of the abdomen. These swellings are perplexing, and are constantly mistaken for serious abdominal tumors. The history of the case, the absence of grave constitutional symptoms, the most frequent occurrence of the tumefaction in females, especially in hysterical females, and the usually coexisting constipation, furnish us with valuable signs of distinction. But I believe the use of anæsthetics to be the most important means of diagnosis. I was first led to employ them a number of years ago, in a case which had baffled the skill of several eminent surgeons, one of whom had proposed to the patient an operation as the only means of relief from what was considered an ovarian disease. The patient was thirty-one years of age, a widow, and evidently of highly hysterical temperament. She was very subject to constipation; and the swelling of which she complained was of irregular outline and occupied the centre of the abdomen, extending some distance on each side of the median line. It was hard and resisting to the touch, but, on strong percussion, yielded a tympanitic sound. Whenever it was touched she shrank. Thorough relaxation was produced by the administration of ether; the hand could be pressed almost against the vertebral column, and all signs of the tumor disappeared. A complete recovery took place; and thus terminated a case which had lasted for fully one year, and in which it is highly probable, from the fact that the patient was fond of having her urine drawn off by the catheter, and had shown other manifestations of a similar type of hysteria, that the swelling was, in part at least, artificially produced. But in any of the phantom tumors I would recommend the use of anæsthetics

for purposes of diagnosis ; nay, they may be most advantageously employed, for similar reasons, in all cases of abdominal swelling in which the rigid state of the abdominal walls interferes with accuracy of investigation.

In soldiers we at times observe one or several small movable tumors, yielding a tympanitic sound on percussion, in the epigastric or at the upper part of the umbilical region. They are, probably, small portions of intestine which have been pushed between the fasciculi of a ruptured rectus muscle, similar to umbilical hernia.

*Umbilical Region.*—Tumors which are found in this region form, as a rule, merely portions of a swelling that is principally seated in the epigastrium or in the hypochondria, such as cancer of the stomach, of the liver, of the pancreas, or of the omentum, and dilatation of the gall-bladder. The only two affections which are apt to occasion a swelling solely, or at least principally, limited to and perceptible in the umbilical region, are tuberculous disease of the mesenteric glands and a movable kidney.

The symptoms of the former malady, or *tabes mesenterica*, are much the same as those of tubercular peritonitis. Indeed, unless the enlarged mesenteric glands can be felt through the abdominal parietes, the discrimination is uncertain. The abdomen is prematurely large, is slightly tender on pressure, and has often a doughy feel ; the child loses flesh, the digestion is impaired, the evacuations are frequent, liquid, and offensive. It often presents signs of scrofulous or tubercular disease elsewhere ; and under such circumstances we cannot be at a loss in determining the nature of the tumefaction in the umbilical region. The disease is very rare in adults, though it occurs.\* Its simulation, especially in young women, by *pseudo tabes mesenterica*, has been described in reviewing the affections of the stomach.

When the *kidneys* are not firmly held by their attachments, they become displaced, and are apt to give rise to serious errors in diagnosis. The dislocated organ is perceived under the margin of the ribs on the right flank, or in the umbilical region, and sometimes extends across the median line. The mass is easily moved, may be, by careful and methodical pressure, returned to the renal region,

---

\* See case reported by Gairdner, Lectures to Practitioners.

and presents, on palpation and on percussion, the outline of the kidney. The lumbar region yields a tympanitic sound on percussion, and we find less resistance and a slight depression over the usual seat of the organ, which depression is effaced by pressing the tumor into the lumbar region. There is in some instances sensitiveness over the displaced organ, especially after fatigue, or a blow, or strong pressure; and pressure in examining the part gives rise to the same sensation as when the renal region of the non-affected side is pressed; but we never find any disturbance of the urinary functions, nor, in fact, except a disagreeable feeling in walking, does any real inconvenience result from the accident, save in those cases in which the movable kidney has become painful, or, by compressing the vena cava or portal veins, occasions dropsy. Yet we meet with exceptions to the rule that the disorder gives rise to no decided symptoms. Sometimes dyspepsia, especially nervous dyspepsia, is pronounced, as well as intercostal neuralgia, and so-called gastric crises occur marked by vomiting, with severe abdominal pain and fever, or there are attacks simulating renal colic. Further, we may find intermitting hydronephrosis.\* There seems to be a special connection between the disorder and hysteria, gastric dilatation, enteroptosis, and membranous enteritis.

The disorder is most apt to occur after violent exertion, or after many pregnancies, or may be due to attacks of congestion of the organ, or to tight lacing. It is rare in men. The right kidney is oftener movable than the left. A movable kidney is best detected by palpation with both hands while the patient takes a deep breath.

The affection may, of course, be mistaken for any form of abdominal tumor, and if the kidney should have become adherent the diagnosis is uncertain. Generally the disorder can be distinguished by the absence of signs of constitutional disturbance; by the history of the case; and by the physical phenomena mentioned. To these may be added the comparatively slight dullness or rather the tympanitic character of sound elicited, except on very strong percussion, over the seat of the tumor. This is an important fact as regards the discrimination of a movable and *displaced spleen*, in which, as the organ is generally enlarged, there

---

\* Knight, Lancet, Oct. 1893.



is considerable and extended dullness on percussion. Moreover, the history of the splenic disorder, which not uncommonly can be traced to a malarial affection, the usually great tenderness, the nausea, dyspeptic symptoms, and hemorrhagic tendencies which attend the displacement of the spleen, and the notch which can be felt in it, will assist us in our diagnosis.\* A movable kidney may be simulated by *malignant disease of the colon*.†

Yet another of the abdominal organs is occasionally displaced and movable,—the liver. Now, a *movable liver* would be often mistaken for a movable spleen, were it a more common affection. But very few well-authenticated cases are on record.‡ In these the peritoneal attachment of the organ had become lax, usually in consequence of pregnancy; in the hepatic region there was a tympanitic sound on percussion; and in the umbilical region and toward the right flank a solid body was discerned, the upper border of which presented a convex outline, the lower border was in the inguinal region. The displaced organ was easily pushed about, and could be replaced in its proper situation. The spleen was found in its usual seat; the symptoms were merely those of weight and uneasiness in the abdomen. The movable or wandering organ may be painful or painless. It has the physical characters of the liver, and the most certain sign is the detection, on palpation, of the notch between the right and the left lobe and of a zone of tympanitic resonance between the swelling and the lung. The diagnosis is, however, always difficult and doubtful. New growths of the kidney, as a case of Legg's proves, are particularly confusing. In most recorded cases autopsies are want-

---

\* Cases of displaced spleen are recorded by Dietl, Wiener Med. Wochenschrift, No. 23, 1856, also in Archives Générales, 1858, tome ii.; Rokitsansky, quoted in Brit. and For. Med.-Chir. Rev., Oct. 1860; see, too, Clarke, Dubl. Hosp. Gaz., Aug. 1860; Med. Times and Gaz., Nov. 1869; G. Engel, Centralbl. f. Gynäk., Leipz., 1886, x.; Brown, Pacific Med. Journ., San Francisco, June, 1892; Dorsett, Med. Review, St. Louis, Dec. 1891.

† Henry Morris, Lancet, April, 1895.

‡ See Cantani, Ann. Univers di Medicina, 1866; and Meissner's article in Schmidt's Jahrb., 1869, No. 1; also *ib.*, No 2, 1871; Blet, Le Foie mobile, Thèse de Paris, 1876; Legg, St. Bartholomew's Hospital Reports, 1877; Arini, Anales del Círculo Méd. Argentino, quoted in Amer. Journ. Med. Sci., July, 1884; H. W. Seager, Brit. Med. Journ., Lond., 1885, ii.; L. Landau, Deutsche Med. Wochensch., Berlin, 1885, ii.; Richelot, L'Union Médicale, Paris, Aug. 1893.



ing ; and the whole subject is very obscure. The affection is more usual in women than in men, and, besides pregnancy, tight lacing and chronic inflammation of the peritoneum are said to lead to it.

*Lumbar Region.*—Tumors in this region, or on either flank, are occasioned by some morbid growth of the kidney, or by an abscess in it or its surroundings, or in the psoas muscles. Again, they may be due to fecal accumulations ; or, if on the right side, to very considerable increase of the liver ; if on the left, to a greatly-enlarged spleen. To discriminate between these conditions, we have to determine whether the swelling fluctuates or not ; we must also analyze the urine, and inquire minutely into the circumstances preceding and attending the tumefaction. It is thus only that we can attain the necessary data for a diagnosis, which has, indeed, often to be reached by the process of exclusion.

Tumors behind the peritoneum may give rise to a visible prominence in either lumbar region, extending to the upper part of the iliac region. The most common cause of these tumors is *cancer of the lymphatic glands* lying by the sides or in front of the vertebral column. The disease is very difficult of detection. Still, we may suspect its existence if, in a patient who is evidently cachectic, and who is steadily losing flesh and strength, we discover, on deep palpation on one side of the linea alba or in the flank, a tumor which, owing to its being surrounded by intestine, returns a tympanitic percussion sound. In some cases the swelling communicates the beat of the aorta and simulates an aneurism, or it presses on the vena cava and gives rise to enlargement of the abdominal veins and of those of the lower extremities, and to œdema of the legs. The disease may involve the iliac glands and the tumor extend into the pelvis, or it may reach upward to the diaphragm ; and, by the cancer spreading to the posterior mediastinum, it may finally open the aorta, producing hemorrhages precisely like those coming from an aneurismal sac.\*

*Iliac Regions.*—Tumors in either of these regions may be due to many different causes. They are, as we have elsewhere discussed, principally owing to ovarian affections ; to fecal accumulations ; to disease of the large intestine, such as intussusception or cancer ; and to pelvic abscess. Sometimes they are caused by displacement

---

\* Case reported by Haldane, Edinburgh Medical Journal, Aug. 1868.

of the kidney, by enlargement of the spleen, and in women by retro-uterine hæmatocele, or by extra-uterine pregnancy.

The *ovarian tumors* are, as a rule, distinguished from the other disorders mentioned by their more or less globular form, by their movability from side to side or in an upward direction, by their seeming to spring out of the pelvis, and their evident attachment below, by the displacement of the womb, by the comparatively unimpaired general health, and by their indolent and generally painless nature. These remarks do not apply to the very slight swelling occasioned by ovarian inflammation, for here the tumid spot is often the seat of severe pain. The healthy ovary is not sensitive to the touch. To examine the ovary with exactness, the abdominal muscles must be completely relaxed; the patient is placed in the attitude recommended by Marion Sims,—on her back, with the shoulders supported, the legs drawn up so that the heels are a few inches asunder and the thighs fall easily apart.

As ovarian tumors grow and spread upward they give rise to difficulties in diagnosis, which we have already examined into. We may here again mention the manner in which ovarian may simulate *renal growths*. Spencer Wells dwells particularly on the absence of fluctuation in the vast majority of instances of enlarged kidney; on the renal tumor being first detected between the false ribs and the ilium; on the signs in the urine, and on the absence of those changes in the quantity and regularity of the menstrual discharges which are common in ovarian disorders. Moreover, the ovarian growth usually displaces the intestine backward; in the renal growth it is pressed forward; and large tumors of the right kidney ordinarily have the ascending colon on their inner border, while tumors of the left kidney are generally crossed from above downward by the descending colon.

Among the causes of a tumor in either iliac fossa, *retro-uterine hæmatocele* has been mentioned. The tumor, commonly of rounded shape, rises above the brim of the pelvis, but is traceable into it. It forms quickly, and an examination through the vagina detects a boggy swelling in Douglas's cul-de-sac, and at times the grating of the blood coagula; faintness and collapse attend its production. Much the same physical phenomena are presented by the swelling due to *pelvic cellulitis*. But the slow way in which the tumor forms, the presence of that hot, puffy, thickened, brawn-

like condition of the vaginal wall, so dwelt upon by Simpson, the usually greater tenderness of the swelling felt through the walls of the vagina, and the feverishness and constitutional symptoms attending the gradual formation of the abscess, are distinguishing marks, except where the contents of the hæmatocele suppurate, when for a differential diagnosis we may have to rely on the history of the case.

*Hypogastric Region.*—Distention of the bladder and enlargement of the uterus, whether produced by air, by liquid, by a morbid growth, or by pregnancy, are the most usual sources of a swelling in this region. If due to any one of these causes, the outline of the tumor is regular and rounded; and by the aid of the catheter, of explorations through the vagina and the rectum, of the history of the case, and of the attending symptoms, we are generally enabled to arrive at a correct diagnosis.

A tumor in the hypogastrium may also have its origin in splenic enlargement, in diseases of the peritoneum, or in hæmatocele. In the latter case it is apt to be uniform and to extend to the iliac fossæ.

In concluding this sketch of abdominal tumors, we shall briefly glance at those which are likely to occupy more than one region, and sometimes even the whole or the greater part, of the abdomen. In rare instances, a cancer of the liver, or hydatids of that organ, or a fibrous tumor of the uterus, or a solid ovarian growth, or an enlarged spleen,\* or a kidney the pelvis of which has become enormously distended in consequence of obstruction of the ureter, may lead to the formation of a swelling which occupies nearly the entire abdomen. But the most usual cause of so diffuse a tumor is *carcinoma of the peritoneum*.

This affection, when very extensive, may give rise to a uniform swelling stretching across the abdomen, and equally marked on both sides of the median line, or to several small tumors, which are evidently unconnected with any organ beneath. It is, moreover, apt to occasion a peritoneal friction sound, to exhibit a varying resistance to pressure at different points, to lead to ascites, to loss of flesh and appetite, and chiefly, by the peritonitis it sets up, to

---

\* As in the case reported by Porter, Philadelphia Medical Times, June, 1875, in which the spleen weighed twenty-one pounds.



the occurrence of fever. Much the same symptoms may be produced by *hydatid disease of the peritoneum*, though here there is less fever, the swelling may be uniform or even more irregular, the abdominal enlargement greater and painless, and we may be able to detect the hydatid fremitus and the hooklets in the evacuated fluid.\* Yet as regards the hydatid thrill we must bear in mind that a similar sensation is obtained from large parovarian cysts† or from *colloid cancer of the peritoneum*, a sensation of peculiar and very superficial fluctuation,‡ associated, however, here with grave symptoms of cachexia, and generally with a rapidly-spreading growth. *Peritoneal abscesses* enclosed by adhesions will also, if large, give rise to several of the signs of a cancer; but the history of an antecedent local or general peritonitis, the swelling not being influenced by changes in the posture of the patient or by the acts of respiration, the indistinct fluctuation of the tumefaction, and its acute course, will ordinarily enable us to distinguish the non-malignant from the malignant affection. In rare instances the tumor may be enormous, increase rapidly, yet be simply *fatty*. There are no means of positively distinguishing the affection.§ *Sarcoma* cannot be told from carcinoma; it is more common in advanced age.

In some cases the malignant disease is closely simulated by *dilatation of the colon*, caused ordinarily by fæcal tumors. This, though it may present but a single swelling, generally occasions several, which are commonly seated at the middle third of the abdomen, are apt to appear on both sides, to be movable and painless and to bear handling without pain, to change their position slightly at intervals, and to become occasionally less in size. Then, after the case has been for some time under observation, we may be able to notice large and characteristic discharges; though we must not forget that a mere sluggish state of the bowels, or even diarrhoea, may exist while the colon is dilated and perhaps filled with fæcal accumulations.|| Sometimes the mass may be seated

---

\* See the cases of Bright, in *Clinical Memoirs on Abdominal Tumors*, republished from Guy's Hospital Reports by the New Sydenham Society.

† Bristowe, *St. Thomas's Hospital Reports*, vol. xi.

‡ As in the instances recorded by Albert Robin, *Bull. de la Soc. Anat.*, 1873, and Vidal, *Bull. et Mém. Soc. Méd. des Hôpit.*, 1874.

§ See *St. George's Hospital Reports*, vol. v., 1870, p. 253.

|| For several interesting cases of the disorder, see Kennedy, *loc. cit.*



above the symphysis and be mistaken for a pelvic tumor. Like a cancerous growth, it may lead to occlusion of the intestine and complete intestinal obstruction. The tympanites and the dilatation it occasions, which may be idiopathic, produce at times fatal results.\* The dilatation may be enormous, though there is no constriction in any part of the bowel.

*Cancer of the intestine* has symptoms similar both to faecal accumulation and to cancer of the peritoneum. The marked cachexia and the signs of persistent and increasing narrowing of the bowel, as shown by the flattened faeces, the blood and pus in the stools, the frequent attacks of colicky pains, and the vomiting, distinguish it from the former affection. The limitation of the swelling, the absence of dropsy, the character of the stools, the frequent change in the position of the tumor and in its distinctness,† and, if it affect the duodenum, the decided jaundice, separate it from peritoneal cancer.



## SECTION V.

### ABDOMINAL PULSATION.

**Aortic Pulsation.**—By far the most frequent cause of a pulsation visible in the abdomen, and especially at the epigastric region, is a throbbing of the abdominal aorta. It is common in hysterical persons. Some women are liable to it immediately before their menstrual periods or during the earlier months of pregnancy. In men it is seen most often in those who suffer from inveterate dyspepsia, and is apt to come on in severe paroxysms, which are alarming to the patient, but which generally disappear under brisk purging. In hypochondriacs whose abdominal walls are thin, the beating at the epigastrium may become a source of continued distress.

The increased action of the aorta, or, as happens in emaciated persons, the greater distinctness with which the beat of the artery

---

\* Gee, St. Barthol. Hosp. Rep., vol. xx.; A. Money and S. Paget, Clin. Soc. Transact., 1888; Formad, Trans. Coll. Physicians, Phila., 1892.

† Leube, Ziemssen's Cyclopædia.

is perceived, without there being abnormal throbbing, may be distinguished from an enlarged and somewhat displaced heart by the circumstances of the case and the absence of the physical signs of cardiac disease ; and from an aneurism, by the mode of invasion, and by the want of those signs which characterize an aneurism.

**Abdominal Aneurism.**—Aneurism of the abdominal aorta is a disease of middle life, and of males. Its most frequent cause is excessive muscular exercise ; sometimes it is produced by a blow on the abdomen, or by syphilis. Its duration is very uncertain : occasionally six or seven years elapse from its earliest indications until the fatal termination ; not unusually the patient lives twenty to thirty months after the outbreak of the complaint.

The chief *symptoms* are pain, and an absence of dropsy, of fever, or of any considerable constitutional disturbance. The pain is generally felt in the back, or in the right hypochondrium, or shooting down the sciatic nerves to the lower limbs. It may be constant and dull, or occur in protracted and violent paroxysms ; ordinarily there is a persistent pain which has periods of fierce exacerbation. The disproportion between its violence and the otherwise almost unimpaired health is a striking and common feature of the disease, and is apt to continue until the aneurism becomes very large and occasions displacement of important organs.

The *physical signs* of an abdominal aneurism are : an impulse communicated to the hand when placed over the swelling ; a systolic blowing sound ; a thrill ; and in some instances a distinct prominence and alteration in the form of the abdomen. The impulse corresponds, with rare exceptions, to the beat of the heart, is single, and ordinarily very forcible. Generally it cannot be felt from behind ; it is a beat discerned only anteriorly and on either side of the pulsating sac. Corresponding to the throbbing of the tumor, we often hear a short blowing sound, to be detected both posteriorly and anteriorly, sometimes perceived in the recumbent posture only ; or a dull, muffled sound ; rarely are there two sounds. A thrill felt at the same time as the pulsation is not unfrequently noticed ; still, it may be absent, even in large-sized aneurisms.

Aneurism of the abdominal aorta may be confounded with—  
RHEUMATISM ; NEURALGIA ; COLIC ;  
DISEASE OF THE SPINE ;

## AORTIC PULSATION ;

## LUMBAR AND PSOAS ABSCESS ;

## NON-ANEURISMAL PULSATING TUMOR.

The first four of these affections are likely to be mistaken for an abdominal aneurism, on account merely of the pain ; the others, because of the presence of pulsation, or of a swelling, or of both pulsation and swelling.

*Rheumatism ; Neuralgia ; Colic.*—The pain caused by an aneurism may closely simulate rheumatism of the lumbar muscles, or sciatica, or abdominal neuralgia, or colic. There is nothing in the pain itself which will lead to the detection of its origin : this can be effected only by a recognition of the physical signs of the aneurism. When these are not well defined, the diagnosis is doubtful. Yet, even when they are slightly marked or absent, if the pain be very obstinate, and we have excluded the affections named or cannot trace them to their usual causes, we shall often be right in attributing the pain to an aneurism. This is especially true as regards abdominal neuralgia occurring in males,—a disorder which ought always to make us examine for an aneurism, and which is not unfrequently found to be due to it.

*Disease of the Spine.*—Patients who are suffering from aneurism often complain of pain in the spine, and present sometimes an obvious spinal curvature. But a careful examination, by detecting the physical signs of an aneurism, will enable us generally to distinguish the source of the difficulty. The constant boring pain so much complained of in cases of aneurism is usually thought to be due to absorption of the vertebræ ; but, as Stokes proved, it has no necessary connection with this lesion.

*Aortic Pulsation.*—Simple abdominal pulsation, such as we observe in hysteria, in dyspepsia, and in pregnancy, or excessive epigastric pulsation due to an enlarged right ventricle or to insufficient aortic valves, may be readily mistaken for an aneurism. But in the former case the history will generally lead us to a correct conclusion, especially if taken in connection with the facts that the pulsation is not heavy and slow, as in an aneurism, but jerking and sudden ; that there is no thrill ; no tumor with corresponding dulness on percussion, if we except pregnancy ; no systolic murmur audible in front of the abdomen or along the spine ; and no pain.

The pulsation due to disease of the heart is discriminated by the physical signs in the thorax. Regurgitation at the aortic orifice, which is the cardiac affection most liable to be confounded with an aneurism, on account of the marked pulsation it may occasion in the left hypochondrium or at the scrobiculus cordis, is distinguished by the single or double blowing sounds, which are heard not only over the thorax, but also over so many arteries of the body, and by the character of the pulse.

*Lumbar and Psoas Abscess.*—In some cases, soft, fluctuating, deep-seated tumors, which are really produced by an aneurism, may arise in the lumbar region; nay, they may seem to point, as happens in psoas abscess, at Poupart's ligament. But, unlike an abscess, the effusions of blood give rise, with rare exceptions, to impulse and to murmur.

*Non-aneurismal Pulsating Tumors.*—When a tumor of any kind presses upon the aorta, a distinct pulsation is communicated, and the similarity to an aneurism is heightened by the circumstance that the morbid growth may produce a murmur. The tumors which most usually occasion the phenomena mentioned are: enlargement of the left lobe of the liver, cancer of the pylorus, disease of the pancreas, or of the omentum, or of the mesentery, and, in rarer instances, enlargement and distention of the kidney, fecal accumulations, and cancer of the lumbar glands.

Now, to avoid error, we must pay close attention to the history of the disorder; we must trace, by percussion, the outline of the solid mass, and see if it correspond with any viscus; we must lay stress on the presence of digestive disorders, and on the amount of constitutional disturbance,—both of which are so slight in abdominal aneurism; we must examine the urine carefully, and find out whether there are renal symptoms in the case. Then, in non-aneurismal tumor the patient has almost always been in bad health before the tumor is detected, and the swelling rarely causes pain of such severity as is observed in an aneurism; moreover, the transmitted aortic impulse is, as a rule, lessened by placing the patient on his hands and knees, thus taking away the pressure from the artery. A varicose state of the epigastric veins and the existence of ascites will also decide against an aneurism; while, on the other hand, the lateral as well as the forward direction of the impulse, violent neuralgic pains in the loins or shooting down the back, and



an immovable tumor, are in its favor. Still, there are cases in which a morbid growth lying across the aorta occasions symptoms so nearly like those of an aneurism that the most skilful diagnostician finds himself in doubt.

There are cases of aneurism in which the physical signs are absent, and in which the affection affords no indication of its existence, beyond, perhaps, pain. Under these circumstances we can only suspect its occurrence.

But supposing that, from the combination of the physical signs and symptoms, we know that we are dealing with an abdominal aneurism, can we be sure that it is aortic? We cannot; for, although this is generally its seat, an aneurism of the splenic or the celiac artery, of the superior mesenteric artery, or of the renal artery, may produce the same phenomena.\*

When an aneurism bursts, it gives rise to symptoms which vary with the seat of the rent. The accident is always fatal, but death may not follow for several days; usually great tenderness of the abdomen and changes in the physical signs are at once produced.

---

\* See Ballard, *Physical Diagnosis of Diseases of the Abdomen*, p. 217.

## CHAPTER VII.

### ON THE URINE, AND ON DISEASES OF THE URINARY ORGANS.

BEFORE discussing the diseases of the urinary organs let us notice the urine in its pathological and clinical aspects.

#### URINE.

The main function of the kidneys is to remove water and nitrogen from the system, at the same time that they take from the blood many of its salts. The excreted liquid contains a variety of substances, and by its study we are enabled to arrive not only at the condition of the organ which prepares it, but also at the state of the circulating fluid, and often indirectly at that of several viscera, the disorders of which give rise to impurities in the blood, which the kidneys endeavor to eliminate. Hence the urine, besides being the most accurate index of the condition of the urinary organs, becomes a fair indication of that of many other important secreting glands in the body; and, further, throws light on the workings of the nervous system.

To glean the full benefit from an analysis of the urine, we must explore it not merely qualitatively, but quantitatively, and examine its deposits with the microscope. Modern chemistry is especially endeavoring to find means which will bring it within the power of every one to determine, by apt volumetric processes, the exact proportion of the ingredients as accurately and as easily as hitherto we have detected their presence. This is a subject which cannot be more than indicated in these pages: only such of these investigations will be noticed as have furnished results which may be made readily available for the exigencies of professional life.

It is customary, in quantitative analyses, to use the French system of measures, and to employ instruments on which cubic centimetres are marked. One thousand cubic centimetres are equal to one litre, or 2.1 U. S. pints, or to a thousand grammes of

water; and one gramme is equal to 15.434 grains; one centigramme to .1543 of a grain.

Urine, in its normal state, is of acid reaction, of amber-yellow color, and of specific gravity of 1018 to 1025 as compared with distilled water at 1000. On standing from eight to twelve hours, a slight cloudy deposit takes place, consisting mainly of mucus, epithelial cells from the urinary passages, and a few crystals. Healthy urine freshly voided contains no bacteria, and is aseptic.

Ordinarily, urine soon undergoes decomposition, which renders the results of analysis valueless. It is advisable, therefore, to examine every specimen promptly, but, as this cannot always be done, the addition of some preservative may be needed. Chloroform seems to be the most suitable; six or eight drops added to each fluidounce, the mixture to be well shaken, will preserve samples for months, even in hot weather. Chloroform gives a strong reaction similar to sugar with Trommer's test, but does not reduce bismuth subnitrate nor interfere with the phenylhydrazin test. It arrests the fermentation of sugar and of urea.

In the examination of sediments great advantage, both as to time and complete collection of the suspended matters, is gained by the use of a centrifugal machine, several forms of which are now procurable. Where a motor current is available, the electric centrifuge is the best. The method tends to exaggerate the amount of material, as compared with the old method of sedimentation. We may also obtain casts and other free elements which otherwise would escape. In addition to its usefulness in urine-examination, a good high-speed centrifugal machine is of much use in other clinical work, especially in examining sputum and blood. Purdy's percentage tubes add much to the advantage of the instrument.

The manner of obtaining a specimen of urine is not unimportant. We should instruct our patient, as is so strongly recommended by Sir Henry Thompson,\* to pass the first two ounces into one vessel, and the remainder into another. We thus procure a specimen of the renal secretion, in addition to anything in the bladder, separate from any urethral products, and avoid the error of confounding prostatic or urethral with vesical or renal disease. When it is essential to obtain a specimen of urine absolutely pure

---

\* Clinical Lectures on Diseases of the Urinary Organs.

and unmixed with products of the bladder, the same authority recommends the drawing off of the urine by means of a soft gum catheter, while the patient is standing. The bladder should then be carefully washed out by repeated one-ounce injections of warm water. The urine is now to be permitted to pass, as it will do, drop by drop, into a small glass vessel. The bladder contracts around the catheter, and the urine percolates direct from the ureters, through their virtual prolongation,—the catheter,—into the receptacle. The urine passed in the morning, immediately after rising, will be found to represent with sufficient accuracy the general process of disassimilation; but, if greater accuracy be desirable, a specimen of the mixed urine of the twenty-four hours should be used.

As regards the *quantity* of urine daily voided, the mean average of healthy persons is 1500 cubic centimetres (fifty fluidounces). In summer, when the skin is acting freely, less fluid passes off by the kidneys than in winter. The more liquid that is taken into the system, the greater is the secretion of urine, unless the other organs that eliminate water, as the skin, the lungs, and the intestines, are excreting with unwonted activity.

The quantity is diminished in all cases in which the specific gravity is increased, with the exception of diabetes; it is diminished in acute diseases, in fevers, in cholera, and in the early stages of dropsies; in some forms of Bright's disease, particularly the acute forms, through their entire course, and often in the last stage of all forms of that disease. It is, on the other hand, augmented in cardiac hypertrophy and whenever the specific gravity is diminished; in hysteria; in contracted kidney, and in polyuria. In almost all vesical and renal affections frequent micturition is a marked symptom,—not always, however, associated with increased quantity of urine.

The *ingredients* of urine are numerous. The principal are: urea, sulphates, phosphates, chlorides, uric acid and urates, kreatinin, hippuric acid, mucus, coloring-matter, and a large proportion of water.

The following table, by Parkes, shows the composition of normal urine, the figures representing the amount passed in twenty-four hours by a male adult weighing sixty-six kilogrammes (one hundred and fifty pounds).



Water .....	1500.00	grammes.
Urea .....	33.18	"
Uric acid.....	.55	"
Hippuric acid .....	.40	"
Kreatinin.....	.91	"
Pigment and minor organic matters.....	10.00	"
Sulphuric acid.....	2.00	"
Phosphoric acid.....	3.16	"
Chlorine.....	7.00	"
Ammonia .....	0.77	"
Potassium.....	2.50	"
Sodium.....	11.90	"
Calcium.....	.26	"
Magnesium.....	.27	"

In the present state of our knowledge of the condition of substances in solution, it is not possible to arrange these ingredients in definite combinations. Besides the elements mentioned, the quantities of which fluctuate with the food-supply and with the activity of tissue-metamorphosis and vary especially when the system is deranged, we meet, in morbid states, with substances that do not exist at all in healthy urine, or the presence of which is doubtful, such as various forms of albumin, sugar, blood, bile, fats, oxalate of lime, and certain pigments. Most of these are dissolved in the urine, and are not to be detected except by delicate tests; others form in sediments after the urine has been discharged, and may be recognized by the microscope.

Having thus, in a general manner, mentioned the constituents of the urine, normal and accidental, let us, in the same general manner, look at the points of clinical interest to be decided by an analysis; in other words, let us ascertain what the physician, not the professed chemist, is in quest of.

Usually, we endeavor to fix all these waymarks: the color, the specific gravity, the quantity, the reaction, the presence or absence of such important abnormal ingredients as albumin and sugar, and the character of the deposits. Frequently, too, we extend our examination until we have determined approximately, if not accurately, the increase or diminution of the main constituents of the urine, especially of the urea, uric acids, chlorides, phosphates, and sulphates, and the distribution or non-distribution of bile and other unusual constituents through the fluid. Let us examine these points more in detail.

**Color.**—The color of the urine is much affected by food and medicine, as well as by various morbid processes; so rapidly, indeed, affected, that we must be chary of drawing conclusions from the appearance of the secretion alone. Yet we suspect the presence of certain substances, or are nearly positive of their absence, by the appearance of the fluid. Thus, a smoky or a red aspect is apt to be owing to admixture of blood; a very light color denotes generally an increase of water, and is commonly found in diabetes, in hysteria, and in kindred nervous affections. In febrile diseases the urine is of dark hue. A greenish-yellow or brownish tint of the discharge is indicative of bile; but a similar tinge may be present when rhubarb has been taken. A dirty-blue urine happens from an indigo sediment; it is alkaline, and occurs chiefly in typhus and in cholera. Strong coffee darkens the urine; turpentine darkens and imparts a violet odor to it; carbolic acid, tar, and creasote render it black; so do disintegrated blood and melanotic cancer. Santonin, logwood, and senna discolor it. The first-named substance gives it a bright yellow color, which on the addition of an alkali becomes crimson. Senna may impart to it a brownish or a deep red color, which, however, like that due to rhubarb, is lightened on the addition of mineral acids, and is thus distinguished from the hue of urine containing blood. The altered appearance is mostly due to the coloring-matter of these articles being excreted with the urine.

The chemistry of the coloring-matters of the urine is still incomplete, and the clinical significance of the color-changes still obscure. The principal normal coloring-matter is *urobilin*, which is an oxidation-product from blood and bile-pigment. In febrile conditions a less oxidized product is excreted, which MacMunn has named *pathological urobilin* and declares to be identical with the coloring-matter of the feces, *stercobilin*. He further states that the presence of this body in the urine is to a certain extent an indication of the absorption of fecal matter and ptomaines which have not been destroyed by the liver. Other pigments have been described, among which may be named *uroerythrine* and *urochrome*. The employment of the spectroscope is one of the means of distinguishing between these colors, but a description of their minute differences would be beyond the scope of this work.

**Specific Gravity.**—We take the specific gravity of urine to

judge of the solid matter it contains. The readiest means is the urinometer. For the implement to yield trustworthy results the fluid should be brought to the temperature at which the urinometer has been graduated. A difference of  $7^{\circ}$  F, corresponds to about 1 degree of the urinometer. Most instruments are graduated for use at  $60^{\circ}$  F., and the cheaper forms are often inaccurate. Squibb makes a urinometer adapted for use at  $77^{\circ}$ , which is convenient for office work. It is easy to obtain instruments sufficiently accurate for clinical use. More accurate than the urinometer is the specific gravity bottle, or the Westphal balance.

If there be but a small quantity of urine for examination, we note the amount and how many volumes of distilled water it takes to fill the vessel to the height required to float the urinometer. We then multiply the number above 1000 that the instrument shows, by the total number of volumes of the mixed fluid. This is only approximate.

From the specific gravity we may calculate approximately the quantity of solid matter passed by multiplying the number above 1000 by 2.33. This may be done whether we estimate in grammes or in grains. For instance, in urine of specific gravity of 1010 there will be 23.3 grammes of solid matter in each 1000 grammes of urine; in urine of 1030, 69.9 grammes. This information obtained, it is easy to find the whole amount of solids contained in the urine of twenty-four hours, after ascertaining the quantity passed in that time. To take the first illustration: if 1000 grains yield 23.3 grains of solid matter, how much would be yielded by 20,000 (the quantity passed, let us say, in twenty-four hours)?

$$1000 : 23.3 :: 20,000 : x. \quad x = 466 \text{ grains.}$$

This method is not very precise; when exactness is required, the urine must be evaporated until a dry residue is left, which should then be carefully weighed.

The amount of solids in healthy urine is variously estimated. Beale places it approximately at from 800 to 1000 grains in twenty-four hours; Hofmann and Ultzmann fix it at 60 to 70 grammes,—about 920 to 1080 grains,—and in persons who are fasting, or have taken little food, as in fevers, at 30 grammes, in the twenty-four hours. As a rule, the proportion is greatest in persons of heavy weight: if, therefore, we wish to make nice

comparisons, the weight of the body should be always stated. To ascertain how much of the solid matter consists of mineral matters, the organic substances must be burned off at a red heat.

In disease, the solids, and with them of course the specific gravity, fluctuate very much. We find the specific gravity decidedly increased, rising to 1030 or higher, when sugar or an excess of urea is present, and when the urine is concentrated and of deep color. A low specific gravity is met with in certain forms of Bright's disease, in many cases of hysteria, and in pale urine except that of diabetes. But to be accurate—and, indeed, accuracy in regard to the other physical and chemical properties is unattainable without attending to the same rule—we must not lay stress on the specific gravity without taking into account the measure of urine passed in the twenty-four hours, as well as the quantity of drink and of food swallowed; all of which of necessity influences the specific gravity. So, too, does the activity of the tissue-metamorphosis.

**Reaction.**—Normal urine usually reddens blue litmus-paper. The *acidity* depends upon acid salts, especially acid sodium phosphate. The degree of acidity is, even in health, not always equal, and is much influenced by digestion. If no food has been taken for hours, the discharge is highly acid; that passed after a meal, and while the process of digestion is going on, is but faintly so, or neutral, or even alkaline. In about three or four hours after meals the alkaline tide turns, and the acidity of the urine slowly increases until food is again taken. There seems, however, to be a limit to the increase of acidity, for Bence Jones found that continuing to fast for twelve hours beyond the usual meal-time did not intensify the acidity of the urine. The alkalinity of the urine after meals is rarely detected at the bedside. For, although the urine may be alkaline when secreted by the kidneys, it is generally mixed in the bladder with that which collected before or after the alkaline tide, and the mixed urine when passed may have an acid reaction.

The acidity of the urine is augmented by the administration of the vegetable or the mineral acids; yet they do not cause, even in large doses, as great variations as does digestion. We find the urine very acid during a meat diet. We find acidity of the urine



strongly marked if any acid be present in it which sets the uric acid free, or if this be in decided excess.

For determining reaction, litmus-paper is used. Solution of litmus is divided into two parts ; to one part nitric acid is added, drop by drop, until the color is wine-red. This is then mixed with the other half. Slips of filtering-paper are dipped in this and dried. They have a purple tint, and are very delicate, responding to a trace either of free acid or of alkali. We thus avoid the use of two colors. Litmus-paper is best kept in a closed dark bottle.

We may estimate the amount of free acid in the urine by a solution of sodium hydroxide (caustic soda) containing 4.0 grams to the litre. This solution is added drop by drop to 100 cc. of urine, which has been measured off in a beaker glass. After the addition of each half cubic centimetre, a drop of the mixture is placed, by means of a glass rod, on well-prepared litmus-paper. When the paper is no longer reddened, the analysis is finished ; and by noting how much of the standard solution has been used, we can determine the acidity of the urine, which it is customary to express as equal to so many grains of oxalic acid, the value of the sodium hydroxide solution in terms of oxalic acid having been previously ascertained.

Urine, when voided, remains ordinarily acid for at least a day ; but it may lose its acidity much sooner. This is always a significant fact, having much the same meaning as if the fluid had been discharged in a neutral or an alkaline state.

Now, an *alkaline* reaction may result from several causes : from the effect of digestion, as already mentioned ; from the presence of sodium or potassium carbonate ; or from the decomposition of the urea into ammonium carbonate. In the former case, heat does not restore the color of the red litmus-paper,—it remains blue ; in the latter, a gentle heat soon brings back the original red tint. Moreover, in either case, the earthy phosphates are precipitated, the fixed carbonate causing the precipitation of the amorphous calcium phosphate ; while by the ammonium carbonate ammonium and magnesium phosphates, in conjunction with the calcium phosphate, are thrown down, and the triple phosphate is abundantly formed, and can be easily recognized under the microscope by its prismatic crystals.

Alkalinity of the urine from fixed alkali is not inconsistent

with health. We have adverted to the effects of digestion and to the fact that alkaline urine results from the use of certain articles of vegetable food, or of the salts of sodium and potassium. Urine owing its alkalinity to ammonium carbonate is always to be viewed as pathological. The disturbance is generally long continued, and the urine loses its acidity in the bladder, in consequence of a disease of the mucous coat of the viscus, or from being long retained there, as in cases of paraplegia, or from admixture with pus, which acts as a kind of ferment and leads to decomposition of the urea.

**Changes in the Quantity of the more Important Constituents of Urine.**—*Urea.*—The amount of urea excreted by well-nourished, healthy, adult males in the twenty-four hours is estimated, in round numbers, by Roberts at  $3\frac{1}{2}$  grains per pound weight of the body, and by Neubauer and Vogel at 25 to 40 grammes, or 0.37 to 0.6 gramme for every kilogramme of weight of the body. Purdy places the mean excretion of urea in healthy adult males between the ages of twenty and forty years at 33.18 grammes (512.4 grains) in twenty-four hours.

Urea is the principal product of the change of nitrogenized substances. Its proportion fluctuates, therefore, with the food partaken of, as well as with the activity of the transformation of the structures of the system: hence it becomes the most important index of the waste and repair of tissues. Exertion of body and of mind leads to the discharge of a larger quantity of urea. If this is replaced by a nourishing diet, nothing is lost; the body retains its health. But when the requisite amount of nitrogenized aliment is not taken, or, if taken, cannot be assimilated, owing to a disturbance in digestion, the person wastes. We notice, too, in acute febrile states, until their height is reached, hand in hand with the emaciation an increase of this significant urinary constituent,—a proof, then, of the rapid and unsupplied disintegration of the tissues. We see the same increase during paroxysms of intermittent fever, in inflammations, and in some cases of nervousness; also from a predominant animal diet, and in certain forms of indigestion, in which the food is speedily passed off in the shape of urea instead of acting its part in the nutrition of the economy. Degenerative changes in the liver may be accompanied by a diminution of urea-excretion.


A lessened quantity of urea is excreted during fasting, from an almost exclusive vegetable diet, in dropsies, and in many long-continued organic diseases which gradually undermine the general nutrition and diminish tissue-change, or in states attended with diminished oxidation. But the diminished amount in the urine may also be due to a want of secreting power of the kidneys. The urea, or the products of its decomposition, then act as a poison in the blood; and the symptoms indicative of uræmic poisoning are encountered. Urea is sometimes not found in the urine at all, or only in traces, having been replaced by leucine and tyrosine.

Quantitative estimations of urea are almost exclusively made by the use of solutions of sodium hypochlorite or hypobromite, which decomposes the urea, liberating nitrogen and carbon dioxide in amounts proportional to the urea present. The carbon dioxide is kept in solution by using excess of sodium hydroxide or carbonate, and the volume of nitrogen is measured. The most accurate results seem to be obtained with the hypobromite, but this does not keep well, and its extemporaneous preparation is troublesome and annoying. Sodium hypochlorite is readily obtained, being the common Labarraque's solution. It keeps in good condition for a long while, and, according to Squibb, who has investigated the matter carefully, gives good results. It must be prepared according to the U.S.P. 1890,—that is, contain a marked excess of sodium carbonate. Several observers have reported that improved effect results from the addition of potassium bromide (1 gramme to 25 cc. of *Liquor sodæ chloratæ*). To avoid the annoyance of using pure bromine, when it is desired to employ the hypobromite solution, Charles Rice suggested the use of a solution of bromine in potassium bromide. This keeps well, and is more convenient to handle. In this method the solutions used are as follows: (a) 10 grammes of potassium bromide are dissolved in 80 cc. of water, 10 grammes of bromine added, and the liquid shaken until the latter is dissolved; (b) 10 grammes of sodium hydroxide are dissolved in 25 cc. of water. For use, equal quantities of the two liquids are mixed and slightly diluted with water. Sodium hypobromite may also be prepared by adding directly 1 cc. of bromine to 25 cc. of the above solution of sodium hydroxide; but the liquid must be used within a few hours, or accurate results cannot be expected. The mixture must be made

in a well-ventilated place, as bromine is exceedingly irritating and corrosive.


For collecting and measuring the nitrogen evolved, many forms of apparatus have been devised. That of Hufner is a standard form, but simpler and less expensive forms are now usually employed in clinical work. Fig. 56 shows a form devised by Doremus, which is much used. The apparatus is filled with solution of hypobromite or hypochlorite, so that when the graduated tube is upright the bulb is about half filled. A large watch-glass or shallow dish should be placed beneath, to catch any overflow.

FIG. 56.



Doremus's ureometer.

FIG. 57:



Greene's ureometer.

A measured quantity of the urine (1 cc.) is introduced by means of the dropping-tube, the opening being pushed well into the bend of the upright tube, and the apparatus being tilted forward to prevent any escape of bubbles or urine into the large bulb. After about twenty minutes the volume of gas is read off; 1 cc. of nitrogen may be taken to represent .0028 of urea, but the tube is usually graduated so as to read directly the percentage of urea, a definite volume of the urine being taken for each test. Squibb furnishes a simple and satisfactory apparatus. Fig. 57 shows Greene's ureometer, also a simple instrument.



Fowler's method has been endorsed by several careful observers. It depends on the fact that the decomposition of urea greatly reduces the specific gravity of the urine. It may be performed as follows. The specific gravity of the sample is carefully taken, and then 25 cc. added in a large beaker to 175 cc. of solution of chlorated soda (U.S.P. 1890), and, after mixing well, allowed to stand for a few hours, when the specific gravity is again taken. Multiply the specific gravity of this residual liquid by 7, add the specific gravity of the original liquid, and divide the sum by 8, subtract from this quotient the specific gravity of the residual mixture, multiply the remainder by 0.77, and the product is the percentage of urea. In case the urine is of high gravity it is better to use 12.5 cc. diluted with an equal volume of water and then add the 175 cc. of solution of chlorated soda. The result must be multiplied by 2.

A method for fixing the quantity of urea approximately is that proposed by Haughton. It consists in the use of tables showing how many grains of urea are excreted in the urine, of which the amount daily passed and the specific gravity are predetermined. On the following page is the table, as abridged by Roberts. Its main use is as a standard of comparison, and it cannot be depended on when sugar or albumin is present.

*Uric Acid.*—Uric acid, like urea, is a product of the metamorphosis of tissue. It was supposed by Liebig that the acid is an early stage of the transformation of urea. Hofmann teaches that uric acid is deposited owing to the decomposition of the urates by the acid phosphate of sodium. Under ordinary circumstances, the deposition of uric acid occurs subsequent to the expulsion of the urine; but should the acid sodium phosphate be in excess, the uric acid may be precipitated before the secretion is voided, and thus give rise to gravel and calculi. This may also happen through too great concentration of the urine.

The amount of uric acid passed in twenty-four hours varies from 0.5 to 1.0 gramme. It corresponds in general to the amount of urea in the proportion of 1 to 33. In normal urine the presence of uric acid cannot be detected without the addition of a strong acid, since it exists in the form of soluble urates, which must be first decomposed. The uric acid is gradually thrown down in small red grains, which, should it be desirable to deter-


Haghton's Table for the Estimation of the Daily Excretion of Urea from the Specific Gravity.

Fluid-ounces.	SPECIFIC GRAVITY.																			
	1003	1004	1005	1006	1007	1008	1009	1010	1011	1012	1013	1014	1015	1016	1017	1018	1019	1020	1021	1022
20	35	36	43	57	71	85	100	103	106	119	130	136	142	151	160	196	233	241	249	257
22	38	40	47	62	78	95	110	113	116	130	143	149	156	166	176	215	257	265	274	282
24	42	43	51	68	85	101	120	123	127	142	156	163	170	181	192	235	280	289	299	308
26	45	47	55	73	92	110	130	134	137	153	169	176	184	196	208	254	303	318	324	334
28	48	50	59	79	99	118	140	144	148	165	182	190	198	211	224	274	326	337	349	360
30	52	54	64	85	106	127	150	155	159	177	195	204	213	226	240	294	349	361	374	386
32	55	57	68	90	113	135	160	165	169	188	208	217	227	241	256	313	378	385	398	411
34	58	61	72	96	120	144	170	175	180	200	221	231	241	256	271	333	396	409	428	437
36	61	64	76	102	127	153	180	185	191	212	234	244	255	271	288	352	419	433	448	462
38	65	68	80	108	134	161	190	195	201	224	247	258	269	286	304	372	442	457	478	488
40	69	72	85	114	142	170	200	206	212	236	260	272	284	302	322	392	465	481	498	514
42	74	75	89	119	148	178	210	216	222	247	273	285	298	317	336	411	489	506	523	540
44	76	79	93	125	156	186	220	226	233	259	286	299	312	332	352	431	512	530	548	566
46	80	82	97	130	163	195	230	236	243	271	299	312	326	347	368	450	535	554	573	592
48	84	86	101	136	170	203	240	246	254	283	312	326	340	362	384	470	558	578	598	618
50	87	90	106	142	178	212	250	257	265	295	325	340	355	377	400	490	581	602	623	644
52	90	94	110	147	185	220	260	267	276	307	338	353	369	393	416	509	605	626	648	669
54	94	98	114	153	192	229	270	277	286	319	351	367	383	408	432	529	628	650	673	695
56	96	100	119	159	199	238	280	288	297	331	364	380	397	423	448	548	651	674	698	720
58	100	104	123	165	206	246	290	298	308	343	377	394	411	438	464	568	674	698	723	746
60	104	108	128	171	213	255	300	309	319	355	391	408	426	453	480	588	697	722	748	772
62	108	110	132	176	220	263	310	319	329	366	404	421	440	468	496	607	719	746	772	797
64	110	114	136	182	227	271	320	329	339	378	417	435	454	483	512	627	742	770	797	823
66	114	118	140	187	234	280	330	340	351	390	431	448	468	498	528	646	766	794	822	849
68	116	122	144	193	240	288	340	350	361	402	443	462	482	513	544	666	790	818	847	875
70	120	126	149	199	248	297	350	361	372	414	456	476	497	528	560	686	814	843	872	901
72	122	128	153	204	255	305	360	371	382	425	469	489	511	544	576	705	838	867	896	926
74	126	132	157	210	262	314	370	381	393	437	482	503	525	558	592	725	861	891	921	951
76	130	136	161	216	269	323	380	391	404	449	495	516	539	573	608	745	884	915	946	977
78	134	140	165	222	276	331	390	401	414	461	508	530	553	588	624	765	907	939	971	1002
80	139	144	170	228	284	340	400	412	425	473	521	544	568	604	640	785	930	964	996	1028

mine the quantity of the acid, are washed, dried, and carefully weighed. The quantity of uric acid may be approximately estimated by mixing 100 cc. of the urine with 5 cc. of strong hydrochloric acid and allowing it to stand in a cool place for two days. The separated crystals are then collected, washed with a little water, and weighed. A correction of .0038 of a gramme for 100 cc. of liquid used in washing should be added to allow for solubility.

The characteristic reaction of uric acid is furnished by the *murexide test*. A few drops of nitric acid are mixed with the suspected deposit in a capsule, and the mixture is slowly evap-

FIG. 58.



Crystals of uric acid, magnified about 200 diameters. Most of these forms are seen in the urine of acute rheumatism.

orated to dryness, best on a water-bath; a drop of ammonium hydroxide is then added, which produces instantly a rich purple.

But both uric acid and the urates can be easily and quickly discriminated by the microscope. The crystals of uric acid are readily discerned, notwithstanding that they vary both in size and in form. Rhombic plates with rounded angles are frequent. To obtain the crystals rapidly, where they are not passed as uric acid, a portion of the suspected deposit is dissolved in a drop of potassa, and the alkaline solution treated with an excess of acetic acid: after the lapse of a few hours crystals of uric acid will be formed.

In disease, the fluctuations in the quantity of uric acid are great; as a general rule, they correspond to the rise and fall of



urea. We find the acid diminished in hydruria and affections in which the eliminating power of the kidneys is interfered with, as in the more advanced stages of Bright's disease and in anæmia and chlorosis. An increase is encountered in acute inflammations, in fevers, in functional disorders and many of the structural affections of the liver, in heart and lung diseases attended with dyspnœa, in leukæmia, and in acute rheumatism. In the latter malady the little red granules, visible to the naked eye, form a deposit in the urine soon after it is voided.

We must, however, be careful not to suppose the uric acid to be in excess because it is readily precipitated. It may or may not be in larger amount: the sediment merely proves an augmentation of acidity in the urine sufficient to take away the base from the uric acid. This happens often as the result of acid fermentation of the urine. Frequently urates are separated along with the uric acid; we find then generally a dark urine of high specific gravity and of very acid reaction.

Persons who habitually pass urine of the character described are subject to gastric or hepatic disorders. They are also often gouty, or lithæmic, and frequently consumers of a large amount of animal food, or intemperate or indolent in their habits.

Uric acid or urates are never found as sediments in freshly-voided healthy urine. Occasionally precipitates of uric acid or urates occur in the urinary passages. Now, these sediments may concrete and form the nuclei of calculi; or they may be passed in small particles, commonly spoken of as "gravel."


*Urates.*—The pathological conditions in which the urates are changed are much the same as those in which alterations in uric acid occur. They are principally the sodium, potassium, and ammonium urates. The deposits formed by their precipitation are of pink color, sometimes brown, or like brick-dust, or yellowish, or even white. From pale urine of low specific gravity a white sediment is apt to settle. All the deposits are dissolved with readiness by heat. Acids decompose them and separate uric acid. They are all more soluble in warm water than in cold, and the neutral salts are more soluble than the acid ones.

Under the microscope, the urates are seen to be either irregular, amorphous particles, needle-like crystals, dumb-bells, or round globules of various sizes, from some of which fine needles project.



The latter, like the dumb-bells, are commonly supposed to be sodium urate; the globules and crystals, sodium urate and ammonium urate; the granular, amorphous powder, mixed urates, more especially sodium urate and potassium urate. These amorphous urates may, under the microscope, be mistaken for calcium

FIG. 59.



Mixed urates.

phosphate. The differential test consists in their behavior with acids: the phosphate is dissolved by acetic or hydrochloric acid; the urates are gradually transformed into crystals of uric acid. Then, a deposit of calcium phosphate is often more cloudy than the urates, and, unlike them or uric acid, is not soluble in liquor potassæ. From calcium carbonate, which also occurs in a granular form, both the urates and the calcium phosphate are distinguished by the effervescence of the carbonic acid which happens on the addition of a strong acid.

Urine containing a sediment of urates is generally markedly acid, or soon becomes so, either from an absolute increase of the uric acid, or in consequence of changes in some of the constituents of the fluid—as of the pigment—which take place either before or shortly after emission. Not unfrequently, too, it is scanty, and the urates are deposited as soon as the urine cools to the temperature of the atmosphere. Their precipitation may be, and indeed often is, due to there not being water enough to hold them in solution. We may judge of this being the case by ascertaining the amount of urine passed in twenty-four hours. If the quantity

be about normal, the deposit is in all likelihood due to an excess of urates. In cold weather these deposits occur more quickly and more extensively than in warm.

Sediments of urates are at times met with in pale urine, and without either diminution of water or excess of acidity. The urine yields but a faintly-acid or a neutral or an alkaline reaction, and under the latter circumstances calcium phosphate, or even triple phosphates, may be observed to accompany the urates. The urate present is acid ammonium urate.

*Phosphates.*—The phosphates are derived in part from the food, in part from the disintegration, or rather the oxidation, of the disintegrated albuminous substances, and especially of the nerve-structures. They occur either as calcium and magnesium phosphates, the *earthy phosphates*, which exist in small amounts, about 1 gramme in twenty-four hours, and as sodium phosphate, about three times as abundant, forming the greater part of the *alkaline phosphates*.

In health the phosphates are kept in solution by their acidity; but as soon as the urine ceases to be acid they are deposited. Hence the appearance of phosphates bespeaks a neutral or an alkaline condition of the urine, with the exception that calcium phosphate may occur in acid urine. Often the fluid, as we have already seen, becomes alkaline from the decomposition of the urea into ammonium carbonate. This acts upon the phosphate, forming ammonio-magnesium phosphates, which crystallize commonly in transparent prisms or in feathery-looking bodies, easily distinguished from the amorphous powder or small round globules of calcium phosphate. Yet there is, as Roberts has pointed out, a crystalline form of calcium phosphate which might be mistaken for one of the stellar forms of uric acid, but it may be distinguished by its being invariably colorless. These earthy phosphates are all readily soluble in acids, even in weak acids like acetic acid, and this at once distinguishes them, even under the microscope, from calcium oxalate, which some forms resemble. In many specimens of urine they are precipitated by heat; but the addition of an acid soon dissolves them, and thus prevents the turbidity from being mistaken for that due to albumin.

The triple phosphates are often met with in heavy deposits mixed with pus, especially in the alkaline purulent urine resulting

from chronic vesical catarrh. They are also seen in cases of retention of urine due to temporary or permanent paralysis of the bladder, as in low fevers, in hemiplegia, or in paraplegia. They are found, too, in many affections in which the vital powers have

FIG. 60.



Earthy phosphates; the granules are chiefly calcium phosphate, the rest triple phosphates.

been seriously lowered and the acidity of the urine diminished, as during convalescence from acute disease. Under the latter circumstances, and in fact whenever the urine has become alkaline from the presence of a fixed alkali, the phosphatic deposit is apt to show a large excess of the amorphous phosphates, if, indeed, it do not altogether consist of them.

Urine alkaline from fixed alkali, and depositing phosphates, is, unless this condition have been brought about temporarily by fruit or other food, a matter of serious import. We encounter it in persons laboring under great general debility and indigestion associated with an impaired tone of the nervous system, and with aching pains in the lumbar region and a tendency to boils,—in fact, in those of whom it is customary to speak as exhibiting the phosphatic diathesis, or as having *phosphaturia*. Such a morbid state is not uncommon in men depressed by mental toil or anxiety, and may become associated with harsh, dry skin and enormous flow of urine and marked emaciation, giving rise to so-called *phosphatic diabetes*.

In spite of the distinct sediment of the phosphates, it is some-

times doubtful if the latter are really increased in quantity. The want of the acidity of the urine permits their precipitation, and causes them to become readily apparent. On the other hand, the phosphates may be actually in excess, and yet this excess be concealed from view. This happens especially with the alkaline phosphates, the proportions of which change in disease much more than do the earthy phosphates, and indicate much more clearly the variations of the phosphoric acid.

Now, a real, not merely an apparent, increase of the phosphates occurs, according to Bence Jones, in acute inflammatory diseases of the nervous structure, and in fractures of the skull when an inflammatory action takes place in the brain. It also occurs after mental strain. We find the phosphates also augmented by the abundant use of animal food, and by very active exercise. The earthy phosphates are markedly increased in rickets and in extensive bone disease; the phosphoric acid, as well as the sulphuric acid, the urea, and the sodium chloride, is excreted in less amount than in health during the course of a maniacal paroxysm, in epilepsy, in melancholia.\* In gout as well as in Bright's disease, too, the excretion of phosphoric acid is diminished.

To determine the proportion of the *earthy* phosphates, a few drops of ammonia are added to the urine; soon a whitish precipitate is produced, which is not removed by heat. From the quantity of the deposit, after settling, we may form a rough estimate of that of the earthy phosphates. In an ordinary-sized test-tube a deposit 1 c. high represents a normal amount. But if the amount is to be accurately ascertained, we must employ a graduated glass, separate the precipitated phosphates by filtration, ignite them in a platinum capsule, and weigh the ash. The *alkaline* phosphates are not thrown down by alkalis, and, unlike the earthy phosphates, are very soluble in water. They are procured by taking the fluid from which the earthy phosphates have been carefully removed by filtration, and adding to it a saturated solution of magnesium sulphate. Or we add to the urine about one-third as much of the magnesium mixture, and if the precipitate be copious, giving the fluid the appearance of cream, then the alkaline phosphates are in excess; if there be merely a milky turbidity, they are normal.

---

\* Adam Addison, Brit. and For. Med.-Chir. Rev., April, 1865.



From the deposit obtained in testing for the phosphates, some idea may also be formed of the quantity of *phosphoric acid* in the urine. The average quantity passed by an adult male in twenty-four hours is, according to Vogel, about 3.5 grammes, or nearly 53 grains. For the volumetric processes by which the amount of the acid may be determined, I refer to special treatises on the chemistry of the urine.

*Chlorides.*—The chlorides in the urine are derived from the food; they correspond closely with the amount of salt ingested. In consequence, the sodium chloride—the main chloride in the urine, for it contains but little potassium chloride and calcium chloride—is, even in health, liable to great fluctuations; the mean in twenty-four hours is estimated by Vogel and Parkes at 11.5 grammes, or about 177 grains. Bischoff states the average at 14.73 grammes. Large quantities of chlorides are excreted after active bodily or mental exercise, smaller quantities when the body is at rest, as at night. In disease, very various amounts are eliminated with the urine. In cases of chronic indigestion, of dropsy, and during an ague-fit, the chlorides are diminished. In typhus fever and in acute inflammatory affections they sink to a low level, and rise again in convalescence: an increase after a diminution is thus always a favorable sign. We may study these changes in pleurisy and pericarditis, but especially in pneumonia. At the period of hepatization the chlorides are absent from the urine, and appear in increased quantity in the sputum; during resolution they reappear in the urine; between these stages there is, probably, a determination of the salt to the inflamed organ.

Sodium chloride is detected by acidulating the urine with nitric acid and adding a solution of silver nitrate; a dense white precipitate of silver chloride quickly appears, insoluble in nitric acid, but soluble in ammonia. The amount of the chloride is approximately estimated by comparison with healthy urine, or by employing the method of Hofmann and Ultzmann. According to this method, if in using a solution of silver nitrate of definite strength, 1 to 8, we find curdy masses of silver chloride falling to the bottom, which on shaking the glass do not separate, we judge the chlorides to be in normal amount. If the precipitate of silver chloride be small,  $\frac{1}{10}$  per cent. or less, a simple milky turbidity arises and no curdy mass deposits; whereas if the

chlorides be entirely wanting there is neither milky cloud nor turbidity. If the urine contain much albumin, it should be coagulated and filtered off before the test is applied.

*Sulphates.*—Sulphates are found in the urine in large quantities. They consist of potassium sulphate and sodium sulphate, the former in excess. Like the alkaline phosphates, they are soluble in the urine. To detect them, a few drops of nitric acid are added to urine, and subsequently from fifteen to twenty drops of a saturated solution of barium chloride, when a white precipitate insoluble in acids occurs. If there be merely an opaque milky cloudiness, the sulphates are in normal quantity.

The sulphates are obtained in part from the food, in part from the oxidation of the sulphur entering into the constitution of the albuminous substances of the body and the subsequent union with a base of the sulphuric acid which is formed. They are enhanced by an exclusively animal diet, after violent exercise, in acute rheumatism, in pneumonia, and in all acute febrile processes with a large excretion of urea; in fact, their increase is apt to go hand in hand with that of urea. The administration of potassium raises in a striking degree the proportion of the sulphates. The sulphates show decrease during an exclusively vegetable diet and in urine of low specific gravity.

The average daily quantity of sulphuric acid passed in the urine is about 2 grammes. Vogel gives an easy method of determining approximately whether it is increased or diminished. After ascertaining the whole amount of urine in twenty-four hours,—say it is 2000 cc., and then each 100 cc. would contain 0.10 gramme of sulphuric acid,—100 cc. are rendered acid, and as much of a test-solution of barium chloride\* is added as corresponds with 0.05 gramme of the acid. The mixture is now filtered, and if the filtered liquid be not made turbid by the barium chloride, we may infer that the patient has secreted less than 1 gramme of sulphuric acid in the twenty-four hours. If the liquid, however, be rendered turbid by barium chloride, a further quantity of this agent, corresponding with 0.5 gramme of sulphuric acid, is added; and if the filtrate be still rendered turbid, it is evident that the

---

\* Made generally by dissolving 30.5 grammes of crystallized barium chloride, powdered and air-dried, and diluting the solution up to 1 litre; 1 cc. of it then equals 10 milligrammes of sulphuric anhydride.

quantity of sulphuric acid is greater than normal. In addition to the sulphates proper, the urine contains small quantities of derivatives of sulphuric acid, known as the ethereal sulphates, one of which is phenylsulphuric acid. The origin of these bodies is believed to be in some way connected with the action of putrefactive processes dependent on micro-organisms, but as yet no definite information as to their exact clinical significance is at hand.

*Kreatin and Kreatinin.*—These substances found in the urine are purely excrementitious, and are derived from a disintegration of the muscular tissue. Kreatinin is the product of the change of kreatin. From 0.5 to 1 gramme is excreted daily.

But few observations have as yet been made on the increase of kreatin, or on its significance in showing the activity of nutrition in the muscles in health or in disease. Active muscular exercise augments the quantity; and the same effect is probably produced by all spasmodic affections, and, as Munk has shown, at the height of acute disease, while kreatin is diminished during convalescence, and in advanced degeneration of the kidneys.

Both kreatin and kreatinin are generally included, in analyses, under the head of nitrogenous bodies. Under the microscope the crystals of kreatin are colorless and beautifully transparent. Their appearance, as well as that of kreatinin, is faithfully represented in Robin and Verdeil's plates.\*

**Presence of Abnormal Substances in the Urine.**—Here may be mentioned the ingredients, such as bile and blood, observed in the urine in disease only; and along with them I shall notice those constituents the occurrence of which in healthy urine is occasional, but of which it is certain that their presence in any marked degree is abnormal.

*Oxalate of Lime, Calcium Oxalate.*—There can be no doubt that the crystals are not found in large numbers except in a morbid condition. Some pass habitually a considerable quantity. They are generally persons weighed down by care and anxiety, or who overtask their brains by incessant application to study, or weaken their nervous power by excessive sexual indulgence or by masturbation. Sometimes they are troubled with frequent seminal emissions and irritation of the bladder, or they are dyspeptic,

---

\* *Traité de Chimie anatomique*, Paris, 1853.



and suffer from uneasiness after meals ; but the appetite may be good and the digestion unimpaired. They are always languid, and either very irritable or very dejected. Frequently they complain of loss of memory, and of a sensation of weight or of a dull pain across the loins. They are liable to boils and carbuncles, grow thin, and evidently are generally out of health. The urine is of high specific gravity, shows an increase of urea, and ordinarily a cloudy deposit consisting of mucus and the crystallized oxalates. Not unfrequently traces of albumin are associated with small amounts of calcium oxalate in urine apparently otherwise normal.

This is the disorder called by Golding Bird *oxaluria*, and is generally combined with tissue-changes and increased excretion of urea. Its existence as a separate affection has been denied ; but I believe the clinical association of a considerable number of oxalates with the symptoms mentioned to be undoubted. The presence of uric acid and of oxalates is not uncommon in lithæmia. The origin of the oxalic acid is not certain. It is generally the product of incomplete oxidation of organic matters in the body, as well as of sugar, of starch, and of the salts of the vegetable acids. Probably in the first class of cases alone are the constitutional symptoms described present. In the others we may at times detect evidence of the irritation of a calculus, or of disease of the bladder or the kidneys. Acid fermentation of mucus in the urinary passages also occasions it.

Calcium oxalate may be detected in the urine when articles which contain it, such as sorrel and the rhubarb plant, have been eaten, or after the free use of carbonated drinks. It may be also found in the urine of those recovering from severe acute maladies, and is encountered, but only in very small quantities, in the urine of healthy persons. But in neither instance is it at all permanent, nor can the presence of a few crystals be looked upon as of the least importance.

The microscope is incomparably the readiest means of detecting the salt. This appears in the urine in well-defined octahedra of most varying size, and in dumb-bell bodies. The former are much the more common and characteristic, for the dumb-bells are not frequent, nor is this formation peculiar to calcium oxalate. Occasionally, long or pointed octahedra or prismatic crystals are observed. All forms are unaffected by acetic acid.



The oxalates are often mixed with deposits of urates or uric acid ; a fact which some use as an argument that oxalic acid is but the direct transformation of uric acid, just as others regard it as a lower degree of oxidation than is necessary to form the products of disintegration into urea. Sometimes—Beneke says constantly—

FIG. 61.



Calcium oxalate crystals.

the earthy phosphates coexist in large amount with the oxalates. Occasionally the irritation from the passage of the crystals gives rise to tube-casts. A case came under my observation years since in which a patient suffering from a protracted attack of oxaluria voided for weeks, along with the oxalates, hyaline, exudative, or small waxy casts. Neither heat nor nitric acid detected albumin. Under treatment, the crystals disappeared from the urine, and with them the casts. The gentleman recovered perfectly. The urine examined ten years afterwards showed not the slightest signs of degeneration of the kidneys.

*Leucine and Tyrosine.*—Both these substances are the result of the decomposition of highly nitrogenous animal matter, are very similar, and are usually associated. They replace urea, and have been found in the urine only in disease, as in yellow atrophy of the liver, in typhoid fever, in smallpox, in phosphorus-poisoning, in cancer of the liver.\* They are either spontaneously deposited, or

---

\* Vaughan and Beringer, Contributions from the Chemical Laboratory of the University of Michigan, vol. i., 1882.

form a deposit if a small quantity of urine be evaporated. Tyrosine is readily detected by the microscope. It crystallizes in long, very fine, shining needles, which may congregate in globular bodies.

Hofmann has proposed the following delicate chemical test for tyrosine. A solution of mercuric nitrate, nearly neutral, is to be treated with the solution suspected to contain tyrosine: if it be present, a reddish precipitate is produced, and the supernatant fluid is of a very dark rose-color. Leucine crystallizes in granular masses, consisting of roundish globules, sometimes of concentric form, and for the most part of yellowish color, and resembling oil-drops, but, unlike oil, is not dissolved by ether. The chemical test for leucine is to place the suspected deposit on platinum foil and then to evaporate it with nitric acid. The residue is moistened with caustic soda, and this mixture is carefully heated over a spirit-lamp. It is gradually condensed into oily-looking drops,—a property which Scherer has pointed out as characteristic of leucine.

*Bile.*—The occurrence of bile in the urine imparts to it a very dark color. Its presence is a proof that the bile passes into the blood, and that the kidneys are performing a function forced on them by the deranged action of the liver, or by an impediment in the biliary passages. All the constituents of the bile may appear in the urine, or only the pigment, without the acids or their salts. The pigment is sometimes found transiently, and in small quantities, without yellowness of the skin: its more permanent and marked occurrence is, however, always attended with jaundice. It may be discerned both before the discoloration of the skin is noticeable, and after it has lost its yellow hue. The biliary acids are not of necessity present in the urine of icterus.

The detection of the coloring-matter of bile is effected by pouring a small quantity of urine on a white plate; a drop of the yellow fuming nitric acid of commerce is then permitted to fall on the thin layer of fluid. Soon a play of color takes place, beginning with green and blue, passing to violet and red, and often finally to yellow or brown; the green is the predominant and the most characteristic of the colors. According to Frerichs,\* this reaction may fail in cases where the other symptoms of jaun-

---

\* Diseases of the Liver, Sydenham Soc. Transl., vol. i. p. 100.

dice are undoubted, owing to the bile-pigment having already passed through stages of transformation. When this is the case, the urine is at one time of a brown or brownish-red color, and becomes red on the addition of nitric acid ; at another time it is of a deep red, which is converted by nitric acid into a dark bluish-red. Murchison has made a similar observation\* in rare cases where jaundice has resulted from a blood-poison, and he has frequently found the urine to present these characters where there has been no jaundice, yet obvious derangement of the liver.

Heller's test is also very easily performed. In a small beaker glass containing about 6 cc. (1.62 fluidrachms) of pure hydrochloric acid mix enough urine to discolor this, then allow nitric acid to trickle along the sides and form a layer underneath. A beautiful play of colors takes place at the point of contact, and, on stirring up the mixture with a glass rod, throughout it.

The following is also a delicate test for bile. Add to the urine some calcium chloride solution, and then solution of sodium carbonate. The precipitate will contain any bile-pigment, and may be collected by agitating the liquid with chloroform. The chloroform solution should be agitated with water and acidulated with acetic acid. Any bilirubin will color the chloroform yellow, which will become green on adding the acid.

If the urine contain only altered biliary coloring-matters (biliofusin), they may, according to Hofmann and Ultzmann, be recognized as follows. A piece of clean white linen is dipped into the urine and then allowed to dry ; it is discolored brown. Further confirmation is found in a very dark reaction for urophæin (by adding about double the quantity of urine to strong sulphuric acid), the urine appearing not garnet-red, but black. A similar reaction is produced only by the presence of sugar and of blood-coloring matter, both of which can be excluded by the appropriate tests.

The *biliary acids* are sought for by *Pettenkofer's test*. It consists in adding a few drops of a solution of sugar to a small portion of urine contained in a test-tube or in a china dish, placed in cold water. To this mixture an excess of concentrated sulphuric acid is added, drop by drop. The fluid assumes a yellowish-

---

\* Clinical Lectures on Diseases of the Liver.

red color, which, if bile be present, passes into a crimson or violet. The test is not applicable to albuminous urine, unless the albumin be first coagulated and separated. And it is inconclusive; for urine containing an excess of indican may display, when thus treated, a reaction exactly similar to that caused by the bile acids. Moreover, Neubauer and Vogel state that oleic acid and albumin give analogous reactions.\* The spectrum, which shows lines by F and near to E, affords, according to Schunck, the most certain test of bile acid; indeed, minute distinctions between the different coloring-matters cannot be attained except through spectroscopy.

A delicate test very generally used for biliary acids is *Oliver's test*. The test solution consists of half a drachm of pulverized peptone, four grains of salicylic acid, half a drachm of acetic acid, and distilled water to make eight ounces. The fluid is made transparent by repeated filtering. Twenty minims of urine are added to sixty minims of the test solution: if bile acids are in excess, a distinct milkiness quickly appears.

*Sugar*.—This substance is not a normal ingredient of urine, or exists only in traces too minute to be detected by the ordinary tests. When met with in normal urine it is probably due to the decomposition of the indican. Sugar may be found occasionally in the urine of those who live exclusively on a starchy diet, or who take large quantities of sugar; but the proportion even then is very small. Sugar appears in the urine after inhalation of carbon monoxide, and, as this is a common ingredient in illuminating gas, cases of light chronic poisoning giving rise to apparent slight diabetes are probably not uncommon. The urine secreted while under the influence of turpentine, ether, chloroform, chloral, or amyl nitrite is found to respond to the copper tests for sugar, but this may often be due to the medicaments themselves. Bordier† has grouped together many observations which led him to conclude that saccharine urine may be considered as an almost normal occurrence in the stage of recovery from acute diseases. Measles, pneumonia, erysipelas, all inflammatory fevers, are likely to exhibit it during convalescence. It may be detected in certain lesions of the brain and spinal cord. At Guy's Hospital the urine

---

\* Neubauer and Vogel's Analysis of the Urine.

† Archives Générales de Médecine, 1868.



of a large number of patients, laboring under various complaints, was found in several instances, particularly in cases of phthisis, to give a more or less marked reaction of sugar.\* A large and persistent amount occurs only in diabetes.

Urine holding sugar in solution is light-colored, of high specific gravity, and of peculiar smell. It rarely deposits sediments, and the excess of water in it may be large.

To detect the presence of sugar, several tests have been proposed, nearly all of which are easy of application. When albumin is present, it should be first separated by boiling and filtering.

*Trommer's Test.*—A few drops of a solution of copper sulphate are dropped into the test-tube holding the urine. Solution of caustic soda is now added in excess. If the fluid be saccharine, the faint greenish tint is changed to a deep blue, the precipitate which is formed when the alkali is first added being soon redissolved. On heating the blue mixture it becomes brownish, then yellow, and finally a reddish-brown mass of copper suboxide is thrown down, very different from the flocculent or greenish sediment noticed when no sugar exists. A very small quantity of sugar can be detected by this process: but, good as the test is, it has its drawbacks; for sugar is not the only substance which possesses the power of reducing the salts of copper. Chloral, cellulose, kreatinin, and to some extent uric acid and the urates, share with it this property. Furthermore, Beale has shown that the presence of ammonium salts will prevent the precipitation of the suboxide in urine containing but little sugar.

For the quantitative determination of sugar, Fehling's solution is generally employed. This may be made by the following formula, in which, in accordance with the recommendation of Allen, the quantity of Rochelle salt is rather greater than ordinarily given. 34.64 grammes of pure crystallized copper sulphate are dissolved in pure water, and the solution is made up to 500 cc. 70 grammes of caustic soda in sticks and 180 grammes of pure Rochelle salt are dissolved in 400 cc. of water, and this solution also is made up to 500 cc. The two solutions should be kept in separate well-stopped bottles. For use equal quantities are mixed

---

\* Researches on Diabetes, by F. W. Pavy.

as required. To determine the proportion of sugar in a sample, five cc. of each solution are mixed, diluted with about an equal volume of water, and brought to the boiling-point, in a porcelain basin. The porcelain dish with handle, called a casserole, is very convenient for this purpose. No precipitate nor loss of color should result from the boiling of the solution. The sample of urine is then added by small portions at a time, boiling between each addition, and watching the liquid so as to note the point at which all the blue color is removed. The condition is best determined by withdrawing the basin from the flame from time to time, inclining slightly, and allowing the red precipitate to settle. Any trace of blue color is easily seen. Ten cc. of the solution require .05 gramme of glucose to reduce them completely; the amount of urine used, therefore, contains this amount of glucose, and a calculation of percentage can easily be made. To get accurate results, the urine should be quite dilute, and if the qualitative tests indicate considerable sugar it will be necessary to dilute the liquid to five or even ten times its bulk. This dilution must, of course, be allowed for when making the final calculation.

Allen recommends the following test for cases in which there may be doubt as to the presence of sugar. Heat, to boiling, about ten cc. of Fehling's solution, and add a nearly equal quantity of the urine; heat for a few minutes, and then set aside to cool. If no turbidity is produced as the liquid cools, the urine is free from sugar, or, at most, contains less than  $\frac{1}{40}$  per cent. Fehling's test can also be used for peptone and propeptone. It gives at the point of contact in the test-tube a rose-pink or purple color.

Purdy recommends the following solution as a substitute for the ordinary Fehling's solution. The ingredients can be mixed in one bottle.

Crystallized copper sulphate.....	4.742	grammes.
Potassium hydroxide.....	23.50	"
Ammonium hydroxide (U.S.P., sp. gr. 0.9).....	450.00	cc.
Glycerol.....	38.00	"
Water sufficient to make.....	1000.00	"

All the materials should be pure. The mixture is prepared by dissolving the copper sulphate and glycerol in 200 cc. of the water by a gentle heat. In another 200 cc. of water dissolve the potas-

sium hydroxide. Mix the two solutions, and when the mixture is cold add enough water to make the volume 1000 cc. Of this liquid 35 cc. are exactly reduced by .02 gramme of dextrose. The presence of the ammonium hydroxide prevents the precipitation of the cuprous oxide, but does not interfere with the reduction. The test is applied by measuring into a flask 35 cc. of the solution, adding about 70 cc. of water, and bringing the mixture to full boiling, when the diluted urine is added by small portions at a time until the liquid is decolorized and clear.

*Boettger's Test.*—Add to the filtered urine about half its volume of sodium hydroxide solution and a pinch of pure bismuth subnitrate, and boil the mixture. Sugar will be indicated by a *black* precipitate. If sugar is not present, the precipitate will be white, or, at most, somewhat gray. This test is very delicate and tolerably free from fallacy. Dark-colored urines of high gravity may produce a gray precipitate, but it does not settle so rapidly nor so completely to the bottom of the tube. Only a pure finely-powdered preparation of the bismuth compound should be used for the test.

*Phenylhydrazin Test.*—Phenylhydrazin is a coal-tar derivative which possesses the property of forming crystalline compounds not very soluble in water with bodies of aldehydic or ketonic type, to one or the other of which classes the sugars belong. It is generally used in the form of phenylhydrazin hydrochloride, a soft, very light, brown crystalline powder with a geranium flavor. It is said to cause a persistent eczema when much in contact with the skin. The test may easily be conducted without danger. The following method seems, according to some comparative experiments made by Dr. Leffmann on the different published processes, to be the best. Fifty cc. of the urine are mixed with 0.75 gramme of phenylhydrazin hydrochloride and 1.0 gramme of sodium acetate, and the mixture is heated for one hour at least in a test-tube placed in boiling water. Very small amounts of sugar will produce a marked yellow precipitate—a compound of sugar with the reagent—which under moderate magnifying power exhibits either brush-like branchings or more decidedly radiate crystals, somewhat like chestnut-burs. A flocculent brownish precipitate or small brown globules should be disregarded. The precipitate is almost characteristic of sugar, but cannot by the microscope be distinguished

from a similar precipitate by glycuronic acid, a rare substance which is closely allied to dextrose in structure. The distinction can be made only by collecting the precipitate and determining its melting-point. The test, however, is principally of value in distinguishing those cases in which very limited reducing action is exhibited by a sample of urine when tried by the ordinary tests. As a delicate reaction for true sugar it does not seem to possess the great advantage over Boettger's test that has been claimed for it.

Further tests, though now not much employed, are *Moore's test*—boiling the urine with an equal part of potassium hydroxide—and the *fermentation test*.

Other forms of sugar, such as *sugar of milk*, may be found in the urine. Sugar of milk has hitherto been detected only in the urine of lying-in and of nursing women.

*Acetone*.—Ralfé gives the following test. About four cc. (one drachm) of sodium hydroxide solution containing a gram (fifteen grains) of potassium iodide are placed in a test-tube and boiled. An equal volume of urine is then poured in cautiously, so as to float on the surface of the alkaline liquid. At the point of contact a ring of phosphates will be formed, and after a few minutes will be colored yellow and studded with crystals of iodoform. Alcohol and lactic acid also give this result.

On adding a very dilute alkaline solution of sodium nitroprusside to a fluid containing acetone, a ruby-red color is produced which in a few minutes changes to yellow.

*Diacetic acid*, a body somewhat similar to acetone, is occasionally present in urine. It is generally recognized by the red color produced by solution of ferric chloride.

*Inosite*.—This is a substance not belonging to the sugars, but having some of their properties, and at times found in the urine. Inosuria is a symptom rather than a disease.\* The characteristic reaction of inosite is exhibited when a solution of the substance is evaporated with nitric acid nearly to dryness on platinum, and the residue, moistened with a little ammonium hydroxide and a solution of calcium chloride, is again evaporated to dryness: a marked rose-color appears,—which is not the case when true sugars are treated in the manner described.

---

\* Gallois, De l'Inosurie, 1864.



The presence in the urine of the *blood-extractives* indicates merely the escape of blood-material, and proves the existence of congestion or inflammation of some part of the urinary surfaces. Rees has pointed out \* that in Bright's disease the extractives can be found in the urine before albumin is met with, and also that they exist after the albumin has disappeared,—thus warning us, on the one hand, of the approach of albuminuria, and, on the other, against too early a belief in convalescence; for, as he justly observes, so long as the blood is losing its extractives so long is the patient in peril. The presence of the extractives also enables us to diagnosticate nephritic irritation from renal calculus before albumin, blood, or pus has appeared. To the delicate test by guaiacum for the crystalloids of the blood, which has been used to detect the prealbuminuric stage of Bright's disease, we shall presently more particularly refer.

*Albumin and other Proteids.*—The study of the various proteids occurring in urine is a matter of difficulty; and much uncertainty and confusion still exist with reference to them. The genito-urinary tract being a mucous area of great extent, abnormal secretions are frequent, and it is in many cases impossible to determine the boundary between health and disease. Thus, much discussion had been held as to the occurrence of albumin in *normal* urine, without any clear definition as to what is meant by the term normal. Efforts have been made to secure tests of extreme delicacy, but, while some of these have value in physiological investigation, they are often too delicate for practical work.

By the term albumin, unqualified, clinicians generally mean serum-albumin, and that meaning will be understood in this work. A proteid, derived from the mucous tissue, is generally present in urine. This has been designated *mucin*, but seems to be identical with a body called nucleo-albumin and also obtained from bile. Fibrin and hæmoglobin may appear, and also all the products of the transformation of proteids under the influence of digestive ferments, that is, the various proteoses and peptones. Recent research has shown that what is often designated peptone is an intermediate product,—an albumose.

---

\* Guy's Hospital Reports, 3d Series, vol. xiv. p. 431.

Albumin appears sometimes for a short time and then for a period is not found. Egg-albumin, it is stated, may show itself in the urine after the free use of eggs as food.

The tests for albumin depend on coagulation. The most important are :

*Heat ;*

*Nitric acid ;*

*Picric acid ;*

*Potassium ferrocyanide ;*

*Trichloroacetic acid.*

*Heat Test.*—Albumin is coagulated by heat of about 150° F. (65° C.). The application of heat to normal urine often causes a precipitate of phosphate. To avoid this fallacy a small amount of acid, nitric or acetic, is added. The test is best performed as described by Purdy : Mix a portion of the sample with about one-eighth its volume of a saturated solution of common salt, filter, and fill a test-tube nearly full with the mixture. Add two or three drops of acetic acid, and boil the *upper* stratum of liquid. The contrast between the two layers of liquid will be sufficient to indicate very small amounts of albumin. The salt solution prevents the interference of mucin, which is not precipitated under these conditions. This is a satisfactory method for the detection of minute amounts of albumin. Small quantities may be also found by thorough boiling of urine to which a few drops of acetic acid have been added, without admixture with the common salt solution. For the detection of notable quantities the following tests will be ample.

*Nitric Acid, Heller's Test.*—Fifteen drops of commercial nitric acid are placed in a somewhat narrow test-tube, and some urine poured slowly down upon it, the tube being considerably inclined. Another method is to put the urine in first and introduce the acid by means of a pipette, so as to form a clear layer at the bottom of the tube. A white ring forms at the point of contact. Urine in which this test does not show albumin may, for practical purposes, be regarded as not containing it.

Tests by this so-called underlaying method are conveniently made by the use of the albumin-test glass designed by J. A. Kyner. The precipitating substance, *e.g.*, nitric acid, is put in proper quantity in the tube, and the liquid to be tested is allowed

to flow through a filter folded in the usual way and placed in the funnel-shaped top of the glass.

In testing filtered urine it must be borne in mind that many forms of filter-paper will furnish enough soluble vegetable albumin to give distinct reactions with the more delicate tests for proteids. To avoid this error it will be best to use the centrifugal machine to secure a clear liquid.

Urine rich in urea sometimes forms a precipitate of urea nitrate. It may be distinguished from albumin by its crystalline character, especially after standing a few hours, and by its solubility when the liquid is warmed. Excess of urates may also produce a precipitate that might be mistaken for albumin, but the ring is irregular and will in a few hours become distinctly crystalline and can be easily determined under the microscope.

Resinous bodies administered as medicines are occasionally excreted by the urine, and are precipitated by the addition of nitric acid. They may generally be recognized and distinguished from albumin by their strong odor and by their solubility in alcohol.

According to Hofmann and Ultzmann, the precipitate produced by nitric acid in the cold may be taken as a rough quantitative approximation. If the white zone has a depth of from one-tenth to one-eighth of an inch, and appears clearly defined only when placed against a dark ground, the amount of albumin is less than one-half of one per cent.; if the zone is somewhat deeper, and visible without a dark ground, the amount of albumin is about one-half per cent.; while if the precipitate is flocculent and separates in lumps, it amounts to over one per cent. In urine containing alkaline carbonates an effervescence will occur when any acid is added, but this will soon cease and the coagulum will be formed. More accurate ways of determining the quantity of albumin in urine are by Esbach's albuminometer and Purdy's centrifugal method.

Sometimes urine is encountered on which neither the heat nor the acid test yields the customary result. This is owing to its containing modified albumin, *albuminose*. Such a case was pub-

FIG. 62.



Albumin test-glass.

lished by Bence Jones.\* No coagulation was produced by heat, and none by nitric acid, unless the urine was subsequently heated and permitted to cool. The solid that formed on cooling disappeared on heating. The patient was laboring under mollities ossium. Basham recommends the tincture of galls as a test for this modified form of albumin. There are several albuminous forms in urine in different conditions,—some, like the peptones, the result of incomplete digestion; others, like paraglobulin, derived from the blood.† Senator has taught us to view both peptones and albumoses as due to altered blood-composition.

*Picric Acid Test.*—This acid occurs in the form of bright yellow crystals, which may be used as a test by simply dropping them into the filtered urine, in falling through which they form a coagulum. The saturated solution may also be employed in the manner of the contact nitric acid test. The solution, being lighter than most urines, will form the upper layer. Picric acid makes a very delicate test, but shows about the same fallacies as the other acid tests. It also forms a slight precipitate with mucus, stains the skin yellow, and is somewhat explosive.

*Potassium Ferrocyanide Test.*—Twenty-five drops of strong acetic acid are thoroughly mixed with three times that amount of a solution of potassium ferrocyanide (1 in 20). A considerable volume of the urine is then added. Albumin if present will form a precipitate. Purdy states that the test thus applied is absolute evidence of albumin. The directions must be carefully followed.

*Trichloroacetic Acid.*—This is a solid, highly deliquescent body. It is corrosive, and should be handled with care. It is employed in the strongest possible solution, best obtained by allowing the solid to absorb water from the air until a solution is just formed. It is an extremely delicate reagent. D. D. Stewart investigated its action on many samples of presumably normal urine, and found that almost all would give some reaction. It is rather too delicate for general clinical work. It reacts with all proteids.

Halliburton sums up the reactions for the different *proteids* as follows :

---

\* Philosophical Transactions for 1848.

† Senator, Virchow's Archiv, Bd. lx., 1879; Brunton and Power, St. Barth. Hosp. Rep., 1877; Neubauer and Vogel, *op. cit.*, 7th ed., p. 384, Amer. transl.



If no precipitate forms on boiling after acidulating, albumin and globulin are absent. A precipitate may indicate both.

If no precipitate is produced after neutralizing the original liquid and saturating with magnesium sulphate, globulin and hetero-proteose are absent. If the urine gives no precipitate by the boiling test for albumin, nor with nitric acid in the cold nor when saturated with ammonium sulphate, peptone is the only proteid that can be present. Peptone may be detected by the so-called biuret reaction, which depends on the red color produced by adding solution of sodium hydroxide to the liquid to be tested and then a small amount of a dilute solution of copper sulphate. Other proteids react with this test, but give a reddish violet color.

The complete removal of all the other proteids from a mixture containing peptone by means of ammonium sulphate is difficult. Peptone occurs physiologically only during the puerperal state.\* It occurs pathologically during many varying conditions, especially where there is tissue degeneration. It is frequent in general paralysis.†

*Mucin, Nucleo-albumin.*—The reactions of this substance have been studied especially by D. D. Stewart,‡ who employed a solution of nucleo-albumin obtained from bile and believed to be identical with that occurring in urine. He found that solutions of citric acid both dilute and concentrated, used by the underlaying method, as noted in the cold nitric acid test for albumin, gave distinct contact rings. Picric acid associated with citric acid also gave such precipitates, but picric acid alone gave, with solutions containing not more than .02 per cent. of nucleo-albumin, only a tardily appearing haze.

If urine be diluted with water and then strongly acidulated with acetic acid, mucin is precipitated, and may be collected, re-dissolved in water by the aid of alkali, and again precipitated by acetic acid.

*Diazo-Reaction.*—Under this title, Ehrlich introduced some years ago a method intended to detect certain nitrogen compounds produced by febrile conditions, especially typhoid fever. The

---

\* Robitzschek, Zeitschr. f. Klin. Med., xxiv.

† Arch. Gén. de Méd., March, 1894.

‡ Medical News, July, 1894.

test is performed by mixing a solution of sulphanilic acid in hydrochloric acid with the urine, adding a small amount of solution of sodium nitrite (not nitrate), and then rendering the liquid strongly alkaline with ammonium hydroxide. On shaking the test-tube a carmine-red color appears in the foam, if the reaction takes place. Several observers have stated that this reaction occurs in typhoid fever after the fourth day, and, when faint, indicates a mild case. The reaction also occurs in phthisis pulmonalis when rapidly tending to death. There is, however, need of further demonstration of the value of this test.

*Blood.*—The passage of blood with the urine constitutes hæmaturia. The urine is of a red color, or of a more or less dingy or smoky hue, and deposits, on standing, a reddish-brown or a dark coffee-ground sediment. If much blood be present, small, irregular masses are seen at the bottom of the vessel.

The appearance of urine containing blood is therefore not uniform. The diagnosis is at once rendered certain by the use of the microscope. And only by this means can it be rendered certain; for urine may be red or black, from the admixture of various pigments derived from substances swallowed as food or medicine or belonging to the economy. Thus, beet-root, some kinds of strawberries, logwood, and rhubarb impart a deep red color, which may be the cause of groundless alarms; or urine deeply tinged with bile, or discolored by fever, may be thought to signify the occurrence of hemorrhage from the urinary passages.

The chemical tests for blood are, on the whole, inferior to the microscopic examination. We may have sometimes to resort to them. I have found a rough test in the addition of carbolic acid, which not only coagulates the albumin, but also changes the color of the fluid. It does not produce the same peculiar reddish tinge with bile, or, so far as I have tried, with any other substance. The guaiacum test is very accurate. It is said by Mahomed to detect infinitesimal traces of blood, or rather its characteristic crystalloids, when neither the microscope nor the spectroscope nor the nitric acid test for albumin affords any indication of their presence. It is especially valuable in the recognition of the prealbuminuric stage of Bright's disease, in which hæmoglobin appears in the urine before albumin.\* The test, as modified by Stevenson,

---

\* Medico-Chirurgical Transactions, 1874.

consists in adding to a few drops of urine in a small test-tube a drop of tincture of guaiacum and then a few drops of ozonic ether. The mixture is agitated, and as the ether collects at the top it carries with it the blue color produced by the hæmoglobin, leaving the urine colorless below. If saliva or a salt of iodine be present, the test is fallacious. The spectroscope affords a very delicate test. The characteristic bands of hæmoglobin of yellow and green are seen between D and E. If the hæmoglobin be in a state of destruction or reduction, only one broad band appears.

But the microscope, as already stated, is the means most employed. The corpuscles we detect with it are not always of uniform appearance, yet they are never collected in rouleaux. But, after having found blood-corpuscles to indicate the true nature of the changed hue of the excretion, the questions remain to be solved, at what point has the blood been poured out? Is it really from the urinary organs? and if it be from them, whence?—from the kidneys, from the bladder, or from some other portion of the tract? Again, what morbid state lies at the root of the hemorrhage?

Now, the first of these questions must always be answered at the onset. Blood may flow from the vagina or uterus and become mixed with the urinary secretion, or it may have been added for purposes of deception. In the former case, a careful inquiry into the state of these organs, or, if necessary, a digital examination, will eliminate the source of error; in the latter, drawing off the urine by the catheter will detect the imposture. When we have fully satisfied ourselves that the blood is derived from the urinary organs, the next point to be ascertained—and clinically its importance cannot be overrated—is, whether it proceeds from the kidney or from the bladder. To determine this, we have not only to study the character of the fluid excreted, but also to investigate closely all the conditions of the accident.

If the blood come *from the bladder*, it is not equally diffused through the urine; the fluid discharged is at first clear or nearly so, but at the end of the act of micturition is much more deeply colored, or pure blood, in a liquid form or in clots, is voided. Then, too, there is usually pain over the bladder, with a frequent desire to pass water, and a stoppage in doing so; the urine is generally alkaline.

When the blood is derived *from the kidney*, we discover, on

the one hand, pain in the lumbar region, and other symptoms pointing to the affected organ, the existence of albumin in considerable quantities in the urine, or the passage of gravel. Clots are not encountered in renal hemorrhage, except when the blood coagulates in the infundibulum or the ureter and is gradually forced downward. Such clots are of a whitish color, and generally of cylindrical shape. In their passage toward the bladder and out of the urethra they become often the source of distressing pain. They are very significant, yet they are not absolutely pathognomonic of renal hemorrhage; for coagula formed in the bladder may be retained there for some time, and lose their color before they are expelled. Sometimes we meet with little solid or gelatinous fibrinous coagula which bespeak simply localized fibrinous exudation from some part of the urinary passages.

Aid in diagnosis may be derived from the study of the shape of the clots, which for this purpose should be floated out in water. According to Hilton,\* they will oftentimes be exact moulds or casts of the cavity in which the blood was effused. Thus, coagula formed within the bladder have a somewhat irregular, circular outline, and are flattened in shape, with bevelled and serrated edges. The use of the microscope, furthermore, is very valuable in the differential diagnosis. The epithelium which is mixed with the blood from the kidney is not flat and in scales, like that from the bladder, but small and more or less round or columnar; nor are there fibrinous shreds to be seen. Hofmann and Ultzmann direct attention to the various size of the corpuscles as significant of the hæmaturia which attends parenchymatous affections of the kidney; besides the normal disk-formed corpuscles, which are, however, frequently quite colorless, very small, even dust-like blood-corpuscles are met with. Sometimes the blood-globules are seen to be collected on casts that have been moulded within the renal tubes. These blood-casts warrant an absolute conclusion as to the source of the hemorrhage. But they do not always occur; and their absence, therefore, is not so valuable a proof as their presence.

*Renal hæmaturia.*—When of renal origin, the hæmaturia is often due to congestion or an acute parenchymatous inflammation

---

\* Guy's Hospital Reports, 3d Series, vol. xiii. p. 19 *et seq.*



of the kidneys in infectious maladies, such as scarlatina, small-pox, malignant measles, and typhus. Here we have the history of the malady, and the presence of tube-casts, of blood-casts, and of a considerable amount of albumin, to explain the meaning of the hemorrhage. The blood is derived from the engorged and ruptured Malpighian corpuscles. But as regards the large amount of albumin present, we must not lay too much stress on this as indicating marked kidney implication. I have known it to happen where the kidneys were not affected. Irritant medicines, too, such as turpentine and cantharides, may cause congestion and bloody urine; and so do strains and blows on the back. In all these varied circumstances, a careful survey of the history and the symptoms will establish the diagnosis.

It has been recently stated \* as a diagnostic sign that in renal hæmaturia the blood-corpuscles show fragmentation, while this does not happen in vesical hemorrhage.

Renal hæmaturia of chronic character is generally due to cancer of the kidney; to cystic degeneration; to ulceration within the pelvis of the organ; or to irritation, with or without ulceration, set up by a calculus. In the first of these affections there is nothing in the urine to point out the source of the hæmaturia until the disease is far advanced, when pus, and sometimes disorganized cancerous tissue, may be discerned in the sediment. The manifestations of cystic degeneration are uncertain unless we can detect a large tumor; the signs of a non-calculous pyelitis are not definite, but hæmaturia is a rare symptom.

The existence of a calculus—the most common of the agents producing chronic hæmaturia—is indicated as the source of the hemorrhage by localized pain, and by the bleeding having followed active exertion, or a jar of the body from a fall, and by its recurring from time to time under circumstances like those just mentioned, favorable to the disturbance of a calculus lodged in the kidney. The presumption of this being the reason of the repeated bleeding is converted almost into certainty if there be also pus-cells present and if on any occasion a stony concretion have been expelled. Simon has catheterized the ureters and thus determined renal calculi; but this is not a procedure

---

\* Gumprecht, Deutsch. Archiv f. Klin. Med., liii., 1894.

easy to imitate. We find also hæmaturia in tubercular disease of the kidneys ; as in cancer, it is apt to be intermittent. Hæmaturia is at times met with in interstitial nephritis.

*Hæmoglobinuria*, or *intermittent hæmatinuria*, differs from ordinary renal hemorrhage ; the urine, although coagulable by heat and nitric acid, exhibits very few or no blood-corpuscles, but shows much granular pigment ; there is blood dissolution, and only the blood coloring-matter is found in the urine. We may use the guaiacum test to develop the presence of the dissolved hæmoglobin ; the hæmin crystals of Teichmann can be produced, and with the spectroscope we find the oxyhæmoglobin bands between D and E, occasionally also the methæmoglobin bands in the red. The urine voided is generally of a deep blood-color, and within an hour or two, perhaps, changes suddenly to a pale straw-color. It shows an increased proportion of urea. According to Greenhow,\* crystals of calcium oxalate are constantly passed during a paroxysm, and are absent at other times. The affection is unattended by any permanent lesion of the kidneys. It is paroxysmal in form, and not of malarious origin. It is ushered in by a chill, which is followed by an imperfect hot stage, and more rarely by sweating ; in some instances immoderate yawning and stretching of the limbs are the initiatory symptoms, and urticaria, great thirst, and local cyanotic appearances are observed. There is, indeed, a close association with Raynaud's disease. The temperature may be that of health, or may rise to be high. Transitory albuminuria may precede the attacks ; between them the urine is normal. Pain in the loins is not unusual. In the blood during the attack a marked diminution of red corpuscles is observed, as well as masses of granules and spindle-shaped bodies and other products of destructive change ; and it is very likely, as Ponfick maintains, that the blood condition is primary and the hæmoglobinuria secondary. The etiology of the disease is unknown. It often happens in syphilitic subjects. In those predisposed, brain-worry brings on attacks ; rest and food may prevent them. The influence of cold seems to be a very potent cause.†

There is an intermittent hæmaturia which is malarial. This *malarial hæmaturia* has been frequently observed in our Southern

---

\* Transactions of the Clinical Society, 1868, vol. i.

† Rosenbach, Berlin. Klin. Wochensch., 1880 ; Mackenzie, Lancet, Feb. 1884.

States, especially in Florida, Louisiana, and Texas, and in the East and West Indies. The hæmaturia may occur in daily paroxysms, or at longer but regular intervals. The bleeding sets in suddenly, but the urine soon clears up, though in some instances it remains persistently bloody. The urine is albuminous, contains casts, hæmoglobin, and generally only few blood-disks; it shows, then, a hæmoglobinuria rather than a hæmaturia. The attacks are mostly preceded by coldness of the extremities; some elevation of temperature may follow. When there are distinct fever and yellowness of skin, the hemorrhage from the kidney forms part of the disease known as hemorrhagic malarial fever. Malarial hæmaturia is more common in men than in women.\* It differs from ordinary paroxysmal hæmaturia above described in the decided malarial history, in the greater regularity of the paroxysms, and in the influence that quinine generally exerts on them. Malarial organisms have been found in the blood.†

There is also a form of hæmaturia which is endemic and depends upon the presence of a *parasite*, *Bilharzia hæmatobia*. It prevails in the Mauritius, certain parts of Cape Colony, Natal, Egypt, and Brazil. The parasite inhabits mainly the small vessels of the mucous membrane of the urinary passages and the kidneys, and it gains chiefly access to these parts during the act of bathing in rivers. Persons affected with the *Bilharzia hæmatobia* are often observed to pass small renal calculi of calcium oxalate having for their nuclei the ova of this parasite;‡ they may also present chylous urine. A similar *parasitic hæmaturia*, due to the *Filaria sanguinis hominis*, is met with in India.

Further, there is a hæmaturia peculiar to infants. This has been described by Parrot,§ under the name of *renal tubal hæmaturia*, and is characterized by hæmaturia and the accumulation in the tubules of the kidney of the red globules of the blood, and by a bronze discoloration of the skin, and cephalic symptoms.

---

\* Tyson, System of Pract. Med. by Amer. Authors, vol. iv.; see also Baker, Prize Essay, North Carolina Med. Journ., 1887; J. A. Stamps, Therap. Gaz., 1888, 3d S., iv.

† Stefanowicz, Wien. Klin. Wochensch., No. 20, 1893.

‡ Geo. Harley, Med.-Chir. Transact., vol. xlvii. p. 55, and vol. lii. p. 379; Handford, Brit. Med. Journ., 1887; Allen, London Practitioner, April, 1888, and Hill, London Lancet, May, 1888.

§ Archives de Physiologie, Sept. 1873.

Besides these causes, renal hemorrhage may occur from rupture of the kidney, of which it is the most prominent sign. It may also result from an altered state of the blood, as in purpura and in scurvy or in leukæmia; or we may find hæmoglobinuria in these states. We also find the dissolved blood in the urine after extensive burns and excessive doses of potassium chlorate.

*Vesical hæmaturia.*—One source to which this may be owing is a congestion of the bladder, as witnessed in fevers of a low type; another is irritant diuretics; another is blood-effusion from purpura or the hemorrhagic diathesis. Yet another is inflammation, whether acute or chronic, and whether of traumatic origin or brought on by a stone. In most of these contingencies the history of the case and the local symptoms establish the diagnostic distinctions; in arriving at which we are often materially aided by the introduction of a sound into the bladder.

Hemorrhage from the bladder, dependent upon tumor or malignant growths on its mucous coat, is usually attended with pain, with a constant desire to empty the viscus, and with considerable emaciation and a general cachectic condition. The fluid which is passed contains pus, and, as the malady advances, from time to time large quantities of blood. Yet the appearance of the blood in the excretion may be the first sign of disturbance.\*

Vesical hæmaturia, more frequently than renal, occurs as a vicarious discharge. Persons who are subject to bleeding piles lose blood occasionally from the bladder instead of from the rectum. But, in obscure cases of this kind, before arriving at a conclusion it is necessary to bear in mind that true vesical hemorrhoids are not uncommon.

Blood may be discharged from other parts of the urinary apparatus; it may come from the *prostate gland* or from the *urethra*. Now, in either case the bleeding is usually profuse, and large quantities of blood are passed pure, or unmingled with urine. Besides, there are local signs of disease of these parts, furnishing important points of discrimination.

Such, then, are the various conditions under which hæmaturia may be noticed. As regards its gravity, it is evident that this depends less upon the hemorrhage itself than upon the disorder

---

\* See case by Todd, Case XI., Lectures on Urinary Diseases.




of which the hemorrhage is a symptom. The flow of blood in itself is very rarely fatal. One of the worst consequences it may entail is the retention of a clot which serves as a nucleus for the formation of a calculus.

*Pus.*—Urine containing pus deposits an opaque creamy sediment or a glairy mass, is generally alkaline, and always slightly albuminous. If the deposit be agitated with a strong solution of caustic soda it becomes gelatinous. This is the chemical test for pus. But it is a clumsy one, compared with the rapid and absolute diagnosis of the pus-corpuscles by means of the microscope; this is especially valuable where the amount of pus is very small.

A deposit of phosphates may be mistaken for pus; a few drops of acetic acid clear it up, but do not influence pus. Sometimes a large amount of mucus is mixed with the purulent sediment, or a deposit due wholly to the former ingredient is so considerable that

FIG. 63.



Pus-corpuscles; those at the lower part of the field exhibit the action of acetic acid on the corpuscles.

it is mistaken for pus. Yet the mucous deposit shows distinct points of difference: it is less dense, and collects more in clouds at the bottom of the vessel; and it does not under any test show albumin. Again, the microscope is a valuable means of discrimination. In place of pus-corpuscles, quantities of epithelium are always seen to be entangled in the transparent mucus, and the

action of acetic acid develops the filaments of mucin. Sometimes, also, there are thin flakes of cylindrical bodies, unlike any appearance exhibited by pus. Yet, when the urine is strongly ammoniacal, even the microscope does not furnish a certain test; for the salts of ammonia obliterate the distinctive pus-globules and convert pus into a slimy mass, in which nothing but the nuclei may be distinguishable.

The occurrence of pus in the urine is a sign of suppuration somewhere in the genito-urinary system, or a proof that an abscess has opened into it. But as to the exact seat of the formation of the pus, its existence in the urine affords no clue. To some extent, however, we can judge of this by the microscopical appearance of the corpuscles. When these are round and well developed, with their characteristic nuclei readily brought out by acetic acid, they generally have their origin in a catarrhal inflammation of the mucous membrane of the bladder, and are apt to be associated with triple phosphates. On the other hand, as Vogel points out, pus-corpuscles of irregular contour, exhibiting irregular nuclei when treated with acetic acid, or an ill-defined granular mass, consisting of irregularly-shaped pus-corpuscles and partly-destroyed cells, indicate the probable existence of deep-seated suppuration, ulceration, or tubercular disease. The sudden appearance in the urine of large quantities of pus points to the bursting of an abscess; an abundant deposit of pus in acid urine is chiefly noticed in pyelitis.

*Fat.*—Fatty matter may occur in the urine in various forms and in different conditions. It may be found in the shape of globules, when oil or milk has been added to the urine for purposes of deception, or when the former article has been swallowed for some time in considerable quantities, as for instance during the administration of cod-liver oil. Fat is also encountered in globules of varying size, either free, in cells, or in tube-casts, as in fatty degeneration of the kidneys. Fat, too, may be found in the urine in phosphorus-poisoning and after fractures.

The tests for fat are its solubility in ether, and its microscopical characters. Lee and Atlee have pointed out\* an illusory detec-

---

\* Amer. Journ. Med. Sci., April, 1869, p. 357.

tion of fat. They found, in testing a specimen of urine, that the ether rose to the top so charged with matter as to resemble a half-liquid pomade. Separated by a pipette and spontaneously evaporated, it left a dirty-whitish greasy mass. A careful examination of this residue showed that, instead of consisting of fatty acids, it contained nothing but the normal constituents of the urine, for it was soluble in water, reappearing as normal urine. It was then ascertained that almost any urine will form an emulsion when violently agitated with ether, especially if the ether contain a small amount of alcohol. When, therefore, ether appears to dissolve out fatty matter from urine, the ethereal solution should be separated, and allowed to evaporate spontaneously, and if the residue be soluble in water it cannot be held to contain fat.

There is no certainty of the presence of fat unless the sediment be examined chemically and microscopically. The opalescence of urine caused by a sediment of urates has been mistaken for that from oily matter, and so also has been the pellicle which often forms on urine, and which consists not of fat, but of vibriones, fungi, and crystals of the triple phosphates. The "kystein" pellicle observed in the pregnant state is of similar kind, though some oily matter may enter into its composition.

In some cases fat is met with in a very finely divided state, imparting to the urine a milky look, which disappears on its admixture with ether. This condition, to which the name *chylous urine* has been given, does not depend upon any permanent morbid change in the kidney; the chylous character of the urine is intimately connected with the absorption of chyle, but precisely how the urine acquires that character is uncertain. It may be absent in the day urine and very marked in the night urine; there are at times small quantities of albumin present. The affection may continue for years without impairment of the general health, being always perceptibly increased by exercise. In the tropics chylous urine is found often in connection with the *Filaria sanguinis hominis*. The parasite has been detected in the blood of persons in the southern part of this country by Guitéras.

A urine which spontaneously coagulates soon after being voided, owing to *fibrin*, a fibrinuria, is very uncommon except in the Isle of France and in Brazil. A thick urine may be due to pus dis-

solved in alkalis, as in certain bladder affections. But the thick matter is at once greatly thinned by water, and on the addition of acetic acid a white precipitate of alkaline albuminate falls.\*

**Sediments.**—In connection with the ingredients of the urine, the nature of the urinary sediments has been discussed, and it has been insisted that they cannot be accurately determined save by a microscopical examination. I shall here group together only their general characteristics :

1. A light and flocculent cloudy deposit is commonly mucus, entangling epithelial cells, bacteria, or spermatozoa.
2. A dense, abundant, white deposit is generally composed of urates or phosphates ; but it may be pus or extraneous matter.
3. A yellow or pink deposit is almost always due to urates.
4. A granular or crystalline deposit, of reddish or dark-brown color and small in quantity, is uric acid.
5. A dark, sooty or dingy-red deposit is blood.

The following table may serve a useful purpose, in showing how both the sediments and the soluble urinary ingredients are affected by the reagents commonly employed :

TABLE EXHIBITING THE SIGNIFICANCE OF THE MAIN CONDITIONS AND THE ACTION OF THE MAIN REAGENTS EMPLOYED IN THE EXAMINATION OF THE URINE.

SPECIFIC GRAVITY .....	{	High .....	{	Urine high-colored.....	{	Increase of urea, uric acid, etc.
		Low .....		Urine pale.....		Diabetes.
HEAT.....	{	Throws down deposit .....	{	Urine high-colored or normal	{	Certain forms of Bright's disease.
				Urine pale.....		Excess of water.
	{	Dissolves deposit	{	Soluble in nitric acid .....	{	Phosphates.
				Insoluble in nitric acid .....		Serum-albumin.
				Does not dissolve deposit .....		Serum-globulin.
						Urates.
						Uric acid.
						Phosphates.

\* Hofmann and Ultzmann, *op. cit.*



TABLE EXHIBITING THE SIGNIFICANCE OF THE MAIN CONDITIONS AND THE ACTION OF THE MAIN REAGENTS EMPLOYED IN THE EXAMINATION OF THE URINE.—*Continued.*

NITRIC ACID....	Precipitates.....	Quickly .....	{ Albumin.
		More gradually..	{ Uric acid. Urea nitrate (crystalline).
	Dissolves.....	Earthy phosphates. Alkaline phosphates. Oxalates.	
HYDROCHLORIC ACID.....	Produces play of color.....	{ Bile-pigment.	
	Precipitates.....	{ Uric acid.	
	Transforms.....	{ Urates into uric acid.	
SULPHURIC ACID.....	Changes color of urine .....	Change of color to violet.....	{ Uroxanthin or indican.
		Brown.....	{ Urohæmatin.
		Crimson or violet (if sugar have been added)....	{ Biliary acids.
ACETIC ACID...	Precipitates deposit (not soluble in excess of the acid).....	Violet.....	{ Indican.
		Mucin (nucleo-albumin).	
	Precipitates with potassium ferrocyanide.....	{ Albumin and albumoses.	
PICRIC ACID...	Precipitates.....	Albumin, albumoses, peptones.	
		Red deposit—blood.	
		Slowly-developed haze—mucin.	
SODIUM HYDROXIDE..	On boiling, turns urine brown....	{ Sugar.	
	Dissolves .....	{ Uric acid.	
	Forms gelatinous mass .....	{ Deposits of urates Pus.	

TABLE EXHIBITING THE SIGNIFICANCE OF THE MAIN CONDITIONS AND THE ACTION OF THE MAIN REAGENTS EMPLOYED IN THE EXAMINATION OF THE URINE.—*Continued.*

AMMONIUM HYDROXIDE..	Precipitates .....	{ Earthy phosphates.
	Dissolves .....	{ Cystin.
BARIUM CHLORIDE....	Precipitates.....	{ Deposit, soluble in free acid. { Phosphates.
		{ Deposit, insoluble in acids. { Sulphates.
SILVER NI- TRATE.....	Precipitates.....	{ Yellow deposit, soluble in nitric acid and ammonia. { Alkaline phosphates.
		{ White deposit, insoluble in nitric acid, but soluble in ammonia. { Sodium chloride.
COPPER SUL- PHATE AND SODIUM HY- DROXIDE ....	Precipitates with heat yellowish-red deposit ....	{ Sugar.
	Turns violet.....	{ In cold..... { Peptone.
		{ With heat..... { Serum-albumin.
ETHER.....	Precipitates.....	{ Albumin.
	Dissolves.....	{ Hippuric acid, soluble in alcohol.
		{ Fat.
	Does not dissolve	{ Uric acid.

## URINARY ORGANS.

### Diseases of the Kidney of which Pain is a Prominent Symptom.

The group embraces acute inflammation of the kidney, and those painful affections classed under the term *nephralgia*.

**Nephritis.**—Acute inflammation of the kidney is not a frequent disease; it is chiefly observed in old persons and in damp climates. It may be occasioned by an attack of acute rheumatism,

by direct violence to the organ, or by the irritation of a calculus ; but probably its most frequent cause is exposure.

It begins with a chill, soon followed by fever of moderate degree ; there are nausea and vomiting, and at times diarrhœa with tenesmus. The urine is voided drop by drop ; it is red, and may contain blood. The patient complains of pain in the renal region, sometimes dull, at other times sharp and lancinating, and augmented by pressure and by moving. The pain is not limited to the kidney, but radiates to the diaphragm and to the bladder. With it are often associated numbness of the thigh of the affected side and retraction of the testicle. The disease may occur in both kidneys ; yet it rarely affects more than one. It lasts from one to three weeks, and generally terminates in resolution. But it may lead to suppuration and disorganization of the organ.

The disorder is recognized by the pain, the fever, the retraction of the testicle, and the appearance of the urine. It differs from an attack of colic by the signs of disturbance of the urinary organs, by the seat of the pain, and by the fever ; from rheumatic pains in the back, by the former of these symptoms. Then, in lumbago we rarely find much febrile excitement, nor are there nausea and vomiting, or numbness along the course of the anterior crural nerve ; but, on the other hand, the pain is much more influenced by movements, especially by stooping and such other motions as call the muscles of the back into play. Congestion of the kidneys is distinguished from inflammation by its affecting both sides, by the absence of protracted or severe pain, and by the comparatively slight derangement of the urinary functions. Further, the congestion is not idiopathic, and we can generally trace it to the swallowing of some irritating substance, or to the poison of a febrile malady, such as smallpox or typhus. From the passage of a renal calculus acute nephritis differs by the steady, less paroxysmal and less violent pain, which does not, as in renal colic, begin suddenly and end suddenly ; by the fever ; by the bloody urine, and by the absence of a history of previous attacks.

**Nephralgia.**—Severe pain in the kidney, unconnected with inflammation of the organ, is ordinarily caused by the passage of a calculus. There is no fever, although passing elevations of temperature may occur. Nephralgia exhibits a great similarity to

colic ; but this has already been discussed ; and in particular cases we are often much aided by the knowledge that in “renal colic” the patient has on a former occasion passed renal concretions.

The amount of pain varies according to the magnitude of the stone and its character. As a rule, calculi composed of *oxalate of lime* give rise to most pain. We may distinguish them by their roughness and irregularity and their brown or dark-gray color ; those of *uric acid* and *urates* are reddish and much softer, and not jagged, and, unlike calculi consisting of the salts of lime, are combustible on platinum foil, leaving a mere trace of residue, while the oxalate of lime calculus leaves considerable residue, and is soluble in mineral acids without effervescence. Calculi of the *mixed phosphates* are white, very brittle, soluble in acids, insoluble in alkalies, and fuse in the blow-pipe flame. The mixed phosphates rarely form a stone entirely, being often only an incrustation around a blood-coagulum or a foreign body, or having a kernel of uric acid. Indeed, the majority of phosphatic stones have uric acid centres, while calculi of uric acid or its salts possess, as a rule, the same composition throughout ; calculi of oxalates have often a nucleus of uric acid and a crust of phosphates. *Xanthine* and *cystine* are the rarer constituents of stones. The former, like uric acid and the ammonium and sodium urates, is consumed by heat, and burns without visible flame, but the murexide test exhibits an orange-yellow color ; cystine burns with a bluish-white flame, emitting an odor like that of burning fat, and the powder is soluble in dilute ammonia. The crystallization of the ingredients of the urine forming a calculus is very apt to take place around particles of mucus.

As already stated, we have in the severity of the pain a sign indicative of the nature of the case. Still, there are states in which *paroxysms of pain* referred to the neighborhood of the kidney are attributable to other causes than the passage of a calculus. Leaving out of consideration that doubtful disease, pure *neuralgia of the kidney*, we find a few affections—very rare, it is true—which closely simulate the passage of a renal calculus.

The first of these is the pain occasioned by an *inflamed and ulcerated ureter*. Todd relates a case of the kind.\* The patient

---

\* Clinical Lectures, Lecture II., on Diseases of the Urinary Organs.



had severe attacks of lancinating pain, referred to the right loin, lasting for weeks, and accompanied by constant and intractable vomiting. The urine contained pus in varying quantity, but neither blood nor calculous matter could be detected. At one time he continued free from any paroxysm for four years. After death the most careful search was made for a calculus, but none could be discovered. The ureter of the right side was thickened throughout the greater part of its course, and deposits of lymph adhered to its mucous membrane. A somewhat similar train of phenomena may occur from an irritation or inflammation of the ureter caused by the poison of rheumatism or gout, although the paroxysms of pain are apt to be neither so severe nor of so long duration.

Another morbid condition closely resembling the passage of a renal calculus may result from *malarial poison*. How close this resemblance may be, the following case will show :

A soldier, twenty-four years of age, of fair complexion, and evidently of strong constitution, was seized rather suddenly with pain over the left kidney. The loin was sensitive to the touch, and appeared somewhat red and swollen. The skin was hot ; the pulse 100. The urine was not found to be abnormal, though containing a reddish coloring-matter. The pain continued for several days, becoming more severe, notwithstanding that by direction of Dr. Hilborne West, under whose charge the man was, and with whom I saw him, six ounces of blood were drawn from near the affected part. On the fourth day of the disorder the patient was assailed with excruciating pain along the course of the ureter, attended with the voiding, at short intervals, of a high-colored urine. The attack lasted from six o'clock in the evening until five o'clock the next morning, leaving the patient much exhausted ; the only relief throughout its duration being obtained from the inhalation of chloroform. At six o'clock that evening another seizure, of equal violence, set in ; and, after the lapse of twenty-four hours, again another. Seeing the recurrence of the paroxysms at about the same time of each day, and learning from the patient that a few months before he had had a remittent fever, which had left behind an irregular intermittent, we resolved upon the administration of large doses of sulphate of quinine in the interval between the paroxysms. The seizure did not take place

that night; but, the remedy being a day or two afterward suspended, the fourth night was again a night of anguish. The antiperiodic was resumed, and continued, in lessened doses, for three weeks. The patient remained under observation for about six weeks after the last attack, gradually recovering his health and spirits. When he was lost sight of, there was still a dull pain in the left lumbar region, with inability to stand erect; but no return of the excruciating intermittent pains.

In a case of this kind it is evident that nothing but a knowledge of the history of the patient, and the noting of the regularly-recurring onsets of the pain, could have led to a correct appreciation of its cause. We sometimes meet with a so-called neuralgia of the bladder, of similar origin, and having much the same symptoms, except that the distressing pain is referred to the bladder. As in the case just detailed, the attacks occur at night.

These remarks are all based on the assumption that the renal pain is very severe and paroxysmal in its character. Let us now briefly inquire into the significance of a steady and less acute pain, premising that we have excluded from consideration abdominal aneurism, affections of the muscles of the back, of the spine, and of the tissues surrounding the kidney, in which diagnosis, of course, we are materially assisted by an examination of the urine.

An ingenious application of the effect of cocaine has been made in the study of pain. If a urethral injection of a twenty-per-cent. solution of cocaine immediately relieve pain in the kidneys, E. H. Fenwick points out,\* we may recognize a transient and unimportant cause for it, such as congestion, uric-acid urine, pressure of the colon. If the renal pain be uninfluenced, a stone or renal cancer may be suspected. Injections of cocaine into the bladder are not always, however, free from risk.

We meet with *persistent pain* referable to the kidney itself, in inflammation of the organ, especially in that variety of inflammation affecting the infundibula and pelvis, termed pyelitis. We also encounter it in malignant disease of the kidney; sometimes, although it is not then of long duration, from the irritation of concentrated and highly-acid urine; much more generally from the presence of a stone lodged in the kidney. The pain in the

---

\* *Lancet*, May 5, 1888.

latter complaint often extends along the course of the ureter to the testicle, which is retracted and swollen. Not unfrequently there is also tenderness on pressure over the affected kidney, and the pain is greatly increased by active exercise; and it is not uncommon to find, associated with these exacerbations of pain, nausea and vomiting, and the appearance of blood in the urine.

There is yet another point in the diagnosis of the *passage of calculi* which we must not overlook,—namely, that the pain may be referred to other parts than to the region of the kidney and the course of the ureter. It may be felt near or at the sacrum, and not merely on one side; it may extend to the bladder and become associated with a painful spasm of this viscus and with the voiding of urine drop by drop; or to the testicle, which becomes sensitive and swells; or to the thigh, which feels numb; or it may be referred to the right hypochondrium and extend downward, but not be perceived in the loin. Under the latter circumstances there may be, with pain of great intensity, coexisting distention of the colon, vomiting, and constipated bowels, and the symptoms so closely resemble those of the passage of a biliary calculus that only the detection of blood in the urine prevents error.\* Again, as happened in two cases which came under my notice, the pain may be referred to the left hypochondrium or along the course of the colon, may be associated with soreness to the touch and with digestive disorders, and may closely simulate an organic lesion of the stomach or intestine. Nothing but careful and repeated examinations of the urine, and observing the irregular and whimsical course the supposed intestinal malady pursues, will enable us to arrive at a knowledge of the truth.

Nor must we be unmindful that a calculus may be months in passing, and that as it changes its position the seat of the pain changes. I had a case of the kind under my charge in a lady about fifty years of age. She suffered for weeks at a time from excruciating pains, beginning in the left kidney, then felt somewhat below it, and finally localized in the neighborhood of the left ovary. She was occasionally free from pain for five or six days. But it was only after fully nine months of recurring suffering that the passage of a calculus the size of a plum-stone

---

\* Case of Owen-Rees, Guy's Hospital Reports, 3d Series, vol. x.



followed by a discharge of large amounts of a gritty substance and a soapy-looking urine, removed her distress. The stone consisted of urates.

The symptoms of renal calculus may, after having existed for a longer or shorter time, entirely cease, owing either to the calculus becoming encysted and thus remaining innocuous, or to its obstructing the ureter, causing retention of the urine, and, by pressure, producing gradual atrophy of the cortical and tubular structures, the kidney being finally converted into a mere bag.

In concluding the subject, it will be useful to group together the symptoms by which we may infer the existence of a *calculus in the kidney*. They are: frequent micturition, often attended with pain at the end of the penis; pain in the loin, with or without accompanying soreness, occasionally passing suddenly into a violent paroxysm, with a tendency to shoot along the course of the ureter to the testicle and the hip of the aching side; and in some cases the discharge of pus due to coincident pyelitis. These symptoms become positive evidence if the blood-extractives be present in the patient's urine, or if this, when examined microscopically, be found to contain blood-corpuscles; or if we know that attacks of hæmaturia have previously happened, and that gravel or small urinary concretions have at any time been discharged. The presence, too, in the urine of microscopic calculi, as Beale shows, points to the existence of larger concretions in the pelvis or in the structure of the kidney. But all these indications are far from being always present. The renal stones may be so large that they cannot leave the kidney; we may have nothing but the symptoms of a pyelitis, which we suspect to be calculous, and even these symptoms may be wanting. To determine whether both kidneys are implicated in the calculous disease, we must examine the urine during the passage of a renal calculus. If the urine become perfectly healthy, when previously it has been abnormal, we conclude that it comes from a healthy kidney, and that the secretion from the diseased one is temporarily blocked up.



## Diseases marked by an Albuminous Condition of the Urine, associated with more or less Dropsy.

The chief of these diseases is Bright's disease. At the present day we hold that the disease which bears Bright's name consists of a group of maladies having the common feature of a more or less albuminous state of the urine. But, though I believe this view to be the correct one, I shall in this sketch prefer to consider the disorder in the main as it is seen separated by broadly-drawn lines into an acute and a chronic form, and then examine the further differences these present.

**Acute Bright's Disease.**—In this form of the affection the symptoms are of an acute character. Especially so is the dropsy, which is quickly developed and soon becomes the most marked token of the malady. The history of a large number of cases is as follows. After exposure to wet or cold, a fever sets in, accompanied by nausea, and by a dull pain in the region of both kidneys, extending along the ureters. The eyelids and face become puffy and swollen, and soon a general œdematous condition of the skin is observable, showing itself very plainly in the extremities, scrotum, and abdominal parietes. Subsequently dropsical effusions often take place into the interior cavities.

The same symptoms are noticed in the acute Bright's disease, or the acute parenchymatous nephritis, which so constantly attends scarlatina, except that, following as it does an exhaustive disease, there are from the onset much greater pallor and general debility. Acute parenchymatous nephritis is also met with, though far less frequently, in other infectious diseases, as in smallpox, measles, diphtheria. It may follow a lightning-stroke,\* or sewer-gas poisoning.†

The urine in the acute malady is of high specific gravity, and dingy from its admixture with blood. There is a frequent desire to void it, although the whole quantity passed is rather below the natural average. The urine contains a large amount of albumin; a microscopical examination brings to light casts, lined here and there with blood-corpuscles. As the malady progresses, these

---

\* Medical and Surgical Reporter, July 23, 1887.

† Lancet, March, 1894.

“blood-casts” disappear, and we find casts coated with epithelium, which may be normal or slightly fatty, and with free nuclei; or we observe slightly granular or hyaline casts; or we may discern leucocytes and long cylindrical ribbon-like mucous casts. Furthermore, crystals of uric acid, of urates, even of oxalates, and a considerable amount of renal epithelium, are often seen in the sediment. The normal constituents of the urine are considerably changed. The chlorides may have disappeared altogether; the phosphates are diminished; the uric acid and the pigments are increased. The amount of urea fluctuates much: it is generally diminished.

There is moderate fever, with a temperature of about  $101^{\circ}$ ; the pulse, however, may be quick, tense, and full. The skin is

FIG. 64.



Epithelial casts and epithelial cells from the kidneys found in a case of acute Bright's disease (*acute desquamative nephritis*); magnified about 460 diameters.

generally harsh and dry; nausea and vomiting are of common occurrence.

The urgent symptoms last ordinarily for several weeks. When recovery is about to take place, they abate; the temperature becomes normal, the skin moist, and hand in hand with a diminution of the dropsy the quantity of the urine largely increases and the albumin gradually disappears. But this, although fortunately the common, is not the invariable issue. The disease may gradually lapse into a chronic form. Or a certain amount of albumin may remain in the urine; and after exposure this increases, and

the dropsy and most of the acute symptoms return. In some instances of the malady, not in many, there are numerous tube-casts in the urine, but little albumin.

There is a form of acute Bright's disease due to a bacillus. Letzerich \* describes it as "*nephritis bacillosa interstitialis primaria*." It occurs in children, runs its course with a moderate fever in from two to six weeks, and generally ends in recovery. The urine contains red blood-corpuscles, a few leucocytes, only small amounts of albumin, but great numbers of bacilli, shorter and thicker than the tubercle-bacilli, and easily stained with methyl-violet. An infectious nephritis also has been described due to the *bacillus coli communis*.†

Whatever the attending circumstances, the risk to life, when an attack of acute Bright's disease has been prolonged, is greatly increased by the supervention of local inflammations,—as of the pleura, lungs, peritoneum, or pericardium; or by the sudden effusion of fluid into the pulmonary structure; or by the retention of urea in the blood and consequent uræmic intoxication.

The recognition of the disease is readily effected. The puffy, pale face; the general dropsy; the albumin in the urine, associated with tube-casts,—form a combination of signs so remarkable that it is difficult to mistake their meaning. Many of the same phenomena are encountered in the chronic form of the malady: therefore, what is about to be said of the differential diagnosis of the acute complaint may be in the main applied with almost equal correctness to the chronic ailment.

The chief disorders with which acute Bright's disease is apt to be confounded are:

ACUTE NEPHRITIS;

SUPPURATIVE NEPHRITIS; PURULENT URINE;

HÆMATURIA;

SIMPLE ALBUMINURIA;

PULMONARY ŒDEMA;

PLEURISY AND PERICARDITIS;

DROPSY;

COMA; CONVULSIONS.

\* Neurol. Centralbl., 1887, quoted in Sajous' Annual, 1888, p. 483.

† Fernet et Papillon, Bull. et Mém. de la Soc. Méd. des Hôp., 1892.

*Acute Nephritis.*—This differs from acute Bright's disease by its affecting generally only one kidney, by the much greater pain and tenderness in the lumbar region, by the retraction of the testicle, and by the higher degree of febrile excitement. Then, too, the deeply-colored urine which is voided contains little or no albumin.

*Suppurative Nephritis; Purulent Urine.*—In rare cases the suppurative process may coexist with Bright's disease. But, on the whole, the two disorders are distinct and may be readily discriminated. We find pus of renal origin in the urine, in consequence of pyelitis or of abscess of the kidney. The former is very generally linked to the irritation of calculi, or is an infective process; the latter shows a fever of a remittent type, and often a well-defined swelling is felt in the lumbar region and extending far downward. All this is very different from Bright's disease. Then, we detect blood and pus in the urine of cases of pyelitis or of renal abscess, and any casts that are found are apt to be covered with pus corpuscles.

*Hæmaturia.*—In hæmaturia, if we can speak of it as a separate disease, there is albumin in the urine; and, on the other hand, traces of blood as well as of pus may be present in the urine of Bright's disease. But, as in purulent urine, the quantity of albumin met with in hæmaturia is small; in fact, it is in exact proportion to the amount of blood or pus the urine contains; whereas, on the contrary, if the secretion from a Bright's kidney be mixed with pus or blood, the amount of albumin is generally large.

*Simple Albuminuria.*—By this is meant an albuminous urine unconnected with any marked structural lesion, except congestion,—such an albuminuria as is sometimes observed as a transient phenomenon in the course of several diseases; as, for instance, in the exanthemata, in typhus, in cholera, in hectic fever, in chronic congestion of the liver, in oxaluria, or as a consequence of surgical diseases and operations.\* An albuminuria of similar kind is met with when the kidneys become congested from interference with the circulation, as in disease of the heart, or from the pressure of a gravid womb. Albumin in the urine may also be encountered in erysipelas, in diphtheria, in pneumonia, in acute

---

\* Henry Lee, Lectures on Practical Pathology and Surgery, 3d ed., London, 1870, vol. ii. p. 380.



rheumatism and in gout, consecutively to a burn, to a blister or a large mustard-plaster, or to the use of salicylic acid or of turpentine or of carbolic acid. But in all these conditions the quantity found is small and transitory, very unlike what it is in the persistent albuminuria of Bright's disease, and the urine is usually dense and high-colored. Then the constitutional symptoms and the general clinical features in the morbid states referred to are so dissimilar to those of Bright's disease that they become a safeguard against error.

Moreover, there is really often more than mere congestion; there is present a parenchymatous inflammation to a limited degree, and of a transitory kind. In all these cases of simple albuminuria the amount of albumin is apt to be small, and there are few, if any, casts. When found, these are generally of the epithelial or hyaline variety, and not highly granular or fatty.

In addition to these forms of simple albuminuria there is one of great importance to recognize, where the albumin happens in persons who in every respect seem healthy, and occurs shortly after partaking plentifully of food, especially of albuminous food, or after severe exercise, particularly in young persons at or near the age of puberty. Some of these cases are cyclic, occurring only at certain times of the day; in much fewer, the albuminuria is persistent. In the great majority of cases, indeed, there is a time in every day in which the urine is free from albumin. It is normal in quantity, normal or slightly increased in specific gravity,\* normal in the amount of urea it contains, and no tube-casts are found in its sediment, but uric acid and oxalates are not uncommon. The amount of albumin in these functional albuminurias is small, and there are no cardio-vascular changes; indeed, there is no symptom except the albuminuria to suggest disease. The bearing this form of albuminuria has on life assurance is leading to its very careful study.

There is a form of albuminuria that stands in close connection with excessive uric acid formation and oxaluria, to which I not long since called attention. The amount of albumin is generally small; hyaline and epithelial casts are found, though they are scanty. The specific gravity of the urine is high, and this, as

---

\* Purdy on Diseases of the Kidney.

well as the urates or the oxalates in the urine, is of much significance. Violent exercise increases this albuminuria of uric acid and oxaluria. The cases may be of short or of long duration; recovery is the rule.\*

*Pulmonary Œdema.*—Bright's disease is one of the most frequent causes of dropsical effusion into the air-cells; oppression in breathing, inability to lie in the recumbent position, cough, frothy expectoration, are therefore common among the symptoms attending the renal affection. And to distinguish this œdema from that produced by other morbid states we have only to examine the urine carefully,—a matter, indeed, which ought not to be neglected in any case of œdema of the lungs. Yet we must not forget that small amounts of albumin may be found in urine from any stress of breathing, and from diseases that, like those of the heart, congest the lungs and kidney and are themselves among the causes of pulmonary œdema.

*Pleurisy and Pericarditis.*—The tendency to inflammations of the serous membranes is a remarkable peculiarity of Bright's disease. We may discriminate pleurisy or pericarditis complicating the malady from either of these affections of other origin, by noting the far greater amount of dropsy than is found in these disorders, and by detecting persisting albumin and tube-casts in the urine.

*Dropsy.*—By an examination of the urine, too, may be distinguished the dropsy of the complaint under consideration from that produced by other causes. And we also see very often the evidences of the true nature of the dropsy in its beginning with swelling of the face and then becoming universal, and in the striking and characteristic physiognomy which it has a share in developing.

*Coma; Convulsions.*—A dangerous complication of Bright's disease manifests itself by drowsiness and convulsions. Now, it is very important to distinguish the cases produced by uræmic poisoning from epileptiform convulsions and kindred states in which there is no appreciable change of structure in the kidneys. Let us see how they differ.

*Uræmia*, or uræmic intoxication, is commonly preceded by a

---

\* Amer. Journ. Med. Sci., January, 1893.

diminution in the urinary secretion. There is headache, with indistinct vision, great drowsiness, and vertiginous sensations; the pupils are sluggish and usually dilated; the hearing is impaired; the countenance is dusky; the skin is cool, with short exacerbations of heat; and the patient suffers from constipation, nausea, and obstinate vomiting. Paralysis of sensation may be observed in the extremities, and various kinds of cutaneous eruptions. The dullness of mind is apt to deepen into stupor or coma, or convulsions set in as precursors of the coma, which terminates in death unless the urinary secretion be freely re-established. The coma may at one time be so profound that it is impossible to arouse the patient, whilst at another time he rouses himself and acts with intelligence. The convulsions generally succeed one another rapidly.

In some cases the marked phenomena set in with a chill; in other cases there is no such obvious beginning. And as regards the decided lessening, or even suppression, of the urinary secretion, though this is the rule, it is not constant. I have known many an error in diagnosis to be committed, and the symptoms of uræmia many a time to receive an erroneous interpretation, from supposing that uræmia could not exist, as the quantity of urine passed was about normal. We must test for urea and the other urinary ingredients, which may be profoundly changed in amount, notwithstanding the seemingly healthy aspect of the secretion, and notwithstanding, too, that it may be found free from albumin. In addition to the great decrease in the urea, the uric acid is reduced; the specific gravity is generally low; casts are mostly found in the urine.

Cases of uræmic coma differ from ordinary comatose conditions, as witnessed in apoplexy, in fevers of a low type, or following narcotic poisoning, by the dissimilar symptoms ushering them in. The coma is much more suddenly developed than that in fevers; far less suddenly than that of apoplexy or narcotic poisoning.\* Then, the stertorous respiration, to adopt the observation of Addison,† is peculiar: the loud sounds of the expired air are of much

---

\* There may, however, be exceptions to this rule, as was the case in a curious instance reported by Moore in the *London Medical Gazette*, 1845, in which a person became comatose after taking laudanum, yet his death was found to have been caused by contracted kidneys.

† Guy's Hospital Reports, 1859.

higher key, not like the low, guttural tones of apoplexy. Furthermore, we may have in the general dropsy a clue to the nature of the case; but of course the most certain light is thrown on it by the analysis of the urine.

The same remarks apply to the delirium or to the epileptiform convulsions of uræmia. Here the difficulty in diagnosis is increased by the first seizure often happening unexpectedly,—so much, in truth, increased, that, unless we are aware of the history of our patient and have previously examined the urine, the true explanation of the symptoms is not to be reached. *Uræmic delirium* is rare, but I have met with it under circumstances in which nothing preceded it to indicate its nature.\* Cases of *acute uræmic mania* may also originate thus suddenly. Cases of *uræmic convulsions* may occur in pregnant women; in them, however, the tendency to disorder of the kidney is so great that we are rarely in error in concluding convulsions to be of uræmic origin. We must, however, here, as in all convulsions, be certain that we do not mistake effect for cause. A slight amount of albumin may follow violent convulsions in epileptic seizures. The temperature in uræmic convulsions is variable. It is generally stated to be low; but this is denied by Bartels, who notes it as considerably elevated,† and by Hughes.‡

Among the other marked nervous manifestations of uræmia may be persistent headache, anæsthesia, temporary blindness, and palsies of uræmic origin.§

The cause of uræmia is still undetermined: a contamination of the blood by retained poisonous urinary ingredients or poisonous substances that have formed from them always happens, though these toxins may be of different kinds. Uræmia has been attributed to the salts of potassium accumulating in the blood.|| But this is not proved.

**Chronic Bright's Disease.**—An acute attack of Bright's disease may gradually pass into a confirmed malady, or the com-

\* Case at the Pennsylvania Hospital, April, 1865.

† Ziemssen's *Cyclopædia*.

‡ Philadelphia Hospital Reports, 1893.

§ Laguel, *Journ. de Méd. et de Chir. Prat.*, October, 1888.

|| Feltz et Ritter, *De l'Urémie expérimentelle*, Paris, 1881.



plaint may come on insidiously and develop itself slowly. In either case we have a dangerous chronic affection established.

The transition from the acute to the chronic disease is indicated by the disappearance of blood from the urine, by its lessened specific gravity and the smaller amount of albumin it contains, by the temperature becoming normal, and not uncommonly by a temporary diminution of the anasarca and an increase in the quantity of urine voided.

When the disease runs a more or less chronic course from the beginning, its initiatory steps are obscure. We generally find such cases in persons who are poorly fed and half clad, who live in damp, ill-ventilated houses, who are intemperate, or who have been subject to great grief or worry, or who are saturated with malaria, or whose constitutions are ruined by syphilis or by scrofula. The first symptoms they notice may be frequent desire to urinate; swelling of the extremities or of the face; increasing pallor and general debility; and headache, especially occipital headache. An examination of the urine reveals at once the cause of their protracted indisposition. Yet the renal disease may lead suddenly to a fatal termination without the patient having previously experienced any manifest signs of ill health. And even after the malady has been fully recognized, it is difficult to predict its course. We meet in many with the same phenomena as those encountered in the acute variety, except the fever. But in others the signs are dissimilar,—the dropsy, for instance, is slight or is wholly wanting. The only constant and characteristic manifestations are the profound and increasing anæmia, and the presence of albumin and tube-casts in the urine.

Generally, the urine is of unchanged specific gravity, though this is lowered as the urinary solids and the urea are lessened. Commonly, also, the urine is not so abundant as in health, and its reaction is less acid. The albumin is very variable in amount; its quantity may, indeed, fluctuate much in the same patient, and even change from day to day. It is persistent; yet it may, in some cases, disappear for a short time.

The tube-casts, too, are not uniform,—not nearly so much so as in the acute variety of the affection. We meet with casts almost or quite homogeneous, and small or large; with casts besprinkled with shrivelled degenerating epithelium; with casts covered with

granules or with oil-drops. In the progress of a particular case, nearly all these forms may be encountered, although, as we shall hereafter see, the preponderance of any one of them affords an indication as to the exact state of the kidneys. There is only one kind we do not find in the chronic disorder: the one covered with well-developed epithelial cells or blood-corpuscles. The apparent absence of casts from albuminous urine is not absolute proof of the non-existence of renal degeneration. In some cases their absence is only temporary, while in others they are small and few in number and easily escape detection. This is especially the case in the contracted kidney.

From these remarks, it is obvious that a great diversity of phenomena is witnessed in chronic Bright's disease: so great, in truth, is this diversity that the opinion is generally adopted that there are several distinct affections embraced under the one term, and attempts have been made to define accurately the train of symptoms significant of each. But, notwithstanding that a means of separation is also afforded by the very varied aspect of the organ, it is best to consider the differential diagnosis of chronic Bright's disease continuously, pointing out, after having done so, the clinical features that are indicative of the various forms of the malady.

Leaving out of consideration those affections for which both the acute and the chronic disease may be mistaken, and which have been already discussed, chronic Bright's disease may be confounded with—

ANÆMIA ;

NEURALGIA ;

CHRONIC RHEUMATISM ;

CHRONIC BRONCHITIS ;

ASTHMA ;

DISEASE OF THE HEART ; CARDIAC DROPSY ;

GASTRO-INTESTINAL DISORDERS ;

CANCER ; TUBERCULOSIS ; CYSTS OF KIDNEY ;

CHRONIC CONSECUTIVE NEPHRITIS ;

RENAL INADEQUACY.

*Anæmia*.—There are few diseases which alter the blood so completely as does chronic Bright's disease. The blood-corpuscles go on steadily diminishing, while the fibrin holds its own, and the quantity of albumin fluctuates considerably, being ordinarily much

reduced. Besides these changes, the blood often retains its effete ingredients, since the kidneys are incapable of performing their function. The alteration and gradual impoverishment of the blood make themselves manifest by the increasing debility, and by the pallor and waxy look of the countenance.

We may discriminate this well-marked anæmic condition from that unconnected with renal disease by the existence of albumin and tube-casts in the urine, and often also by the prominence of the dropsical symptoms. But it is essential to know that some of the phenomena—certainly albuminous urine and dropsy—may attend the anæmia following profuse or frequently-repeated hemorrhages, without the structure of the kidneys having been impaired. It is difficult to distinguish these cases from true Bright's disease, except by taking into account the diminution of the albumin as the hemorrhagic tendency is lost, and the absence of the tube-casts. The dropsy, unless it be considerable, can hardly be looked upon as a valuable differential index, for a slight or moderate amount of dropsy, or even none, may be encountered in either morbid state.\*

The ophthalmoscopic appearances presented by the retina, and described in a previous part of this work, afford help in distinguishing between the anæmia of Bright's disease and that produced by any other cause. Albuminuric retinitis is not limited to any form of Bright's disease. It generally happens in both eyes, and, though in the chronic variety of the malady it may greatly improve, it does not disappear. The sight itself deteriorates; and we have attacks of blindness, uræmic amaurosis, which come on suddenly and pass off suddenly.

*Neuralgia.*—As this is not infrequent in the chronic form of Bright's disease, we must always, in obstinate cases of neuralgia, examine the urine. Neuralgia of renal origin may affect the fifth nerve, or other nerves; sometimes it takes more the form of hemi-crania, and it is often associated with disordered vision, or with impairment of other special senses; or it may coexist with persistent headache or with strange and anomalous nervous symp-

---

\* The occurrence of marked albuminuria after hemorrhage, to which attention was here called, has been since studied by Fischl, Arch. f. Klin. Med., Bd. xxix., and by Quincke, *ib.*, Bd. xxx., No. 4.

toms. Headache from Bright's disease may also be present without neuralgia; it may be of the nature of megrim, and occur in paroxysms attended with nausea and vomiting.

*Chronic Rheumatism.*—Frequently patients affected with chronic Bright's disease complain of muscular pains. The pain is dull, not increased on pressure; sometimes shooting, more like that ordinarily termed neuralgic, to which we have just called attention. The pain is oftenest met with in those instances in which the dropsy is slight or wholly wanting, and an examination of the urine is then the only means of determining its real significance.

*Chronic Bronchitis.*—This is one of the most common complications of Bright's disease,—so common, indeed, that Rayer observed it in seven-eighths of his patients, and Wilks\* states it, from an extensive analysis of cases, to have been more universal than any other single symptom, albuminous urine alone excepted. It is hardly necessary to add that the last-mentioned sign is the one that distinguishes this secondary pulmonary affection from all other forms of bronchial disease.

*Asthma.*—Whether or not there be coexisting bronchitis, attacks of shortness of breath, like paroxysms of asthma, occur as the result of Bright's disease. This renal asthma is most common in the chronic contracted kidney. It has no features by which it can be recognized from ordinary asthma, except that the wheezing and the rales are not so marked, and that it does not subside by copious expectoration. It indeed more resembles cardiac asthma, and is most frequent at night.

*Disease of the Heart; Cardiac Dropsy.*—In very many cases of chronic Bright's disease there is disease of the heart, particularly hypertrophy. This manifests itself by the ordinary physical signs of hypertrophy. The vessels become more tense and rigid, especially in contracted kidney, and there is marked accentuation of the second sound of the heart. With these physical signs, dropsy may show itself, and frequently does so in chronic parenchymatous nephritis. With kidney and heart both affected, it is often difficult to say which is the primary lesion, and to which the dropsy is to be especially referred; either may occasion albumin in the urine.

---

\* Guy's Hospital Reports, 2d Series, vol. viii.



Let us suppose that in cases of so-called cardiac dropsy we find albumin in the urine : is this a proof of coexisting Bright's disease? Not unless the amount of the abnormal ingredient be considerable and more than occasional tube-casts accompany the albuminuria. Mere congestion of the kidneys, resulting as it does from an obstruction to the flow of the venous blood along the vena cava, may occasion albuminuria ; but the presence of albumin is temporary, and its quantity small, and the specific gravity of the urine is generally high. A large amount of albumin, persistent and conjoined with characteristic tube-casts, shows that changes are present in the renal textures. Disease of the heart and disease of the kidney are often combined. It is the disease of the kidney which produces or coexists with the disease of the heart. The cardiac affection does not give rise to the renal affection nearly as often as supposed.\*

*Gastro-Intestinal Disorders.*—These are among the most common consequences of the renal malady. They manifest themselves in various ways: by flatulency and indigestion; by diarrhœa; by nausea and vomiting. The latter symptoms are apt to occur when uræmic intoxication is developed. They may be, however, also met with at any period of the disease without the concurrence of other urgent symptoms, and become so prominent as to throw into the background most of the other signs of the renal affection. I have seen cases of Bright's disease which first manifested themselves by apparently causeless nausea and vomiting; the tongue was clean.

*Cancer; Tubercle; Cysts of Kidney.*—These morbid products affect the kidneys but rarely,—at all events, rarely in a form so marked as to give rise to conspicuous clinical phenomena. In all of them there may be albumin present in the urine, but it is generally in small amounts, and mixed with some ingredient having a more specific meaning. Thus, in *cancer* of the kidney we may find blood with the albumin; indeed, hæmaturia is a very important symptom, and in some instances we discern with the microscope cells like those observed in any cancerous growth; often the

---

\* See proof in Middleton-Goldsmith Lecture, 1888, or Relation of the Diseases of the Kidney, especially the Bright's Diseases, to Diseases of the Heart, by J. M. Da Costa.

hemorrhages are profuse and frequently recurring, are preceded by severe pain, and we detect a palpable tumor in the flank, passing upward into the hypochondriac region and downward to the iliac region, or even forward, not affected by the act of breathing, and sometimes causing bulging posteriorly. In cases of melanotic cancer, whether it have its seat in the urinary apparatus or elsewhere, Eiselt and Bolze\* have noticed that the urine on standing assumes the color of porter, and that on the addition of concentrated nitric acid it instantly presents the same dark color. In children, cancer of the kidney is not a rare disease,† and when we can exclude as the cause of the renal tumor cystic degeneration and hydronephrosis—in them congenital affections—we can diagnosticate the case with some confidence. In adults the diagnosis is always doubtful, at least when the disease is primary. A rapid and irregular growth of the one-sided renal tumor, severe pain, bloody urine, and cachexia are the most certain signs. Sudden and rapidly-growing varicocele is stated to be a symptom of malignant tumor.‡ *Syphilomata* of the kidney may be suspected from the history, but cannot be recognized with certainty; they rarely cause pain or produce a tumor large enough to be detected, but mainly give rise to the ordinary manifestations of chronic Bright's disease,§ most often of the amyloid form. In *sarcoma* of the kidney the swelling in the abdomen attains, in children especially, very great size; hæmaturia is comparatively rare, and the peripheral lymphatic glands do not become implicated.||

In *tubercle*, little yellow cheesy masses of degenerated tubercular matter collect as a sediment, as in the cases referred to by Frerichs in his work on Bright's disease. The constant presence of this sign is, however, doubtful. The tubercular matter is generally derived from the pelvis of the kidneys. The deposit it forms in the urine is insoluble in acetic acid. Pus and other signs of chronic pyelitis are present, and there is no other assignable cause for the existence of the suppurative disease than tubercle.

\* Prager Vierteljahrschrift, vols. lix. and lxvi.

† Braidwood, Liverpool Reports, 1870.

‡ Guillet, Tumeurs malignes des Reins, Thèse de Paris, 1888.

§ Wagner, Archiv f. Klin. Med., Bd. xxviii, 1881; Mauriac, Arch. Gén. de Méd., Oct. 1886; Jaccoud, Gaz. des Hôp., 1888.

|| Neumann, Archiv f. Klin. Med., Bd. xxx., 1882.

The disease may be primary, or the infection take place from the bladder, the prostate, or the ureters. We may be assisted in the diagnosis by finding tubercles in other organs. Rayer tells us that scrofulous disease of the vertebræ has repeatedly been observed to be associated with tubercular kidneys. In tubercle of the kidney, extreme pain, occurring in paroxysms like those of nephritic colic, is a very important sign. This pain, as I have had occasion to observe, is associated with frequent micturition, and is temporarily relieved by the flow of water. The urine is, however, scanty. A moderate amount of hæmaturia may happen; the patient passes at times little fibrinous shreds, and emaciates steadily. The bacillus of tubercle may serve as a means of diagnosis in the urine; it has been detected in a number of instances.\* In some cases the kidney most diseased enlarges sufficiently to form a tumor discernible through the abdominal walls.

In *cysts* of the kidney—those at least enclosing echinococci—small vesicles containing the characteristic structures of the parasites may perhaps be detected. Ordinary cysts are not to be recognized with any certainty during life; nor can they be distinguished from Bright's disease, since they are very frequently developed in the chronic varieties of this disorder. When the cysts attain decided dimensions, they give rise at times to the discharge of highly-bloody urine, and to albuminuria, and to large tumors, which may be detected through the front walls of the abdomen. They may affect one or both kidneys, producing slow cachexia and enormous abdominal swelling. Cysts of the kidney and liver often coexist.†

*Chronic Consecutive Nephritis.*—In consequence of affections of the bladder, of stone in the bladder, of strictures of the urethra, of disease of the ureters and of the prostate, indeed of various surgical affections of the urinary organs, we may have a kidney disease established which is rather a form of slow inflammatory change than Bright's disease. It may affect only one or both kidneys, and the diseased organs are tough and hard, large or small, and show great increase of fibrous tissue. The source of

---

\* R. Shingleton Smith, *Lancet*, i. 942, 1883; Irsai, *Wien. Med. Presse*, p. 1173, 1884; and others.

† Sabourin, *Arch. de Phys.*, ix., 1882.

irritation which has led to the secondary inflammation is at times in the kidney itself, in the shape of a large calculus in the pelvis.

In another form of this consecutive nephritis suppuration takes place, affecting first especially the pelvis of the kidney, a suppurative pyelo-nephritis,—the condition often called *surgical kidney*. It is difficult to distinguish these consecutive forms of nephritis, especially where pus is found in the urine, either from the condition last mentioned or from coexisting bladder disease, except by the history. Very often there is pain along the course of the ureter; and the urine, when passed free from pus, contains neither albumin nor casts, or only a small amount of albumin and a few hyaline casts. The urine is apt to be copious and of low specific gravity. When it contains pus from the kidneys, and the bladder is comparatively unaffected, the purulent urine is generally acid. The heart rarely becomes disturbed, though hypertrophy has been occasionally noticed in the non-suppurative form.\*

*Renal Inadequacy.*—There are patients who pass the ordinary amount, or less than the ordinary amount, of urine daily, of low specific gravity, from 1002 to 1008, not containing more than two per cent. of urea, though the uric acid may be normal, and who in consequence of this insufficient action of the kidneys are always ailing and weak, take cold easily, and suffer from headache and nervousness. Even if they drink water freely, they do not pass more urine; this does not contain albumin or casts, differing in this respect from Bright's disease. But dropsy, as Sir Andrew Clark, who first described the complaint,† states, with puffy face and dry, glossy skin, may happen, and a state similar to myx-œdema be gradually developed.

Having now treated of chronic Bright's disease as *one* affection, I shall briefly refer to the distinctions between its forms. In so doing, I shall follow the classification based on the diversified anatomical aspect of the kidneys.

First there is the chronic *enlargement* of the organ, of which several kinds exist:

---

\* Fagge's cases, in *Practice of Medicine*, 1886, vol. ii. p. 483.

† *British Medical Journal*, vol. i., 1883.



1. *The fatty kidney.* The kidney is very large and fatty. The deposit may occasion yellow scattered granulations, or the enlarged organ is pale, and mottled by red vascular patches. The convoluted tubes are filled with oil, accumulated in their epithelial cells. The fatty disease is recognized by the numerous oily casts, fatty cells, and free oil-cells which appear in the highly-albuminous urine. It is a fatal complaint, generally very chronic in its course, and attended with persistent dropsy. This morbid condition must not be confounded with a simply fatty kidney, such as is sometimes found in phthisis, or oftener in drunkards, and which is not associated with albuminous urine. A certain amount of

FIG. 65.



Fatty casts and epithelial cells filled with fat, as seen in the discharge coming from a highly-fatty kidney.

fatty casts and fatty cells may appear in the urine and not be persistent or indicate the real, dangerous fatty kidney. It is thought by several, by Dickinson especially, that the fatty kidney may follow a high degree of inflammation in the acute form of Bright's disease, particularly in that form brought on by exposure to cold. The acute form attending scarlet fever is more apt to pass into the large white kidney.

2. *The enlarged chronically inflamed kidney.* I allude to the chief form of the large white kidney so often mentioned by English physicians, the chronic parenchymatous inflammation of the kidneys of the German writers. The organ is white, enlarged, dense ;

its tubes are filled with exudation-matter, their walls thickened. The cortical portion of the kidney is pale and increased in breadth, evidently full of an inflammatory deposit; the medullary cones retain their vascularity. This variety of the malady may or may not follow acute Bright's disease. It may last for a few years, but generally terminates unfavorably before that time. The urine is diminished in urea and pigment and in chlorides; it contains large amounts of albumin and granular and epithelial casts, with some hyaline casts and a few slightly oily casts. The dropsy occasioned is extensive and persistent, and there is usually little difficulty in tracing it to an acute attack. Sometimes the dropsy lessens materially, then actively recurs, and there seems to be rather a series of subacute attacks than a continuous chronic malady. The large kidney is not supposed ever to contract; but this is not a settled point. Grainger Stewart holds that it does, as does the waxy kidney, yet believes that both in a stage of atrophy are distinct from the so-called cirrhotic or contracting form of Bright's disease.\* The large white kidney may pass into the fatty kidney. Dilatation of the heart is common in chronic parenchymatous nephritis, more common even than pure hypertrophy, which is more usual in contracted kidney.

3. The *waxy* or *amyloid kidney*, an affection in which the enlarged organ is smooth, of firm look, and of pale-yellow color, and is the result of a general disease involving the kidneys in common with other organs. It originates in the exudation from the minute arteries of a waxy material which infiltrates the tissues. This condition generally follows upon protracted suppuration from whatever cause, either wound or disease. The urine is increased in quantity in the earlier stages, and of low specific gravity; it contains much albumin, but not many casts. Those which are seen are pale, and for the most part hyaline, or highly-refracting, structureless moulds of the tubules of large diameter; they may or may not give the characteristic amyloid reaction, the red color when treated with a watery solution of iodine and of potassium iodide.† Methyl-green colors amyloid substances an intense green. It is used for staining in the form of a one per cent.

---

\* On Bright's Disease of the Kidneys, 1871.

† Curschmann, Virchow's Archiv, vol. lxxix., part 3.

aqueous solution. Methyl-green colors hyaline casts *in situ* ultramarine blue, so that these also can be readily distinguished in sections of the kidney from the green-colored tissues around, in which they may lie. Blood is rarely present in the urine of the amyloid kidney, and the urea is but slightly diminished in quantity. Diarrhœa frequently coexists, and the liver and spleen are apt to be enlarged; but the heart is not affected. The dropsy is absent or trifling in amount, yet its persistence while the urine is

FIG. 66.



Hyaline or waxy casts, magnified about 460 diameters. On some of them are scattered a few shrivelled epithelial cells and oil-drops; the large cells to the left are epithelial cells from the bladder.

The kind of casts here depicted may be found in any form of Bright's disease, acute as well as chronic. In the waxy kidney, however, they vastly preponderate, and are of large size,—many much larger than those in this figure.

increased in quantity is peculiar to this form of renal disease, and it may exist markedly as a late symptom; the patient is sallow-looking and emaciated; his disease may last for years.

In laying stress on the hyaline and waxy casts we must be careful not to confound them with those still larger mucous moulds of the uriniferous tubules, or *mucous casts*. They are also smooth, but of enormous length, subdividing into smaller ones, and of cylindrical shape. They are met with in acute parenchymatous nephritis, but are particularly apt to occur in consequence of transmitted irritation from the bladder, and are then perhaps associated with small amounts of albumin and of pus. Yet unless the latter ingredient be present there is no albumin, or the merest trace.

Further, flask-shaped hyaline bodies and cylinders may be moulds of the vesicles and smaller ducts of diseased prostates.\*

4. Then we have the small *contracted kidney*, the granular kidney or interstitial nephritis, which is viewed as the last stage of Bright's disease by those who believe in the various appearances

FIG. 67.



Granular casts, or casts covered with disintegrating epithelium and granules. Casts of this character are chiefly found in the chronic inflammatory forms of Bright's disease. They are not seen in the acute complaint, except when it is assuming a chronic form.

being only successive stages of the same morbid process, but which is more generally held to be an independent disease. This form of disease is frequently found in gouty persons, or after prolonged mental anxiety and distress,† or as the result of lead poisoning. The urine contains but an inconsiderable amount of albumin; the tube-casts are granular, or simple fibrinous moulds, generally small, sometimes large; here and there a little oil is observed. Dropsy is absent in a certain proportion of cases, and when present is generally slight. It often disappears for a while and returns. The urine is increased in quantity, although toward the termination it may become scanty or even suppressed. Dyspepsia, puffy eyelids, chronic bronchitis, increased arterial tension, hyper-

\* Sir Andrew Clark, Transactions of the Clinical Society of London, vol. xix., 1886.

† Clifford Allbutt, British and Foreign Medico-Chirurgical Review, Oct. 1877.



trophied ventricles, albuminuric retinitis, headache, and disorder of the nervous system are common symptoms. The malady runs a very chronic course. It is chiefly characterized anatomically by an affection of the fibrous tissues surrounding the Malpighian corpuscles and lying between the tubes, a slow increase, followed by a slow contraction, of the intertubular fibrous tissue and atrophy of the tubules, connective-tissue changes in the renal plexus,\* and fibroid changes in the small vessels of the body. The sphygmograph shows marked pulse-tension, and this, with altered specific gravity, has been noticed before albumin is present in the urine. In the uric acid or gouty nephritis, uric acid deposits may be found in the straight tubes of the medullary substances. A chronic interstitial nephritis may also be associated with deposits of lime, which take place very generally in the uriniferous tubules in the cortex. These lime deposits may be, as Virchow points out, calcareous matter washed into the kidney from diseased bone.

In contracted kidney, especially in the earlier stages, albumin may be absent from the urine, and we may have to recognize the malady rather by the hypertrophied heart and thickening of the vessels, the high arterial pressure, the accentuation of the second sound of the heart, the headache, vertigo, nausea, breathlessness, retinal changes, and the anæmia. The urine may be of low specific gravity and copious, but there are many exceptions to this; it is generally deficient in urea. Occasionally a few hyaline or granular casts are found; and the albumin may not be entirely absent, but appears every now and then in traces. There may be even chronic general œdema present without albuminous urine,† and various nervous and mental symptoms.

The different kinds of albumin have been above mentioned. Of these serum-albumin and serum globulin are by far the most important. Indeed, as yet there has been nothing of clinical value gained by the study of the other varieties.

In the following table the clinical differences between the various forms of Bright's disease are set forth :

---

\* Da Costa and Longstreth, *Amer. Journ. Med. Sci.*, July, 1880.

† As in Case 31 of Mahomed's paper on Chronic Bright's Disease without Albuminuria, *Guy's Hospital Reports*, 3d Series, vol. xxv.

TABLE EXHIBITING THE CLINICAL DIFFERENCES BETWEEN THE PRINCIPAL FORMS OF BRIGHT'S DISEASE.

*Acute Cases in which Dropsy occurs quickly and is extensive.*

Acute Bright's disease; acute desquamative or tubal nephritis; acute parenchymatous nephritis; acute diffuse nephritis; acute renal dropsy .....	<p>Caused mostly by exposure, or scarlet fever.</p> <p>Dropsy extensive, generally begins in the eyelids or in the feet; usually febrile symptoms; uræmia may be met with. Disease most common in childhood and among young adults.</p> <p>Recoveries frequent; but disease may terminate in acute parenchymatous nephritis.</p>	<p>Urine usually scanty, deep-colored, of high specific gravity, containing much albumin, often blood; also blood-casts; casts, many of large size, covered with epithelium, and a few hyaline casts; and free epithelial cells, cloudy and granular; urea diminished.</p>	<p>Kidneys enlarged, congested or mottled, shedding epithelium; cortical substance increased; cones usually redder than cortical substance. Dilated convoluted tubes, distended with swollen, cloudy epithelium; at ends of tubules also blood or plugs of fibrin. Tubules darker and denser than normal.</p>
--	--	--	---

*Chronic Cases in which Dropsy is variable in amount and may be absent.*

Chronic inflammatory form; chronic tubal nephritis; chronic parenchymatous nephritis; chronic diffuse nephritis; large white kidney .....	<p>History often of antecedent acute inflammatory attack; dropsy a prominent symptom. Uræmic phenomena not uncommon; among them at times uræmic coma, with its usual symptoms. Inflammations of serous membranes also not uncommon.</p> <p>Hypertrophy of heart, especially of the left ventricle, or dilatation.</p> <p>Recovery possible, but doubtful; may pass into fatty kidney.</p>	<p>Urine in normal or in increased quantity; albumin generally in considerable amount; granular casts; at times compound granule cells and partially fatty epithelium; casts with fragments of epithelium or a little fat; no blood-casts; leucocytes; specific gravity below normal; urea diminished.</p>	<p>Kidneys enlarged, capsules easily stripped off, cortical substance increased in volume, cones may be of natural color; tubes often irregularly distended, and filled with granular epithelium here and there slightly fatty, and with detritus. Thickening of intertubular matrix.</p>
---	---	--	---

TABLE EXHIBITING THE CLINICAL DIFFERENCES BETWEEN THE PRINCIPAL FORMS OF BRIGHT'S DISEASE.—*Continued.*

*Chronic Cases in which Dropsy is variable in amount and may be absent.—  
Continued.*

Fatty Bright's kidney.....	<p>Persistent and obstinate dropsy, coming on gradually; face pale and puffed; hypertrophy of heart affecting often both sides.</p> <p>Always fatal.</p>	<p>Urine contains much albumin, fatty casts, fatty epithelial cells, free oil.</p> <p>Spec. grav. variable, usually from 1015 to 1030.</p> <p>Quantity variable, generally moderate or diminished; urea diminished.</p>	<p>Kidneys enlarged, and very fatty; sometimes have a mottled look.</p> <p>The tubes, especially the convoluted ones, full of highly-fatty epithelium, and free oil.</p>
Waxy kidney; lardaceous or amyloid degeneration of kidney.....	<p>Follows usually exhausting and wasting diseases, syphilis, caries, and long-continued suppuration.</p> <p>Rare in very early and in advanced age.</p> <p>Dropsy trifling, except late in disease; great emaciation; striking sallowness of face; liver and spleen enlarged; diarrhoea; much thirst; heart not affected; nervous symptoms infrequent.</p> <p>Unfavorable prognosis.</p>	<p>Urine increased, contains much albumin, but few casts, which are pale and transparent or highly refracting. The casts may or may not give the mahogany-red reaction with a watery solution of iodine.</p> <p>Spec. grav. low, yet usually above 1010; urea normal or slightly diminished.</p>	<p>Kidneys enlarged, smooth, and waxy-looking; capsule easily detached; cortex pale, anæmic; reddish-brown discoloration on testing with watery solution of iodine; cones often dark and congested. Morbid process at first chiefly along renal vessels.</p>

TABLE EXHIBITING THE CLINICAL DIFFERENCES BETWEEN THE PRINCIPAL FORMS OF BRIGHT'S DISEASE.—Continued.

*Chronic Cases in which Dropsy is variable in amount and may be absent —Continued.*

Chronic contraction of the kidney; cirrhosis of the kidney; interstitial nephritis; granular kidney; gouty kidney.	Dropsy slight, frequently absent; face sallow, yet not so much so as in the waxy disease; often headache and retention of urea, tendency to coma, and to convulsions; impoverished blood; hypertrophy of heart; epistaxis; liver may be cirrhotic; retinitis. Most common between forty and sixty years of age. May exist for years unsuspected; is a very chronic disease, and incurable; may lead to death by apoplexy.	Urine more copious than in health, yet extremely small amount of albumin, this at times temporarily absent; hyaline and large finely granular casts; altered epithelium; a little oil. Spec. grav. low; rarely above 1010, much oftener below; urea decreases gradually; marked decrease late in disease.	Kidneys waste slowly, become dense and contracted; capsule very adherent; surface often granular; thickness of the cortical substance diminished; cysts common. There is hypertrophy of connective tissue; compression and atrophy of gland-elements and of tubules. Cardio-vascular changes. Tissue-changes in renal ganglia.
--	---	---	--

### Diseases associated with Purulent Urine.

In every case in which pus in any quantity is detected in the urine, it becomes of great importance to ascertain primarily that it is not derived from the urethra, from the vagina, or from an abscess that has opened into the urinary passages. The first point we may decide by examining into the history of the case, and, if necessary, by an exploration of the parts, as well as by an examination of the urine procured in the manner recommended in the first part of this chapter; the second, by the same means, and by determining that a discharge takes place equally when no urine is voided; the third is more difficult to make out, but there is generally something in the symptoms and in the history of the case furnishing a clue to its interpretation,—such, for instance, as the sudden appearance of a large quantity of pus in the urine.



Having excluded each of these morbid states as the source of the purulent urine, we next turn to see which of the maladies that are its most common cause is before us. They are :

**Acute Cystitis.**—Acute inflammation most frequently affects the mucous membrane at or near the neck of the bladder. The inflammation may spread from the mucous membrane to the muscular coat ; but it rarely reaches the peritoneal covering. In some cases it is propagated along the uterus, and even to the kidneys. The morbid action is not often of idiopathic origin, although sometimes it follows exposure to cold and damp ; much more usually is it due to the extension of an attack of gonorrhoea, to disease of the prostate, to traumatic causes, to protracted retention of urine, or to the irritation produced by medicines or stimulating drinks. Sometimes it is owing to the poison of rheumatism or of gout. It is frequently due to a bladder infection by pyogenic germs.

Acute cystitis is much more commonly encountered in men than in women, and in adults than in children. Its main symptoms are a feeling of weight and pain in the hypogastric region, augmented by movement and by pressure. The pain does not, however, remain confined to the region about the bladder, but is felt also in the iliac and sacro-lumbar regions. It is attended with considerable febrile disturbance and extreme irritability of the affected viscus. The urine is voided drop by drop, and its passage is usually accompanied by straining and a scalding sensation at the neck of the bladder ; it is high-colored, cloudy from vesical mucus, and contains blood and pus and sometimes shreds of lymph. At first the urine is acid. The acute disease generally terminates within a week, leaving often an irritable bladder or a chronic inflammation.

The symptoms of acute cystitis are similar to those of *acute nephritis*, and the exciting causes are much the same. But acute inflammation of the bladder differs from acute inflammation of the kidney by the greater severity of the pain, its much lower position, and the distress in voiding the urine. *Neuralgia*, or spasm, of the bladder may be distinguished from acute inflammation by the absence of fever, and by the sharp, lancinating, but paroxysmal pain, each onset of which lasts hardly longer than from two to six hours, and is attended with difficulty in making water, which disappears as the pain subsides.

*Metritis* exhibits several of the traits of cystitis: we find the same hypogastric pain shooting to the thighs or to the anus and loins, the same feeling of weight in the peritoneum, and the same signs of irritation of the bladder and of fever. As it, however, generally occurs in the puerperal state, we have the history, and the character of the discharges from the vagina, to guide us, as well as the knowledge to be gained by a local examination.

**Chronic Cystitis.**—This affection, often called chronic vesical catarrh, is common in advanced age. It generally comes on in an insidious manner, and is excited by some obstacle to the evacuation of urine, such as a stricture, or by the presence of a stone in the bladder, or by an enlargement of the prostate gland. A paralysis of the viscus leading to retention of its contents, or a serious structural disease of its coats, whether malignant or non-malignant, may, however, also establish the morbid process.

The symptoms are partly those of constitutional debility, partly those of local disease. The most usual of the latter, indeed in every way the most characteristic, are the dull pain, a frequent desire to make water, and the passage of a large quantity of mucus or pus with each act of micturition. The urine, which is alkaline, on standing deposits a glairy, viscid sediment, in which, under the microscope, vesical epithelial triple phosphates, large pus-corpuscles, extremely regular both in contents and in shape, and pathogenic germs, especially the *bacillus coli communis* and the *staphylococcus pyogenes*, may be detected. The urine usually contains more albumin than is found in acute cystitis.

The diagnosis of the disease in males is easy. The only affection with which it is liable to be confounded is abscess of the kidney. In females, uterine disorders may so closely simulate it that it may require a local examination to tell the difference.

But, having decided the case to be one of chronic cystitis, it is always more difficult to discover its exciting cause. We have to depend, to a great extent, upon the history of the malady; its association with a stone can be determined only by the use of the sound.

**Abscess of the Kidney.**—This dangerous condition is the result of suppurative inflammation of the kidney, or of abscesses forming in connection with pyæmia, or with embolism. The suppurative inflammation is sometimes traceable to an acute attack

of nephritis brought on by exposure or by external violence, to retention of urine, or to the impaction of a renal calculus; but at other times it originates without any assignable cause, and in an insidious way. The association of suppurative nephritis with erysipelas has engaged much attention, and the renal affection is even thought to be erysipelatous in its origin.\* Abscess of the kidney also arises from the acute interstitial nephritis followed by suppuration that occasions the so-called surgical kidney.

Abscess of the kidney is a rare disease. It has much the same symptoms as pyelitis. There is a fulness on one side of the spine, associated with tenderness on deep pressure in the lumbar region, and with more or less constant pain, the pain and tenderness being increased by lying on the affected side; there are also fever and occasional rigors, digestive disturbances, and blood and pus in the scanty urine. In some cases a marked tumor is found in the loin, extending toward the iliac fossa. If the abscess bursts into the calices, there occurs, simultaneously with a subsidence of the tumor, a sudden and copious discharge of pus with the urine, or, if it breaks into the intestine, with the fæcal evacuation.

The disease almost never affects more than one kidney: hence so-called uræmic symptoms are rarely met with, since the healthy kidney enlarges and becomes capable of performing a double amount of work. Ebstein† has, however, observed that chronic abscess in one kidney may produce amyloid disease of the other. The disorder gradually leads in most cases to a fatal issue, from the irritation, the vomiting, the diarrhœa, the wasting discharge, and the protracted hectic; sometimes paralysis of one or both legs happens, adding greatly to the distress. There is a possibility of recovery, if the patient have strength enough to withstand the purulent drain until the abscess empties itself. It may do this through the urinary passages, through the colon, through the lumbar muscles, through the diaphragm, and be evacuated by coughing, and the cavity of the abscess then cicatrizes; or the abscess may burst into the peritoneal cavity and cause rapid death.

The diseases for which the malady is most apt to be mistaken

\* Goodhart, *Guy's Hospital Reports*, 3d Series, vol. xix.

† Ziemssen's *Cyclopædia*.



—leaving out those extremely rare cases in which abscesses from diseased vertebræ break suddenly into the urinary tract—are chronic cystitis, perinephritis, and pyelitis. From *cystitis* it may be distinguished by the dissimilar local signs and the different appearances of the urine. Thus, in the affection of the bladder the quantity of pus constantly discharged is far greater,—for in abscess of the kidney there are times when little or no pus is voided; on the other hand, the urine of the vesical disorder is less albuminous. In the renal malady we can detect casts and other renal products in the sediment.

*Perinephritis* unconnected with inflammation of the kidney is a very rare disease. When primary, it may result from exposure; but it is more generally due to contusion or strain. I saw an instance of it which occurred in a young man who, returning home from a long walk, strained his back in jumping a fence. An abscess gradually formed, giving rise to a slight fulness in the left lumbar region, and severe pain, which disappeared as matter was discharged through the integuments. The function of the kidney was not affected.

But an external opening may be established when the process of inflammation and suppuration has begun in the kidney and thence spread to the loose tissues surrounding it. Under these circumstances, the appearance in the urine of pus prior to its discharge through the muscles of the back would be the only certain means by which we could judge where the suppuration had primarily taken place. The inflammation may travel upward from the pelvic viscera or from the head of the colon or the appendix; it has been also noticed after irritation of the testicles and of the spermatic cord. Secondary perinephritis has been observed in pyæmia, and after typhoid and typhus fevers and smallpox. The disease is not uncommon in childhood.\*

The prominent symptom in perinephritis is pain, which at times is so severe as to confine the patient to bed with his knees flexed, with a sense of fulness and dragging weight, with tenderness in the region of the kidney, and with lameness owing to the interference with the play of the psoas muscles. The urine is generally unaltered, or only full of urates; the bowels may be

---

\* Gibney, Amer. Journ. of Obst., reports twenty-eight cases, April, 1876.



constipated, owing to the pressure of the tumor on the intestine. A rounded, doughy, and generally indolent swelling, uninfluenced by the respiratory movements, is usually found in the lumbar region or a little lower. The abscess may cause pulmonary or pleuritic complications, but almost never gives rise to jaundice. As the disease advances, severe chills, with high fever and copious night-sweats, occur, as well as emaciation and marked debility, and the thoracic symptoms may mask the renal; fluctuation may at times be detected, and, before the abscess breaks externally, a phlegmonous appearance of the skin where the abscess points is not unusual. Great relief follows the discharge of the pus.

From *inflammation of the psoas muscle* we distinguish perinephritis by the absence of marked sensitiveness over the renal region in the former complaint, and by flexion of the thigh in it producing pain.

**Pyelitis.**—Inflammation of the mucous membrane of the pelvis of the kidney is almost never idiopathic, being commonly caused by a calculus which has been arrested at the commencement of the ureter; or by a retention of urine from an obstacle in the ureter, bladder, or urethra; or by an extension upward from the bladder of an inflammation. Bright's disease and diabetes are not unusually, and typhus and the eruptive fevers, pyæmia, scurvy, diphtheria, carbuncle, puerperal septicæmia, are occasionally, complicated with some degree of pyelitis. Pyelitis may be also catarrhal or rheumatic. Under these circumstances, and in all the infectious diseases, pyelitis is apt to show itself in an acute form.

The symptoms of the chronic malady are in part those produced by the morbid states exciting it, especially those denoting a calculus lodged in the kidney or arrested in its transit toward the bladder; partly those directly traceable to the inflammation of the pelvis and infundibula. The manifestations of the latter disorder are a constant dull pain in the loin, felt also in the course of the ureter, and the passage of pus and occasionally of small quantities of blood with the urine; in cases from retention and decomposition of urine there are recurring chills, sweats, vomiting, headache, delirium, and fever. In most cases of pyelitis the urine is acid, albuminous, very abundant, and offensive. It may be acid even if it abound in triple phosphates. If detained any

length of time in the bladder it becomes ammoniacal. Bacteria are a frequent cause of pyelitis, as well as of abscess of the kidney, by migrating from a diseased bladder. In some instances of pyelitis an eruption like rubella is noticeable. Pyelitis not unfrequently affects only one kidney.

The most difficult point connected with the recognition of pyelitis is the ascertaining that the purulent discharge does not proceed from the bladder. And there is no positive sign to guide us, except the existence in the urine of epithelium from the pelvis of the kidney, distinguishable by its oval or fusiform shape, and by the frequent occurrence, in a cell, of clearly-defined, dark-colored, round granules, and of two nuclei. But this epithelium will not be always found, and we have then to fall back upon the history of the case, upon the attacks of renal pain, upon the hæmaturia caused by a calculus, and upon the combination of signs as pointing more to one disease than to the other. In some cases there is a perceptible swelling in the loin; at times, too, owing to coexisting degeneration of the cortex of the kidney, the amount of albumin is wholly disproportionate to that contained in pus, and this becomes a valuable indication of the affection not being vesical. But if there be a coincident disease of the bladder, the differential distinction may become impossible. Under these circumstances, too, the acid state of the urine, on which in uncomplicated cases much stress may be laid, is not apt to be a feature to aid us. Pascallucci\* has brought forward a sign of pyelitis which he regards as certain. It consists in taking note of the manner in which nitrate of urea crystallizes when nitric acid is added to the urine. If the catarrh be limited to the bladder, the microscope shows the crystals arranged in the form of hexagonal rhomboidal blades; in pyelitis the blades are irregular and set at angles, and some of them are in the shape of small feathers.

Supposing the point settled, and the vesical origin of the pus disproved, the diagnosis is limited to an inflammation of the ureter, to an abscess in the substance of the kidney, and to pyelitis. Here again the history of the case comes into play. Furthermore, in the former of these affections—a very rare one, unless associated with pyelitis—the amount of pus in the urine is very

---

\* Il Morgagni, quoted in *Lancet*, June, 1873.

trifling; in the second, too, it is less than in pyelitis, except when the abscess empties itself. The pus is also, as already indicated, not constant, alternately appearing in and disappearing from the urine; there is usually more obvious swelling, although this is by no means always discernible or even present in abscess, and the abscess is attended with much greater constitutional disturbance. Still, here again we must admit that the disorders are sometimes very obscure and difficult to distinguish, and it may be impossible to discriminate between them should the morbid states coexist, or a typhoid condition and uræmic fever be induced by the retention of the urine and its decomposition.

*Catarrhal or rheumatic pyelitis* is generally a short disease which ends favorably; so does the idiopathic pyelitis of the puerperal state, which rarely lasts more than from five to eight days. The pyelitis with retention and decomposition of urine is a much more serious complaint, and, although it usually runs a rapid course, not having a duration of more than a week or two, it may become a protracted state. Pyelitis due to the *irritation of calculi* is apt to develop into a chronic condition.

In *tuberculous pyelitis* the symptoms are the same as in the ordinary form. The association with tuberculosis in other parts, and the detection of tubercle-bacilli in the urine, establish the diagnosis.

In those cases of pyelitis in which there is a very decided obstruction to the flow of urine through the ureter, caused by a calculus, a clot of blood or viscid pus, or other débris, the discharge of pus is suddenly arrested and the cavity of the pelvis dilates greatly; gradually the gland-tissue is compressed, and a large pus-containing sac is formed, giving rise to a condition known as *pyonephrosis*, and to a distinctly limited swelling in the side. Tumors of this kind are ordinarily not painful to the touch, are sometimes very indolent, and do not materially affect the general health, certainly not, as a rule, nearly so much as might be supposed. They not unfrequently subside gradually by free discharges of pus, and the patient recovers.\* Sometimes they become much reduced, and then swell up again from time to time. They have been known

---

\* See, for instance, Cases XLVIII. and L. in Todd's Clinical Lectures on the Urinary Organs.

to occur in both kidneys ; but this is of great rarity. The urine generally contains albumin and considerable pus ; it is acid and of low specific gravity.

Pyonephrosis cannot be distinguished from suppurative nephritis and ordinary *abscess of the kidney*, except it be by the history. The more constant and larger discharge of pus may be also made a point of diagnosis, as well as the obvious variations in the swelling and, as a rule, the slighter constitutional symptoms. But too much stress must not be laid on these points ; and the fact should not be overlooked that abscess of the kidney may be latent, or be present almost without fever, or with very obscure manifestations of pain, irregular attacks of fever, and vomiting, coming on at intervals for months or years.

When the changes resulting from an impediment to the flow of urine are unassociated with suppuration of the mucous membrane of the pelvis of the kidney, although the pelvis dilates extraordinarily from the accumulating urine and the kidney-tissue in time disappears, we have the condition designated by Rayer as *hydronephrosis*. It is due to mechanical obstruction from retroflexion or cancer of the womb, or from morbid growths or abscess of the bladder, or to congenital malformation of the ureter. Sometimes it is double. The swelling to which it gives rise may subside simultaneously with a sudden and copious discharge of urine. When this symptom is absent, the diagnosis must be based on the existence of a fluctuating renal tumor and on the absence of signs of suppuration.\* There may be attacks of renal colic due to the passage of clots of blood. The urine is at times copious, at times scant. The disease may lead to temporary, but entire, suppression of urine. Accurate percussion enables us to distinguish hydronephrosis from ascites ; in the former the dulness is generally one-sided, and it is uninfluenced by change of position. Ovarian cysts are more difficult to discriminate. Careful examinations by the rectum and by the vagina, and an investigation of the fluid after an exploratory puncture, are alone of value ; and even they may mislead. Urinary constituents, for instance, have been found to be absent in rare cases of hydronephrosis.

*Hydatid tumor* of the kidney is of comparatively rare occur-

---

\* See Schroeder, *Diseases of the Female Sexual Organs*, p. 385.



rence, and is very apt to be confounded with hydronephrosis. When the urine contains no hydatid vesicles or their débris and the hydatid fremitus is absent, the diagnosis is extremely difficult, and must rest chiefly on the history of the case.

Ordinary *renal cysts*, when large enough to occasion a tumor, cannot be distinguished from hydronephrosis save by the history, and by the albuminous and decidedly bloody urine which the cysts give rise to, while in hydronephrosis the urine presents nothing peculiar, or occasionally only small amounts of pus and of blood. Then, renal cysts are double-sided, preserve the shape of the kidney, and do not rapidly change their size.

Pyelitis may be connected with fibrinous clots due to repeated hemorrhages from *multiple aneurisms of the renal artery*. We may suspect this condition if the other more usual causes of pyelitis seem to be absent, and if the affection happen in an old person having repeated attacks of hæmaturia and atheromatous arteries.\*

### Disorders in which a very large Amount of Urine is discharged.

**Diabetes.**—In diabetes mellitus, or glycosuria, the urine is of pale color, decidedly acid, and of high specific gravity, ranging generally from 1030 to 1050. The quantity passed is enormous: seventy pints and upward have been known to be discharged daily. The urea is increased; so are the sulphates and the chlorides, and the earthy phosphates, while the alkaline phosphates vary greatly with the food, and uric acid is generally diminished; so is the coloring-matter. The urine contains from one to eight per cent. of sugar. In a small proportion of cases the flow of urine is not increased.

The symptoms attending the drain of fluid from the system are great thirst, constipation, and generally a dry, harsh skin, a red tongue, and a feeling of constant emptiness and of hunger. To these are added a steadily-progressing waste of the body, muscular feebleness, chills, a somewhat hurried breathing, a peculiar mawkish odor of the breath, peevishness of temper, a tendency to eczema and to boils and carbuncles, and in women pruritus of the vulva. The temperature is subnormal, often not over 96°. Cata-

---

\* Ollivier, Archives de Physiologie, 1873.

ract and other kinds of defective vision are not infrequent. The knee-jerk is generally absent. Galezowski\* has described a peculiar form of retinitis in diabetes; retinal hemorrhage, and palsies of the muscles of the eyeball, have also been noticed. Attention has been directed to diabetic hypermetropia, and with the change of refraction a quantity of sugar in the urine is observed.† Defects in accommodation are common. Diabetic endocarditis also happens, and is more frequent in women than in men;‡ and neurites and neuralgias and periostitis§ and arthritic disorders|| may have their origin in diabetes. Double sciatica is often of diabetic source. There is frequently a connection to be traced between gout and diabetes.

Diabetes is a very fatal disease; yet it is impossible to foretell its exact mode of termination. Some are cut off rather suddenly; others drag out a long existence, and die worn out and dropsical, or of cirrhosis of the liver, or of broncho-pneumonia, or of phthisis. For some days, or even for weeks, before death, the sugar may disappear from the urine.¶ Diabetic gangrene is also a mode, though not a frequent one, of termination of the disease.\*\*

When the disease ends suddenly, it is apt to do so by so-called diabetic coma. The comatose condition is prone to be preceded by vomiting and abdominal pain, rapid pulse, great anxiety and restlessness, labored breathing, depressed body-heat, headache, and drowsiness. These symptoms are attributed to the poisoning of the body by the development of *acetone*, a derivative of acetic acid, in the blood; the acetone can be found in the urine, and may be readily detected on the breath by its odor resembling that of chloroform. The evidence, however, of the decomposition of the sugar into acetone, and of the consequent nervous symptoms

\* *Compte-Rendu du Congrès Ophth. de Paris*, 1862.

† Landolt, *El Siglo Médico*, quoted in *Lancet*, April, 1880.

‡ Lecorché, *Arch. Gén. de Méd.*, June, 1882; *Bulletin de l'Acad. de Méd.*, 1880.

§ *Arch. Gén. de Méd.*, Feb. 1882, and *Amer. Journ. Med. Sci.*, April, 1882.

|| Dyce Duckworth, *St. Barth. Hosp. Rep.*, vol. xviii., 1882.

¶ In a case for a long time under my charge, in which the diabetes lasted for several years, sugar entirely disappeared from the urine as the signs of phthisis became fully developed, for some months before death.

\*\* See cases collected by Hunt, *Transact. Phila. Co. Med. Soc.*, Nov. 1888.

called diabetic coma, is still not conclusive. At all events, it is certain that there may be diabetic coma without acetonuria, and acetonuria without coma and even without diabetes. Diacetone is believed to be more certainly than acetone a forerunner of diabetic coma.

Whence comes the sugar? Is it from the food? the blood? the stomach? the liver? These are questions that cannot be satisfactorily answered. Since Bernard's discovery of the sugar-forming properties of the liver, saccharine urine is thought to proceed from an inordinate formation in this viscus of sugar, which is not fully destroyed in the lungs and is excreted by the kidneys. On the whole, this view has been sustained; but it is impossible to apply it clinically as an exclusive theory. We find sugar in some instances of gastric malassimilation, yet oftener in liver affections, especially in cirrhosis. That the sugar is not derived from the sugar in food is certain; for patients kept even on the most rigorous meat diet still pass sugar. In some cases diabetes has been observed to be associated with paralysis of the tongue, palate, and vocal cord, and other signs of disease in the floor of the fourth ventricle, or of tumors pressing there; or it has been noticed after fractures of the skull involving the base. Much evidence of the frequent connection of diabetes with structural alterations of the nervous system has been adduced. That there is a diabetes of nervous origin can, indeed, not be doubted. Again, diabetes in a number of cases has been found to be linked to a lesion of the pancreas. It also often follows mental emotion and worry. In some instances it is malarial, in many hereditary.

Starchy and saccharine substances increase the quantity of diabetic sugar. Nay, they may be the cause of a little sugar appearing in the urine of healthy persons. Yet those in whom a saccharine state of the urine is readily induced are in great danger of becoming diabetic. If we are in doubt whether we are dealing with a case of diabetes, we may follow Seegen's advice and let the patient eat heartily of saccharine and sugar-forming substances, and examine the urine three hours after the meal; if no sugar then be found in the urine, diabetes may be excluded.

In the aged, sugar may be present in the urine without being attended with distressing symptoms. It is in such cases that we are most apt to meet with the intermitting diabetes to which

attention has been called by Bence Jones.\* When the abnormal ingredient thus disappears from the urine, it is replaced by uric acid and by oxalates. There is still another form of intermitting glycosuria. Sugar is sometimes—Burdell† says uniformly—found in the urine during the paroxysms of intermittent fever; but it vanishes during the intervals.

Sugar is also found in the urine in small quantities after inhaling chloroform or taking chloral. Among the insane, sugar may be present in the urine without there being other symptoms of diabetes, and without grave significance.‡ Indeed, this appearance of sugar in the urine from passing causes or without other marked symptoms has given rise to the distinction made by some between *glycosuria* and *diabetes*, restricting the latter term to persistent saccharine urine with decided symptoms. The passing glycosuria gets well; true diabetes rarely does.

In some instances we have *diabetes with coexisting albuminuria*, and even with other evidence of Bright's disease. In the majority of such instances the degeneration of the kidneys has happened subsequently to the diabetes, and in its more advanced stages from its constant irritation; but I have met with cases in which Bright's disease has preceded the diabetes. Amyloid kidney has also been noticed in connection with diabetes. A small amount of albumin in diabetic urine is not uncommon.

**Chronic Diuresis.**—This disease is otherwise known as *polyuria*, or *diabetes insipidus*. It is characterized by the habitual discharge of a very large quantity of urine of low specific gravity, from 1001 to 1008, containing an excess of water, but no sugar; urea is increased; uric acid is very deficient; inosite is often present; kreatinin may be excreted in increased quantity. The general symptoms are much the same as those of diabetes; the thirst is generally extreme, and it may happen that more water is passed than is drunk. Most cases recover under treatment, except when dependent upon irremediable lesion. They sometimes die of suppression of urine.§

The cause of this singular malady is obscure. We meet with

\* Medico-Chirurgical Transactions, vol. xxxviii.

† L'Union Médicale, No. 139, 1859.

‡ Lailler, quoted in Journal of Mental Science, May, 1871.

§ Case under my charge at the Philadelphia Hospital.



polyuria after cerebro-spinal fever, or in connection with tumors of the brain, or with disease of the medulla oblongata or of part of the floor of the fourth ventricle, or with tumors compressing the abdominal ganglia. Lancereaux tells us that the disorder is not uncommon in syphilitic affections of the nervous centres;\* and Bartholow's experience is that syphiloma of the brain is its most usual cause. I have repeatedly encountered the malady after injuries to the head,† after sun-stroke, or in persons broken down with malaria. At times it is seen in instances simply of great nervous depression without organic disease. It is, indeed, mostly connected with some abnormal state of the nervous system. It has been stated to coexist with marked excess of phosphates, and to be a phosphaturia.

We must take care not to confound cases of chronic polyuria with true *diabetes*. They differ by the low specific gravity of the urine, and the utter absence of a saccharine ingredient. Sometimes a state of diuresis is found to exist temporarily during the removal of dropsical effusions, or when the action of the skin is insufficient. We also meet with apparent cases of diuresis in *hysterical women* and in persons who suffer from incontinence of urine, whether due to an external injury, or dependent upon simple irritability, or upon inflammation or paralysis of the bladder. In all such, however, we can establish the diagnosis by laying stress on the history of the patient, and by measuring the amount of urine passed in the twenty-four hours,—which amount may be large, but is not inordinate. In hysteria it may be temporarily very large after a paroxysm, but is not persistently so. In some instances diabetes mellitus alternates with diabetes insipidus. The discovery of an hydræmic centre in the cerebellum by Eckhard, as well as the well-known points at the floor of the fourth ventricle, which, according to the exact seat of puncture, produce increased flow of urine with sugar or without sugar, gives us the clue in which direction to look for the explanation of such cases. The large flow of urine we sometimes meet with in *contracted kidney* is told from hydruria by the presence of albumin and tube-casts and the other signs of kidney degeneration. An

---

\* Sydenham Society's Translation, p. 77.

† Transactions of the College of Physicians of Philadelphia, 1875.

excessive flow of urine may happen in *hydronephrosis*. But the antecedent history, the previous existence, as a rule, of a fluctuating tumor, and the character of the urine, either normal or containing at times traces of albumin or of blood, will throw light on the character of the malady.

### Disorders in which little or no Urine is discharged.

**Suppression of Urine.**—Suppression of urine, unconnected with degeneration of the kidney, is a rare disorder. Yet it may occur in previously healthy persons, or in the course of fevers of low type, and probably associated with no other morbid state than congestion of the kidneys. It is occasionally met with as one of the freaks of hysteria, or is caused seemingly by the irritation reflected to a healthy kidney from a diseased bladder.

The symptoms it occasions, independently of the absence of the discharge of urine, are drowsiness, nausea, vomiting, coma, sometimes convulsions; in one word, the symptoms of uræmic poisoning. The formidable complaint may give rise to marked urinous smell of the perspiration and of the breath, and to exceeding and very general cutaneous hyperæsthesia.\* The temperature may be low, and remain so even if there be coexisting internal inflammation, or be above the norm.†

Concerning the exact cause of the suppression we are often kept in the dark until the termination of the malady; for, unless familiar with the patient's antecedent symptoms, we are unable to determine, in the absence of the urinary secretion, whether or not a disease of the kidney lies at the origin of the mischief.

Oppolzer tells us that we may diagnosticate *thrombosis of the renal vein* if we have diminution of the secretion of urine and its final suppression preceded by blood, albumin, and casts in the urine. If there be a history of severe injury to the kidney, these symptoms have a much more positive meaning.

---

\* This was the most obvious symptom in a case under my care in 1864 at the Philadelphia Hospital, in which no urine was secreted for many days, the catheter being repeatedly introduced into the bladder. The patient recovered. She had, previously and subsequently to the attack, vesical catarrh. In a case reported by Fuller, St. George's Hospital Reports, vol. v., the difficulty existed for eight days without occasioning convulsions. It was the same in a case of mine that lasted eleven days and got well.

† Bourneville, Gaz. Méd. de Paris, 1872.

**Retention of Urine.**—The urine retained in the bladder distends the viscus and forms a swelling in the hypogastrium, discoverable both by palpation and by percussion. The urine is generally not wholly kept back, for a slight discharge every now and then takes place, or there is a constant dribbling,—a matter which in itself should suggest the introduction of a catheter.

Retention of urine, if soon recognized, is not a dangerous complaint, as it can be at once relieved by the passage of a catheter; but if the ailment escape observation, or be inefficiently dealt with, the bladder may burst,—though Sir Henry Thompson tells us that this is a circumstance of exceeding rarity,—or the patient dies from the absorption of the noxious urinary ingredients.

The causes which lead to retention are various; prominent among them, at least in a medical point of view, is paralysis of the bladder, especially that form of paralysis which occurs in low fevers; retention is also one of the symptoms of paraplegia; then inflammatory swelling of the neck of the bladder, organic stricture, or enlarged prostate may give rise to it; again, retention or incontinence may be due to hysteria. If the urine be long retained in the bladder, it becomes alkaline, and putrefactive changes occur, and fission fungi, especially the *micrococcus ureæ*, develop in great numbers in the ammoniacal urine.

The disorder is readily detected. It may be discriminated from suppression of urine by the existence of the hypogastric tumor, and by the introduction of a catheter,—a means which, in cases of doubt, ought never to be neglected. Sometimes the abdominal swelling is so great as to lead to the belief of the existence of dropsy; and the error is fostered by learning that the patient has been passing his water, and has a constant desire to discharge it, or by seeing that it dribbles from him.\*

The retention from paralysis is distinguished from that due to other causes, as obstruction, by observing that the catheter enters readily, and that the urine flows out in a continuous stream, increasing and lessening with the respiratory movements, but does not come out in jets.

---

\* In a case reported by Schneider, and quoted in *Brit. and For. Med.-Chir. Rev.*, April, 1864, urine was passed; yet when, owing to the peculiar shape of the tumefaction, a catheter was introduced, fourteen pints of urine, and subsequently eight more, were removed.

## CHAPTER VIII.

### DROPSY.

AN abnormal collection of watery fluid in the areolar tissue or in the serous cavities of the body constitutes dropsy. Now, dropsy is but a symptom, and as such we have already examined into it as associated with various disorders; but, though only a symptom, it is one so obvious, and comprises so often apparently the whole complaint, that it will serve a useful purpose to investigate connectedly the clinical meaning of its typical forms.

#### Dropsy, according to its Seat and Extent.

Dropsies may be external, or be confined to internal parts. To the latter variety belong hydrothorax, hydrocephalus, and ascites,—affections elsewhere described, which we shall consider here only so far as they may form part of a general dropsy.

External dropsies are illustrated by *anasarca* and *œdema*: the first, a universal accumulation of serous fluid in the areolar textures; the second, a localized collection in the same structures, differing, therefore, in nothing but extent. Both, as ordinarily met with, exhibit painless swelling of the surface, devoid of redness; a skin often stretched and shining, pitting upon pressure, and retaining for some time the mark of the finger; and in both, the tumid part, if punctured, allows a watery fluid to run out. *œdema* is most commonly perceived around the ankles; the tumefaction of *anasarca* is found generally not only in the lower extremities, but also in the arms and in the face.

*Anasarca* is usually dependent upon disease of the kidneys, or of the heart. The swelling rarely shows itself at all parts of the body at once; it ordinarily begins at the feet and ankles in diseases of the heart, in the face in diseases of the kidney. It is greatest where the areolar tissue is loosest.



*Œdema* may be due to the same causes. Yet a limited collection of fluid is often the consequence of a purely local difficulty unconnected with visceral disease, but of a character interfering with the venous circulation. Thus, the compression or obliteration of a large vein occasions *œdema* below the point of the disorder. We see *œdema* happening if a bandage be applied too tightly, or if swollen glands press upon the main vein of a limb. We also meet with it in the adhesive form of venous inflammation, and in milk-leg, or *phlegmasia alba dolens*. In all of these forms the *œdema* is one-sided; and, the cause being external to the thoracic or the abdominal cavity, there is little difficulty in its recognition. A circumscribed *œdema* also accompanies erysipelatous inflammations of the skin or subjacent tissues; so, too, do we find *œdema* confined to a limb the general nutrition of which has been lowered by paralysis.

When the dropsical effusion is dependent upon a *tumor* seated in an internal cavity and interfering with the passage of the blood, it may be very local and one-sided, as we sometimes find in connection with abdominal cancer; but it is most apt to be found on both sides of a portion of the body, although more particularly marked on one side. The *œdematous* extremities exhibit usually also marked enlargement of the veins.

Another source of a double-sided *œdema* is *anæmia*. This form of dropsy is often seen without there being any disease of an internal organ. The watery state of the blood is highly favorable to the transudation of the serum, and this collects first about the ankles, and subsequently in other parts of the body. The absence of any discoverable organic affection, the pallid countenance, the pearly whiteness of the conjunctiva, and the venous murmurs in the neck, are very significant. A microscopical examination of the blood and a blood-count establish the diagnosis.

A dropsical effusion in part of similar origin, but much more often connected with *internal dropsy*, especially with ascites, is the dropsy we observe in those broken down by *malarial poisoning*. The state of the liver and spleen, added to the condition of the blood, determines the greater extent of the effusion.

### Dropsy, according to its Causation.

Having viewed anasarca and œdema as in the main uncombined with internal dropsies, and as forming the sole signs of the dropsical complaint, let us now look at them when associated with effusions of serum elsewhere. The same remarks will also apply to hydrothorax and to ascites, the meaning of which, when occurring alone, we have inquired into, but which we shall here consider in their relations to *general* dropsy, or that form of the disorder in which anasarca or œdema coexists with dropsy of one or of several of the large serous cavities.

First, let us examine into the causes of general dropsy. The most common are a disease of the heart, of the kidneys, or of the liver; so common, in truth, that in every case of dropsy we must always examine these organs carefully. According as the dropsical accumulation originates in a morbid state of these viscera, it is called cardiac, or renal, or hepatic.

*Cardiac dropsy* arises in consequence of the deranged or enfeebled circulation produced by a disease of the walls and cavities of the heart, associated or not with a valvular lesion. The dropsy begins in the feet and ankles, being much influenced by position, and gradually extends upward; but it is rarely very obvious in the face or upper extremities. The thighs and scrotum are sometimes greatly swollen, and there is a watery effusion into the pleural cavities or into the pulmonary parenchyma.

*Renal dropsy* is usually much more general than cardiac dropsy. It does not, like this, begin in the most dependent parts, but is often first noticed in the face and eyelids. There is hardly a space in the body where, as the complaint progresses, fluid may not accumulate. The proof that the dropsy is renal is furnished by the presence of albumin and of casts in the urine.

Occasionally the dropsy is owing to an affection both of the kidney and of the heart; and the inquiry may arise, which of the organs was primarily disturbed and gave rise to the dropsy? The kidney disorder generally precedes the heart disorder. Should it be of importance, in an individual case, to determine which has occurred first, we may be enabled to arrive at a conclusion by a close examination of the history of the case: did the patient

suffer from palpitation and shortness of breath prior to or coincident with the anasarca condition, and has he ever had rheumatic fever? or did he have an attack of acute dropsy before the persistent swelling of the feet or of the face occurred? If this has happened, there is a strong probability of the renal disease having been antecedent to the cardiac malady.

*Hepatic dropsy* may, like the preceding forms, be more or less general; but it is very rarely so, unless it be of long standing, or unless there be coexisting disease of the heart or of the kidneys. The most usual kind of dropsy depending upon an affection of the liver is abdominal dropsy, and this is so well understood that ascites is frequently looked upon as constituting a proof of hepatic disorder. But it is a mistake so to regard it; for ascites may also be produced by peritoneal tumors or inflammation, by enlargement of the spleen or of the pancreas, or by the pressure of diseased glands,—in fact, by any lesion which occasions a decided impediment to the portal circulation.

Again, it is possible, though it is not a cause which acts often, that mere irritation of the *areolar tissue* will occasion more or less general dropsy. This was a favorite doctrine of the older physicians; and H. C. Wood thus explains the dropsy of arsenical poisoning.\* Another cause of general dropsy, especially of anasarca, is *peripheral multiple neuritis*. I have seen this in obscure cases, in which the electric reactions, the absence of the knee-jerk, the altered sensation, made the diagnosis clear.† Perhaps this is the cause of the dropsy in *beriberi*,—an affection with decided anasarca.‡

Besides these sources of general dropsy, we may find deterioration of the blood, with, perhaps, a simply enfeebled condition of the heart, giving rise to it. But such a state is much more likely to occasion œdema, or, in some instances, anasarca, than general dropsical effusions; and it is thus that, while the former phenomena are not uncommon in exhausting diseases or in marked impoverishment of the blood, the latter are rarely met with unless there be at the same time some cardiac or renal complaint.

---

\* Amer. Journ. Med. Sci., July, 1871.

† As in a case seen with Dr. Lewis Brinton.

‡ Simmons, *Beriberi*, or the “*Kakké*” of Japan, Shanghai, 1880.

There is a disease allied in its symptoms to dropsy which has attracted much attention. It is the disease pointed out by Sir William Gull, and called by Ord *myxœdema*, consisting in the progressive invasion of the body by a mucus-yielding dropsy, unassociated with albuminuria or disease of the heart, but invariably combined with destructive change and decrease of the thyroid gland. It affects chiefly adult women, who present swollen, waxy-looking features, with a circumscribed flush on the cheeks, and are markedly anæmic, having an interstitial development of fibrous tissue, with an excess of subcutaneous fat. The skin is everywhere thickened and rough, is devoid of perspiration, and the puffy integuments do not pit, or pit but slightly, on pressure. The hands are often swollen and misshaped, the nails are brittle; there is loss of teeth and of the hair. The temperature is apt to be below the normal, the excretion of urea to be diminished. The movements of the limbs are slow and languid; the gait is uncertain; sensation is impaired; there is irritability of temper, with increasing hebetude, monotonous voice, slow, drawling speech; finally, melancholia and aberration of mind may supervene. The disease, so similar to a cretinoid state, may be artificially produced by the removal of the thyroid gland. It has been also noticed after the long administration of iodide of potassium.\* In a certain number of cases the atrophy of the thyroid is preceded by hypertrophy.† The disease may be preceded or be attended with intractable uterine hemorrhage without apparent cause.‡

The swelling of myxœdema is distinguished from the dropsy of *acute nephritis* by affecting the forehead as well as the face, and by causing a thickening of the bridge of the nose and of the lips and the tongue; the swelling of the body and the legs is uniform and uninfluenced by gravitation; there is not the swelling of the scrotum that we see in *acute nephritis*.

It is more difficult to distinguish *contracted kidney* from myxœdema. In both we may find excessive flow of urine of low specific gravity, and a few hyaline casts associated with very small amounts of albumin. But in the exceptional cases in which we have dropsy in contracted kidney, it shows itself in the lower ex-

---

\* Stalker, *Lancet*, Jan. 1891.

† Ord, quoted in Sajous' *Annual*, vol. iv., 1891.

‡ Kirk, *Lancet*, Sept. 1893.



tremities, and not on the forehead, the lips, and the tongue. Then, we do not find in contracted kidney the dry desquamating skin, and the loss of hair so common in myxœdema.\*

### Dropsy, according to the Rapidity of its Development.

Dropsy may come on suddenly, or be gradually developed. The first is called *acute* or *active* dropsy; the second, *chronic* dropsy. To the latter class belong the majority of instances of the forms of dropsy just discussed, in which the watery accumulation is thought to arise from defective action of the absorbent vessels, or in which, in other words, the dropsy is passive. Acute dropsy has active symptoms much like those of an inflammatory fever. The effusion takes place suddenly, and in consequence of exposure to cold and wet, or of checked perspiration. In the vast majority of examples it is accompanied by albumin in the urine, and is, in truth, due to an affection of the kidneys. Yet there are cases of acute dropsy which are not of renal origin, and in which the rapid occurrence of universal anasarca is not susceptible of being traced directly to a definite lesion.

---

\* Allen Starr, Med. News, Dec. 1894.

## CHAPTER IX.

### DISEASES OF THE BLOOD-VESSELS.

ONLY a short description of these will here be given, partly because many of the diseases of the arteries and veins have already been mentioned in connection with other maladies, and partly because the knowledge we have of a number of them is still pathological rather than clinical.

#### Diseases of the Arteries.

The principal of these are inflammation and atheromatous changes.

**Arteritis.**—Inflammation may attack the outer coat, *periarteritis*, the inner coat, *endarteritis*, or all the coats, *general arteritis*. All these processes may be the result of rheumatism, of gout, of syphilis, of lead poisoning, or of inflammation spreading from surrounding textures.

In *periarteritis* the last-named is the most common cause. The large arteries are the ones that are pre-eminently affected, and inflammation of the external coat of the thoracic aorta is more often encountered than that of any other artery. It may be acute; occasionally it takes its origin in inflammation of the inner coat. It may lead to suppuration, and, the pus finding its way into the calibre of the vessels, pyæmia and metastatic abscesses are caused. But it is not possible to make a certain diagnosis of the condition.

There is a peculiar disease of the arteries, *periarteritis nodosa*, which, with the signs of acute desquamative nephritis and fever, gives rise to small swellings under the skin, to rapid loss of muscular power with deficient electro-muscular contractility, and to such severe muscular pains that they are readily mistaken for

those of the trichinous affection.\* But the history of the ailment, the signs of the thickening of the vessels, and, if necessary, an examination of the muscles, will throw light on the cause of the muscular distress.

*Endarteritis* is almost always chronic, and chronic endarteritis is most commonly due to rheumatism, to gout, to the poisonous influence of lead or of arsenic, or is seen in connection with contracted kidney. As regards the latter, the question may arise as to whether it has caused the chronic inflammation of the arteries or is a mere coexisting affection owing to the same general morbid process. Arthur V. Meigs† in several able papers urges this view, and I believe it is generally the true explanation. It is certain that chronic endarteritis is found without Bright's disease, or preceding it, and gives rise to symptoms by which it can usually be recognized. It is very commonly described as *arterio-sclerosis*, in consequence of the hardening of the walls of the artery, the result of the endarteritis, and especially of the intima.

The inflammatory thickening of the intima of the arteries and arterioles starts as simple inflammation of the lining, but may extend to some degree into the veins. The symptoms to which chronic endarteritis gives rise are epistaxis, or hemorrhages into internal organs, such as the brain or the lungs; œdema without recognizable cause; attacks of bronchitis or catarrhal pneumonia; and torpor of the liver. An appearance of prominence of the smaller vessels and their greater resistance show the fully-developed disease, and we are then able to find marked nervous symptoms, such as vertigo, loss of memory, general want of power in the limbs, and anæmia. Valve-changes may also present themselves, and albumin and casts in the urine, and other signs of kidney affection. But these do not necessarily occur. Again, there are cases in which they seem to precede the endarteritis. The visceral complications of the malady make statements about the temperature uncertain, but I believe that it is persistently slightly elevated. Endarteritis is at times compensatory in slowing of the blood-current.‡ An accentuation of the

---

\* Kussmaul and Maier, quoted in Schmidt's Jahrb., No. 8, 1868.

† Transact. College of Physicians of Philadelphia, 1888 and 1889.

‡ Thoma, Virchow's Archiv, April, 1888.

second sound, as well as its reduplication, is a very usual feature in arterio-sclerosis.

Extensive inflammation of the arteries, a *general arteritis*, is a very rare affection, and when it happens it is acute. In a few instances of rheumatism we find *acute arteritis* arising, and especially inflammation of the fibrous structures of the aorta. This condition may be suspected should we observe intense general uneasiness and distress, with pain, increased pulsation, a distinct murmur in the course of the vessel, and tumultuous action of the heart without there being obvious signs of disease of that organ present. Still, the diagnosis is never a positive one. The result of the inflammation, particularly of the acute endarteritis which forms part of it,—seen, too, when this alone is present,—is that the blood may clot, and thrombi or emboli, as well as pyæmia, result. The coagula that are formed break away, and occasion embolism, generally in the smaller peripheral vessels; though thrombi may also develop in the larger vessels, and pyæmic fever result. It is generally impossible to recognize the malady until after the coagula occur, and then signs of narrowing of the calibre of a vessel, with a localized murmur, may give us the clue to the cause of the symptoms. Hæmatomata may form after embolic arteritis and consequent perforation.\*

**Atheromatous Changes.**—These are the common results of processes of decay occurring especially in persons of advanced years. But we may also find them in chronic Bright's disease, in syphilis, and in lead poisoning, and they may be so much mixed up with the signs of chronic endarteritis that it is frequently impossible to separate the manifestations of the maladies. In truth, even when the atheromatous changes are predominant they are not always easy of recognition. These alterations, happening in internal arteries, are beyond the accurate discernment of the physician. He may infer that they exist, if a distinct systolic blowing sound be heard in the track of the aorta or its branches, in a person who is not markedly anæmic, who is past middle life,—and therefore at an age at which these kinds of changes of tissues happen,—and in whom no cardiac murmurs, or only faint cardiac murmurs, are perceived. But it is chiefly by the age

---

\* Rushton Parker, Transact. Clinical Society of London, 1881.



of the patient, the general circumstances of the case, the rigid resisting superficial arteries, often irregular to the touch, and the gradual development of cardiac enlargement from the resistance to the circulation, that a conclusion as to the meaning of the physical signs is arrived at. These changes in the arteries are important in themselves, and important also because they lead to thrombi and are often the first step toward the laceration or the dilatation of the vessels,—in other words, toward the establishment of an aneurism, which may be miliary aneurism, as in the brain and lungs, or a large aneurism, as in the thoracic and abdominal aorta. The atheromatous change may be so great as to cause almost complete occlusion of large arteries, as of the common carotid.

### Diseases of the Veins.

The chief affection of the veins in a diagnostic point of view is inflammation.

**Phlebitis.**—This is met with by the surgeon much oftener than by the physician, who encounters it more especially in affections of internal organs, such as the liver, and has to study it in connection with the formation of thrombi, and metastatic abscesses to which it leads; as, for instance, in inflammation of the veins of the liver. But the most common form of phlebitis which comes under the cognizance of the medical practitioner is milk leg, or *phlegmasia alba dolens*. Here we have usually phlebitis with an obstruction by a coagulum of the venous circulation in the affected limb, and in the coagulum cellular elements and connective tissue and fine vessels are found.\* There are also cases of thrombosis without inflammation of the veins. The leg becomes painful, swollen, shiny, and œdematous. The femoral vein is tender and blocked by a clot; sometimes this is in the crural. The disease is seen in puerperal women, or as a sequel of typhoid fever or of chlorosis. The phlebitis or the thrombus that forms may be of septic origin. This is most likely in the puerperal state, probably also in typhoid fever. The disease is, with rare exceptions, one-sided. The pain in the leg may cause it to be mistaken for rheumatism, but the one-sided swelling and the œdema dis-

---

\* Damaschino, Bull. Soc. Méd. des Hôp., 1880.

tinguish it. Among its early and significant symptoms is pain on pressing the calf of the leg on the affected side.

### Diseases of the Capillaries.

Some of the organic diseases of the capillaries belong to the arterio-sclerosis in Bright's disease, or to the waxy degeneration in purpura. It is difficult to say what the functional disorders are, for many of them are regarded as forming part of the peripheral diseases of the nervous system, and the affection of the arterioles and of the capillaries is a mere vaso-motor spasm in connection with the neurosis. This is supposed to be the case in the anomalous localized sensations of cold which some patients have in particular parts of the body, though their persistency is unlike the history of a spasm. The painful flushings of the feet bespeak temporary excessive dilatation of the fine vessels. The so-called "dead fingers" are regarded as a spasm of the arterioles. They are most common in emotional women, but they have also been observed in men.\* The finger-tips become white and numb, warmth and feeling returning after a variable time. An attempt has been made to prove a very close connection with Bright's disease, but the dead-finger symptom is not characteristic of this. The disorder may happen only at night, forming part of a passing loss of power and sensation,—the so-called "night palsy." It may disappear with old age.†

A spasm of the minute vessels of more permanent character may lead to profound disturbance of nutrition in a part, even to its destruction. This is the case in the so-called *symmetrical gangrene*, or "Raynaud's disease." It is a remarkable neurosis, producing purple discolorations, which are very painful and on which bullæ form. These symmetrical patches of blood-stasis or local asphyxia lead to gangrene, which, however, does not terminate fatally, and is generally completed within ten days. The malady is often seen in the hands, beginning in or certainly affecting corresponding fingers. The patches of local asphyxia, believed to be due to spasm of the arterioles, may recur for months again and

---

\* J. E. Squire, *Lancet*, Dec. 6, 1886.

† Donaldson, *ib.*

again without the disorder causing mortification of the skin, or this may take place in any attack. The disease may not happen in the fingers or the toes, but on the exterior surface of the forearm and below the knee. By far the greatest number of cases occur in winter, and the disease is thought to be allied to paroxysmal hæmoglobinuria, which, indeed, has been repeatedly observed in association. Neuritis of the affected part has been found by a number of observers. Henry \* looks upon neuritis as a mere coincidence.

Raynaud's disease must not be mistaken for *chilblains*. These do not appear, disappear, and reappear in the manner in which the discoloration does in Raynaud's disease. In *erythromelalgia*, described by Weir Mitchell,† there are vascular changes, acute congestion, or cyanosis. The disorder manifests itself in the heel or the sole of the foot, and is attended with great pain.

---

\* Trans. of Assoc. Amer. Physicians, 1894.

† Medical News, Aug. 1893.

## CHAPTER X.

### DISEASES OF THE BLOOD.

IN the following sketch I shall attempt to describe only those diseases of the blood which are seemingly, for the most part, idiopathic, and may be recognized by well-marked clinical traits. Prominent among these, and to a certain extent characteristic of all blood disorders, are general debility, a changed aspect of the mucous membranes and of the skin, especially in color, and alterations of nutrition.

In the investigation of diseases of the blood the microscope is of the first importance. It informs us with regard to the relative proportions of the white and red globules, and exhibits the peculiar homogeneous, fibrinous blood-plates or hæmatoblasts. It tells us something as to what part of the blood-making organs the former are derived from; it indicates whether the red globules are of the right color, whether their outline is regular, whether they form rouleaux properly, and whether their number is decreased. In this respect recent research has aided us much by supplying us with accurate means of computation.

The method of determination of the globular richness of the blood introduced by Vierordt, in 1854, was to allow a stated amount of a definite dilution to dry upon a glass slide, and then by the aid of the micrometer to count the number of the globules. Imperfect as it was, by it he ascertained that the normal number of blood-corpuscles in a healthy male adult was between five and six millions to a cubic millimetre, and that in certain diseases this number was much diminished. Clinical observers confirmed these observations, and subsequent improvements have rendered the apparatus more precise and made the results more accurate. The forms of apparatus now mostly in use are the hæmacytometer of Thoma-Zeiss and the hæmacytometer of Gowers. To these has been added the graduated moist-chamber globule-counter of



Malassez. Another form of apparatus for determining the globular richness of the blood is the hæmatokrit of Hedin.

The Thoma-Zeiss, or Zeiss, hæmacytometer is very simple. It consists of three parts: first, a graduated pipette or mixing-vessel, with rubber tube attached; second, a counting-cell on an object-slide made of ground glass; third, a cover-glass with ground level surfaces.


When counting the red corpuscles of the human blood, the tip of the finger should be thoroughly cleaned, the middle finger of the left hand being generally selected. By rubbing the end of the finger with a coarse towel a slight hyperæmia is induced, so that a cut with a spear-pointed needle will permit of the flow of a drop of blood sufficiently large for examination. The tip of the pipette is placed into this drop, and the blood carefully drawn up to the mark 1,—*i.e.*, one cubic millimetre. After this has been accomplished, the tip should be cleaned by means of a soft cloth and the pipette inserted into a carefully-filtered ten per cent. solution of sodium sulphate, or Thoma's substitute of a three per cent. solution of sodium chloride. This is drawn up into the tube until the bulb is filled to the mark 101. The blood and fluid are then thoroughly mixed by shaking the tube, holding the finger over the tip of the pipette, that the liquid may not escape. After the mixture has been thoroughly effected, half of the fluid in the bulb is blown out, and the drop that follows is allowed to flow on to the previously cleaned floor of the counting-cell. The cover-glass is then immediately placed in position, and the apparatus allowed to rest quietly upon a horizontal surface for a few moments, that the corpuscles may be permitted to settle. For the success of this operation perfect cleanliness must be maintained throughout.

In order to make the examination, the slide should be placed in the stand of the microscope and held in a horizontal position, that the corpuscles may not be displaced. The cover-glass should lie accurately; great care should be taken that no liquid flow between the cover-glass and the ring. It is important that the drop of blood mixture shall remain standing in the centre of the cell, and that by the spreading of the cell the under surface of the cover-glass shall be in contact with the mixture for several millimetres. Using a one-fourth or a one-fifth objective glass to

bring into view the divisions cut upon the floors of the cell, we find that upon these lie the red blood-corpuscles. The number of corpuscles in each space is then noted. Through each fifth horizontal and vertical row of the lines an additional line is drawn, for the purpose of fixing more readily the position of the squares counted.

Each field of the net-work contains a surface of one four-hundredth of a square millimetre. The distance of the cell-floor

FIG. 68.



Artificial capillary of Malassez, magnified 100 diameters.

from the under surface of the cover-glass is one-tenth of a millimetre. Each square, therefore, represents the one four-thousandth of a cubic millimetre. The number of corpuscles contained in one of these cells multiplied by the number of times the blood has been diluted will give the amount of corpuscles contained in the one four-thousandth of a cubic millimetre. The amount contained in a cubic millimetre can, therefore, be found by multiplying by four thousand. The surest method is to count at least forty spaces, to take the average of them all, and proceed as above. It is sometimes rather difficult to distinguish the white from the red blood-corpuscles, and this difficulty is obviated by adding the one-third per cent. solution of acetic acid to the diluted blood. Another method for computing the white corpuscles and


their relative number to that of the red is to use, with the salt solution, a few drops of a one per cent. solution of gentian violet; this leaves the red blood-corpuscles unaltered and stains the leucocytes a deep violet.

The following method for differential counting of leucocytes in fresh blood has been devised by Elzholz. After drawing blood into the pipette, a solution composed of seven grammes of two per cent. eosin solution, forty-five grammes of glycerin, and fifty-five grammes of water is added; then, a solution composed of four drops of concentrated watery solution of gentian violet with one drop of absolute alcohol and fifteen grammes of water, by which the polynuclear cells are more deeply stained; the eosinophile cells are reddish violet.

The *hæmacytometer* of Gowers is about the same as that of Zeiss, differing mainly in the number of divisions on the cell, each space being but one-tenth of a millimetre in length. The method of preparing the blood solution is not so convenient as that of Zeiss. A hæmic unit of five millions of corpuscles to one cubic millimetre of blood is assumed.

By the original method of Malassez the blood was diluted with artificial serum so that it represented  $\frac{1}{100}$  or  $\frac{1}{200}$  of the original. A small amount was then introduced into a flattened capillary tube of known capacity and, with the micrometer eye-piece, the globules were counted in the capillary tube of a certain length, say 500 micro-millimetres. The capacity of this length of the tube in parts of a cubic millimetre being already known, the entire number of globules in a cubic millimetre of the undiluted blood was easily determined by calculation. For the purpose of diluting the blood\* Potain's capillary pipette (Fig. 69) is well adapted. It is so constructed as to contain in a part of its extent a reservoir

FIG. 69.




Potain's pipette.

\* Malassez recommends for artificial serum a five or six per cent. solution of sodium sulphate, having a specific gravity of 1020 to 1024.

imprisoning a glass bead, the capacity of this chamber being exactly one hundred times that of the capillary tube leading to it. To the opposite extremity is attached a rubber tube, which, when placed between the lips, causes the fluid to ascend to the desired extent by aspiration, or by blowing through it the tube may be emptied.

Malassez, in describing his new globule-counter,\* recommends, without abandoning his original design, an improved cell for

FIG. 70.



Graduated moist-chamber of Malassez. In the lower figure the compressor is seen attached to the slide.


microscopic work. It consists of a thick glass slide having ground in the centre of its upper surface a ring or circular trench one and a half millimetres in breadth and one millimetre in depth, which leaves a plateau about seven millimetres in diameter separated from the remnant of the surface of the slide by a narrow gutter, so that when the cover is in position water may be placed under it by capillary attraction, but cannot reach the islet in the centre. In this way any fluid may be protected from evaporation while under examination,—an important precaution while counting blood-cells. Outside of this ring, three or four holes pierce the glass slide, from which the points of screws are made to

\* Arch. de Phys., 1880, and Oct. 1882.



project, so as to maintain the cover-glass at exactly one-fifth of a millimetre above the surface. A micrometer scale is engraved upon the object-holder, which obviates the necessity of regulating the microscope in advance. The scale on the object-holder is divided into rectangular spaces one-fourth of a millimetre long by one-fifth broad, representing each one-twentieth of a square millimetre. Each of these is subdivided into twenty little blocks, as shown in the figure (Fig. 71).

FIG. 71.




Blood-mixture as seen with the square micrometer ruling of the moist-chamber of Malassez; magnified 250 diameters.

In order that the cover-glass shall be placed quickly and exactly upon the screw-points and the drop of diluted blood, the cover is attached to a frame moving upon a hinge, which is clamped to the slide. The glass slide is kept perfectly horizontal, and, if it is feared that the object examined will dry, a little water or the blood-mixture may be dropped upon it, so as to surround the circle already mentioned. The number of globules contained in

twenty of the little squares is now to be counted, and if the fluid used be a centesimal dilution it is only necessary to add four ciphers to the number in order to obtain the number in a cubic millimetre, since the large squares represent the ten-thousandth part of this unit. To be exact, several observations should be made. The greatest care is required after each enumeration, in order to insure cleanliness.

FIG. 72.



DALAND'S HÆMATOKRIT.—The central cut represents the complete instrument, with revolving tubes in position. To the left is the tube-holder or frame, with one tube removed so as to show the spring, by which the tube is to be held in place. To the right is a tube containing blood that has been subjected to centrifugal force, indicating ninety per cent. of corpuscular elements, as compared with normal blood.

The use of the hæmacytometer in any form requires skill, patience, and leisure, and even with great care the counts are not accurate, varying between two and four per cent. of error. The

eye-strain in counting the cells in a sufficient number of squares to insure even an approximately correct result is very considerable. In order to obviate this, Hedin,\* in 1890, devised an ingenious apparatus, which he named the *hæmatokrit*, by which the entire mass of the globules in a definite quantity of blood can be rapidly ascertained. The instrument consists of a capillary glass tube, correctly graduated, in which a certain volume of diluted blood is held, while the tube is subjected to centrifugal action, by which the separation of the plasma and the cellular elements is effected. The proportion is determined by the scale engraved upon the side of the glass tube, and the globular richness of the blood is promptly determined. If this be supplemented by a microscopic examination to determine the proportion of the leucocytes to the red cells, and the hæmoglobin also ascertained by the Fleischl hæmometer, the most important clinical data in a case of anæmia are supplied. The original instrument of Hedin has been advantageously modified by Gärtner,† Arnold,‡ and especially Daland.§

The illustration represents the form used by Daland. It is essentially a machine for producing rotation of a metal frame supported upon a spindle and arranged to carry two glass tubes, the outer ends of which fit into small, cup-like depressions, the bottom of which is covered with thin rubber. The glass tubes are held in place by springs at their inner extremities. Each tube measures 50 mm. in length, with a lumen of half a millimetre, and upon it is a scale representing one hundred equal parts; a lens front, by magnifying the column of blood, facilitates the reading of the scale. A single revolution of the large handle causes one hundred and thirty-four revolutions of the frame, so that a speed of ten thousand revolutions in one minute can be readily obtained. Daland instructs us to secure the instrument firmly to a solid table, and to oil it daily when in active use. The method of employing the *hæmatokrit* is simple. To fill the glass tube, a rubber tube is slipped over the end of the capillary pipette, and to the extremity of this rubber tube a mouth-piece

---

\* Scandnavisches Archiv für Physiologie, No. 2, 134; Prager Med. Wochenschrift, 1891.

† Berliner Klinische Wochenschrift, 1893, No. 4.

‡ Medical News, 1894, Sept. 29, p. 348.

§ Transactions of the College of Physicians of Philadelphia, May 2, 1894.

is attached, precisely in the same manner as when the hæmacytometer is used. This glass tube or pipette must be absolutely clean and dry. The finger of the patient is punctured; the blunt point of the pipette is to be placed into the blood, and the tube completely filled by suction. The finger of the operator is then quickly applied to the blunt extremity of the tube, which is next inserted into the frame, and rotated at the rate of ten thousand times per minute. All that remains is to read the percentage volume of blood from the scale. The divisions on the pipette are one-half millimetre apart, so that the scale can be read without difficulty.

The entire procedure need not occupy more than three minutes, and requires no special skill, while the elements of error are so few that the results are trustworthy. In health the volume of red corpuscles is a little over fifty per cent., so that by doubling the number as read from the scale we can get a proportionate expression of the percentage of corpuscles in a specimen as compared with the normal. The amount of hæmoglobin in each corpuscle is approximately obtained by taking the proportion of red corpuscles obtained by the hæmatokrit as the denominator of a fraction whose numerator is the hæmoglobin proportion determined by the hæmometer of Fleischl. The condition of even slight leucocytosis is easily recognized, and when the increase in the white blood-cells rises to thirty thousand or more, the grade of leucocytosis is readily detected. With the hæmatokrit of Daland, when a column of red corpuscles obtained from a healthy man is examined, the white cells present a sharp, clearly-defined, and shallow white band. When the leucocytes are much diminished in number, this white band is imperfect, and in places the red color of the biconcave disks is visible. In the Arnold-Hedin hæmatokrit, diluted blood is used, and the readings are made more accurate by the use of a low-power microscope; the motor runs by electricity.

Exact results cannot be obtained by any method, since the norm of five million red corpuscles is not constant. As shown by Henry, there is a diurnal variation in the number in health, so that all expressions of results of examination are only approximate.

Whatever be the method used in counting the blood-corpuscles,



a number of observations should be always made. Since from any of the instruments but an approximate result can be obtained, the mean of several observations will give us estimates sufficient for all practical purposes. In the several forms of anæmia, it is necessary to obtain a correct knowledge of the state of the blood, not only as to the number of its corpuscles, but also as to the amount of hæmoglobin contained, for in the diagnosis of any disease of the blood it is absolutely required to know the relationship existing between them. Normal blood contains about five million red blood-corpuscles, nearly ten thousand white blood-corpuscles, and two hundred and fifty thousand blood-plaques or hæmato-blasts,\* to the cubic millimetre, and each red corpuscle holds in suspension a certain percentage of hæmoglobin. Any marked variation in the number of corpuscles or in the amount of hæmoglobin is indicative of an abnormal state.

In estimating the number of blood-corpuscles both age and sex must be taken into account. In healthy women the number per cubic millimetre is somewhat less than in healthy men, being about four million five hundred thousand; in new-born infants it often exceeds six million, as both Hayem† and Henry‡ have found by repeated observations. But in the infant the constitution of the blood is remarkable for its variability. The very suggestive observations of John K. Mitchell§ have shown that in adults massage increases enormously for the time being the number of red corpuscles in the blood count.

The white blood-corpuscles in normal healthy blood are in the proportion of about one to six hundred of the red, this varying somewhat in different individuals without being indicative of disease. When the red blood-corpuscles are reduced in number, the proportion of leucocytes is greater, without there being necessarily an increase in the number. The safest method of procedure is to estimate the number of white corpuscles to the cubic millimetre, so that any increase or diminution in their amount will give their true condition irrespective of the change in the number of red disks. The same observation may be applied

---

\* Hayem, Du Sang, Paris, 1889.

† Du Sang et de ses Altérations anatomiques, Paris, 1889.

‡ Amer. Journ. Med. Sci., April, 1890.


§ Transactions of the College of Physicians of Philadelphia, 1893.

to the hæmoglobin, for often this is only relatively diminished, whereas, if the red blood-corpuscles were counted, we should find that each disk had its normal or even an increased amount of hæmoglobin.

The chief apparatuses for *estimating the hæmoglobin* are the hæmoglobinometer of Gowers, Fleischl's hæmometer, Hénocque's hæmatoscope, and the chromocytometer of Bizzozero.

Of these, the hæmometer of Fleischl is the most used. Hénocque's is especially valuable for spectroscopic examination. Gowers's apparatus consists of two glass tubes of exactly the same

FIG. 73.



The hæmoglobinometer of Gowers. A, bottle with pipette-stopper; B, capillary pipette; C, graduated tube; D, tube containing standard tint, fixed in E, a wooden block; F, guarded needle.

size. One contains a standard of the tint, of the dilution of twenty cubic millimetres of blood with one thousand nine hundred and eighty cubic millimetres of water. The second tube is graduated to one hundred degrees, which equal two cubic centimetres. The twenty cubic millimetres of blood are measured by a capillary pipette. This quantity of the blood to be tested is dropped to the bottom of the graduated tube, a few drops of distilled water being first placed in the latter, and the mixture is rapidly agitated, to prevent the coagulation of the blood. The distilled water is then added drop by drop until the tint of the solution is the same

as that of the standard, and the amount of the water added indicates the amount of hæmoglobin.

Fleischl's hæmometer\* consists of a stand to which is attached a reflector made of card-board. On the under surface of the plate there are two grooves, into which slides the frame, holding in position a wedge-shaped glass colored red, the intensity of the hue being graduated from zero to one hundred and twenty degrees. The frame is moved by means of a thumb-screw so that when it is operated the tinted glass passes beneath one of the compartments of the comparing vessel. The horizontal projection of the partition of this vessel should fall directly upon the outer edge of the glass wedge when the instrument is properly adjusted. In operating the instrument, care should be taken to have everything perfectly clean. Accompanying each apparatus, are a glass pipette for dropping the water into the compartments, and several minute capillary tubes for securing the blood.


The compartments—that is, the blood and wedge compartments—are filled almost to the top with distilled water, and the vessel is placed *in situ*. The instrument should then be so arranged and the reflector so adjusted as to secure the full rays of light from either a candle, a lamp, or a gas-flame. Before securing the blood, the tip of the middle finger of the left hand should be carefully cleansed and dried. The automatic blood-pipette, with a capacity of six and a half cubic millimetres, and about eight millimetres long, to which is attached a frail wire for its manipulation, should always be greased, to prevent the blood from adhering to its sides. This is dipped into the blood sideways, to facilitate the flow into the tube: the greatest accuracy is essential to the correctness of the test. With as little delay as possible the tube is then placed into the blood compartment and its contents allowed to escape, aiding by gently moving the tube back and forth along its own axis. The diluted blood remaining in the tube is then washed out by means of the pipette and allowed to flow into the compartment. This is filled, as is the wedge compartment, with distilled water, care being taken not to allow the fluid in the two chambers to run together, and that the upper surface of the water is perfectly level, neither curved nor concave.

---

\* Wiener Med. Jahrbücher, 1885, pp. 425-445: Das Haemometer.

The blood is now ready for examination. In looking at the compartment the eyes should be shaded, that the direct rays of light may not strike in and thus cause error in the observation. The thumb-screw is turned, which slowly moves the wedge from right to left; this movement is continued until the eye can perceive no difference in color between the two compartments:

FIG. 74.



should the difference be imperceptible for a considerable distance, then the point at which the color appears lighter and that at which it appears darker should both be noted and the mean ascertained. The number of degrees—that is, the percentage of hæmoglobin as compared with healthy blood, which is taken as one hundred—will be found on the movable slide.



It is often a matter of great convenience to represent the blood-examinations graphically. An excellent chart for this purpose is in use at the Johns Hopkins Hospital. Fig. 74 shows it, and the manner in which the record is made.

**Anæmia.**—Poverty of blood is met with as a consequence of profuse or frequently-recurring hemorrhages, of insufficient nourishment, of affections which prevent the nutriment taken from being properly absorbed or assimilated, thus impoverishing the blood by depriving it of its most needed constituents, and of profuse chronic discharges, which drain the blood of many of its important elements, and especially of its albumin. Besides these causes of anæmia, we find it occasioned by particular poisons, as by malaria, by syphilis, by uterine complaints, by the retention of noxious ingredients in the blood, or by diseases of certain glands. Again, it is sometimes encountered apparently resulting from constipation, or without our being able to trace it to any obvious source. But under all these circumstances except in the anæmia after hemorrhage, where all the constituents of the blood are diminished together, we have to deal with a watery blood deficient in red corpuscles, and the corpuscles are often badly shaped, and shrunken at their edges. The hæmoglobin may be increased relatively, or it may be diminished, or it may not be materially changed.

Whatever may have given rise to the anæmia, the manifestations of the disorder when well marked are much the same. The patient is weak and pale; his lips and tongue have lost their red color; the eye is pearly; his pulse is feeble, but generally accelerated; the appetite is deficient or depraved; the bowels are apt to be costive. Yet some persons, who are apparently well nourished and are not pale, may have deficiency of red blood cells and of hæmoglobin. Exercise induces great fatigue, shortness of breath, and palpitation; and the disturbance of the heart may be associated with cardiac murmurs or with blowing sounds in the cervical veins, and is sometimes so persistent as to lead to structural changes. In some cases, among the symptoms we meet with obstinate headache and with dropsy, and in many with a persistent pain in the left side, in the region of the spleen.

Anæmia may be owing to the presence of parasites, such as intestinal worms. The very marked form which is common in

Egypt is that due to *anchylostomiasis*. The *anchylostomum duodenale* is taken into the body in the muddy water, or by eating earth containing the embryos of this worm. *Anchylostomiasis* is an insidious, wasting disease, characterized by progressive anæmia and by digestive and nervous deterioration, occurring chiefly in earth and brick laborers of warm climates, caused by the presence in the duodenum and jejunum of a blood-sucking, rhabditic, nematode worm. The blood shows great diminution of red blood-corpuscles and reduction of hæmoglobin.\*

*Chlorosis*.—Here the pallid, wax-like countenance, the very pale lips, and the pearly eye afford unmistakable evidence of the deterioration of the blood, consisting chiefly in great deficiency of hæmoglobin, which is generally much more marked than the reduction in the red corpuscles; these, indeed, may be of almost normal amount. Lloyd Jones† regards the disease as an exaggeration of a change which occurs in the blood of the healthy female at puberty, and which leads to an increase of the amount of blood-plasma associated with a diminution of the amount of hæmoglobin. Van Noorden has shown that the assumption of Bunge that there is increased albuminous decomposition in the intestines in chlorosis is unwarranted by the facts. Meinert has noted gastroptosis in sixty cases of chlorosis, associated with enteroptosis and occasionally movable kidney, to all of which the chlorosis is ascribed, and, as these displacements are produced by wearing corsets, his deduction is that chlorosis is due to faulty wearing-apparel. Clement has found enlargement of the spleen and occasionally phlegmasia alba dolens in chlorosis, which he ascribes to some form of infection. Considerable stress is laid on the fact that in chlorosis there is a greater tendency to inflammation of the optic nerve and retina than in pernicious anæmia, while the tendency to retinal hemorrhage is considerably less.‡

The complaint is especially encountered in young females, and is, as a rule, associated with amenorrhœa. Indeed, many restrict the term to the obvious anæmia combined with suppression of the menses, so often affecting girls about the age of puberty. In

---

\* Sandwich, Proceedings XI. International Med. Congress, Rome, 1894.

† Brit. Med. Journ., July, 1894.

‡ Stephen Mackenzie, Sajous' Annual, 1895, vol. i., L. 10.

pure chlorosis, organic diseases of the gastro-intestinal apparatus of the spleen and lymphatic glands, or of the lungs and kidneys, are absent; the temperature shows a slight rise; the nutrition of the body is fairly well kept up; the urine is pale and abundant, containing but a small amount of phosphates. Forchheimer finds a diminution of urobilin in the urine, which he regards as of considerable diagnostic importance. The nervous system is irritable. Pigmentation about the second joints of the fingers, on their dorsal surface, has been noticed.\* Sometimes these symptoms of chlorosis happen before puberty; or there are relapses of the malady in middle age. Boys about the age of puberty may also develop the manifestations of chlorosis. Virchow has pointed out the frequent association of chlorosis with narrowing of the aorta and of the great arteries, and such cases are distinguished by obstinate relapses. There is a variety of chlorosis in connection with tubercle, at times preceding it. Chlorosis as well as anæmia may be associated with nasal hypertrophies or the *adenoid vegetations* in the vault of the pharynx,† and relieved by their removal. Both the corpuscles and the hæmoglobin may be decidedly decreased in consequence of surgical shock.‡

Fever may occur in chlorosis, though to but slight degree. Jaccoud attributes it to *anoxæmia*, the deficiency of oxygen in the blood acting as a stimulant to the calorific centres. Fever may be also due to local causes, such as *phlegmasia alba dolens*.

**Pernicious Anæmia.**—This is a fatal form of anæmia, which was well known, at least in some of its varieties, to Addison, and which, since the recent researches of Biermer, has actively engaged the attention of the medical world. It is an extreme anæmia advancing steadily, or with remissions, toward a fatal ending; yet no certain cause can be detected for the profound and disastrous alteration the blood is undergoing. To pernicious anæmia belong most of the cases of “essential” or “idiopathic anæmia” which, since the time of Addison, have been reported.

\* Bouchard; also Pouzet.

† F. Oppenheimer, Berl. Klin. Wochenschrift, Oct. 3, 1892; Sajous' Annual of the Universal Med. Sciences, vol. iv., 1894.

‡ Joseph Leidy, Jr., Transactions of the College of Physicians, Phila., 1893, vol. xv. p. 242.



The disorder is most frequent in women, and has been especially observed in child-bearing women after several pregnancies; still, it also often happens in men, especially before the age of forty. It sometimes seems to have its origin in long-continued dyspepsia or diarrhœa, and atrophy of the gastric tubules; or to arise after protracted hemorrhages or incessant worry,—after indeed slowly but steadily-acting debilitating influences; and it has been noted to arise after nervous shock, or to be of parasitic origin, and due to worms, sometimes to a tape-worm,—*bothriocephalus latus*.\* But in the majority of instances it originates seemingly without cause, and, although it has periods of deceptive improvement which may last for months, or, as I have known, even for a year, it progresses relentlessly toward a fatal issue.† It is true that some cases of recovery have been recorded; but of these it is not quite certain that they presented all the characteristic symptoms.

There is an insidious beginning, except at times when the anæmia develops itself in the pregnant state. Pale tongue, bloodless lips, pearly eye, becoming paler, more bloodless, more pearly, from week to week; breathlessness; palpitation of the heart, especially on exertion; weak digestion; constipation, or constipation alternating with diarrhœa; loud systolic murmurs in the heart, and venous hum in the jugulars; vertigo; a marked lemon-colored hue of the skin about the large joints, at times jaundice; finally extreme exhaustion, sluggishness of mind, fainting-fits, and dropsy, without persistent albumin in the urine, or disease of the liver, or enlargement or valvular disease of the heart, to account for it,—are the prominent symptoms. In the later stages, too, hemorrhages from the nose and from the gums are not uncommon; and hemorrhages from the uterus or from the kidneys, or into the skin and into the retina, may also be noticed; the latter especially is very frequent. Yet, notwithstanding all these grave signs, the body appears well nourished; there is certainly no decided emaciation, except in instances in which fever is more than commonly marked. Now, fever is a significant feature of progressive pernicious anæmia; it has been present in every case that I have met

---

\* Schmidt's Jahrb., i., 1891; also *ib.*, No. 10, 1887; and Berl. Klin. Wochens., No. 40, 1886; also Deutsches Arch. für Klin. Med., Bd. xxxix.

† See also case with remissions in Schmidt's Jahrb., No. 4, 1882.



with. It is not an early symptom, belonging to the full development or to the latter part of the disease. It is of very irregular type, and not of high intensity, the temperature rarely exceeding 103° F. It is apt to be continued, or to show occasional exacerbations, followed by remissions, the febrile state lasting for days, or even for a week or two at a time; then there are periods of shorter or longer duration when it wholly disappears, to come on again in an outbreak attended with all the usual signs of a febrile paroxysm for which no cause is apparent. Toward the end of the case it is not unusual for the anæmic fever to have entirely ceased, and for the temperature to have fallen below the normal standard. The disease may run an acute course.\*

The state of the blood in this perilous malady has naturally been made a subject of minute investigation. The red globules are strikingly diminished in number,—to about a million and a half or less; the white corpuscles are not relatively altered, or they may remain normal, and seem to be increased, because the red globules are much fewer. The hæmoglobin is generally increased,† the white corpuscles are normal or diminished, the pale hæmatoblasts are diminished and may quickly assume irregular shapes. The shape of the red corpuscles was stated by Eichhorst to be characteristically changed, in so far at least that the blood contains a quantity of ill-developed, small, spherical, highly-colored red corpuscles. But these are not pathognomonic; for they have been found by Cohnheim in medullary leukæmia, and by Greenfield in lymphadenoma; on the other hand, in a well-marked instance of pernicious anæmia examined by Bradbury‡ they were absent. They are the corpuscles arrested in their growth. Besides this there are giant cells of irregular shape, on which Hayem§ lays great stress, also many very large normal-looking red corpuscles, some of which are, however, nucleated. Nucleated red corpuscles were detected in the blood of all the patients examined by Howard:¶ the blood seems to revert to a lower type. This has been also insisted upon by Henry.¶ A

---

\* Lantener, *Rev. Méd.*, Louvain, 1883, ii.

† Hayem, *Du Sang*, Paris, 1889.

‡ *British Medical Journal*, Aug. 14, 1880.

§ *Op. cit.*

¶ *Montreal General Hospital Reports*, vol. i., 1880.

¶ *Anæmia*, P. Blakiston & Co., 1887, p. 116.

much larger proportion than is found normally of small disks of deep color, is regarded as important by Pye Smith.\* The accompanying cut (Fig. 75), from a well-marked instance of the

FIG. 75.



Blood in pernicious anæmia, illustrating the irregularly-shaped blood cells (Poikilocytosis).

disease, shows the irregular shape of the corpuscles and their varied size and appearance ; some are nucleated.

Of the real cause of the disease we are in ignorance. No constant lesion of the blood-making glands has been found. The structure of the spleen and of the lymphatic glands is not altered ; the marrow of the bones may or may not be,† though cases in which it is are thought to be instances of myelogenous pseudo-leukæmia rather than of pernicious anæmia. Perhaps the most constant lesion is fatty degeneration of the heart, often associated with the same change in the inner coat of the large arteries. Hunter‡ brought forward strong proof that the characteristic

\* Guy's Hospital Reports, xxvi., 3d Series, 1883.

† Pepper, Amer. Journ. Med. Sci., Oct. 1875 ; see also Cohnheim, Virchow's Archiv, Bd. lxxviii., and Waldstein, Arch. f. Path. Anat., Berlin, 1883, xci.

‡ Lancet, London Practitioner, Aug. 1888.

anatomical change is the presence of an excess of iron in the liver, the seat of disintegration of the corpuscles being chiefly in the portal circulation.

Biruli has found, by microscopic examination, hemorrhages in the substance of the cerebral hemispheres, also round structureless bodies resembling corpora amylacea, and fatty degeneration of motor cells, with shrinking and vacuolation of the cells of Purkinje. The existence of some toxic substance in the circulation is highly probable. It can scarcely be doubted that diminished or faulty hæmogenesis may also exist and contribute to the anæmia, as insisted upon by Van Noorden.\*

Lichtheim,† in 1887, directed attention to alterations occurring in the *spinal cord*, especially in the form of *degeneration in the posterior columns*, with slight changes in the lateral and anterior columns, apparently caused by some toxic substance in the blood; Van Noorden‡ reported a case of progressive and apparently causeless anæmia in a woman fifty-nine years of age. The red blood-corpuscles numbered seven hundred thousand, with relative increase but absolute decrease of leucocytes; some small lymphocytes were present, and irregularly-shaped blood-cells (poikilocytes). Death occurred ten months after the onset of the affection. Microscopic sections showed distinct degeneration of the lateral pyramidal and lateral cerebellar tracts, and of the posterior columns. Bowman§ regards the anæmia as primary and probably the cause of the change in the cord. Recently, Charles W. Burr|| has made a careful study of the spinal cord lesions and symptoms of pernicious anæmia. He concludes that the coexistence of pernicious anæmia and of certain lesions of the cord is not accidental, and that in well-marked anæmia the latter changes are constant. The cervical swelling is the principal site of these degenerative changes. The only parts of the cord ever affected are the posterior columns, the lateral columns in and near the crossed pyramidal tracts, and, rarely, the direct cerebellar tracts. The gray matter is seldom even slightly involved. The

---

\* Quoted in Sajous' Annual for 1895, vol. i. L. 8.

† Congress für Innere Medizin.

‡ Charité Annalen, 1891.

§ Brain, 1894.

|| University Medical Magazine, April, 1895.

degeneration of posterior peripheral roots is ordinarily insignificant. There is usually marked posterior, with less marked lateral, degeneration. Almost invariably a band of normal white tissue separates the diseased area in the posterior columns from the gray matter. Burr favors the view that both the anæmia and the cordal lesions are due to a common cause,—a poison or poisons, as in diphtheria or ergotism.

The diagnosis of pernicious anæmia is never an easy one, for the reason that it is difficult to be quite certain that no obscure and latent disease exists which would account for the exhaustion and the progressive impoverishment of the blood. Indeed, it is only after the most careful and repeated examinations of all the organs of the body and the most searching inquiry into the history of the case that we are justified in making the diagnosis of pernicious anæmia. I have more than once known ill-developed *organic disease of the stomach*, especially gastric cancer, where the tumor could not be discerned, or *contracted kidney*, with but little albumin in the urine, regarded as a typical illustration of the malady, until the autopsy revealed the true cause of the fatal exhaustion. With reference to the former affection the error is all the more likely to happen because symptoms of gastric disorder are not unusual in progressive anæmia; with reference to disease of the kidney the misleading part is that a trace of albumin is occasionally present in progressive anæmia. But it is not persistent; and microscopical examination of the urine will tell us the real amount of kidney affection. The cachectic pallor of subjects of *malignant disease* may be mistaken for the straw-colored appearance of the skin in pernicious anæmia.

*Diseases of the heart* may be mistaken for pernicious anæmia. A fatty heart, in an elderly person, with or without valvular disease, with failure of strength, and with the peculiar pallid, sickly look occasioned by the malady, may mislead. But the long duration of such cases, and the absence of fever, are strong points in the case. Indeed, the error is apt to be the other way,—that, overlooking the symptoms of profound anæmia and general failure, we regard the murmurs and the other cardiac symptoms which are associated with the fatty heart of pernicious anæmia as pointing to a disease of the heart alone. The physical signs will not always assist: the murmurs may be very distinct and loud.



A number of trophic and vascular disturbances have followed surgical extirpation of the thyroid gland, and have also been noticed after atrophy of the gland has occurred. To this condition the name of *cachexia strumipriva* has been given. It is distinguished from anæmia by the occurrence of signs of myxœdema often with cretinism and circulatory disturbances, with local asphyxia and transient or intermittent albuminuria. In some cases epilepsy is developed, in others pulmonary phthisis.

If we have excluded any organic disease that could account for the anæmia, we turn to the diseases of the blood itself to obtain an explanation of the symptoms. And here we find first that pernicious anæmia differs from *ordinary anæmia* by the absence of the history of the causes that commonly give rise to the anæmic state, such as acute diseases, malaria, tubercular or cancerous cachexia, loss of blood, and the like, but above all by its relentless course and the little influence the most nourishing diet and courses of iron have on it. Moreover, the distinctness of the cardiac murmurs, the slight emaciation, and the irregular outbreaks of fever are significant. The marked accessions of fever, the presence of dropsy, though moderate, the retinal extravasations, the other hemorrhagic symptoms, and the unyielding blood-change, separate pernicious anæmia from the *chlorosis* so common at the age of puberty in girls. The pernicious malady sometimes seems to develop out of a long-standing chlorosis, and then the grave symptoms just spoken of prove its supervention. The same grave symptoms happen also, at least the hemorrhages are as frequent, and the fever and dropsy may happen, in *leukæmia* and in *pseudo-leukæmia*. But the great increase in the white corpuscles, the tumefaction of the spleen, or the affections of other blood-making parts, distinguish the former malady; and pseudo-leukæmia, while the blood microscopically will not differ materially, exhibits the enlarged lymphatic glands, their progressive invasion, the lymphoid tumors, the abdominal pains, and the steadily-increasing emaciation so characteristic of the disease.

There are other forms of idiopathic anæmia of which we cannot clearly recognize the cause, that we shall probably soon be able to separate into groups. But for the present we have to admit that cases may occur which cannot be classified.

**Leukæmia.**—This morbid state consists in a decided increase

of the white corpuscles and a decrease of the red. Under the microscope the white globules of the blood, instead of bearing the normal proportion of about 1 to 500 of the red, are found in the proportion of 1 to 6, or even of 1 to 0.5, and cases have been met with in which near the point of death the white corpuscles have exceeded the red as high as five times. Besides the increase of white corpuscles and the diminution of the red, peculiar, colorless, shining, elongated octahedral crystals have been pointed out by Neumann and by Charcot. Haig has stated that the proportion of uric acid in the blood is increased in splenic leukæmia. Jaksch \* has shown that the blood is rich in peptone, although this substance is rarely met with in the urine in leukæmia.

The abnormal condition exists in connection with hypertrophy of the spleen, "splenic leukæmia," or of the liver, with other diseases of this viscera, and with various malignant or non-malignant affections of the lymphatic glands, "lymphatic leukæmia," or of the thyroid body, especially with an increase of the cellular elements. But none of the blood-glands is so constantly and so markedly affected as the spleen. Owing at times to a large production of lymphoid cells in the marrow of the bones, there is produced also a "myelogenous" or medullary form of leukæmia.

The disorder may occur at all ages ; it is more common in men than in women. Leukæmia is consequent upon obstinate intermittents with decided enlargement of the spleen, syphilis, over-exertion, long-continued mental depression, chronic intestinal catarrh, and blows on the splenic region. The form affecting the marrow of the bones frequently results from injury to the bones. Ebstein reported cases of leukæmia following traumatism, but the causative relationship is not clearly made out. But in many cases of leukæmia no adequate cause can be detected. Its beginning is usually gradual and ill defined ; sometimes it clearly follows other diseases. When fully developed, it often occasions, besides the obvious pallor and the cachectic appearance, exhaustion, diarrhœa, extremely hurried breathing, hemorrhages from various parts, especially from the nose, profuse sweating, slight rise of temperature in the evening, increase of uric acid in the urine, fleeting abdominal pains, and dropsy dependent upon the enlarge-

---

\* Wiener Med. Presse, Oct. 1882.

ment of the spleen or of the liver or upon the leukæmic new formations in the latter. In some cases a swelling of the glands on both sides of the throat, attended with inflammation of the mucous membrane of the mouth and the pharynx, and followed by swelling of the axillary and the inguinal glands, precedes the enlargement of the liver and of the spleen.\* Indeed, glandular tumors are often present; the glands of the groin are, as a rule, enlarged. There is disturbance of vision, connected with retinal changes, also melancholy, and in some instances deafness, and peritoneal or pleural inflammations. Pain in the bones, too, particularly in the sternum, is observed. The medullary or myelogenous variety is especially marked by pain, which is increased or developed by pressure over the sternum and ribs and over other affected bones.†

The diagnosis of leukæmia is possible only by the microscopical examination of the blood, which detects the decided increase of the white corpuscles. Mathes has examined the blood in leukæmia to determine the existence of peptone and allied bodies. Peptone was not found in the blood, but there were deutero-albumoses in the blood and serum. In the serum of one case there was also nucleo-albumin. This probably resulted from destruction of blood-corpuscles. The excretion of uric acid was only moderately increased. In the most common variety, *splenic* leukæmia, we may be also able even early to discern the enlargement of the spleen, and to find the evidences of cachexia in the appearance of the patient, and in recurring epistaxis. But it is the microscopical examination of the blood alone which enables us to distinguish leukæmic swelling of the spleen from its other affections. And to have a definite diagnostic meaning the white corpuscles must be decidedly and permanently increased; for a mere transitory, slight increase may occur in other diseases of the spleen. Some corpuscles are larger, some smaller, than normal, and many show fatty changes; but in splenic leukæmia the white corpuscles are mostly large. In both varieties the red corpuscles are badly shaped. *Lymphatic* leukæmia is chiefly recognized by

---

\* Mosler, in Virchow's Archiv, xliii.; Dunn, Amer. Journ. Med. Sci., March, 1894, describes a case with growths in the orbits.

† Mosler, Berlin. Klin. Wochenschrift, xiii., 1876; and Schmidt's Jahrb., No. 10, 1877.



the marked swelling of the lymphatic glands, while the spleen is less obviously affected. In the blood the white corpuscles derived from the lymphatic glands are smaller than those coming from the spleen, and have a well-developed nucleus. Yet it is very difficult to judge a case by these traits. Large round corpuscles containing granules which by ether and chloroform are found to be fatty are stated to be derived from the marrow of the bones, and, if abundant, to bespeak *medullary* leukæmia.\* Karyokinesis has been noted in some red corpuscles, as well as free nuclei in the blood. Hayem† found that nucleated red corpuscles were habitually present, and that the very large white cells were destitute of amœboid movement.

*Hypo-leukæmia*.—A condition in which there is an actual or seeming diminution in the proportion of leucocytes has been termed *hypo-leukæmia* by Héricourt and Richet. Maurel has more recently shown that the diminished number of leucocytes is not real, but is due to their arrest and incarceration in the finer capillaries. He therefore suggests that the name *false hypo-leukæmia* is more appropriate. This arrest of the white cells may be due either to vaso-motor constriction of the small capillaries or to direct action of poisonous agents upon the leucocytes, giving them a spherical shape and sluggish amœboid movement. Goldscheider and Jacob express the same view based upon experimental work which seems to indicate the pulmonary capillaries as the point of arrest of the leucocytes. Temporary *hypo-leukæmia* results from injections of plain bouillon, as well as of various organic extracts, and occurs during the course of many of the infectious fevers, as pointed out by Everard and Demoor.‡

*Lymphadenoma*.—As regards the symptoms, the closest similarity to leukæmia is presented by the affection described as lymphadenoma, pseudo-leukæmia, or *Hodgkin's disease*. It consists in an enlargement of the lymphatic glands of the body, often with lymphoid growths in other parts, which soon becomes complicated with extreme anæmia, with weakness and signs of cachexia, with diarrhœa, with dropsy, with cardiac palpitation, shortness of breath, and attacks of suffocation, with tendency to profuse bleed-

---

\* Schmidt's Jahrb., No. 10, 1877.

† Op. cit.

‡ Quoted, Diseases of the Blood and Spleen, in Sajous' Annual for 1895.



ings and to bed-sores, and leads usually in the course of not many months, or, at farthest, of a few years, to death. There is often a sense of fulness in the abdomen, attended with violent pains; the temperature in advanced cases shows mostly an evening rise. Some of the superficial lymphatics are first affected, others follow; the disorder then extends more decidedly, the spleen and the liver increase in size, other organs, too, may become involved, and lymphoid tumors develop in various parts of the body; but among the internal organs the spleen is the one most constantly disturbed.

The disease generally begins in the cervical glands; far less frequently does it show itself first in the inguinal or in the axillary glands; still less frequently in the bronchial or in other internal glands. The affection occurs much oftener in men than in women. It mostly happens in males between the ages of ten and thirty-five and of fifty and sixty; it is not very uncommon in young children. Its cause is unknown; it certainly has no definite connection with either scrofula or syphilis.

The chief anatomical lesion is found to be an augmented formation of the structure of the glands. The spleen is either simply hypertrophied or is the seat of numerous disseminated lymphoid growths; in neither case is it apt to attain to any very great size. At times the follicles at the base of the tongue, in the tonsils, and in the intestines share in the morbid process; changes in the bone-marrow are rare. The blood shows deficiency in red globules, but otherwise no constant alteration. Slight increase of leucocytes has been occasionally noticed, especially during the later stages; but even then the white corpuscles are small.

It is this difference in the state of the blood that makes the chief difference between pseudo-leukæmia and *leukæmia*, in which there may be glandular enlargements. Further, leukæmia is a disease, as a rule, of longer duration, and the splenic enlargement is generally much more marked. Rare cases of diffused *lymphatic cancer* closely resemble Hodgkin's disease; so closely that they are undistinguishable, except by the history of the case and by a microscopical examination of any of the tumors that may have been removed; the spleen is not involved, while the organs contiguous to the glandular cancer are likely to be more rapidly implicated. In *sarcoma of the lymphatic glands* the disease is at first

strictly local, and then, if it spread, invades not the lymphatic tissues specially, but any part of the body ; the enlarged glands do not move freely on each other as they do in lymphadenoma. *Local gland lymphomas* are separated from Hodgkin's disease by their local character, by their want of extension, and by the absence of marked cachexia. *Scrofulous* or *tuberculous glands*, unlike lymphadenoma, enlarge rapidly, have thickened tissue around them, and are apt to undergo cheesy degeneration, or to soften and suppurate. Moreover, they are associated with the presence of tubercle-bacilli, and mostly affect the submaxillary glands. The anterior cervical glands are the ones chiefly and primarily affected in Hodgkin's disease. In some cases of Hodgkin's disease fever is a prominent symptom, and this may be of intermittent type, giving rise to the belief that we are dealing with a *malarial affection* ; recurring chills make error still more likely.

In the early stages of lymphadenoma a diagnosis is impossible, and we are at a loss to account for the increasing signs of cachexia, until the involvement of the lymphatic glands in rapid succession, and their quick growth, or the speedy formation of other lymphoid tumors under the skin or in other parts of the body, clear up all doubt. There will also be great uncertainty in all those instances in which the growths happen first in internal glands or structures,—as in the bronchial glands and the mediastinum, producing severe bronchitis, extreme dyspnoea, and signs of venous stagnation in the veins of the upper part of the body ; or as in the glands around the biliary ducts, giving rise to jaundice ; or as in growths in the spinal cord leading to paraplegia,—until the external swellings explain the case. The kidney is not an organ that often suffers primarily ; the occurrence of more than a mere trace of albumin shows that it has become implicated from parenchymatous changes or disseminate lymphoid growths.

There is a strong probability that in the future we shall find that there are no primary anæmias, and that all forms will ultimately come to be recognized as secondary and be traced to the lesion that determines them. The diagnosis will be directed rather to the affections with which they are associated than to the special types of the anæmia. The same will be the case with the lymph-

adenopathies, which clinically manifest themselves as leukaemia, Hodgkin's disease, or purpura hæmorrhagica.

**Addison's Disease.**—While seeking for the explanation of puzzling cases of anæmia, Addison discovered that a peculiar anæmia always occurs in connection with a diseased condition of the supra-renal capsules, and is characterized by distressing languor and great general prostration, remarkable feebleness of the heart's action, loss of appetite, obstinate vomiting, and a singular alteration of the skin. This consists in a dingy or smoky hue of the surface; or the color may be of a deep amber or chestnut brown, or the altered skin may have a bronzed tinge. The change of color begins on exposed parts, such as the face and neck and the back of the hands, and deepens first there; but we also soon find it marked in parts which are naturally the seat of much pigment, such as the axillæ, the groins, and the areolæ of the nipples. It is also marked around the umbilicus, on the penis, and on the scrotum, and is dependent upon a layer of pigment in the rete mucosum. The skin remains soft and smooth, and becomes in large portions uniformly discolored, gradually deepening, and often presenting a hue on the face and hands like that of a mulatto. Any irritation of the skin is followed by dark streaks. Discoloration in patches is both less constant and less significant than extensive alteration of hue; yet the darkening in undoubted cases may occur in patches, which are usually most obvious on the face or the superior extremities. The patient may seem at first sight to be jaundiced; but the pearly whiteness of the conjunctiva soon dispels such an idea. The nails are pale and bluish; the tongue may have patches of dark color; the body and breath at times exhale an offensive odor; and the blood has been found to contain an excess of white corpuscles and a slight decrease of the red, although it generally does not undergo any important or characteristic change.\*

With reference to the other symptoms, the most conclusive of them are remarkable prostration, generally without any marked waste of the body, feebleness of heart's action and of pulse, and obvious anæmia. In most cases, but far from in all, these symptoms precede the discoloration of the skin; and they are not

---

\* Greenhow, Addison's Disease.

unfrequently associated with pain in the back and with gastro-intestinal irritation, with breathlessness upon exertion, with vertigo, and with dimness of sight or impaired hearing. A peculiar odor of the body, like that perceived in the colored race, was observed in two cases placed on record by Mr. Hutchinson. In the last stages of the malady the temperature falls below the norm.

Death may take place gradually from the constantly-growing asthenia; or it may occur suddenly, and where the amount of prostration does not appear so excessive as to foreshadow it. According to the elaborate researches of Wilks, the destruction of the capsules is dependent upon a peculiar scrofulous degeneration; and this view of the tubercular nature of Addison's disease is now very generally held.\* Should this prove to be correct,—should it appear, in other words, that the nature of the disease of the adrenals influences the symptoms more than the mere fact of their being diseased,—it would explain why in some cases of absence of the glands, or of their cancerous degeneration or supuration, no signs of Addison's disease existed. With reference to the nature of the affection, however, tuberculous disease of the adrenals has been found without bronzing;† and tubercle-bacilli have been detected in the caseous glands. Many of the symptoms of the fully-developed malady may be due to the implication of the nervous branches, derived from the sympathetic and the pneumogastric, which go to the glands. Indeed, the idea of the primary seat of the disease in the abdominal sympathetic nerve is strongly advocated by some observers.

Now, in the diagnosis of Addison's disease the alteration of the color of the skin plays so important a part that we must inquire whether it or something very like it may not happen in other conditions. In persons *long exposed to the sun* a bronzing of the face and neck and arms occurs; but it is extremely uniform; there is a striking contrast between it and the parts that are not exposed, including such as we find greatly affected in Addison's disease, the flexures of the joints, the scrotum, the textures around the nipple and the umbilicus. Moreover, there is often robust

---

\* See, for analysis of cases, Gilman Thompson, Transact. of Assoc. of Amer. Physicians, 1893.

† As in the case of Ballenghien, Journ. des Sci. Méd. de Lille, 1888.



rather than impaired health. In persons who, in addition to exposure, are of uncleanly habits and infested with vermin, especially in elderly persons, a discoloration of the skin happens at various portions of the body, often deepest on the chest, the abdomen, and the back, which is readily mistaken for the bronzing of Addison's disease. But in this *vagrants' disease* the discoloration is in the superficial, not in the deeper layers of the epidermis, and the dark cuticle is harsh and raised, not soft and smooth. Then alkaline baths and washing with soap will greatly diminish the deepened hue. A similar bronzing of long standing, though of doubtful origin, is sometimes met with.\*

During *exhausting lactation*, or in *pregnancies* attended with much constitutional disturbance, there may be marked discoloration of the skin; yet it is not most obvious on the face, and the circumstances of the case are important aids in the diagnosis. So is the history in those instances in which a bronze hue is *hereditary*,† or in which a very deceptive discoloration follows *yellow fever*, or the *malarial fevers*, or *chronic disorders of the liver*. In these diseases, too, the discoloration is not so great, and it is not marked at the sites most affected in Addison's disease. Greenhow has pointed out how certain very long standing instances of *phthisis* exhibit an appearance exactly like that of the earlier stages of Addison's disease. Yet the abnormal pigmentation does not deepen or increase, and the symptoms remain only those of the pulmonary malady. Stains on the skin from *pityriasis versicolor* or from *syphilis* have not the characteristic seats of Addison's disease, and they are in patches and surrounded by healthy skin, and certainly the syphilitic affection coexists with other significant eruptions or signs. Malcolm Morris‡ has called attention to the mistake of pronouncing a case of *acanthosis nigricans* one of Addison's disease. The fact that the processes in the former are confined to the upper layer of the skin, and characterized by an abnormal development of the younger not yet cornified elements of the upper layers,—the so called prickle-layer,—will serve to separate it pathologically from the secondary and relatively unimportant changes of the skin that attend the

\* Crocker, Transact. Clin. Soc. Lond., vol. xiv., 1881; also Carrington, *ib.*

† Medical Times and Gazette, May, 1871.

‡ Medico-Chirurgical Transactions, 1894.

course of the latter affection. A chocolate-colored discoloration of the whole surface of the body has been observed in a case of psoriasis in which the patient continued to take arsenic during a period of two and a half years.\*

One of the confusing points connected with the diagnosis of Addison's disease is that cases occur without bronzing, or with the discoloration of the skin so slight as to be a matter of doubt. Such cases are generally in persons who die before they have had the disease any length of time. If the altered hue of the skin be wanting, the complaint is undistinguishable from *pernicious anæmia*, though we may lay some stress on the comparative absence of febrile phenomena. *Other diseases of the suprarenal capsules*, such as cancer and waxy disease, are also not to be separated from the peculiar affection of the gland occasioning Addison's disease, if bronzing of the skin be not present.

The malady, as Greenhow proves, is very rare except in persons employed in manual labor. In some instances it seems to arise from grief or protracted anxiety. The disorder is a chronic one, generally lasting for years; but it almost invariably destroys life. Yet cases have been recorded in which most of the symptoms of Addison's disease existed and which recovered; and certainly long remissions in the symptoms have been not unfrequently observed, and in these remissions the discolored skin has lightened. The disease is occasionally met with in young persons. Dyson reports a fatal case in a girl thirteen years of age.†

**Pyæmia.**—Purulent contamination of the blood is an affection much more likely to be met with by the surgeon than by the physician; yet the physician must be familiar with its symptoms. These are, great depression of the vital powers, profuse sweats, rapid pulse, and the formation of purulent deposits in different portions of the body. The symptoms may be of gradual development; but often they set in suddenly with a chill, to which a fever of low type soon succeeds; or the shivering is followed by copious sweating, and the febrile phenomena subsequently appear. A transient erythematous blush on the skin is not unusual.

The pyæmic fever rarely lasts longer than a week, and during

---

\* Carrier, Medical News, Feb. 3, 1894, p. 127.

† Quar. Med. Journ., vol. iii., Part I.

its continuance it usually presents the most marked variations in temperature. Yet the disease is not always alike in this respect ; for we find, as Heubner has proved, not only cases in which the most decided increase of heat is constantly followed by an equally decided decrease, but also cases in which there are febrile attacks followed by marked intervals during which the temperature is almost normal, and cases in which continuous fever exists with striking intercurrent rises in temperature.\* Still, in all the maximum temperature is apt to be very high, ranging from  $106^{\circ}$  to  $108^{\circ}$ .

The disorder may arise after injuries and operations ; or where sinuses or abscesses exist that have no free vent for the pus ; or in consequence of the contamination of the blood which happens in phlebitis or arteritis ; or in inflammation of the external coat of arteries, with suppuration, especially in the periarteritis of the thoracic aorta ; or in ulcerative endocarditis ; or the pyæmia results from the breaking down of coagula in the blood-vessels ; or it may supervene upon diffuse cellular inflammations, or upon puerperal fever : in fact, it will be found under many dissimilar circumstances. Micro-organisms play an important part in its production, especially the several varieties of the streptococcus pyogenes and the staphylococci. They render the pus septic, and, under conditions favorable to their development, diffuse the process. This they do by producing a suppurative infecting phlebitis or by inducing coagulation of the blood and disintegrating thrombi. But, without stopping to explain the sources of origin of pyæmia, let us look at its diagnostic traits.

Now, there are several complaints with which pyæmia is likely to be confounded, the chief of which are typhoid fever, rheumatism, acute glanders and færy, and acute affections of the liver.

It is liable to be mistaken for *typhoid fever*, on account of the adynamic character of the fever, and, it may be, the occurrence of diarrhœa and of cerebral symptoms. But the history of the case is very dissimilar : there is no eruption, or, if there be an eruption, it consists, as Bristowe so particularly points out, of sudamina surrounded by a zone of congestion, and is therefore not the eruption of the typh-fevers ; and, on the other hand, we find in typhoid fever neither the profuse sweating nor secondary

---

\* Archiv der Heilkunde, ix., 1868.



deposits of pus, and the thermometry of the disease is very different. Pyæmia may, however, happen as a complication of the febrile malady.

The pain in the joints and their swelling in succession, the fever, and the perspirations, resemble much at times *rheumatic fever*. But the difference consists in the greater severity of the constitutional phenomena caused by the poisoned blood, in the marked exhaustion, in the rigors, and in the history not being that of acute rheumatism. Moreover, the frequent signs of formation of abscesses in internal organs or around the joints, the development of pustules on the skin, and the striking redness of the tumid joints assist materially in the diagnosis.

*Acute glanders or acute farcy* is a disease scarcely distinguishable from pyæmia, since it occasions, for the most part, the same manifestations. The knowledge that the patient who has apparently pyæmic symptoms has been working among horses, the ulceration of the mucous membrane of the nose, and the fetid discharge proceeding from it, which occurs in acute glanders, and which is apt to be associated with nasal hemorrhages, with an offensive breath, with enlargement of the lymphatic glands in the vicinity of the affected mucous membrane, and with hurried breathing, or sometimes with gangrene of various parts, afford us the only means of discrimination. Then we find a peculiar tuberculated or pustular eruption, resembling smallpox, upon the skin, and in farcy the lymphatic glands and vessels specially suffer. But more significant than all is the distinct history of the contagion; for the grave coryza does not happen in all forms of equinia,—certainly not in farcy.

*Acute affections of the liver* resemble pyæmia on account of the jaundice that may attend the latter disorder; the history of the case, the rigors, the sweats, and the purulent deposits distinguish it. Yet it must be remembered that suppurative inflammation of the portal veins and metastatic abscesses of the liver happen.

In conclusion, let us inquire where and how the secondary deposits, or *metastatic* or *embolic abscesses*, are formed. They may take place in the parenchymatous organs, particularly in the lungs and the liver; in the synovial sacs, in muscles, or in areolar tissue, especially in that under the skin. The secondary deposits are mostly due to fragments of septic thrombi filled with pus



organisms which have become impacted in the vessels of distant parts and there become centres of suppurative change and of fresh infection.

If the altered blood coagulate in the arteries, or if from disintegration of fibrin in the arterial system the infected masses occasion deposits in solid organs, as in the liver or the spleen, we may have, with the similar pathological states, symptoms arising like those of ordinary pyæmia. Indeed, in the *arterial pyæmia*, as it has been called, rigors, febrile symptoms and sweating, and pains in the joints are observable. In connection with the obscure febrile condition, the liver and the spleen are often observed to increase in size slowly.\* The heart may or may not be affected; ulcerative endocarditis is often present. Hayem has pointed out that there may be capillary embolism in pyæmia, not to be recognized except by the microscope. It may be one of the causes of the so-called *idiopathic pyæmia* in which the source of infection is not apparent.

There is a form of pyæmia, called by Leube† *spontaneous septico-pyæmia*, which comes on without obvious cause, or is perhaps preceded by a fall or a slight skin wound, in which the symptoms of pyæmia become developed with pain and tenderness in joints and muscles, ecchymosis of the conjunctiva, vesicles in the skin containing blood, extremely high temperature, swelling of the spleen, albuminous urine, pleurisy or perhaps signs of endocarditis or pericarditis, stupor, delirium, cramps, and finally involuntary discharges and coma. The disease, resembling typhus or ulcerative endocarditis, is to be distinguished only by the general association of the symptoms.

The description of pyæmia given represents it as an acute affection, and so it almost always is. Yet there are cases much slower in their course, and extending over months. These *chronic* or *relapsing* instances of the disease have been described by Paget.‡ The symptoms presented are the same as in the acute disorder; but the local evidences of the complaint are more often seated in different parts of the same tissues, and less frequently in internal organs. The malady is not nearly so perilous as the acute disease.

---

\* Samuel Wilks, *Guy's Hospital Reports*, vol. xv., 3d Series.

† *Archiv für Klin. Med.*, xxii., 1878.

‡ *St. Bartholomew's Hospital Reports*, vol. i.

**Septicæmia.**—This is a poisoned state of the blood, produced especially by animal poisons, such as the bites of venomous serpents, or the absorption of putrid matters that have been generated in the economy, or by their inoculation. It may be seen after injuries and wounds, or in the puerperal state. The continued exposure to the breathing of foul air and of septic gases will also occasion septicæmia. There are no discoverable foci of suppuration, but the bacteria occasioning the sepsis are in the main the same as those of pyæmia. Toxines and ptomaines have much to do with the process.

The symptoms of the blood-poisoning vary somewhat with the individual poison that has occasioned it. They are, as a rule, the symptoms of pyæmia, except that secondary pus-formations belong to the former rather than to the latter; and the same may be said of embolism and its results. Rigors are frequently observed. In many instances the altered condition of the blood leads to great prostration, to hemorrhages from internal organs, to petechiæ, to delirium and coma, to extreme rapidity of pulse, to rapidly-developed fever with high temperature, to enlargement of the spleen, to cough and bronchial catarrh, and to gastric and intestinal disorders. The blood shows the white corpuscles almost always in marked excess, although not altered in character as they are apt to be in leukæmia; the red globules are diminished.\* The bacterial types characteristic of the form of septicæmia are generally demonstrable by microscopic examination and by culture experiments in the bacteriological laboratory.

*Malarial Septicæmia.*—Since the discovery of Laveran that malaria is due to the presence in the blood of a micro-organism, the *plasmodium malarix*, this well-known disorder has entered the list of infectious diseases and become recognized as a form of septicæmia. Klebs asserts that he has found flagellate protozoa in the febrile stage of *influenza*, but this observation has not been confirmed.

*Typhoid Septicæmia.*—Several observers, especially in Italy,† have reported cases of typhoid septicæmia without the accustomed

---

\* See the valuable report of the Committee of the Pathological Society of London, Transactions, 1879.

† See Guido Bariti, *Riforma Medica*, 1887.

alterations in the intestinal tract. The diagnosis was based upon the bacteriological examination of the organs and the characters of the bacilli. Dogliotti also reported a case which had fever, enlarged spleen, and copious eruption of typhoid roseola extending over the entire body. There was profuse diarrhœa. Cultures of blood from the finger and from a vein in the arm remained sterile; but cultures taken from the blood of the papules developed bacilli which corresponded with all the characters of the typhoid-bacilli. At the post-mortem examination no ulcers or cicatrices were found in the intestine. The conclusion is that, besides the familiar form of abdominal typhoid, there is another, identical in every way except that it has no intestinal or lymphatic localization. The bacilli select the skin in preference to the mucous membrane or the lymph cells. Septicæmic typhoid then presents the following characters: an irregular fever, not typical as that of ileo-typhoid; the absence through the entire illness of visceral complications and of symptoms pertaining to the digestive system; the presence in the circulating blood, and in the blood extracted from the rash, of a bacillus presenting the characters of the typhoid-bacillus.\*

Pyæmia and septicæmia have shifted much from their old significance. We know much more of direct infection and how foci of suppuration are set up in various parts of the body. Another set of symptoms is occasioned when the products of the micro-organisms only, the so-called *toxines*, are absorbed by the blood; here fever, prostration, and various nervous phenomena are caused. This condition has been termed *sapremia*, to distinguish it from septicæmia, in which the infective agent is actually present in the blood. Occasionally pathogenic micrococci may be present in the body without giving rise to either suppuration or septic disease. The tissue-cells, especially those of the spleen and the kidney, play a very important part in the destruction, and the leucocytes are also active in the warfare.

**Thrombosis and Embolism.**—Although in connection with endocarditis, with obstruction of the cerebral arteries, and with diseases of the kidney, the phenomena of embolism have been de-

---

\* Translation in the Pacific Medical Journal, vol. xxxviii. p. 203; also Dogliotti, Gaz. Med. di Torino, 1894.

scribed, it may serve a useful purpose to view here connectedly, though chiefly in their diagnostic bearing, some of the results of the formation of the clots in large vessels or in the heart, and of their being carried along with the current of the blood and driven into remote vessels,—the results, therefore, of thrombosis and of embolism. Of these embolism is the subject which more particularly concerns the physician in its immediate practical bearing.

The embolus may produce manifestations in the venous system, either in the peripheral veins, or in the venous trunks of the great internal cavities of the body; or a portion of the clot may have been washed into the pulmonary artery from the right side of the heart; or it may have become impacted in the arteries of the general circulation, in the larger arteries, or in those of fine calibre; or it may have been propelled into the very structure of organs through these arteries, as into the liver-structure through the hepatic artery, into the splenic parenchyma through the splenic artery. Let us examine a little more closely some of the symptoms thus occasioned, premising that arterial embolism is of much more frequent occurrence than the other forms.

In the *veins* thrombi may form, which, so long as they do not produce obstruction of the canal, give rise to no marked signs. A slight hardening and pain on pressure if the coagulum be in one of the more superficial veins, their enlargement if the clot be in a deeper vein, are apt to be the only evidences of the disordered condition. But when the occlusion is considerable, and especially when the collateral circulation is insufficient, œdema is developed, which may be attended with very great tenderness of the swollen part, and, if the impediment be of long duration, with changes in the nutrition of the structures sufficient to produce phlegmonous inflammation. These phenomena are encountered in phlebitis, or milk-leg, *phlegmasia alba dolens*. This condition, occurring after parturition, is now regarded as an evidence of puerperal septicæmia due to infection of the system through the uterus. In some cases profuse hemorrhages happen as a consequence of the stoppage in the vein,—as cerebral hemorrhages produced by *thrombosis of the sinus*, or, as in a case referred to by Virchow,\* as enormous hemorrhagic infiltration of the subperitoneal and sub-

---

\* Pathologie und Therapie, p. 172.



cutaneous tissues, as well as of portions of the muscles of the abdominal walls, the result of a coagulum in the external iliac vein, the epigastric, and the crural vein. Thrombosis may also occur in the cerebral sinuses, without causing hemorrhage, but giving rise to pressure symptoms, pain, prominence of the eyes, and œdema; it may be followed by complete recovery.

In *exhausting and wasting diseases*, blood may clot in the veins, or even in the heart, without any clearly-marked cause. Trousseau called attention to the occurrence of milk-leg as a symptom of gastric cancer. Gout may cause phlebitis and clotting in the veins of the body, as Sir James Paget has pointed out. Again, we may have *chlorosis* give rise to thrombosis in the cavities of the heart and the larger veins, such as the femorals, without phlebitis preceding the morbid condition.\*

Now, portions of the clot, situated in any part of the venous system, however remote from the heart, may become, by being broken off and driven onward with the circulation, sources of great danger. When the blood clots in veins connected with the portal system, the detached fragments may be washed into the liver, and there lead to secondary abscesses. But when coagula occur in the venous system and are wholly or in part carried away with the circulating blood, if we exclude those which, from their situation, could only reach the liver, we generally find the manifestations of disturbance arising in the heart or the lungs. Arriving at the *right side of the heart*, the concretion, if at all large, or if it become so by serving as a nucleus for a larger clot, occasions symptoms of exhaustion and collapse; an intermitting, feeble pulse; irregular and confused beating of the heart, and cardiac sounds enfeebled or lost over the right side of the organ; rapidly-developed distress in breathing, referred, by the sufferer, to the heart,† and signs of asphyxia, though all the time the patient is taking deep inspirations; great agitation; and a swollen state of the veins of the body. Death may then take place suddenly if a portion of the clot separate and obstruct the pulmonary artery.‡

But the mode of death, and the symptoms preceding it, in

---

\* Tuckwell, St. Bartholomew's Hospital Reports, vol. x., 1874.

† B. W. Richardson, Medical Times and Gazette, Nov. 1868.

‡ As in a case recorded by Druitt, Med. Times and Gaz., July, 1862.

embolism of the *pulmonary artery*, are not always the same, and depend much upon the size of the embolus and where it is arrested. A large-sized clot, whether it be merely part of one occupying the right heart, or be washed at once into the pulmonary artery, will occasion the same signs as those mentioned as indicative of a large clot in the right side of the heart; the craving for air is particularly intense, and this craving is increased by every movement of the body; the muscular debility, the lowered temperature, the cyanosed look, the turgid veins of the neck and their undulations, the increased, irregular cardiac impulse, though the heart's action is not sufficiently deranged to account for the disturbed respiration and disordered general circulation, are also noticed; and in some cases a systolic blowing sound, and, where the case is at all protracted, vertigo, albuminuria, and œdema of the limbs, may be observable. The intellect remains clear. As regards the pulmonary phenomena proper, collapse of the lung, hemorrhagic effusions or so-called infarctations, œdema, or capillary bronchitis are likely to happen, except in those instances in which the principal trunks of the pulmonary artery are blocked up and almost instantaneously asphyxia ensues. If the fragments be very small, the amount of dyspnœa is not of necessity great, nor are the symptoms of asphyxia marked; and inflammations of the parenchyma of the lungs may take place, occasioning often secondary obstructions and metastatic abscesses in the lungs. These forms of metastatic abscesses are observed in pyæmia, and are not unusual in puerperal fever.

Blood coagulates in the *arteries* as a consequence chiefly of gangrene and of ulceration. Again, inflammation or atheromatous disease of the coats of the arteries may lead to the local development of thrombi; so may feeble action of the heart and increased coagulability of the blood.\*

Still, the most important phenomena connected with obstruction of arteries are those of coagula being washed into them; the phenomena of *embolism*, therefore, rather than those of thrombosis. The manifestations of embolism are distinguished from those of the mere formation of clots by what is always the most significant sign of either arterial or venous embolism,—the suddenness of the

---

\* Liddell, Amer. Journ. Med. Sci., July, 1873.

manifestation of the abnormal state. And in point of fact the symptoms arise less often as the result of any of the conditions alluded to that occasion coagulation, than in consequence of deposits, fibrinous concretions, and excrescences which are seated on the valves of the left side of the heart, portions of which deposits are carried away by the circulating blood into remote parts. When these bodies become impacted in a vessel the calibre of which is such that it does not permit them to pass on, we find rapid changes taking place in the portions of the body supplied by the obstructed artery,—coldness, pallor of the parts, a diminished functional activity, a shrinking; and if the first obstruction be followed by others, and the collateral circulation cannot be established, local death and gangrene ensue.\*

All these changes are, of course, discernible only in external parts, especially in the extremities; the disturbances of function are the most obvious signs where the internal organs are the sufferers. If the emboli be driven to the *brain*, we have often, as has been mentioned, softening as the final result, and this may be preceded by disorder of intellect, without motor disturbances, and by severe attacks of vertigo, in cases in which merely the smaller arteries supplying the surface of the cerebral hemispheres are obstructed. But where, as is indeed the most common seat of emboli, the arteries of the fissure of Sylvius are clogged, the phenomena are those of apoplectic hemiplegia, and the palsy affects the whole of one side of the body. The brain may also suffer from the seat of the obstruction being in the carotids; indeed, of all organs the effects of embolism are most plainly perceptible in the brain. The presence of emboli in the splenic, renal, hepatic, and mesenteric arteries is generally only to be inferred from the history of the case, and does not occasion any clearly-discernible signs. But tenderness, enlargement of the spleen, and pain in the splenic region in *splenic embolism*, or disordered secretion of urine and pain in the loins in *embolism of the renal artery*, or jaundice in *embolism of the vessels of the liver*, may be very marked.

The occurrence of pain in these cases of internal embolism must not be overlooked; and in embolism of the arteries of the

---

\* As regards the anatomical lesions, see Litten, Zeitsch. f. Klin. Med., 1880; and Cohnheim, Allg. Path., 2d edit., Berlin, 1882.

extremities pain is a symptom of still greater prominence. It may be like a violent neuralgia, or so constant that it is mistaken for rheumatism ; and, as happened in a case of embolism of the right iliac artery, under the charge of Dr. James H. Hutchinson,\* which I saw, it may recur in paroxysms of intense severity, and be referred to the foot, though this be already in a condition of sphacelus. Besides the pain, we are apt to find extreme hyperæsthesia in some parts of the affected limb ; and pricking sensations, formication, and loss of tactile sense, followed by complete anæsthesia, in others. Then painful spasms of the muscles, and a more or less perfect paralysis of motion, may occur. If we join to these symptoms an absence of pulsation in the arteries below the occlusion until the collateral circulation is decidedly established, a strong beat of the vessel on the cardiac side of the obstruction, the coldness of the limb below this obstruction, and the signs of defective supply of blood, we have a group of phenomena which, taken in connection with the history of the case, render the diagnosis a positive one. In reviewing the history of the case the state of the heart and the cardiac symptoms must always be carefully examined into ; and a close inquiry often shows that the sudden manifestations of arterial obstruction were preceded by an attack of palpitation and of irregular action of the heart.

A change in the physical signs of the diseased organ, as of its murmurs, may not be evident ; but, should it be evident, it is a sign of utmost moment. Indeed, any change in what may be viewed as the centre from which the embolus may be detached is of great significance. And this holds good quite as much for venous as for arterial emboli. Thus, in a case of coagulum in a vein, a sudden disappearing of swelling and œdema of the affected limb, with the supervention of signs of embarrassed circulation and respiration, would at once tell what had taken place.

In regard also to the diagnosis of embolism we must always bear in mind the causes which are likely to give rise to it. Several of the causes of arterial embolism have already been mentioned ; those of venous embolism are the same as of venous thrombosis, or, to speak more explicitly, the breaking up of the clots and their transportation may occur in any of the conditions which have oc-

---

\* Amer. Journ. Med. Sci., Oct. 1863.



occasioned them. Now, these conditions, too, will produce arterial clots, and indeed some are more apt to lead to coagulation in the arteries than in the veins. Prominent among them are a narrowing of the calibre of the vessel, as by pressure; dilatation of the vessels and of the heart; failure or great diminution of cardiac power, with consequent retardation of the blood-stream,—a state which is more likely to occasion venous than arterial thrombosis; a breakage in the continuity of the vessel, as when it is torn or cut; changes which take place in the coats of the vessels, especially inflammatory changes; and contact of the blood within the vessels with foreign bodies. Then it is very likely that special states of the blood also, by altering the cohesion of the globules, predispose to, if they do not absolutely cause, the clotting.

Another cause of embolism is that due to *accumulations of pigment in the blood*, the result of malarial fever. The pigment may obstruct the capillaries in the brain and thus occasion capillary apoplexies; or be driven to the liver and there produce signs of disturbance of its circulation, and abscesses. As in all forms of capillary embolism, the symptoms are obscure: the suddenness of their development, generally so characteristic of the other forms of embolism, is wanting; and the diagnosis, as throughout in capillary embolia, is always nothing more than a matter of conjecture, based on a close study of the general phenomena, including the microscopic examination of the blood, and on the history of the case. Similar symptoms occurring after fractures of bone point to emboli derived from the marrow, to *fat embolism*.

The blood in malignant disease undergoes changes indicative of denutrition and wasting. It has been asserted that sporozoa can be found in the tissues invaded by carcinoma, and that these parasitic organisms are also present in the blood, but these statements are not yet fully established. With regard to other forms of malignant disease, observations of this character have also been made. Motz's researches concerning the blood of sarcomatous subjects reveal in the cultures obtained a very small micrococcus, simple or double, anaerobic, and staining with difficulty. The sarcomatous affection localizes itself in the venous walls or the capillaries, and causes a thrombosis of the affected vessels. This infection of the blood may serve as a diagnostic sign in doubtful cases.

*Acute endarteritis* may be the cause of embolism as well as of pyæmia. *Air in the blood* produces great disturbance of the circulation, which may be thought to be due to embolism. The air may be the result of decomposition, and get into the venous system and thence into the general circulation. Jurgensen \* has reported a case in which the air passed into the circulation through the splenic vein. Irregular contraction of the heart, pallor of the face, a peculiar systolic cardiac murmur, faintness and the signs of cerebral anæmia, and slow breathing, are the common symptoms.

In conclusion, the *subsequent changes of the thrombus* must be adverted to. It may organize and be converted into connective tissue and yield an impaired passage to the blood; and perhaps the collateral circulation may be freely established; or, what is not so favorable a result, it may soften and undergo fatty metamorphosis. But even when larger portions are not detached and occasion the marked symptoms of embolism, small ones of an infected thrombus may be wafted into capillaries and there lay the foundation of abscesses. It is thus that in a case of thrombus or embolus we may have the secondary results of pyæmia to deal with,—metastatic abscesses caused in the manner described, and attended with a blood profoundly altered and vitiated by the decomposing products circulating in it.

**Scurvy.**—This disease is not often met with in civil practice; but it is one familiar to the military and the naval surgeon. It consists in a deterioration of the blood, produced by living for a long period upon the same kind of food, and especially upon salted meats, without the requisite supply of fresh vegetables being taken. Now, the potent influence of vegetables is attributed to the large quantity of potassium they contain; and, as there is a deficiency of the salts of potassium in scorbutic blood, it was concluded that this deficiency is the real cause of scurvy. But this theory has not been positively proved. Another cause of scurvy is the want of proper assimilation of food, as in prison scurvy.†

The existence of scurvy in childhood is now recognized as of not infrequent occurrence, and it is probably frequently mistaken for rachitis, acute rheumatism, or possibly for purpura. The

---

\* Archiv. f. Klin. Med., Bd xxxi., 1882.

† See Medical Memoirs of the U. S. Sanitary Commission, p. 278.

concurrence of marked anæmia with joint-swellings in a bottle-fed infant, or in older children, should suggest the possibility of scurvy being present. Northrup and Crandall\* found, in over sixty-three per cent. of the cases of infantile scurvy they investigated, that the diet consisted of proprietary foods and condensed milk. The evidence also indicates that milk sterilized for a long time is capable of causing scurvy when used as an exclusive diet.

Babes,† in studying three cases of scurvy bacteriologically, found a thin, long, wavy bacillus, prone to occur in clusters, in the gums, the lungs, and other viscera. There were also streptococci in the gums.

Scurvy is usually slow in its development. The patient becomes low-spirited, easily fatigued, and is loath to exert himself. The appetite is impaired; there is a craving for acids and for vegetable food; the tongue is flabby; the breath fetid; the pulse feeble; the skin dry. The bowels are usually constipated; but a tendency to diarrhœa may exist, and indeed generally occurs as the disease advances. Neuralgic pains, referred chiefly to the lower extremities, to the bones, and to the back or thorax, are common. The face is pale, or has a yellowish tinge; the eyes are surrounded by a dark ring. During the progress of the ailment, or in severe cases almost from the onset, we find swollen, spongy gums, bleeding on the slightest touch; hurried breathing; a rapid pulse; weakened eyesight, sometimes night-blindness; epistaxis; painful swelling and hardness about the joints of the extremities and in the calves of the legs; and purple spots and bruise-like stains on the skin. Should the malady remain unchecked, the symptoms heighten in severity, ulcers form which have a fungoid look and a great tendency to bleed, hemorrhages take place from internal organs, old sores and wounds reopen, well-knit fractures become disunited, there is a constant tendency to swoon, and the patient perishes miserably exhausted, and with his blood in a state of dissolution. Scurvy may be the cause of epidemics of pericarditis.‡ In some cases death takes place from diarrhœa or dropsy, which may be suddenly developed. Recovery from scurvy is slow.

---

\* Proceedings of New York Academy of Medicine, Feb. 1894.

† Quoted in Sajous' Annual, vol. i., 1895.

‡ Von Dusch, Herzkrankheiten.



**Purpura.**—Scurvy is not a disease difficult to recognize ; only one affection resembles it at all closely,—*purpura*. In this disorder also red or purple spots or livid blotches, uninfluenced by pressure, and passive hemorrhages from the mucous membranes, happen. But there is this difference between the two complaints : *purpura* is common in fruit seasons, and often attacks persons who have not been in any way deprived of vegetable food. The gums are not soft and spongy as in scurvy, nor do we find the same weakness of mind and body. Then, the stain of the skin in *purpura* is apt to be more generally diffused, and the purple blotches are smaller, or, at all events, the large patches of discoloration consist clearly of an aggregation of very many small spots. Moreover, the disorder is not controlled, like scurvy, by fresh vegetables, and by lemon-juice,—in fact, by decided antiscorbutics.

From a clinical point of view we find several forms of *purpura*. In the mildest, the purpurous spots are apt to appear only on the legs. They come in crops, which fade, and there are no constitutional symptoms, except a little lassitude, and perhaps aching of the limbs and pain in the back. In the graver cases, “*purpura hæmorrhagica*,” we find, in addition to the cutaneous hemorrhage, epistaxis, hæmatemesis, hæmaturia, or other internal hemorrhages, and extravasations of blood may happen into the substance of the muscles. The amount of pain attending the malady is very different. There may be none, or it may be trifling ; or deep-seated pains in the cavities of the body, or extended neuralgic pains, may accompany the purpurous complaint. In some instances the pains are chiefly felt in and around the joints, and the apparently rheumatic aches subside in a few days, and spots of extravasated blood become visible. This “*purpura rheumatica*,” a variety particularly described by Schönlein, is usually met with in the strong and healthy. It is, indeed, one of the peculiarities of any kind of *purpura*, that it may come on in the midst of seemingly excellent health ; for while it is true that the disorder may be preceded for some time by general debility, or occur in the course of disease of the liver, of Bright’s disease, or as a sequel to the exanthemata and rheumatic fever, it most often happens where, from the previous history, we should least expect it. Its production as the result of a sudden shock to the nervous system, such as fright, and its occasional intermittent character, have been repeatedly noticed.



It has appeared after the administration of quinine, as first observed by Vépau,\* and since by Gauchet † and by Woodbury.‡

The duration of the malady is very variable: only a week may elapse, or several months may pass, before the spots disappear. Its pathology is unknown. It is clearly, however, not merely a disease of the blood; the capillaries lose their retentiveness and allow the corpuscles to migrate. In some cases purpura presents an *acute* form. It is ushered in by a chill, and by intense pain in the back and limbs, but is generally unattended with fever or severe constitutional disturbance. The purple spots usually first appear on the legs, and are wholly uninfluenced by pressure. They last five or six days, or somewhat longer, then gradually change their color and fade. The patient feels languid, but, unless from loss of blood, his strength is not materially impaired. The effusion of blood happens in some cases into the loose connective tissues of the body, or blood is lost from the lungs, and still more frequently from the bowels or the urinary organs. Under these circumstances the pulse, which is apt to preserve its normal frequency, becomes very rapid; but until exhaustion begins to tell on the nervous system—not, as a rule, long before dissolution—the mind remains clear, and cerebral or spinal symptoms are absent. It is thus that we are able to distinguish severe cases of acute purpura, which may indeed prove fatal in forty-eight hours,§ from cerebro-spinal meningitis. Some of these acute or fulminating cases occur in young children, and it is a question whether or not they were previously subjects of infantile scurvy.

The distinction between *hæmophilia* and purpura is generally simple. It is true that the bleeding in a member of a bleeder's family may happen into the skin, or from any of the parts from which it may take place in purpura; but the family history, the congenital proneness to frequent hemorrhages from the slightest cause, their danger and protraction, the functional excitement of the heart, followed perhaps even by cardiac hypertrophy, the attacks of rheumatoid joint-inflammations, especially after exposure

---

\* Gazette Méd. de Strasbourg, 1865.

† Bulletin de Thérapeutique, vol. cl.

‡ Philadelphia Medical Times, 1886.

§ Harrison Allen, Amer. Journ. Med. Sci., Jan. 1865.

to cold and damp, associated with hemorrhage, and the hemorrhagic diathesis exhibited, stamp hæmophilia with distinctive features. There is in this disease diminished coagulability of the blood; in rheumatism the coagulability is increased.

Microscopic examination of the blood of a case of hæmophilia showed a very largely increased number of very small corpuscles or microcytes, with no marked increase of the leucocytes, but the percentage of hæmoglobin was reduced, and also the red blood-corpuscles, proving the presence of both quantitative and qualitative anæmia.\* Henry has directed attention to the changes occurring in the arterioles, consisting mainly in wasting or absence of the middle muscular tunic of the vessel. Vaso-motor influences undoubtedly play their part in bringing about an attack. This is shown by the flushing of the face which so often precedes an attack, and also by the fact that the attack may follow emotional excitement. *Leukæmia* may be accompanied by subcutaneous extravasation of blood, but it cannot be mistaken for either hæmophilia or purpura, if an examination of the blood be made. Hæmophilia is almost exclusively restricted to the male sex. It may be associated with arthritis.

---

\* Daland, College of Physicians of Phila., Jan. 9, 1894.

## CHAPTER XI.

### RHEUMATISM AND GOUT.

RHEUMATISM AND GOUT are affections having a strong tendency to change their seat, and are dependent upon the presence in the blood of some poisonous material which probably accumulates there in consequence of malassimilation. The rheumatic poison has a singular predilection for the fibrous, serous, and muscular textures. Hence we find it attacking principally the joints, the fasciæ, the endocardium and pericardium, and the muscles in various parts of the body. According to its main forms, it is sometimes divided into articular and muscular; but the more usual division into acute and chronic is simpler, and will answer our purpose best.

**Acute Rheumatism.**—Here the poison gives rise to the symptoms of an acute, active disease, and attacks especially the larger joints. These swell, become hot, red, tense, tender, and the seat of pain aggravated by the slightest movement; an effusion also takes place into the surrounding structures, or into them and the synovial membranes of the joint itself. The rheumatic inflammation may either remain confined to the joints first affected until the disease is over, or, what is more common, it shifts from joint to joint, implicating most of the large ones in succession, yet often invading fresh joints before the swelling has subsided in the parts first attacked. The articular disorder is ushered in and accompanied by high fever, soon attended with a full, bounding pulse, profuse, sour perspirations, a deeply-coated tongue, a scanty, turbid, highly-acid urine, and a countenance expressive of suffering. The fever is generally in proportion to the number of joints involved. The temperature runs up to about 102° or 103° Fahr. very soon after the outbreak of the malady, and remains steady, with slight evening exacerbations and morning remissions when the joint-affection is yielding, but with renewed rises when fresh

joints are being implicated. As the disease disappears, the fever temperature gradually subsides.

There is little difficulty in recognizing the complaint. The pains in the joints, their tumefaction and tenderness, the shifting character of the disorder, the fever, the acid sweats, form a group of phenomena eminently characteristic. In truth, excluding acute gout, the only affections at all likely to be confounded with acute articular rheumatism are pyæmia and glanders, acute synovitis, and milk-leg. The diagnosis of the former has been discussed in connection with diseases of the blood; it only remains to point out the marks of similitude and contrast between acute articular rheumatism and the other maladies mentioned.

*Acute synovitis* resulting from an injury, or from cold, occasions, like articular rheumatism, pain and heat in the joint, with distention. But the disorder, except, perhaps, if it happen in a rheumatic constitution, does not affect more than one joint; and, as there is scarcely any or no effusion into the surrounding tissues, the outline of the joint can be distinctly discerned, and fluctuation is readily detected. Often, too, the accumulation of fluid reaches an extent far greater than in rheumatic inflammation; moreover, the febrile and constitutional derangement is not so severe as in acute rheumatism, and the affection has no tendency to change its seat. Still, acute synovitis may be rheumatic.\*

*Milk-leg*, or phlegmasia alba dolens, occurs most usually in women after delivery, or as a sequel of continued fevers, and is commonly a phlebitis of septic origin in which a thrombus forms. Generally, only one leg swells, and this becomes throughout, or sometimes only around the calf, preternaturally white, firm, hot, and shining. The tumefaction is uniform, and very painful, especially so when touched. It does not pit, or pits but slightly, upon pressure, except at the lower part. There is in some cases tenderness with a sense of hardness in the course of the femoral vein, though this is by no means constant; and we are apt to find signs of much debility and of altered blood, and febrile symptoms. The history of the case and the local signs are very different from acute rheumatism. Among the latter, two giving rise to striking dissimilarity may be mentioned: the almost

---

\* See Adams, Medical Times and Gazette, Feb. 1869.



entire loss of power in the affected limb in phlegmasia alba dolens, and the much higher temperature it shows by the thermometer than the other members. And, while alluding to its heat, we may remark that an increase of general temperature corresponds to an increase of pain and swelling in the limb, and of constitutional distress.\* Phlegmasia dolens has been noted in association with chlorosis.†

Rheumatism may be modified in its manifestations by happening in connection with, or consequent upon, other disorders. For instance, the febrile phenomena may be of an adynamic type when the disease occurs consecutively to typhoid or typhus fever; or we may find the local signs of acute rheumatism strangely mixed with the symptoms of puerperal fever, and in some of these cases pus may fill the tumid joints; or the presence of the syphilitic poison or of gonorrhœa may imprint peculiar features upon the complaint; and in most of the instances mentioned the rheumatism is probably of different blood-origin.

In *gonorrhœal rheumatism* there is usually less febrile distress; the articular pain is not so severe or acute; the integument covering the affected joint is apt to retain its normal color; there may be but one joint—and there are not generally many—implicated; the inflammation is confined to the synovial membrane, and a copious sero-fibrinous exudation occurs; the joint-affection, which is pre-eminently an affection of one knee, shows a tendency to shift, and resembles rather an acute or a subacute rheumatoid arthritis than acute rheumatism; the eye, too, unlike what happens in ordinary acute rheumatic fever, is often attacked. There is no copious sweating, and no disturbance of the heart; and often there has been a running from the urethra, which diminishes when the gonorrhœal rheumatism sets in, but which does not cease. The disorder does not come on early in a case of gonorrhœa; and the joint-affection appears really to be of pyæmic origin. It disappears only very slowly, and is uninfluenced by salicylic acid.‡ It is by all these signs that we judge of the malady with much more certainty than by the mere presence of gonorrhœa with the symptoms of rheumatism, for the former may be a mere coincidence.

---

\* Elliott Richardson, *Pennsylvania Hospital Reports*, vol. ii.

† Perret, *Lyon Médical*, 1888.

‡ German edition of this work.

Gonorrhœal rheumatism may run an acute course.\* Purulent effusions into joints may be mistaken for acute rheumatism. The history of the case, the frequent association with an infectious malady, and the location of the swelling, distinguish these *pyæmic joints*.

The traits of an attack of acute rheumatism are frequently altered by certain complications in internal organs which the contaminated blood is apt to occasion. Prominent among them are the cardiac disorders, which are in fact so common that they may be looked upon as forming part of the rheumatic manifestation. Their signs we have investigated already, while examining endocarditis and pericarditis. Certain cardiac phenomena, such as extreme pain without evidence of recent valvular affection, pain which may shoot to the neck and shoulder and be associated with signs of great irritability of the heart or of heart-failure, have been by some observers, as by Peter and Letulle,† attributed either to rheumatic myocarditis, or to an abnormal excitement of the cardiac plexus, of rheumatic origin.

Other complications are inflammations of the lung, of the bronchial tubes, and particularly of the pleura; an affection of the kidney which is generally a parenchymatous nephritis with some albumin and tube-casts, but which may be due to pyæmic or embolic infarction;‡ and—though not often—cerebro-spinal disturbances, exhibiting themselves by headache, violent delirium, convulsions, and coma, and occurring either in connection with a thoracic disorder, or solely in consequence of the action of the vitiated blood on the nervous centres, or in consequence of Bright's disease or of multiple capillary embolism, or of the sudden exhaustion of the nervous centres. This explanation§ has been more particularly applied to the cases in which an excessive temperature attends the rapidly-developed signs of cerebral disturbance, a temperature of 107° or more. But, speaking from a bedside point of view, we must remember that such cases are comparatively rare, and that rheumatic delirium is far from

---

\* Davies-Colley, Guy's Hospital Reports, 1883.

† Archives Générales de Médecine, June, 1880.

‡ Chomel, Recherches sur les Reins dans le Rhumatisme, Paris, 1868; also Schmidt's Jahrb., No. 2, 1871.

§ Weber, Transactions of the Clinical Society of London, vol. i.

always of the same nature. It may be of the kind just mentioned. It may develop itself with or without the signs of cardiac complaint. It may come on early in the disorder during the violence of the fever; or late, and clearly from debility and impoverished blood, yielding to nourishment and stimulants. It is rarely the result of meningitis. The delirium which attends *cerebral rheumatism* may be marked by great talkativeness, or, on the other hand, the patient may be extremely taciturn.\* Insanity may follow the brain symptoms of acute rheumatism. In some instances, whether due to rheumatic inflammation or to mere disturbance of the medulla and lower half of the pons, we find in rheumatic hyperpyrexia nervous symptoms that simulate multiple sclerosis,—exaggerated knee jerks, ankle clonus, scanning speech, nystagmus, and tremor. Foxwell† has reported such a case in which the temperature reached 111°.

The occurrence of *nodules* in connection with rheumatism, especially among children, has attracted a good deal of attention. They are met with chiefly in the neighborhood of joints, especially of the elbow. These fibrous nodules may appear at once in any form of rheumatism, or come out in crops. They are not tender. They most often occur in cases of rheumatic endocarditis or pericarditis.

In a few instances of rheumatism we find *acute arteritis* arising, and especially inflammation of the fibrous structures of the aorta. This condition may be suspected should we observe intense general uneasiness and distress, with pain, increased pulsation, a distinct murmur in the course of the vessel, and tumultuous action of the heart without there being obvious signs of disease of that organ present. Still, the diagnosis is never a positive one.

Acute rheumatism rarely ends fatally; its cardiac consequences are more to be feared than the acute attack. Cases occur not unfrequently in which the inflammation in the joints is lingering, and in which the febrile symptoms are not intense. These cases form an intermediate grade between acute and chronic rheumatism, and are spoken of as *subacute*. The disorder is more apt than the acute variety to affect the muscles as well as the joints; nay, the

---

\* Some of these points are more fully detailed in my paper on Cerebral Rheumatism published in the Amer. Journ. Med. Sci., Jan. 1875.

† Lancet, May, 1886.

former may be alone attacked. It may be witnessed in the joints of one extremity, or in one joint, and might then be mistaken for synovitis. But the dissimilar history of the complaint will guard against error: no accident has happened to account for the swelling of the joint, and often the patient will tell us that he has had previously an attack of rheumatism. The subacute form of rheumatism is more likely to be confounded with rheumatic arthritis: we shall presently refer to their distinction.

**Chronic Rheumatism.**—This may either be a sequel of the acute disease, or the disorder may from the onset assume a lingering form, the constitutional symptoms being slight. The affection may show itself in the joints, giving rise to stiffness, dull aching, pain produced by motion, but without heat or very obvious swelling, tenderness, and febrile excitement, or marked sweating; or it may implicate the muscles in various parts of the body, occasioning stiffness, as well as pain when they are moved; or it may attack both joints and muscles; or it may be seated chiefly in the sheaths of nerves, leading to rheumatic neuritis, of which sciatica often affords a striking example. In any case the occurrence of the pain furnishes the starting-point in diagnosis; and we must ascertain, by careful examination, whether it be augmented by motion, whether it be more or less shifting, whether it be not combined with stiffness either of the muscles or of the joints, whether it be influenced by changes of temperature, whether it be not neuralgic, or associated with a disturbance of some viscus, such as of the liver or the kidneys, before we conclude that the complaint is really rheumatic.

This is especially necessary in the most common form of chronic rheumatism,—*muscular rheumatism*. All kinds of pains in the muscles or their surroundings, the cause of which is not at once apparent, are apt to be pronounced rheumatic. And indeed it is not always easy to say whether they are or are not of that character. We may distinguish them from the anguish of *neuralgia* by the pain in the latter complaint being ordinarily confined to the distribution of one nerve and not being increased by movement or by pressure, nor is it so steady, or attended with soreness, except over a few spots at some distance from one another in the course of the affected nerve.

As regards the pains caused by *organic structural disease*, we



can generally discriminate them from those of rheumatism by close attention to the history of the case, and by a careful exploration of the internal organs. Thus, for instance, we shall find pain radiating from the right hypochondrium to the shoulder to be dependent upon hepatic disease; or pain shooting down to the groin, thigh, and testicle to be caused by a disturbance of the kidney; or a bearing down and an aching near the sacrum to be probably due to uterine disorder.

Muscular rheumatism may affect the neck, the scalp, the muscles of the face, and the parietes of the chest or of the abdomen. It may be not only chronic in any of these situations, but also acute; or, what is more frequent, when it occurs with fever and is transient, it is a sudden acute exacerbation in persons who are rheumatic and suffer more or less persistently from rheumatism, though perhaps in a different part of the body from the one in which the acute affection has happened. Muscular rheumatism has been noticed in an epidemic form.\*

One of the most common seats of muscular rheumatism is in the loins. It then constitutes the disease known as *lumbago*. The patient is unable to stand erect, and finds it nearly impossible to stoop forward, on account of the severe pain occasioned when the muscles of the back are called into action. Unless the attack be very severe or acute, there is no constitutional disturbance; but the disorder is often obstinate. It is easy of recognition. We distinguish it from pain in the loins due to disease of the kidneys, chiefly by an examination of the urine, and by the different way in which movement affects the rheumatic pain; from lumbo-abdominal neuralgia, by the two or three sore spots in the course of the affected nerve; from rheumatism of the vertebral articulations, by the absence of tenderness and swelling around the spinous processes; from lumbar abscess, by the want of local bulging or fulness, of fluctuation, and of fever. Then, we must be careful not to consider as lumbago the pain in the back caused by disease of the spine, or by disorder of the uterus, or by the passage of abnormal urinary constituents, such as oxalate of lime, or consequent upon strains, or blows, or scurvy, or malaria, or anæmia, or a general or local muscular debility.

---

\* Schmidt's Jahrb., No. 12, 1872.

Thus there are many causes of pain in the loins, and where the case is of any duration or of any doubt we must be careful to exclude these causes from consideration before we assume the disease to be really rheumatism of the muscles and fasciæ of the back. This caution is very necessary in investigating the cases of "weak back" so prevalent among soldiers, which, though commonly spoken of as rheumatic, are really, for the most part, due to strains or injuries which have perhaps produced a weakness of the muscles and a persistent cutaneous hyperæsthesia; or to an impoverished blood, to neuralgia, to scurvy; or to digestive disorders attended with the passage from the kidneys of large amounts of urates or of oxalate of lime.

The remarks made with reference to this form of muscular rheumatism and the states which simulate it are also applicable to pains apparently muscular affecting other portions of the body. We may have pain and soreness of the muscles developed by strain or overwork and attended both with muscular and with cutaneous hyperæsthesia,—a condition very different from rheumatism, and designated by Inman "myalgia." This soreness of the muscles is always in direct proportion to their debility, and is chiefly caused by long-continued exertion beyond the power of the muscle, or by an ordinary amount of action when it or the individual himself is extremely debilitated. The morbid state is most marked during the convalescence from scarlet fever, where it may be looked upon as due to over-exertion of the weakened muscles. The soreness of the muscle is almost constantly accompanied by heightened sensibility of the skin over it; and this coexisting cutaneous tenderness may be in any case regarded as an important diagnostic sign. Myalgia is chiefly found in the muscles of the trunk, and is very rarely general.

Another form of muscular rheumatism which we may here mention is wry-neck, or *torticollis*. This depends chiefly upon contraction of the sterno-cleido-mastoid muscle of one side, and occasions the ungainly appearance with which most persons are familiar. But we must be careful not to consider every case as of rheumatic origin. The disorder may be spastic, or may depend upon nervous injury, and when chronic may lead to alteration in the muscular structure. Injections of atropine, hypodermically, may generally be used, not only for their good therapeutic effect,

but also because, even in chronic cases, they may show us, by the difficulty or impossibility of relaxing the muscle, how much of it is really changed.

There are forms of pain in muscles and tendons that are often mistaken for muscular rheumatism. *Achillodynia* is one; the slight swelling about the insertion of the tendo Achillis, with pain on standing or walking but without much tenderness, marks an affection that is frequently not rheumatic. In *Morton's disease* the pain in the metatarsal phalangeal articulation occurs in seizures, yet only when the foot is moved as in walking; there is neither heat nor swelling.

Pain in the muscles and stiffness may be caused by still other conditions than those described, and be mistaken for muscular rheumatism,—the muscular pains of *trichiniasis*. But the marked exhaustion and the signs of gastro-intestinal catarrh are of such significance that they save us from error.

A form of chronic rheumatism which also may be briefly mentioned is that affecting chiefly the fibrous membranes, such as the *periosteum*. This becomes thick, and tender on pressure; its thickening may even be very perceptible to the touch as well as to the eye. This kind of rheumatism happens in those who have syphilis; but it also occurs where no such taint exists. The pains are generally much more severe at night; and this is sometimes assumed to be a proof of the syphilitic character of the disease,—but incorrectly so; for many varieties of chronic rheumatism are aggravated by the warmth of bed. Indeed, the only really diagnostic signs of syphilitic rheumatism are the obvious evidences of constitutional syphilis, or the history of the infection. Still, to cases in which several nodes exist, and in which the pains more particularly affect the long and flat bones, and in which iodide of potassium speedily modifies the pains, we shall be rarely wrong in attributing a syphilitic origin.

Chronic rheumatism is often feigned, especially by malingerers in the army and the navy, and the deception may be difficult of detection. They pretend to be scarcely able to walk, or hobble around with a cane, and complain much of the pain and stiffness in their joints. Yet there is not the least sign of deformity or real stiffness; the pain is always stated to be the same; and their general health is excellent. Their way of using the stick, too, is

characteristic: they move it each time they move the seemingly crippled leg, but, as a rule, not immediately, thus not employing it as a support. Anæsthetics are of great value in enabling us to decide as to the real amount of immovability of the limb.

**Gout.**—This disease may be, like rheumatism, either acute or chronic. Instead of describing its phenomena, I shall at once point out the marks of difference between the two kindred maladies. In gout, the small joints are chiefly or alone affected; in rheumatism, the large. The gouty inflammation is accompanied by more local pain and redness than the rheumatic, and by œdema, enlargement of the veins, and desquamation of the cuticle, and implicates, at least at first, only one or a few joints, especially the joint of the great toe; while rheumatism attacks the joints of the upper as well as of the lower extremities. In gout there is a tendency to disease of the kidneys, with a moderate febrile disturbance, and no profuse sweats; but we meet rarely with a cardiac complication, at least a valve affection, as constantly happens in rheumatism. Gout is more decidedly hereditary than rheumatism; its early attacks are apt to recur with a certain amount of periodicity, and last about a week,—therefore a much shorter time than those of rheumatic fever. During the paroxysm of gout the urine is scanty, and both before the attacks and during the first days the uric acid is strikingly diminished.

Gout occurs generally in those who live high or who drink large quantities of malt liquor, especially in men about middle age, or is seen in those whose systems have been impregnated with lead; while rheumatism is usually seen in the weak, is excited by cold and damp, is as common in females as in males, and is oftener found in the young and before middle age. Gout is frequently combined with a deposition of chalk-stones in the joints; rheumatism never. Then, as shown by Garrod,\* we possess an absolute means of diagnosis in the examination of the blood. Uric acid is always present in large excess in gout, and absent in rheumatism. This test will render easy of discrimination even those cases which, with the usually employed means now at our command, are very perplexing to distinguish. Nor is the method of detecting the uric acid difficult, if we make use

---

\* Gout and Rheumatic Gout, 2d edit., London, 1863.



of Garrod's ingenious plan. It consists in obtaining the crystals of uric acid, crystallized on a thread placed in a mixture of the serum of the blood or of the fluid from a blister with acetic acid, in the proportion of six minims of the acid to each fluid-drachm of the serum. The mixture of the serum and acid with the thread in it is placed in a shallow watch-glass and allowed to stand from twenty-four to forty-eight hours, protected from the dust. In the blood of gouty patients are many polynuclear colorless cells.

The remarks just made apply more especially to the distinction between acute gout and acute rheumatism. The chronic disorders are more difficult to separate. Indeed, unless there be external deposits or chalk-stones, their discrimination may be impossible. In these obscure cases, however, the history and an examination of the blood may throw considerable light on the diagnosis. In many subjects, too, the exploration of the external ear will assist us in arriving at a correct diagnosis: we find one or several spots of deposit of urate of sodium on the helix.

Gouty persons are subject to indigestion, flatulency, pains and cramps, or palpitation of the heart,—phenomena which are due to the gouty poison, and which are generally ameliorated by a fit of gout. The teeth of those of gouty diathesis are remarkably well enamelled, enduring, and free from decay; but there is great proneness for tartar to collect upon them.\* Violent fits of sneezing may be a most annoying symptom,† and so are deep-seated pain in the tongue and a sense of burning.‡ In chronic gout there are often knotty finger-joints and tophaceous deposits in fingers and toes. Gouty endarteritis is not uncommon; and the frequent association of contracted kidney with gout is universally recognized.

The gouty inflammation of the joints may retrocede during an attack, and severe epigastric pain, nausea, vomiting, flatulence and acidity, faintness and a feeling of sinking, and a quick, feeble pulse show that the morbid action is transferred to the *stomach*; or it flies to the *head*, and apoplexy or maniacal symptoms occur; or to the *heart*, and there is violent palpitation, with dyspnoea, and

---

\* Dyce Duckworth, Transact. Odontol. Soc. of Great Britain, 1883.

† Schmidt's Jahrbücher, No. 8, 1881.

‡ Dyce Duckworth on Gout, London, 1889, p. 87.

intense anxiety ; or it attacks the *spinal cord*, and a sense of constriction around the thorax and abdomen, and piercing pains in the limbs, like those of locomotor ataxia, are encountered, and the spinal dura mater and the roots of the spinal nerves are found to be incrustated with uric acid and urate of sodium.\*

Closely connected with gout is *lithæmia*. Indeed, the excessive formation of lithates and the dyspeptic symptoms, with the heart-burn and eructations, the signs of functional derangement of the liver, the vertigo, the mental gloom or the listlessness and indisposition to exertion, the cramps in the legs and muscular twitchings, the neuralgic attacks, the restless nights, the palpitations of the heart and its irregular beat, are in many but the precursors, although, it may be, the long precursors, of a regular outbreak of gout ; while in many more this half-dyspeptic, half-nervous condition, with the faulty assimilation, the imperfect oxidation, the excessive discharge of lithates at times and their disappearance at other times, will go on for years without ever developing into an attack of gout.† Still, in years the same local lesions may follow in internal organs ; we may have the same form of contracting kidney, arterio-sclerosis, and the heart-affection with hypertrophy, and the accentuated second aortic sound of the lithæmic state.

Lithæmia sometimes manifests itself in attacks of pain in the *stomach and bowels*. The pain is associated with tenderness, and is most common when the stomach is empty. Among the symptoms of lithæmia that are very liable to be mistaken and mis-treated are *disorders of vision*. As Risley‡ has recently stated, lithæmia is both a primary and a modifying factor in many of the discomforts and more serious disorders of the eye. It stands second only to syphilis in the frequency with which it causes *iritis*. In adults, *obstinate conjunctivitis* and *episcleritis* are apt to own lithæmia as a cause, and it often gives rise to pain and to photophobia. It may lead to ulceration of the cornea and *errors of refraction* and attendant *eye-strain* and *headache*.

**Rheumatic Arthritis or Rheumatic Gout.**—Gout is rare in this country. But the same cannot be said of that distressing

---

\* Ollivier, Archives de Physiologie, 1878.

† See paper on Lithæmia, by the author, Amer. Journ. Med. Sci., Oct. 1881 ; and University Medical Magazine, May, 1894.

‡ Proceedings of the State Medical Society of Pennsylvania, 1895.

disorder known as *rheumatic gout*, but which is neither rheumatism nor gout, but a distinct affection. The disorder may be acute or chronic. It is not often the former; many of the acute cases, indeed, being rather subacute than acute. Even in those belonging to the *acute* form there is little febrile disturbance; and though we observe pain and aching in the joints, and some discoloration, we find less redness than in acute rheumatism, and certainly the tongue less furred, the pulse not so bounding, much less profuse perspiration, no such heavy deposits in the urine, and an utter freedom from cardiac complication. The acute arthritic disease has rather inflammation of the pleura and of the eye as its attendants, and is often accompanied by a sallow skin, yellowish conjunctiva, and discolored, costive stools. It implicates the large and small joints equally, thus differing from gout, and causes very great swelling, due to an effusion, not around the joint, but into its capsule. It fastens upon several joints, and, though it may pass from joint to joint, it shows but little migratory tendency; the joints first attacked remain the seat of disease. Unlike gout, it is apt to affect the smaller joints of the hands without a previous affection of the toes, and exhibits no periodic paroxysms or exacerbations. Moreover, an acute attack is of very much longer duration. Unlike subacute rheumatism, it does not affect the muscles, and is, both in the suffering at the time and in its ultimate results, a much graver malady.

The great danger in rheumatic arthritis is from the effects of the inflammation on the joints. The changes there produced are obvious in the *chronic* form, for each joint attacked is apt to be much damaged. The chronic complaint may follow the acute, or it may begin without any febrile symptoms, with pain and stiffness in the joints. These soon become much distended with fluid, which is gradually absorbed, and the structure of the joint alters, the cartilages become, sooner or later, implicated, and gradually waste, and chronic changes and permanent deformity are produced. The alterations may go on getting worse and worse in consequence of repeated attacks, until complete immobility ensues, and, the joints becoming permanently affected, the ends of the bones are dislocated and enlarged. But, though there is much swelling, no deposits of urate of sodium are found in the joints. Occasionally, especially in men, the disease is only found

on one side of the body, and may show itself first in a large joint, as in the hip. Among its peculiar, though less constant symptoms, are very rapid pulse, sweating, and pigmentation of the skin, like freckles.

Charcot has pointed out that in *paralysis agitans*, in addition to rigidity of the muscles, deformities of the fingers result resembling closely those of chronic articular rheumatism. But the likeness to the deformities caused by rheumatic arthritis is still closer, and to distinguish them we must take into account the whole history of the case, the tremor, the fixed look, the peculiar gait, the indistinct speech, the tremulous handwriting, the sensation of excessive heat. Moreover, the disfigured joints are not stiff, and do not crack. The *arthropathies* of *locomotor ataxia* may be mistaken for rheumatoid arthritis, but, irrespective of the history and of the characteristic pains, the absence of the patellar tendon reflex distinguishes them. All these joint affections following nervous diseases, and sometimes classed together as *spurious arthritis*, differ from joints attacked by rheumatism or by rheumatic arthritis in the absence of marked swelling and of pain, except on forcible movement; stiffness is the prominent feature.

Rheumatic arthritis is more common in females than in males; like rheumatism, it may be excited by cold and damp, and is very apt to occur in the weak and unhealthy. It generally, even in cases that recover, persists for months. Nor will it yield to the remedies usually administered in acute rheumatism; nor to colchicum and the alkalies, so beneficial in gout.

I shall here add a short description of a disease of nutrition of dissimilar character to those described, but having this in common, that it markedly affects the organs of locomotion,—rickets.

**Rickets.**—In this country rickets is a comparatively rare affection, certainly rare as compared with its prevalence in England, in Holland, in Germany, and in some other Continental States. It is a constitutional disease of early childhood connected with impaired nutrition, and is chiefly characterized by increased growth of the epiphyses and periosteum, and imperfect ossification, producing softening of the bones with curvatures and distortions. The changes are most manifest in the long bones; and the amount of organic matter in them is more than doubled, while the earthy matter is scarcely above one-third of the normal quantity. Be-



sides the osseous changes there is evident cachexia ; and the liver and spleen become enlarged and indurated from overgrowth of the glandular elements and interstitial development of fibroid tissue. A similar process may also happen in the kidneys and in the lymphatic glands.

Insufficient and improper food is a powerful cause of rickets. The malady may show itself as late as the seventh or eighth year ; but it most generally sets in during the first or second year of life. When it leads to death, it does so generally by gradual exhaustion, by impairment of the digestive functions, by thoracic complications, such as extensive bronchitis, pleurisy, collapse of the lungs associated with bleeding of the thoracic walls, by spasm of the glottis, by convulsions, or by chronic hydrocephalus. As a marked disease it does not usually last longer than a year, though the results of the osseous changes may long persist, and, affecting the thorax or the pelvis, prove eventually very injurious. Yet in time the bones may lose their rickety condition and become strong and dense, although some curvature and deformity remain.

The beginning of the disease is insidious. The child makes no attempt at walking, or ceases to walk if it have commenced. It is languid, irritable, its face pale, its tissues flabby. The appetite fails, there are thirst and irregularity of the bowels, or the marked signs of a gastro-intestinal catarrh. Restlessness at night, a disposition to throw off the bedclothes, profuse perspiration about the head, neck, and chest, while the rest of the body is hot and dry, attend an irregular febrile condition which soon shows itself ; while fear of being touched, or general soreness or tenderness of the body or actual pain, bespeaks the local process that is going on in the bones and their covering. The changes in the bones now become more and more distinct. The joints appear swollen, especially at first the wrist-joints, and when these are examined the lower extremities of the radius and the ulna are found to be enlarged ; similar changes are perceived in the tibia and fibula, and in the elbow. There is tenderness along the ribs, and, should the affection continue, nodules are felt at the junction of the ribs with their cartilages ; the sternum protrudes, a pigeon-breast results ; then the limbs show contortions, the clavicles are bent, the spine may be curved, the pelvis deformed. The head is large and square, the forehead high, the anterior fontanel remains unclosed,

the sutures are open and thickened on the sides. A blowing sound is frequently to be perceived over the cranial sutures. Dentition is delayed, or the teeth decay and fall out. The urine is copious, and contains lactic acid and an excess of phosphates. In advanced cases the symptoms of cachexia are very marked; the flabby muscles, the wan, anæmic aspect, the large abdomen contrasting with the small face, the enlarged liver and spleen, the persistent tenderness over the bones, and at times the marked fever, give sad evidence of altered nutrition and of suffering; yet even then the little patient may recover, though most likely with part of the osseous system irretrievably damaged. Of course we have all kinds of gradations in the malady, and the general symptoms attending the morbid process may be slight, just as the rickety condition of the bones may be limited.

The *diagnosis* will have been made apparent from the description of the symptoms. In advanced cases there can be no doubt. The changes in the bones, the curvature, the distortions, the appearance of the patient, the evidences of cachexia, clearly stamp the malady. Earlier in the disease it may be confounded with the manifestations of *hereditary syphilis*. But this affection comes on even sooner than rickets, almost from birth; there are other signs of the constitutional taint, including early enlargement of the spleen, syphilitic coryza, and, at a later period, the notched teeth; a distinctive history may perhaps be obtained; and the enlarged bones not unfrequently suppurate, the swollen epiphyses become detached, and osteophytes form,—changes not met with in rickets.

*Mollities ossium* produces deformities which may be mistaken for those of rickets. But the softening of the bone is the result of its disease, and not of its want of proper ossification. There is considerable difficulty in locomotion, and the bones bend or break, after having been affected with deep-seated pains. The malady lasts for years, and is not one of childhood, being most common between the ages of twenty-five and forty, and attacking chiefly women. The pelvic bones are often implicated; it is doubtful if the phosphates in the urine are increased, but, as in rickets, the urine contains lactic acid. But there are not the characteristic signs at the cranial bones, the open fontanel and sutures, nor the swelling of the epiphyses, which this malady so strikingly presents.

Some of the local deformities that result and the diseases with which they may be confounded, as of the thorax and of the head, have been elsewhere discussed. Besides the alteration of the skull in chronic hydrocephalus, the condition described by Elsaesser and others as *craniotabes* may be mistaken for ordinary rickets. It consists in thinning of the bones of the cranium, especially of the occipital bone, which becomes perforated, allowing the membranes of the brain to come in contact with the under surface of the scalp, and convulsions may be induced by undue pressure over the points of perforation of the bone. The malady, though regarded by some as a separate affection, is by others, by Virchow among them, looked upon as due to a rachitic diathesis; we certainly often find evidences of this in conjunction with the peculiar alteration of the bones of the skull.

There are cases described as acute rickets which are a combination of rickets and of scurvy.\* They are most common in infancy, and generally present the spongy gums only about the teeth that have been cut. They sometimes show, in addition to periosteal hemorrhages, a sudden protrusion of one eyeball. In the early stages rickets may be mistaken for *acute* or *subacute rheumatism*; the fever, the pain, the sweats, and the swelling near the joints mislead. But the age, the size of the epiphyses, the absence of redness of the joints and of heart-lesion, the "beading" of the ribs, the signs of beginning cachexia, the faulty dentition, and the pale urine full of phosphates, tell the true meaning of the symptoms. Moreover, the apparent joint-affection is apt to show itself at the wrist-joints, always a suspicious circumstance in delicate young children.

---

\* St. Louis Courier of Medicine, 1883, p. 453; also Barlow, British Medical Journal, 1883, i. p. 1029, and "Bradshaw Lecture," *ib.* 1894.

## CHAPTER XII.

### FEVERS.

FEVER is either a symptom of some strictly local malady or constitutes the only obvious affection present. It is only in the latter case that the disorder merits the name of essential fever. The first step, therefore, when fever has been recognized, is to determine whether it is symptomatic or idiopathic; whether, in other words, it is but a complement to a disease, or, as far as can be ascertained, the disease itself. This is not generally a difficult matter. The history of the case, the absence or presence of the marked peculiarities of serious local disturbances, soon determine the scale of evidence to rise on the one side or sink on the other. And it is astonishing, with the progress of medicine, how many affections have been passed over from the domain of fevers to the narrower circle of inflammation of individual organs; with what a different eye, for instance, the brain-fevers of the olden times are regarded. While thus the group of idiopathic fevers has been very considerably winnowed, some of their broad traits have been prominently brought forward. It is now well understood that, with some exceptions, they are characterized by the want of definite and invariable anatomical lesions. That in all constant changes occur in parts of the nervous system, or in the blood, is highly probable. But there is certainly no invariable injury perceptible in the organs of the body: sometimes one, sometimes another, suffers; sometimes nearly all; at times, none. When we contrast this with symptomatic fever, the difference is striking. The visceral lesions, then, of an idiopathic fever are not the starting-point of the fever, but rather secondary and uncertain complications influenced by and subordinate to the profound disturbance of the whole system. In idiopathic fever, the fever controls the lesions; in symptomatic fever, the lesions control the fever.



Most fevers run a definite course, showing a strong tendency to a spontaneous termination at a given time. At their beginning, too, they are for the most part similar. There is a prodromic state, marked generally by unsound sleep, pain in the back, and lassitude. This is followed by chills, which are succeeded by heat of skin, arrested secretions, quick pulse, and evident fatigue upon the least exertion. The fever has now reached its full development. Its precise character becomes evident; the symptoms caused by disorders of individual organs stand forth. After a while the disturbance declines, or speedily ceases under the influence of critical discharges. The functions are re-established, and a convalescence, more or less rapid, sets in. An unfavorable termination, on the other hand, may take place at any period after the system has been fairly invaded.

The marked features impressed upon the fever either by the course it runs, or by the specific nature of the symptoms, go to form what is called its *type*, and may be made the basis of the classification of all febrile disorders. But as opinions have been and are still singularly diversified as to what really constitute the most palpable characteristics, so the classification of fevers is as yet, to a great extent, a matter of speculation. In the following table no attempt is made at an exhaustive or strictly scientific classification. Some disorders, such as cholera, epidemic dysentery, and puerperal fever, considered by many eminent pathologists to belong to idiopathic fevers, have no place assigned to them; while others, such as influenza and yellow fever, the claims of which to be here mentioned are undoubted, might have their positions impugned. But from a diagnostic point of view the arrangement adopted is convenient, and is sufficiently accurate to be free from grave objections.

## FEVERS.

CONTINUED FEVERS.....	{	Simple continued fever.
		Catarrhal fever, or influenza.
		Typhoid fever.
		Typhus fever.
		The plague.
		Cerebro-spinal fever.
		Relapsing fever.

FEVERS.—*Continued.*

PERIODICAL FEVERS. ....	{	Intermittent fever.
		Remittent fever.
		Congestive fever.
		Yellow fever.
ERUPTIVE FEVERS.....	{	Scarlet fever.
		Measles.
		Rubella.
		Smallpox.
		Varicella.
		Miliaria.
		Dengue.
	{	Erysipelas.

## Continued Fevers.

All continued fevers are characterized by a steady progress of the febrile movement, without either decided exacerbation or relaxation, the rise and fall observable being too slight to modify the impression of a sustained action.

**Simple Continued Fever.**—Simple fever sets in with feelings of lassitude and chilliness; to these succeed hot skin, excited pulse, thirst, headache, pain in the limbs. The bowels are generally confined, the urine high-colored. The fever is soon at its height; it then either gradually declines, or is more suddenly relieved by copious perspiration or by a critical discharge from the bowels. Generally it runs through all these stages in a few days; but it may be protracted for upward of a week or longer. On the other hand, a day may witness both its beginning and its termination. The convalescence is almost always rapid.

The exciting causes of this form of fever are fatigue, errors in diet, change in mode of life, exposure to cold and moisture, or to the sun. When brought on by mental overwork or by anxiety or grief, it is not uncommonly attended with increased sensibility of the skin, and with considerable prostration, simulating typhoid fever, but differing from it by the absence of epistaxis, of the peculiar abdominal symptoms, and of the eruption. More frequently the fever has the appearance of one of high action. At times, indeed, it is so intense, and the vascular system is so wrought up, that the distemper assumes what is called an inflammatory type. It then exhibits the characteristics of the fever described by the

physicians of the last century as synocha. A temperature of  $103^{\circ}$  or upward, throbbing of the temporal arteries, severe headache, and delirium are among its symptoms. This variety of the fever is not, however, now encountered, save in tropical latitudes. In point of diagnosis, it is most apt to be confounded with internal inflammations, especially with inflammation of the brain. On the history of the case, and on the full consideration of all the symptoms before us, alone can a trustworthy opinion be based. In truth, in all the grades of what appears to be at first sight simple continued fever, we ought to examine carefully all the organs and see whether the symptoms may not be wholly accounted for by some visceral disturbance. And often, then, under what seems to be a very active or "ardent" fever will, on closer scrutiny, be found lurking the traits of an inflammatory lesion.

**Catarrhal Fever.**—This epidemic malady, which belongs to the idiopathic fevers, is sometimes described as a mere variety of bronchitis, because inflammation of the bronchial mucous membrane constitutes one of its most prominent symptoms. But this is not a just view. With as much reason might typhoid fever be omitted from the list of febrile maladies and described as a variety of enteritis or of diarrhœa.

Catarrhal fever, or influenza, is essentially an epidemic disease, and one which has visited the human race from remote antiquity. Its history is thus not confined to any particular time or to any particular nation. Its cause is believed to be a slender bacillus, with rounded extremities, two or three times as long as it is wide, which is to be found in the expectoration and nasal secretion.\* But its bacillary origin, though very probable, has not been actually demonstrated. Each epidemic does not furnish precisely the same train of symptoms; but they all agree in this: the disorder sets in suddenly and attacks pre-eminently the mucous membranes. Generally it is the mucous membrane of the nose, eyes, and bronchial tubes that suffers most, and we find the signs of coryza and of bronchial inflammation,—a watery eye, sneezing, uneasiness about the throat, and a tormenting cough. But associated with these are great depression of spirits and usually an extraordinary amount of lassitude and impairment of strength,—

---

\* Pfeiffer, *Zeitschrift für Hygiene und Infektionskrankheiten*, March 3, 1893.

much more than the cold in the head, or the laryngitis, or the bronchitis, will account for. The skin is hot, at times covered with perspiration; the thermometric record is peculiar only in its extreme irregularity. The temperature generally ranges between  $100^{\circ}$  and  $102^{\circ}$ , or starts up suddenly to  $104^{\circ}$  or  $105^{\circ}$  and in less than a day subsides almost to normal; the pulse is of moderate volume, the tongue coated; the patient complains of debility, of headache, of aching pains in his back and limbs, and of constriction at the lower part of the chest. Often there is some dyspnoea, as well as epistaxis, hyperæsthesia, especially of the neck and head, and disturbance of the alimentary tract, evinced by loss of appetite, nausea, and vomiting, or by diarrhoea; at times catarrhal jaundice coexists. Commonly after three or four days these symptoms begin to subside, the cough and debility outlasting the other morbid signs. With reference to the cough, we are often struck by the fact that its obstinacy and violence are not associated with adequate physical signs of disorder. It is often very dry and harassing.

But all epidemics do not run precisely this course. In some, the prostration is not so evident, and the febrile signs are more active and of an inflammatory type; in others, the pain and soreness in the limbs and in the joints are the most prominent symptoms; or we may find hemiparesis, or torpor and delirium, or parotitis with salivation, or otitis, or epistaxis, or jaundice, or capillary bronchitis, or pneumonia, or tendency to heart-failure, or meningitis, basilar or spinal, and irregular rashes, as complications. Many complications and sequelæ of influenza have been observed. Among these are inflammation of brain, cord, and nerves, and of the meninges, as well as various psychoses and neuroses, bulbar palsy, acute ascending paralysis, hemiplegia, diabetes, vascular occlusion, gangrene, angina pectoris, and painful and inflammatory affections of tendons, fasciæ, joints, periosteum, and bones.

The lung complication of influenza is striking. It is mostly an intense congestion, with bronchitis and here and there with spots of consolidation. True lobar pneumonia is much rarer. The lung affection may be of long duration, showing the record of a fever with marked rises and remissions. After declining, the temperature may become subnormal and remain so with occasional exacerbations for a long time, as seen in the accom-






Fig. 76.

Temperature in a case of influenza with intense congestion of the lungs.

panying chart of a case in my ward at the Pennsylvania Hospital.\*

Influenza is not ordinarily in itself a fatal disease. It is only so in the very young or the very old. It is also a grave malady in persons with weak hearts. A source of danger is the indurated lung it may leave behind becoming the seat of tubercle.

Catarrhal fever is easily discriminated from other maladies. Its peculiar epidemic character and the prostration prevent us from mistaking an ordinary cold or bronchitis for it. Occasionally the attending debility makes it look like the onset of a low continued fever. But brain-symptoms are present only in rare instances in influenza; and, on the other hand, decided catarrhal symptoms are not common in typhoid and typhus fevers. Before long, too, the occurrence of the eruption of these diseases clears up whatever doubt may have existed. The all but constant absence of an eruption in influenza comes also elsewhere into play: it serves to distinguish this disorder from measles and smallpox.

Catarrhal fever may be mistaken for *hay-fever*. But the local symptoms of irritation of the nostrils, the watery eyes, and the reddened conjunctivæ are very striking, and the febrile movement is generally less than in catarrhal fever. Moreover, there are asthmatic symptoms in hay-fever or hay-asthma in a certain proportion of cases; and the history of the case; the manner in which it comes on as a rose-cold in the latter part of May or early in June, or as autumnal catarrh after the middle of August; the hereditary idiosyncrasy so often seen; the persistence of the attack while exposed to the peculiar vegetable emanations that give rise to it; its almost abrupt cessation on removal to certain localities,—make up a set of features which are very distinctive.

When influenza is prevailing on a large scale, it is often found masked by other diseases, and it may be difficult then to separate its manifestations from those of the malady it accompanies. Other peculiarities of influenza are the long time it takes the patient to regain his strength, and the annoying sweats that attend convalescence. This was very striking in the epidemic of the early winter months of 1890; as was also the tendency to relapses, to irregular heart action, and to alterations of cutaneous sensibility.

---

\* For a further description of the lung complications of influenza, see my paper on the subject in the "International Clinics," Vol. I., Second Series.

**Typhoid Fever.**—In this country and on the continent of Europe a form of continued fever prevails, marked by great prostration and disturbance of the nervous system, and by constant anatomical lesions. To this disease the designations of typhoid fever, enteric fever, and abdominal typhus have been applied.

The disorder either attacks single individuals or shows itself as an epidemic. It occurs at all seasons of the year, but, in this country at least, is most frequent in autumn. In some localities it is thoroughly at home; in others it is only occasionally seen. It avoids both extremes of age, seizing mainly on young adults for its victims.

The distemper may set in suddenly, but more generally it has an insidious beginning. For some days preceding the access of the fever the patient feels weak. He is without animation, and his countenance fully expresses his languor. He complains of soreness and fatigue, of dull pain in the head, of loss of appetite. His sleep is unsound; all exertion is wearisome; something is evidently weakening his nervous energies. A fever now appears, preceded mostly by a chill, or, at all events, by chilly sensations, which alternate with flushes of heat. The muscular prostration accompanying the febrile movement is so great that the patient is obliged to seek his bed. His appetite is entirely gone, the tongue is coated, the bowels are loose, the abdomen is somewhat swollen and tender to the touch. On close inspection, a few reddish spots, resembling flea-bites, are found on its surface.

The malady has now completed its first week. It enters on the second week with fever unabated, and with the signs of disturbance of the alimentary tract and of the nervous system more and more unmistakable. There is sometimes nausea or epigastric distress, often pain in the right iliac fossa, increased by pressure and tympanites. The tongue dries and becomes reddish or brownish; it is often glazed and covered with a light coat; sometimes it has deep fissures; very frequently I have noticed at the tip a wedge of brownish or reddish surface free from coat, but which begins to be covered over as the disease declines; the gums and teeth are lined with dark crusts. The mind is dull and wandering; cough and great restlessness exist; the debility is extreme.

The disease now begins to draw to its close. It has reached the third week, and a change, for better or for worse, may be



looked for. Slowly recovery sets in, marked by a brightening of the countenance and by a gradual increase in consciousness and strength; or deepening insensibility, jerking of the tendons, feeble pulse, and cold, clammy sweats indicate that dissolution is fast approaching.

Thus, in one way or the other, the fever itself is apt to terminate by the beginning or the middle of the fourth week. Yet such is not always the case. Death may take place at an earlier period; or, on the other hand, the malady, by troublesome complications, may be lengthened beyond the second month. Under any circumstances, convalescence is protracted. The nervous system rallies but gradually from the shock it has received.

Among the symptoms enumerated, some tend clearly to characterize the disease. And, first, of the more purely *febrile* symptoms. The heat of the skin is especially perceptible in the evening exacerbations of the fever. Frequently the surface is covered with an acid perspiration, very manifest during the whole course of the disorder, and also encountered long after convalescence has set in. The pulse is accelerated, and remains so after the heat of the skin has left; it is very compressible, and even in intercurrent acute inflammations it seldom loses its compressibility. A jerking, irregular beat, or very great rapidity, is an unfavorable sign. Dicrotism of the pulse is not unusual. Associated with the diminished strength of the pulse is a decided faintness of the first sound of the heart.

The *temperature* is peculiar; in the first five or six days of the disease it pursues an ascending line; that is to say, starting at the normal  $98.6^{\circ}$ , there is apt to be a daily evening rise of about  $2^{\circ}$ , with a morning remission of about  $1^{\circ}$ . From the fifth or the sixth day to the twelfth or a little later,—roughly speaking, we may say from the end of the first week to the end of the second,—the fever is continuous, with a morning remission rarely exceeding  $1^{\circ}$ . From that time on, let us say from the twelfth day, although the evening temperature may remain for a day or two quite or nearly as high, there is an abatement of from  $1^{\circ}$  to  $2^{\circ}$  in the morning. These changes between morning and evening become very evident at the end of the week, and are still more evident in the third week, when the morning and evening temperatures may vary as much as from  $4^{\circ}$  to  $6^{\circ}$ . During this week, too, the evening temperature gradually decreases; but in severe



cases it remains high, and there are no decided remissions, either in the second or the third week. The morning temperature is high,  $104^{\circ}$  or more, and there may be still greater heat of skin in the evening, or else it differs but little from that of the morning. The peripheral temperature, as measured, for instance, in the palm of the hand, becomes during the fever as high as the axillary temperature, but their equalization ceases prior to defervescence.\* In exceptional instances, it is stated, the temperature may be normal throughout;† in still rarer instances it is subnormal.‡ I have never seen a case of either kind. Occasionally the curve may resemble that of intermittent fever.§

The *urine* is acid, high-colored, scanty,—the urine of fever. In severe cases it contains variable amounts of albumin, particularly in the cases with high temperature; hæmaturia is very rare. Ehrlich has stated that the urine of typhoid fever gives a special reaction. This test consists in taking forty parts of a saturated solution of sulphanilic acid in hydrochloric acid, one to twenty, and one part of a one-half per cent. solution of sodium nitrite, and adding them to an equal bulk of urine rendered alkaline by strong ammonia. Normal urine is colored brownish by the test liquid, typhoid-fever urine pink or ruby, with slight frothing. The reaction has not been found in all cases of typhoid fever, and has been obtained in a variety of other morbid conditions, such as tuberculosis, measles, scarlatina, enteritis, malaria, pneumonia, meningitis, septicæmia, uremia.|| It has been thought to be due to the presence of bacterial products.

Among the abdominal symptoms, *diarrhœa* is the most prominent. It is never absent, except when the disease is unusually

\* Couty, Archives de Physiologie, No. 2, 1880.

† Finlayson, American Journal of the Medical Sciences, March, 1891, p. 225; Wendland, Deutsche medicinische Wochenschrift, Aug. 29, 1892; Dreschfeld, Practitioner, No. 298, vol. 50, p. 272; Fisk, Medical News, Nov. 3, 1894, p. 479; MacDougall, Lancet, April 15, 1893.


‡ Raimondi, Gazette des Hôpitaux, 1894, No. 109; Centralblatt für Innere Medizin, 1895, No. 6, p. 152.

§ MacDougall, Lancet, April 15, 1893.

|| Taylor, Lancet, May 4, 1889; Edwards, Medical News, April 2, 1892; Dawson, Dublin Medical Journal, 1894, No. 258; American Journal of the Medical Sciences, April, 1894, p. 448; Nissen, Jahrb. für Kinderheilk., B. 38; Fortschritte der Medizin, 1895, No. 5, p. 192.

mild. Generally it is a very early symptom ; at times it is even seen among the prodromes. The clue to its cause is found in the state of the intestinal glands,—in the enlargement and ulceration

FIG. 11.



Temperature in a case of typhoid fever of decided kind.

of the glands of Peyer and of the solitary glands, with the tumefaction of the mesenteric glands. And in these morbid alterations we find an explanation not only of the occurrence of the diar-

rhœa, but also of its frequency. The stools are thin, of a yellow or dark-brown color, and of offensive smell. When the affection is at its height, from three to four evacuations occur during the twenty-four hours; but the passages may become much more numerous, and with their number the danger rises. If they take place without the knowledge of the patient, his situation is precarious. Sometimes the stools contain blood. Should this be present in considerable quantity, it is a very unfavorable circumstance. Yet intestinal hemorrhage is by no means necessarily fatal. In rare instances there is hæmatemesis.\* Dickinson † has reported a case in which profuse hæmoptysis occurred, in the absence of any pulmonary lesion.

FIG. 78.



Eberth-Gaffky's typhoid-fever bacillus, from a potato culture. The broad ones are really two bacilli lying in juxtaposition. *Zeiss*  $\frac{1}{12}$ , homo. im., Oc. 5.

The poison of typhoid fever is conveyed chiefly through drinking water or defective sewerage. It is now generally thought to be the bacillus described by Eberth and by Gaffky as present in the intestinal lesions (Fig. 78) and found in the

\* Weiss, *Wien. med. Presse*, 1887.

† *Lancet*, Feb. 17, 1894, p. 421.



stools. This is a rather plump organism, from two to three  $\mu$  long, with rounded extremities, and staining with the ordinary aniline colors.

While there is a close resemblance between the bacillus of typhoid fever and the bacillus coli communis, there are certain points of distinction. Thus, the movement of the former is more active and more extensive than that of the latter. The typhoid-bacillus manifests a greater tendency to form threads of two or more than does the colon-bacillus. Colonies of the former grow much more slowly on gelatin and on agar than do those of the latter. Potato cultures of the typhoid-bacillus are scarcely visible, while those of the colon-bacillus appear as distinct, broad, orange streaks. The colon-bacillus is capable of inducing fermentation and of curdling milk, while the typhoid-bacillus is not. Flagella are more readily demonstrable on typhoid-bacilli than on colon-bacilli. Finally, the typhoid-bacillus yields the indol reaction with potassium nitrite, while the colon-bacillus does not.\*

*Enlargement of the spleen* is a very constant attendant upon the fever. In fact, whenever we can be certain that the evident increase in size is not due to some previous malady or to malaria, the extended percussion to dulness in the splenic region becomes an element of importance in our diagnosis. The tympany that often exists interferes with the recognition of the enlargement.

Another abdominal symptom of significance is *pain*. It varies much in severity and character, and is, indeed, not always present. It is often a heavy, aching feeling. In some patients it is of a griping kind, preceding the loose discharges; in others it seems to be called into existence only by pressure. Its most common seat is in the iliac fossæ; yet the testimony of the sick man himself as to its exact situation must be received cautiously. He is too ill to answer intelligently. Still, the expression of suffering on his face when pressed on either side at the lower part of the abdomen is indicative of the pain corresponding, for the most part, to the seat of the irritation. In rare instances the pain is really in the muscles, which may, indeed, suppurate.† Often, while the hand is exploring the abdominal regions, a movement

---

\* Archiv für Hygiene, vol. xix., fasc. 3.

† Ebing, Archiv für klin. Med., viii.



of the fluid and gas in the distended bowel, attended with a gurgling noise, becomes appreciable. This sign is best elicited near the ileo-cæcal valve.

Attention has been called to a yellowish discoloration of the palms of the hands and soles of the feet in cases of typhoid fever, and not, as a rule, observed under other conditions.\*

During convalescence, griping pains are not infrequently complained of. They are colicky pains, produced generally by errors in diet, and may be followed by a return of the diarrhœa or by a relapse of all the other symptoms of the malady. Occasionally during the latter period of the fever a sudden pain sets in, of great intensity, unremitting, and attended by spreading tenderness. Such a pain shows that peritoneal inflammation has been lighted up in consequence of perforation. This complication is not necessarily fatal, as the morbid process may be limited by inflammatory adhesions. In a number of instances the abdomen has been opened, the perforation found and closed, and recovery has ensued.† The peritonitis and intestinal adhesions that follow perforation may be attended with symptoms of obstruction of the bowel.‡

Hardly inferior to the abdominal symptoms in import are the signs of *disturbance of the nervous system*. The fever is, as its old name implies, pre-eminently a "nervous" fever: the nervous symptoms are, in truth, never absent; but, though always present, they are less extensive in some cases than in others, and not the same throughout all the stages of the disease. Thus, early in the disorder, dull headache, mental languor, wakefulness, and a perverted state of the senses, such as ringing in the ears and dullness of hearing, are encountered; while later, great restlessness, delirium, somnolence, or coma, and jerking of the tendons are phenomena more likely to be met with. Occasionally the disease is ushered in by acute mania.§

\* Filipovitch, *Lancet*, Aug. 19, 1893. See also *Medical News*, Oct. 4, 1893, p. 444.

† Van Hook, *Medical News*, Nov. 21, 1891, p. 591; Sifton, *Chicago Clinical Review*, vol. iv., No. 7, p. 368.

‡ Blaikie Smith, *International Clinics*, vol. i., 2d Series, 1892, p. 79.

§ Hare and Patek, *Medical News*, June 20, 1891, p. 681; MacDougall, *Lancet*, April 15, 22, 1893.

The *delirium* sets in generally during the second week, for the most part at night, and terminates with convalescence or ends in coma. It is not a wild delirium, but a confusion of mind associated with rambling thoughts. If the patient's attention be strongly engaged, he may be almost always roused, and does for a time as he is told; but after a short interval his muttering lips indicate that some curious fancy has again taken possession of him. In some cases, not in many, the delirium is attended with great restlessness and agitation, and the sick man, if not prevented, attempts to walk about the room. This kind of frenzy often ends in fatal coma. Equally unpromising is early or unremitting delirium. When contrasted with the mental wandering in other acute disorders, the delirium of typhoid fever exhibits peculiar traits. It is ordinarily more active than that of typhus; far less demonstrative or talkative than the mania of drunkards; as aimless as, but less continued than, the ravings of inflammation of the brain. Great restlessness and tremors, associated with a clear mind, and at times with copious perspirations, have a very significant meaning: they point to deep and extending ulceration.

Other symptoms of grave disturbance of the nervous system show themselves in violent general *convulsions*. These are more common in children than in adults, in whom they may be a late symptom; they may or may not be of uræmic origin. As a rule, the knee-jerks are present, unless peripheral neuritis exists. In severe cases both the reflexes and the muscular irritability are said to be increased.\* On the other hand, it has been observed that in children the tendon-reflexes are often enfeebled during the acute stage of the disease and exaggerated during convalescence.†

In some cases of typhoid fever symptoms not only of cerebral but also of spinal origin appear, and they may, indeed, assume a high degree of intensity. We find extensive cutaneous hyperæsthesia, spinal pain and tenderness, with a sense of pricking along the vertebral column, and, in some instances, cutaneous and muscular anæsthesia, numbness of the extremities, partial paralysis or convulsive contractions of the respiratory muscles, convulsive

---

\* Angel Money, *Lancet*, Nov. 7, 1885, p. 842.

† Albouze, *Journal de Médecine et de Chirurgie Pratiques*, Sept. 10, 1892.

cough, paralysis of the sphincters, contractions of the extremities, and even rigidity of the muscles of the neck.\* These spinal symptoms are more common when the disease is epidemic than when it is sporadic, and are always indicative of a very serious form of the disorder. They sometimes persist after the fever has left, or indeed—and this is especially true of paralysis—may not appear until convalescence. The *palsy*, the most common form of which is paraplegia, mostly begins gradually and disappears gradually. It may be preceded by trembling movements, suggesting the idea of disseminated sclerosis; but the tremor is rather the result of general debility, and, unlike sclerosis, it occurs before, and does not attend or follow, the loss of muscular power in the limbs, and is not associated with difficulty of enunciation.

Hawkins† has reported a fatal case of typhoid fever complicated by intestinal hemorrhage and purpuric spots and the development of right hemiplegia and aphasia due to occlusion of the left middle cerebral artery. He has collected seventeen cases of typhoid fever complicated by hemiplegia. Most often the palsy was right-sided and associated with aphasia. Usually there was recovery from the paralysis, but in three cases this was persistent. The complication was most common at the end of the attack or during convalescence. There is much evidence that the paralysis after typhoid fever is due to neuritis.‡

Two other prominent symptoms of the malady must still be inquired into: one is epistaxis; the other, the cutaneous eruption. *Epistaxis* is not often absent in grave cases. It may happen at any period of the complaint; but it generally takes place before the disorder is far advanced. The quantity of blood lost is rarely considerable; and for this reason the occurrence of the hemorrhage is frequently overlooked.

The *eruption* peculiar to the disease is the rose-colored rash. It appears on or shortly after the seventh day, but occasionally not until the end of the second week. It can hardly be called a papular eruption, as it consists rather of small, red spots, only

---

\* Fritz. Étude clinique sur divers Symptômes spinaux observés dans la Fièvre typhoïde, referred to in Arch. Gén. de Méd., June, 1864.

† Transactions of the Clinical Society of London, vol. xxvi., 1893, p. 50.

‡ Pitres and Vaillard, Rev. de Méd., 1885, t. v.; Ross, Amer. Journ. Med. Sci., Jan. 1889; Bury, Medical Chronicle, June, 1892.

very slightly elevated above the skin, somewhat similar to flea-bites, yet differing from them in lacking the central mark and in their finer, paler color and less obvious outline. The spots are seen upon the abdomen and chest, rarely upon the extremities, almost never upon the face. They disappear totally on strong pressure, yet return immediately when the pressure ceases. They are generally few in number, and not persistent. Each spot does not last for more than three or four days; then it fades, and a fresh one near by replaces it, and runs the same course. Spots thus appear and pass away for more than a week, after which, in most cases, they entirely vanish. During convalescence not a trace of them can be found; but should a relapse take place, they reappear with the other symptoms of the malady. This eruption, although very common, is not invariably present; at all events, it is not invariably found. Beyond doubt, too, it is in some epidemics more constant and marked than in others. Late in the disease another eruption appears, consisting of minute transparent vesicles, scattered plentifully over the body. These sudamina are not so frequently encountered as the rose-rash, and are certainly not so characteristic.

After convalescence has set in, we may have a return of fever. It may be either a transitory and slight return, due to fatigue or to some indiscretion in diet, or a more protracted state, in which most or all of the symptoms peculiar to the disease reappear. Thus, typhoid fever *relapses* usually come on in the second week of assured convalescence, and, according to my experience,\* occur suddenly; soon diarrhoea, furred tongue, and enlargement of the spleen are manifest, and on the fourth or fifth day reappears the characteristic rose-rash, which is often somewhat coarser than in the first attack, and does not show the same disposition to appear in successive crops. With the eruption delirium is apt to come back. The temperature is unlike that of the original attack in quickly reaching a high point of fever-heat; after the first day or two it remains more or less stationary, with a slight morning fall, for five or seven days usually, and then shows the well-known remissions and rises of the zigzag decline. The pulse is often noted to be dicrotic.† The relapse is in its duration usually much

---

\* See article on Relapses of Typhoid Fever, Transactions of the College of Physicians of Philadelphia, 1877.

† Steinthal, Arch. f. klin. Med., Feb. 1884.



shorter than the original attack, and generally, notwithstanding the threatening appearance of the symptoms, ends in convalescence. During its progress intestinal hemorrhage may happen; and after return to apparent health a second relapse or more may occur. Each relapse occasions characteristic markings on the nails, from impaired nutrition, which Longstreth has very fully described.\* Ziemssen specifies the fifth, seventh, and fourteenth days after the cessation of the original fever attack as the days on which a relapse is likely to happen.† The temperature sometimes keeps up a degree or two, while the patient is in every other respect fully convalescent, yet will come speedily to the norm if he be made to leave his bed.

Both during the height of the fever and in convalescence, but more especially during the latter, certain complications or sequelæ may arise, some of which are medical, such as parotitis, erysipelas, noma, laryngeal ulceration or stenosis, milk-leg, thrombosis of the femoral artery, the result of arteritis,‡ cirrhosis of the liver, periostitis, osteomyelitis, and transitory aphasia;§ while others, as dislocations, caries, necrosis of bones, epididymitis,|| abscess, and gangrene, come within the domain of surgery.¶

Sometimes sequelæ appear long after the primary disease has come to an end. Orlow\*\* has reported a case in which five and one-half months afterwards typhoid-bacilli were found in a granuloma of the tibia. Péan and Cornil†† observed a case in which five months after an attack of typhoid fever typhoid-bacilli were found in the lesions of a suppurative periostitis. Sudden death may take place in the course of typhoid fever as a result of disturbances in the circulation, from the formation of blood-clots, from inflammatory and degenerative changes in the muscular wall or in the nervous supply of the heart, or from

\* Relapses of Typhoid Fever, Transact. Coll. of Phys. of Phila., 1877.

† Arch. f. klin. Med., Feb. 1884.

‡ Lucas-Championnière, Journ. de Méd. et de Chir. Pratiques, 1888.

§ Arch. f. klin. Med., Bd. xxxiv., 1, 1883.

|| Girode, Archives Gén. de Médecine, Jan. 1892, p. 43.

¶ See an elaborate discussion of these surgical complications, by Dr. W. W. Keen, Fifth Toner Lecture, Washington, 1877.

\*\* Deutsche medicin. Wochenschrift, Nov. 27, 1890.

†† Bull. de l'Académie de Médecine de Paris, Apr. 14, 1891.

the poisoning of the system that is an essential part of the disease.\* Death has also resulted from profuse sweating.†

After this analysis of the symptoms of typhoid fever, it would be useless repetition to discuss at length how the disease differs from all other idiopathic fevers. The attempt will rather be made to explain its diagnosis from those maladies, whether essentially febrile or not, to which it bears the closest resemblance. And here we find that the disorders with which typhoid fever may be confounded are not the same at all the stages of the complaint. Early in the affection it is most likely to be mistaken for simple continued fever, or for one of the exanthemata. But diarrhœa is not present in these, nor are there marked prodromata; and whatever doubt may exist with reference to simple continued fever is cleared up in a few days, as the temperature-record is different and as the symptoms come to an end at a time at which in typhoid fever they begin to be more and more developed. Still, the exanthematous fevers cannot, before their eruptions appear, be distinguished with absolute certainty; though we may suspect measles by the attending coryza, scarlatina by the sore throat, and smallpox by the lumbar pains and high fever.

At a more advanced period, typhoid fever may be confounded with typhus, and with these morbid states:

GENERAL DEBILITY;

TYPHOID CONDITIONS;

ENTERITIS;

PERITONITIS;

MENINGITIS;

ULCERATIVE ENDOCARDITIS;

ACUTE PULMONARY AFFECTIONS.

*General Debility.*—It does not seem likely that so acute and dangerous a malady as typhoid fever could be mistaken for mere debility; yet such an error may occur when the disease is latent, or so light as not to confine the patient to his bed. In these so-called “walking cases” the debility, however, sets in suddenly, and not gradually, as in weakness from general constitutional causes.

---

\* Dewèvre, Archives Générales de Médecine, Oct., Dec. 1887; Galliard, *ib.*, May, June, 1891.

† Juhel-Renoy, Archives Générales de Médecine, 1886, vol. i. p. 274.

Moreover, the abdominal symptoms are rarely wanting, and there is more or less confusion of mind. Due attention to these circumstances will prevent mistake; but the greatest safeguard against error is to be aware that the disease assumes at times a latent form, and to examine every case of sudden debility, to see if under its mask are hidden the features of typhoid fever.

*Typhoid Conditions.*—No blunder is more common than to misconstrue into typhoid fever a typhoid condition of the system. We may find this condition in many different complaints, both acute and chronic; but more especially are purulent infection, some forms of pneumonia, dysentery, and erysipelas attended with delirium, drowsiness, dry, brown tongue, and extreme prostration, —in one word, with a typhoid state.

Yet a typhoid state is not typhoid fever; it is simply a low condition of the system which may be present in many dissimilar maladies, and which is present in its most perfect form in typhoid fever. But in this complaint we have other signs than those of vital depression: we find joined to it diarrhoea, tympanites, epistaxis, an eruption, and special manifestations of disturbance of the nervous system,—all symptoms bearing no direct relation to the adynamia, and thus serving as valuable distinctive marks. An examination of the urine, too, is often of signal service; though we must not forget that in grave cases albuminuria to a moderate degree is present. And there are cases of Bright's disease and of abscess of the kidney in which the poisoning of the blood that happens occasions a deceptive likeness to typhoid fever, so deceptive that only a minute examination of the urine can fully explain the true meaning of the symptoms. The following case well illustrates this:

A man, about forty-five years of age, was admitted into the Philadelphia Hospital in January, 1863. He was very prostrate, and hardly able to give an account of himself. It was ascertained that he was not a person of intemperate habits, and that he had been attending to his work until within two weeks. He was evidently stupid, and, when questioned about himself, seemed to have great difficulty in remembering, and in collecting his thoughts. He had fever; a pulse above 100; a dry, brown tongue. The heart-sounds were feeble, the heart increased in size. The urine was at times turbid, and contained a slight, whitish sediment,

which was not, however, examined with the microscope. His mind wandered at night; the abdomen was distended and in parts slightly tender; several doubtful red spots were detected on its surface. In fact, he appeared to have almost every one of the more constant symptoms of typhoid fever, except the diarrhœa. A few days after his admission he became comatose, and sank. The intestinal glands were found in a healthy condition; but both kidneys were thoroughly disorganized and filled with pus.

What exactly produces the typhoid state it is difficult to say. Milner Fothergill\* connects it with tissue-waste without increased renal activity, and with the accumulation in the blood of the products of the tissue-waste.

*Enteritis.*—The great difference between enteritis and typhoid fever consists in this: in enteritis the inflammation of the intestine constitutes the disease; there are no symptoms other than those referable to the inflamed intestine. We find no great prostration; no mental wandering; no enlargement of the spleen; no rose-spots; no signs of abnormal processes due to a typhoid dyscrasia. The disorder, too, gives rise to much more abdominal pain, and is of shorter duration. In certain rare cases the follicles of the intestines are inflamed and swollen, and the attending febrile malady may closely simulate typhoid fever, without, however, its characteristic intestinal lesions, or eruption, though with considerable diarrhœa and swelling of the spleen.† Again, I have known fœcal accumulations in the intestine to produce and keep up diarrhœa and continued fever of several weeks' duration similar to that of typhoid, and ceasing only when the large fœcal masses were voided. The absence of eruption, of cerebral symptoms, and of enlargement of the spleen proved the points on which the correct diagnosis of the non-existence of typhoid fever was based.

*Peritonitis.*—The same remarks apply to peritoneal inflammation. Here, moreover, the expression of the face, the constipation, and the great abdominal tenderness serve as marks of discrimination. But we must not forget that acute peritonitis may appear in the course of typhoid fever. Generally this untoward event happens at a late period of the disease, and after the patient has

---

\* Edinburgh Medical Journal, September, 1873.

† Cazalis and Renaut, Archives de Physiologie, 1873.



been under observation for some time ; we are then at no loss to understand the meaning of the spreading tenderness, the rapid, small pulse, the marked tympanitic distention, the sweats, the nausea and vomiting, the collapse, and the pinched features. But the accident may occur in cases that we have not previously seen, or in which the affection has run so latent a course as hardly to have attracted even the patient's attention. The cause of the peritonitis is then commonly first revealed by the autopsy, which shows actual perforation of the intestinal walls, in consequence of ulceration of a solitary or an agminated gland. Whenever, indeed, in typhoid fever the signs of peritonitis can be clearly traced, the exciting cause of the inflammation may be announced to be perforation ; for the evidence on which it has been assumed that peritoneal inflammation may take place without the giving way of the intestine is not so positive as to cause us to abandon this diagnostic rule.

*Meningitis.*—Typhoid fever has some symptoms in common with inflammation of the brain ; but the signs of difference have been fully discussed in connection with acute meningitis, and need not here be examined. And in rare cases we really have meningitis as a complication of typhoid, showing small pupils, vomiting, and rigid neck. The distinction from epidemic cerebrospinal meningitis we shall presently trace.

*Ulcerative Endocarditis.*—In some cases the differential diagnosis between this and typhoid fever becomes of great difficulty, especially if the case be not seen until the endocarditis has led to delirium and the symptoms of collapse. Recurring chills, with high temperature and sweats, as in malarial fever, great rapidity of pulse, with sudden changes and marked irregularity, a generally-diffused roseolous eruption, and the signs of the cardiac lesion, form the most trustworthy points of distinction.

*Acute Pulmonary Affections.*—In the majority of cases of typhoid fever we find cough, dependent upon an affection of the bronchial tubes. The bronchial affection gives rise to extreme loudness of the rales, with a cough disproportionately slight ; sometimes, too, owing to the blood gravitating to the most dependent portions of the lungs, the resonance over the posterior part of the chest is impaired. From these phenomena, added to the abdominal and cerebral symptoms of the fever, and the vital depression, there is

no difficulty in discriminating between idiopathic *bronchitis* and typhoid fever.

Not infrequently we find a dry *pleurisy* combined with the bronchitis, and in some cases, not in many, the cough is associated with exudation into the pulmonary structure. Now, it may be extremely difficult to distinguish a *pneumonia* of this kind from inflammation of the lung setting in amid signs of prostration, until the eruption and the abdominal symptoms solve the difficulty. Generally, however, it is not a matter of much doubt, as the condensation of the lung in typhoid fever does not occur early in the disease,—not, in fact, until the symptoms of the fever are clearly developed. Occasionally a cough remains after the febrile symptoms have begun to decline and the mind is regaining its clearness. The cough increases in severity, and the patient soon loses the strength he may have acquired. On listening to the chest, we find scattered over both lungs many fine, dry and moist sounds. The percussion-note is here and there dull; the expectoration is profuse; there are dyspnoea and excessive sweating. Here is a group of signs bespeaking *acute phthisis*. We may also observe acute phthisis with most of the symptoms of typhoid fever; even the delirium, the stupor, and the enlargement of the spleen may be present; but the eruption never is, and the diarrhoea rarely. In general acute miliary tuberculosis the similarity is even greater, and diarrhoea is not uncommon; the disease is, as a rule, longer. Tubercle-bacilli may or may not be present in the sputum; they have been detected in the urine and in the blood; when present they enable us to make a positive diagnosis. In rare instances the two diseases have coexisted.

In concluding the subject of typhoid fever it will be proper to notice those forms of the affection which run their course in a different manner from that ordinarily pursued by the malady,—the *mild typhoid* and the *abortive typhoid*. The former has usually a gradual beginning, and the disease throughout remains mild; its duration may be, however, the same as, or even longer than, that of ordinary typhoid, or it may be considerably shorter,—in fact, an abortive typhoid, the variety of typhoid to which of late years Jürgensen especially has directed attention.\* Yet the

---

\* Sammlung klinischer Vorträge, No. 61, 1873. See also paper by Johnston, Amer. Journ. Med. Sci., Oct. 1875.

abortive type is not always mild : cases are mentioned\* in which the temperature rose to  $106^{\circ}$ , but in which the duration of the fever was only from seven to twelve days. Indeed, it is the short duration that is characteristic of *abortive typhoid*. As a rule, it begins suddenly, and the temperature reaches its highest point on the second or third day. It often does not exceed  $104^{\circ}$ , and it stays at, or near, the height it has so speedily attained for the greater part of the duration of the fever, and then remissions show themselves, and there is a gradual return to a healthy standard, much in the same way as at the end of ordinary typhoid fever ; or the changes are so marked and rapid that the defervescence is accomplished in a few days. The symptoms of typhoid fever are all met with in the abortive malady, though they are not present with the same constancy ; tenderness in the right iliac fossa is the most frequent ; enlargement of the spleen and the rose-colored spots are very usual ; diarrhœa is often wanting. The disease terminates in sixteen days or less ; but there is great proneness to relapses. It is not apt to be a fatal affection.

**Typhus Fever.**—This is a highly-contagious malady, almost always met with in an epidemic form, and generally among those whose systems are depressed. It prevails in jails and camps, among crowded populations, or in badly-ventilated localities, and has no constant structural lesion. In this country it is a rare disease. It is either preceded by a brief stage of lassitude and dejection, or is ushered in with a chill and pain in the head and back. The skin soon becomes dry and of pungent heat ; the pulse rises much in frequency, and is at first full, sometimes even tense. The patient lies in a state of half-consciousness, dull, drowsy, weak, with evident signs of his nervous and muscular system being overwhelmed by the influence of some fearfully-depressing poison. The face is flushed ; the odor from the body extremely unpleasant.

By the fifth day all these symptoms are plainly marked, and about this time a coarse, red, cutaneous eruption makes its appearance. But it occasions no change in the gravity of the symptoms. On the contrary, the confusion of mind and the stupor increase ; the patient wanders, picks at his bedclothes, and ceases to com-

---

\* Liebermeister, in Ziemssen's *Cyclopædia*.

plain of the pain in the head or limbs. The pulse is frequent and feeble; the tongue dry and dark; sordes collect on the gums and teeth. The bowels remain as they were at the onset,—constipated. The urine often comes away drop by drop, or, as the bladder loses the power of contracting, is retained. The case has now reached its height; the signs of a prostrated nervous system, of deteriorated blood, and of utter loss of muscular strength either begin to pass away, or deepen from hour to hour and clearly show the doom that awaits the fever-stricken patient. From the beginning of the distemper until the unfortunate issue is rarely over thirteen days. If the sick man can withstand the poison until the third week, he is apt to throw it off and recover; but it may be so virulent as to overpower him almost at the onset.

Micro-organisms have been found in the blood in cases of typhus fever. Thus, Brannan and Cheesman\* describe a club-shaped bacillus from 0.5 to 0.8  $\mu$  in diameter, and from 1 to 2.5  $\mu$  long. Dubief and Bruhl† have found an organism that they designate “diplococcus exanthematicus.” Lewaschew‡ describes cocci from 0.2 to 0.5  $\mu$  in diameter.

Let us examine some of the symptoms of this pestilential disease in detail.

The *physiognomy* of typhus is peculiar. The expression is stupid, and coarser than in health. The face shows a deep flush of a dusky-red hue. The eye is injected, the pupil often contracted. The skin is covered with a characteristic *eruption*, from which the disease takes its name of “spotted” or “maculated” or “exanthematic” typhus. The rash is at first slightly elevated and much like that of measles. It is of a dark tint, a “mulberry rash,” and fades but does not vanish on pressure. It makes its appearance from the fifth to the seventh day, and is permanent, consisting not of successive eruptions, but of the same spots, which deepen or lighten with the changes in the disease, and do not pass away before the fourteenth day. Each spot thus lasts until recovery or until death, and no new ones show themselves after the second or third day of the rash. They are numerous on

---

\* Medical Record, June 25, 1892.

† Universal Medical Journal, May, 1893.

‡ Deutsche medicin. Wochenschrift, Nos. 13, 34, 1892.



the trunk and the extremities, but are rarely observed upon the face. Some are much lighter than others, and thus a mottled aspect of the skin is produced, on which Sir William Jenner \* lays great stress. Sometimes the spots are of purple color and uninfluenced by pressure. These petechiæ are the attendants of the worst forms.

The skin of a typhus-fever patient is often sensitive, and, as already stated, generally very hot. In some cases the thermometer indicates a *temperature* of  $107^{\circ}$ , or more; commonly it ranges above  $104^{\circ}$ . The heat is sustained: it does not show the decided differences between morning and evening that are observed in typhoid fever, the daily variations to the middle of the second week being rarely  $1^{\circ}$ ; and from that time onward the morning abatement does not amount to more than about  $1.5^{\circ}$ , until the defervescence is reached. The passing away of the high temperature occurs, however, not, as in enteric fever, by more and more evident remissions, but suddenly. Early in or towards the middle of the third week the temperature falls quickly, and in twenty-four or thirty-six hours a normal standard is reached. In rare instances, it is asserted, the temperature may not rise above the normal, or it may even be subnormal.†

The *cerebral* symptoms of typhus fever are never absent. In some epidemics they constitute the prominent feature of many cases, and dangerous and fatal these cases are apt to be. One of the most frequent proofs of the disturbance of the brain is seen in stupor. The patient lies in a heavy slumber, occasionally muttering some incoherent words; or he is sleepless, his eyes remain wide open, he has *coma-vigil*, he takes no notice of anything going on around him. Either of these states may deepen into coma. In other cases delirium is the most conspicuous symptom. This delirium rarely sets in before the end of the first week. In type it is low and muttering, and unaccompanied by great restlessness; or it may be associated with constant movements and trembling of the limbs, or jerking of the tendons,—in fact, with symptoms resembling those designated as hysterical. Sometimes

---

\* Identity or Non-Identity of Typhoid and Typhus Fevers, London, 1850; and Medico-Chirurgical Transactions, vol. xxxiii.

† Combemale, Gazette hebdom. de Médecine et de Chirurgie, 1893, No. 30, p. 352.

the mental wandering is active and very persistent. The patient tosses about, is constantly talking, and can hardly be restrained from getting out of bed. He has illusions of hearing and of sight; his eyes are injected, the pupils often contracted; there is headache, with intolerance of light. Here we have the true brain typhus, with its formidable cerebral symptoms simulating closely those of *acute meningitis*, and differing only by their union with a cutaneous eruption, by the dissimilar aspect of the tongue, and by the beat of the pulse, which is rarely full, and never so tense as that of meningitis. Then, the nervous excitement is accompanied, or, at all events, soon succeeded, by greater and more rapid prostration of strength, and is often exchanged far more suddenly for coma than is observed in the meningeal disorder.

The headache which has just been mentioned is a very constant symptom: usually it is most severe during the first week, and abates with the appearance of the mental wandering. Often it is accompanied by more or less giddiness, which increases with the progress of the disease.

These head-symptoms of typhus are, like those of enteric fever, sometimes connected with a noisy, shallow, and irregular respiration. This kind of breathing can be clearly traced to the abnormal state of the nervous system, as no signs of alteration in the lungs coexist. Often, as Flint\* has pointed out, it is a forerunner of fatal coma. In one case I found the strange phenomenon associated with great distention of the bladder, and subsiding materially after the introduction of a catheter.

The remarks with reference to the cerebral phenomena of typhus apply to those instances in which there is no inflammatory disorder within the cranium. But we must not overlook the fact that this may ensue. Such cases are difficult of recognition. The pulse, as a rule, is slow and irregular, the pupils are contracted, there is a frown on the forehead, and intense headache, sometimes screaming. Vomiting is not always encountered. We may find with these symptoms acute meningitis, and the morbid appearances may be confined chiefly to the base of the brain.† There are other symptoms referable to the nervous system that are occa-

---

\* Clinical Reports on Continued Fever. .

† Kennedy, Dublin Quarterly Journal, Feb. 1867.

sionally very marked, such as great agitation, rigidity of certain muscles, and convulsions. But as regards the latter, the nervous system is for the most part only secondarily disturbed, for the convulsions are generally of uræmic origin.

The *pulse*, after the disease is fully developed, is generally rapid, and of moderate volume or feeble. As the disorder advances the pulse rises in frequency, while it diminishes in force. As convalescence is established, it falls; if it remain frequent, this is generally indicative of some concealed visceral disorder, often of a disease of the lungs. It does not always correspond closely with the condition of the heart, which is in a state of granular degeneration. The beat may be excited and violent, while the pulse is very weak. Often the cardiac impulse undergoes a singular diminution, and with its change the first sound becomes enfeebled; in fact, it is sometimes almost lost, and only very gradually regains its natural tone. Occasionally, at the height of the disease, it is replaced by a soft, systolic murmur of blood origin.

The *urine* is generally high-colored at first, and deposits an abundance of urates and phosphates. There is an absence of the chlorides, or they are reduced to a trace. The urea, as ascertained by an analysis of Parkes\* in a case in which no medicine was given, is increased, and its augmented excretion is remarkably regular during the height of the malady. During convalescence the urea sinks below the normal standard. The water passed is lessened; the urine is apt to contain a large amount of uric acid, and to preserve its acidity. In eight out of twenty-one cases that I examined during an epidemic,† it contained albumin, but this ingredient was present only in the severer cases. In some instances the microscope exhibits in the deposit, besides the salts of the urine, renal as well as vesical epithelium, and tube-casts, either finely granular or hyaline, or epithelial. Much the same condition of urine is also found in typhoid fever. But the pigment that in typhus fever was detected by Parkes throughout only in small amounts has in typhoid fever been found to be immensely increased.

---

\* The Urine in Disease, p. 258.

† Amer. Journ. Med. Sci., Jan. 1866.

The *complications* encountered during the course of the fever, or during convalescence, are much the same as those of typhoid fever, although they do not in the two diseases occur with equal frequency. We meet with abscesses, with large sloughs on the trunk and extremities, or with gangrene of the extremities,\* with milk-leg, with erysipelas, with inflammation of the parotid gland, with œdema of the glottis, and with pulmonary complaints. The latter are very common, and mostly very alarming. Sometimes they consist merely in affections of the larger or the smaller bronchial tubes, and rales of varying size are heard all over the chest. At times, instead of these signs, or associated with them, may be noticed dulness on percussion and bronchial respiration over the lower lobes of the lungs, depending upon congestion, with consolidation, more or less perfect, of the pulmonary tissue. Here is one of the worst of all the complications,—a low form of pneumonia, often of the broncho-pneumonic type. This must, however, not be confounded with the so-called *pneumotyphus*,† in which the manifestations of pneumonia appear early and become later complicated with those of a typh-fever, though generally of typhoid and not of typhus. During the last stages of typhus fever, or after convalescence has set in, *acute tuberculous deposits* occasionally develop themselves in the lungs, with the same symptoms as during or subsequent to typhoid fever.

To discuss now the differential diagnosis of typhus fever. We find various maladies resembling it, but none so closely as *typhoid fever*. The subjoined table shows both their similarities and their differences :

TYPHOID.	TYPHUS.
Age generally from eighteen to thirty-five.	At all ages; often in persons beyond middle life.
Not contagious; mostly sporadic.	Highly contagious; usually epidemic.
Attack generally insidious.	Attack generally sudden.
Duration fully three weeks; frequently much longer.	Duration somewhat shorter; often not prolonged beyond second week.
Death hardly ever before end of second week; more generally in, or after, third week.	Death not infrequently at end of first week, and often before conclusion of second.

---

\* Estlander, quoted in Amer. Journ. Med. Sci., July, 1871.

† Wagner, Archiv für klin. Med., Aug. 1884.



## TYPHOID.

Cerebral symptoms come on gradually; last longer.

Great emaciation.

Face pale, or flush confined to cheeks.

Skin hot, sometimes covered with acid perspiration.

Characteristic temperature-record, chiefly influenced by the changes in the intestinal glandular lesion.

Abdominal symptoms, such as diarrhoea, tympanites; stools contain characteristic bacilli; intestinal hemorrhage not unusual.

Epistaxis common.

Bronchitis and pleurisy.

Eruption light red, and not on extremities or face.

Post-mortem appearances are: morbid state of Peyer's patches; enlargement of mesenteric glands; ulceration of mucous coat of intestine; enlargement and softening of spleen; ulceration of pharynx.

## TYPHUS.

Delirium or decided stupor comes on soon, sometimes almost from the onset; headache has appeared and disappeared by about the tenth day.

Less emaciation; greater prostration.

Face deeply flushed; eye injected.

Skin of pungent heat; sometimes emitting an ammoniacal odor.

Temperature-record more that of a continuous fever; for the most part sudden and rapid defervescence.

No abdominal symptoms; bowels constipated; meteorism rare; intestinal hemorrhage of extreme rarity; sometimes acute dysentery during convalescence or as a sequel.

No epistaxis.

Intense pulmonary congestion; sometimes bronchitis of finer tubes; broncho-pneumonia.

Eruption darker in color, and all over body; very seldom on face.

No constant post-mortem appearances; the most frequent are the dark-colored, liquid state of the blood, and enlargement of spleen. Softening of the heart common. There are no intestinal lesions.

The points of contrast between the two affections are here so manifest that it would seem impossible to confound them. Yet it cannot be denied that in sporadic cases of typhus it may be very difficult to come to a conclusion; occasionally, also, the symptoms of the two diseases are strangely blended or interchanged. Thus, we may have constipation in typhoid, and diarrhoea in typhus, or the eruption may be curiously mixed. For instance:

A boy, sixteen years of age, was received into the Philadelphia Hospital, with evident signs of a beginning fever of a low type. A day or two after his admission, and corresponding, as nearly as could be ascertained, to the fifth day of the disease, an eruption showed itself all over the body. It was dark-colored, petechial in its aspect, and did not disappear on pressure. Associated with

it were drowsiness and constipation. In a few days more, however, the symptoms changed. The dark eruption faded, and rose-colored spots were perceptible on the chest and abdomen ; diarrhœa set in, and the fever ran its course to a favorable termination with the character of typhoid, just as at the onset it had assumed the character of typhus.

Besides typhoid fever, typhus may be confounded with meningitis, with inflammation of the lungs, with measles, with small-pox, and with the plague. The distinctive marks between the first two and typhus fever have been rendered apparent while discussing the cerebral and pulmonary complications of the latter malady. I shall here only dwell again upon the great value of the eruption from a diagnostic point of view. The symptoms that approximate measles, smallpox, and yellow fever to typhus will be analyzed in connection with these affections. One word here as to its difference from the *plague*.

This pestilent disease, which during several centuries left almost annually its deep indent upon the human race, is hardly known to any but Russian physicians at present, save by description. And the descriptions leave on the mind the impression of an exposition of a familiar malady ; for the authors who have most carefully delineated its traits have produced a picture which, with very slight changes, may be suited to a representation of epidemics of typhus fever. Thus, we read of a highly-contagious fever setting in suddenly, attended with constipation, with a rapid, feeble pulse, with dizziness and delirium, with injected eyes, with a dry tongue, with noises in the ears and deafness, with defective urinary secretion, with starting of the tendons, with watchfulness or stupor, and with red patches and purple spots scattered over the surface of the body. The features that the plague does not share with typhus are nausea and vomiting, pale face, an alarmed, despairing look of the countenance, hæmoptysis, and, above all, the buboes and carbuncles in different parts of the body, and the clearing mind when they happen. Moreover, the disease is of much shorter duration. Death generally takes place between the third and the fifth day, or convalescence sets in on the sixth or the seventh day, or early in the second week. It may, however, be protracted by the long-continuing suppuration of the buboes.

In very severe cases death takes place in forty-eight hours.

These cases are apt to be associated with but slight fever and with clear intelligence.\*

The relations of typhus fever to cerebro-spinal fever will be best discussed with the latter disease.

**Cerebro-spinal Fever.**—This disease is also known as cerebro-spinal typhus, as epidemic meningitis, and as epidemic cerebro-spinal meningitis, and is the affection which has been called in this country spotted fever. It was formerly very prevalent in portions of the United States, but the present generation of physicians had little knowledge of it until about simultaneously with the severe epidemic in Germany in 1863 and 1864 it invaded this country and committed great ravages, especially in some of the New-England States, in New York, and in Pennsylvania. Since that time it has become naturalized here, as Ziemssen states to be also the case in Germany.†

Cerebro-spinal meningitis does not always present exactly the same symptoms. These vary somewhat according to the structures which bear the brunt of the disease. Usually, however, marked cerebro-spinal phenomena preponderate; in some instances the evidences of pulmonary embarrassment or of blood deterioration are very prominent. Again, the signs of spinal disturbance may prevail over those of the cerebral, or the reverse.

The disease may be gradual in its approach, feelings of chilliness, succeeded by headache, by tenderness at the nape of the neck, by nausea, and by pain in the back and joints, preceding its full development. Generally its onset is sudden; a violent chill is quickly followed by intense headache, vomiting, and extreme prostration. However the beginning, the disease usually soon reaches its full development. The excruciating headache is associated with vertigo, and often with delirium and stupor. The headache may remit, but does not cease during the attack. Another symptom of the fully-developed disease is stiffness of the deep muscles of the neck, so that the patient cannot bend the head forward; and the stiffness may pass into marked contraction, and the head be thrown backward and rigidly fixed. The contraction

---

\* Hirsch and Sommerbrodt's report on the epidemic in Astrachan in the winter of 1878-79, Berlin, 1880.

† Cyclopædia of the Practice of Medicine, vol. ii., 1875.

of the muscles may extend along the spine, which frequently is painful, not specially to the touch, but on movement of any kind; sometimes, moreover, severe spontaneous pain occurs. There are also pain at the nape of the neck, and in the loins and shooting to the lower extremities, and pain at the epigastrium, and a feeling of contraction of the chest. The face has a fixed or suffering expression; the patient is extremely restless; he trembles; talks incoherently; when spoken to, does not appear to hear; his pupils are generally dilated, and there may be dimness of sight, or double vision. The skin is dry, generally very sensitive, or in some parts the sensibility is increased, in others diminished, and the cutaneous surface is frequently spotted with a red eruption, erythematous and roseolous,—an eruption which often becomes brownish, and then for the most part rapidly petechial, and wholly uninfluenced by pressure; or the purple spots may be seen from the start. Vesicles, too, are apt to appear on the lips. They show themselves from the third to the sixth day of the disease, while the eruption is seen on the first day, or may at all events be detected by the third day.

The pulse at first is natural or slow; but it becomes rather frequent and irregular, and commonly remains accelerated throughout the disease, showing extraordinary variations in a few hours; the impulse of the heart is at times much augmented. The tongue is moist or dry, and brown; the breathing often hurried and shallow; and the urine I have often noticed to contain large quantities of urates and to be slightly albuminous; hyaline and granular tube-casts have been also found in the urine in severe cases attended with high fever.\* In the malignant cases there may be hæmaturia. The bowels are at the outset constipated, but as the malady advances they become relaxed; in some cases dysentery has been observed, with lanceolate micrococci in the stools.† There are usually persistent irritability of the stomach with great thirst and spasmodic contractions or convulsive movements in the muscles of the extremities. The spleen, early in the affection, is apt to enlarge, but does not continue tumefied. With these symptoms, to which those of exhaustion become plainly added, the disorder progresses to its close, presenting now and then

---

\* Flexner and Barker, *loc. cit.*

† *Ibid.*



strange and delusive remissions, soon followed by distinct exacerbations. In fortunate instances the morbid phenomena gradually lose their violence, and the patient, greatly emaciated, enters upon a tedious convalescence.

But though these are the symptoms which frequently recur in epidemics, yet, as already indicated, they cannot always be taken as the standard expression of the disease. Most of them were observed in the formidable examples of the malady which have of late years been encountered in this country; and they have also been met with in the epidemic cerebro-spinal meningitis prevalent in Germany. The temperature is most variable; it may be scarcely above the norm, or may reach between  $106^{\circ}$  and  $108^{\circ}$ , or even higher, without there being a proportionate rise in the pulse. The irregularity of the temperature was a common feature in our epidemics. High temperature may be interrupted by long-continued normal temperature.\*

In a recent epidemic in a mining centre in the State of Maryland, carefully investigated by Flexner and Barker,† symptoms referable to the cranial nerves were observed, particularly loss of smell, strabismus, nystagmus, inequality of the pupils, photophobia, ptosis, impairment of vision, rigidity of the face, trismus, retarded respiration, Cheyne-Stokes breathing, deafness, and disturbances of speech. The strabismus was almost always divergent, and in many cases affected especially the right eye. The nystagmus was, as a rule, horizontal or vertical, but in one case it was rotatory. A considerable number presented engorgement and tortuosity of the retinal veins; some, optic neuritis; and one each, retinitis and thrombosis of the central vein. The pupils were variable,—sometimes contracted, sometimes dilated, often unequal. The tendon-reflexes were not uniform, but were in many cases diminished. General cyanosis was not uncommon, and epistaxis was frequent. In addition to herpes and purpuric and petechial spots, a common form of cutaneous eruption was an indistinct purplish mottling of the surface. Nearly twenty per cent. of the cases presented articular complications, principally

---

\* Wunderlich, *Archiv der Heilkunde*, No. III., 1865.

† *American Journal of the Medical Sciences*, February, March, 1894. For a report of the ocular findings, see, also, Randolph, *Bulletin of the Johns Hopkins Hospital*, 1893, vol. iv., No. 32, p. 59.

effusions into and around the joints, with redness and swelling. Well-marked leukocytosis was a constant feature at the height of the disease; the red blood-corpuscles were little, if at all, changed in number, while the hæmoglobin was somewhat diminished. No micro-organisms were found in the blood.

The duration of the malady is very various. Patients may become rapidly comatose, and die within twelve hours, before any distinctly febrile action has begun; or may sink in a few days; or, on the other hand, the complaint may pursue a very chronic course, lasting for weeks, and during this time deafness and blindness, convulsions, retention of urine, and local palsies—though these are unusual—may be prominent phenomena.

Of the cause of the formidable disease we know little. It is not a malarial disease; for, though occasionally there is a singular intermission or remission in the symptoms, there is no regularity in this respect. The temperature-record, even of these apparently malarial cases, is different, being irregular; and the affection is unyielding to quinine. Many look upon the disorder as modified typhus; and certainly it occurs epidemically under much the same circumstances as typhus. But, though kindred to typhus, it is a fever due to a different poison, and differs broadly from typhus by being far less contagious, if indeed it can be regarded as contagious at all, and in the inflammatory lesions found in the brain and spinal cord. Bacteriological examinations have in many instances resulted in the isolation from the meningeal exudation of the diplococcus of pneumonia, which, there is reason to believe, bears an etiological relation to cerebro-spinal fever.\*

Cerebro-spinal meningitis attacks children very frequently. It is more common in winter and in spring than in summer; though I have seen it in summer. It is an affection very familiar to military surgeons; it seizes on recruits who have been subjected to unaccustomed fatigue or have been huddled together in unhealthy barracks or camps.

To determine the diagnosis is ordinarily not difficult: the sudden onset of the malady and its epidemic character are safeguards against error. The protracted cases simulate *typhoid fever*. They

---

\* Flexner and Barker, *loc. cit.*

resemble it in its long duration, in several of the cerebral symptoms, and in the occurrence of an eruption, and sometimes of diarrhœa. They differ from it in the more abrupt invasion, or rather in the short time in which the disease reaches an alarming aspect; and in the early stages the violent headache, the constipation, the constant vomiting, the slow or normal pulse, and the cool or but slightly heated skin, are unlike the signs of enteric fever. In those cases in which an eruption appears, it is noticed, at latest, by the third or fourth day, not at the end of a week, as in typhoid fever; nor is the rash, save in extremely rare instances, rose-colored. Later in the malady the traits of distinction become broader and broader. The prominence of the abdominal symptoms in the one disorder; the continued violent headache, the fixed spinal pain, the hyperæsthesia, the facial herpes, the severe twitchings or the tetanic rigidity of the muscles, and the absence of marked enlargement of the spleen, in the other,—are signs the import of which is not easily overlooked.

The suddenness with which the morbid phenomena occasionally develop themselves, and the lulls that take place in the course of the affection, may cause it to be mistaken for the cerebral variety of *congestive fever*. But the remissions are not so marked as in this pernicious malady, nor are the exacerbations preceded by a long, violent chill. Moreover, the temperature-record is different, and congestive fever does not begin with congestive symptoms, but the first attack is like that of an ordinary intermittent or remittent: hence we have the history of the case to instruct us. Finally the detection of hæmatozoa in the blood establishes the diagnosis of the malarial affection.

From *tetanus* cerebro-spinal meningitis may be distinguished by its epidemic prevalence, and by the signs of mental disturbance, which are very slight or wholly wanting in the former disorder. Trismus is common and early in tetanus; rare, if ever present, in cerebro-spinal fever. Generally, too, the sudden and painful spasms, aggravating the tetanoid contractions, and the cognizance of the exciting cause of the tetanic convulsions, such as their following wounds or punctures, aid in interpreting their meaning.

How can we discriminate between *inflammation of the meninges of the cord* and epidemic cerebro-spinal meningitis? Thus: in pure spinal meningitis, as in myelitis, mental symptoms are ab-

sent; their presence in cerebro-spinal fever constitutes one of the marked features of the disease. The history of the case in the former malady points to cold and exposure, or to syphilis. Clonic spasms of the extremities are more common; persistent rigidity of the muscles is a less striking peculiarity. We find no eruption.

*Tubercular meningitis* is distinguished by the much more insidious beginning, the much more protracted course, the absence of eruption, and usually of marked stiffness of the neck, the variations in the pulse according to the stage of the disease, the irregular breathing, and the history of a tubercular taint.

*Idiopathic or sporadic cerebro-spinal meningitis* is a very rare disease. It runs a much slower course than the epidemic malady generally does, and its spinal symptoms are less marked. But it cannot be distinguished with any certainty from isolated cases of cerebro-spinal fever. The absence of an eruption and of the striking variations of temperature presented by the latter is of significance.

There are other diseases with which cerebro-spinal meningitis has been confounded; for instance, owing to the eruption and to the sore throat that may attend it, with *scarlatina*. But the onset and the neck-symptoms are very different; and so is the eruption; certainly it is different in its course. Still, as regards the onset, we must bear in mind that both may be ushered in by convulsions. An extremely rapid pulse would be in favor of scarlatina. Cerebro-spinal fever also resembles at times the onset of *malignant measles*; but the catarrhal symptoms and presently the eruption guide us.

I have known more than once the disease, on account of the congestion of the lungs or the broncho-pneumonia which may accompany it,—and in some epidemics the lung-affection is very marked,—to be mistaken for *pneumonia*. In truth, the diagnosis is sometimes far from easy. The mental symptoms, the intense headache, the variations in the pulse, the hyperæsthesia, the vomiting, the stiffness and retraction of the muscles of the neck, the eruption, are distinguishing traits of value; but when these important symptoms are ill defined, much doubt may exist. So there may if epidemic cerebro-spinal meningitis become intercurrent, as it sometimes does in pneumonia as well as in other acute affections. Supervention of the severe headache, and ap-



pearance of rigidity of the neck, of great restlessness, of hyperæsthesia, and of coma, are the symptoms of most importance.

In some instances of cerebro-spinal fever there is great pain, with some swelling of the joints, and the disorder is thought to be *acute rheumatism*. But the head-symptoms, the state of the muscles of the neck, and the dissimilar course of the malady soon clear up the diagnosis.

The poison may produce so light a case that the stiffness of the neck may be mistaken for *rheumatism of the cervical muscles*. There is, however, even in these instances, an unusual amount of headache, and in a case in which I was consulted it became a permanent condition for several years, and then yielded.

*Uræmia* with contracted kidneys may give us most of the same symptoms as cerebro-spinal fever, especially headache, vomiting, and retraction of the head; the differentiation will depend upon the previous history, the existence or non-existence of similar cases, and the presence or absence of febrile phenomena and of cutaneous eruptions.

Lastly, let us look at the clinical features separating cerebro-spinal fever from the disease it is most like,—*typhus*; let us contrast its phenomena with those of this affection, which in many respects it so closely resembles. Both diseases are apt to prevail at the same time; both attack all classes and ages; both are evidently attended with dissolution of the blood,—but this alteration in the blood occurs much more rapidly and is much more marked in epidemic cerebro-spinal fever than in ordinary cases of typhus;\* the eruption is different from that of the common form of typhus; we find less delirium; a less intense, though more irregular, fever; the affection is generally of much shorter duration; the countenance is not of a dusky hue and stupid, but pale or of a sallow color, and dull or expressive of suffering; and there is the stiffness of the muscles of the neck, with the fixed spinal pain, and muscular contractions and other signs of spinal or cerebro-spinal

---

\* The deterioration of the blood occurs, indeed, very soon in cerebro-spinal fever. In an autopsy of a child that died in twenty-four hours, I found the blood diffuent and black; in an adult patient who had been ill but two days, I detected blowing sounds in the heart, evidently of blood-origin. The poisoned blood unquestionably gives rise to many of the nervous symptoms, and it is on the blood and the nervous centres that the poison mainly acts.

lesion ; and the herpetic eruption on the face. Certainly, therefore, the clinical manifestations of cerebro-spinal fever are very dissimilar to those of the usual varieties of typhus. But they are not so dissimilar to those occurring in some epidemics of malignant cerebral typhus.\*

Cerebro-spinal fever may, during an epidemic, complicate other acute maladies, and mix its symptoms curiously with them. With the attack the difficulty does not pass off, for it may leave all kinds of want of power and local palsies, besides derangement of vision, permanent deafness, impaired intelligence, epilepsy, persistent headache, chronic meningitis, which may be indeed the cause of the headache, and chronic hydrocephalus. In one instance I have known an extraordinary swelling of the whole body to follow ; the skin is hard, tense, and greatly thickened, pits very little on pressure, except around the ankles, and is tightly drawn over the face ; this swelling and thickening, very much like a general sclerema, has now lasted for upward of twenty years, and has been attended with a feeling of numbness in the skin and a moderate amount of anæmia. There is no palsy or albuminuria ; the patient suffers little inconvenience, except from her size. She has a waxy countenance, and looks like a very fat woman.

**Relapsing Fever.**—This is a form of fever characterized by its rapid course and its proneness to relapse. Epidemics of this disease—and it occurs only in epidemics—are frequently encountered in Ireland and in Scotland. In this country it was until of late years almost unknown.

The disorder is decidedly acute. Its invasion is sudden, and marked by rigors, pain in the back and limbs, vertigo, severe headache, and nausea and vomiting. Fever is soon developed, and rises high, it may be to between  $107^{\circ}$  and  $109^{\circ}$ . There are severe muscular pains, particularly in the muscles of the extremities ; the pulse is very rapid ; the temporal arteries throb ; the tongue is covered with a thick white fur. The bowels, as a rule, are constipated. In many cases there is engorgement of the liver, with yellowness of skin ; and in nearly all there are epi-

---

\* An extraordinary case, bearing on the relationship of the complaints under discussion, was under my charge in 1865 at the Pennsylvania Hospital. See Case XII. of a series of typhus fever cases published in *Amer. Journ. Med. Sci.*, Jan. 1866.

gastric tenderness and marked enlargement of the spleen. The matter ejected from the stomach is greenish, or sometimes black and like coffee-grounds. Minute points of extravasated blood are not uncommonly seen upon the integument. The urine is scanty, and contains usually bile-pigment, some albumin, and hyaline casts. On the fifth or the seventh day, though sometimes not until the tenth, the symptoms subside as speedily as they set in, a profuse perspiration preceding their decided abatement, and the temperature falls to the norm or even below. Convalescence is now apt to be rapid, and seemingly complete, the patient being up and going about; but the apparent return to health does not last long. Ordinarily after a week, therefore on the twelfth or fourteenth day from the first beginning,—sometimes sooner, rarely later,—the attack, preceded perhaps by a slight rise in temperature for an evening or two, returns, presenting again the same signs, and again terminating by a critical sweat in convalescence. This second attack may be short and mild; but it may be both longer and of graver character than the first. It is, at times, followed by another, and yet another, relapse. When the patient finally throws off the disease, he is very weak, and his blood is much impoverished. He shows a tendency to dropsy of the extremities; and blowing murmurs, evidently not organic, are perceptible while listening to the heart. These murmurs, however, may also be heard during the paroxysms. The patient is not really well during the intermission; his spleen remains enlarged, the pulse is slow, the action of the heart is weak, and the muscular and arthritic pains do not entirely disappear.

Relapsing fever has an intimate connection with destitution. It is a contagious but far from a fatal disorder, except, perhaps, in the negro. In fatal cases death sometimes happens during the first paroxysm as the result of syncope, of hemorrhage into the brain or from the lungs; or it may occur suddenly during the intermission from paralysis of the heart. But the most common termination of the cases having an unfavorable issue is in consequence of complications or of states which have been induced by the malady, such as lobular or lobar inflammation of the lung, hemorrhagic pachymeningitis, abscess of the spleen or of the kidney leading to pyæmia, chronic diarrhœa, Bright's disease, dropsy, parotitis, palsies. At times the patient perishes in a condition

similar to the collapse of cholera, though the collapse is more protracted and the pulse can be felt, and discharges from the bowels are by no means a constant accompaniment. The extreme prostration, attended with great coldness of the skin, may last for days. It is more particularly met with in the "bilious" or "bilious typhoid" form of the malady,—a dangerous variety, in which severe vomiting, jaundice, and delirium are encountered, and the paroxysm is not followed by a distinct intermission or remission, but often by the signs of collapse alluded to, in which uræmic symptoms have been more particularly noticed.\* The collapse, however, may happen not only at the close of the paroxysm, but in the remission, whether this be distinct or not, or in a subsequent paroxysm; and this may be the case no matter what variety of the disorder we have to deal with, and whether or not the grave symptoms be due to uræmia.

Yet the state of the kidneys and of the urinary secretion has commonly a great deal to do with the graver phenomena of the malady. Acute renal disease with albumin and tube-casts in the urine was discerned by Obermeier† in two-thirds of his cases. It was also, with or without tube-casts, met with in a number of Pepper's cases.‡ The urea, Reisenfeld§ found, was during the first paroxysm always increased, and this increase continued beyond the crisis. The products of the heightened tissue-metamorphosis may be retained, and thus grave symptoms arise. Leucine and tyrosine have been also found.

There is no constant obvious lesion in relapsing fever, unless it be the lesion in the spleen. This organ is greatly enlarged, and presents numerous round or irregularly-shaped bodies, of white or yellowish-white color.|| But myriads of minute organisms, spirilla, are found in the blood just prior to the outbreak of the paroxysm, and at its height. Indeed, since Obermeier's discovery of the spirilla in relapsing fever, there is no doubt that they are the cause of the malady, and their detection in the blood re-

---

\* Hermann, Account of St. Petersburg Epidemic, Schmidt's Jahrb., No. 6, 1865. See also further observations in Meissner's article, *ib.*, No. 2, 1870.

† Virchow's Archiv, 1869, Bd. xlvii.

‡ American System of Medicine, article "Relapsing Fever."

§ Virchow's Archiv, 1869, Bd. xlvii.

|| Pastau, *ib.*




moves all doubt in the diagnosis. In a single field of the microscope we may see but a few or from twenty to thirty spirilla.

The diagnosis of the malady cannot be made positively during the primary seizure. Yet the presence of the fever, while an epidemic prevails, may be suspected from the sudden fierce beginning of the attack; from the fact of the high fever-heat of  $104^{\circ}$  to  $107^{\circ}$  showing itself in less than twenty-four hours, and exhibiting either a morning remission of one to two degrees and the maximum of temperature in the early afternoon or evening, or but little difference between morning and evening, until the rapid and great fall which takes place at the crisis; and from the character of the gastric symptoms. Then the

microscopical examination of the blood is of great importance. Relapsing fever resembles *yellow fever* in its short duration and in some of its manifestations. But there is this evident difference: in yellow fever the paroxysm or febrile stage is usually much shorter; the symptoms in the remission do not subside nearly so completely; this stage is a very brief one as compared with the decided intermission of relapsing fever; the black vomit of yellow fever does not come on until the stage of collapse is reached; and this far more fatal malady presents lesions in the liver and heart which are not found in relapsing fever, while it does not show the extraordinary enlargement of the spleen.

From *typhoid and typhus fevers*, relapsing fever may be distinguished by the shorter prodromata, by the presence of jaundice, by the absence of the characteristic eruptions, and by the short period during which the symptoms last. Again, critical sweats with the rapid cessation of the fever are not likely to be seen in these disorders, certainly not in typhoid fever; and the very high

FIG. 79.



Spirilla of relapsing fever (from Heydenreich). a, single spirillum; b, star-shaped bundle; c, nidus of spirilla, with blood-corpuscles.

temperature, the severe muscular and arthritic pains, the tenderness over the liver and the spleen, the vertigo, and in some cases the early collapse without apparent cause, are characteristic; while, on the other hand, delirium and stupor are rarely encountered in relapsing fever. After the relapse has taken place, the diagnosis is easy, if the case have been watched during the first attack. But, should it not have been under notice before, it may be at times very difficult, without an examination of the blood for spirilla, to say whether we are dealing with relapsing fever or with a relapse of typhoid or typhus fever. And this difficulty is enhanced by the want of uniformity of the symptoms in the second onset of the strangely recurring malady, and the close similarity they occasionally show to those of typhoid or of typhus fever. Another difficulty is presented by the fact that relapsing fever may exhaust itself in the first paroxysm. But this is an unusual occurrence, and the abortive cases are light. In them too the spirilla may be detected in the blood.

### Periodical Fevers.

These fevers are characterized by the distinct periodicity of their phenomena: they exhibit intervals during which the patient is wholly or nearly free from febrile disturbance. With the exception of one,—and its place here is, indeed, doubtful,—they are all owing to marsh miasm, or malaria. This noxious agent gives rise to a group of fevers ever betraying their common origin by their strong family resemblance: alike in occurring in low, swampy localities; alike in most of their symptoms, and in the difficulty of eradication from the system; alike in the secondary lesions, in the enlargement of the spleen and of the liver, and in the altered condition of the blood, which they leave behind them; and also alike in being under the control, absolute and immediate, of cinchona and its various preparations. Along with the forms of miasmatic fever I shall describe yellow fever; not because it is of identical nature, but on account of the similarity of the prominent symptoms.

**Intermittent Fever.**—The paroxysm comes on with a chill: the face becomes pale, the lips bluish; the teeth chatter; the skin is cold; there is a feeling of uneasiness and fatigue. After

a period varying commonly from half an hour to an hour, this cold stage passes off. Now we find decided heat of the surface, with restlessness, thirst, a full, rapid pulse, muscular pains, a scanty secretion of urine; in other words, active febrile symptoms. These continue for hours, for a period always much longer than the first stage: then a sweat breaks out all over the body; the pulse becomes softer and less frequent; the secretions are fully re-established; and this sweating stage terminates the paroxysm.

The patient is now, for the time being, well; but the disease soon recurs: in from twenty-four to seventy hours the paroxysm repeats itself. In the former case we call the fever a *quotidian*; in the latter, a *quartan*. The *tertian* type is before us when the paroxysm sets in again in about forty-eight hours; the *double tertian*, when we find a daily attack, but those of alternate days alone corresponding in time and severity. Even a *quintan* ague may happen.\* The period between the ending of one attack and the beginning of another is spoken of as the *intermission*, or *apyrexia*; while the time between the beginning of the two paroxysms, including the first with its succeeding intermission, is called the *interval*.

The varied types of the fever present marked differences in the character and duration of the several stages. The *tertian* has generally the longest hot stage, the *quartan* the longest cold stage. In the *quotidian* there is a short cold stage, followed by a hot stage which may last for upward of fifteen hours. Occasionally the stages are very irregular and anomalous. Thus, the sweating stage may precede the cold stage, or it may be the only one which shows itself; or, again, the rigor may be altogether wanting. Sometimes there are no distinct stages, but the patient has a "dumb ague," which manifests itself at definite periods by a feeling of great depression, or of a severe pain at some portion of the body, or by chilly sensations, or by headache, or by nausea and vomiting, or, as I have seen, by excruciating pain over the kidneys, and almost entire suppression of urine, or by spasmodic obstruction of the intestine.†

---

\* Case of Henry, Brit. Med. Journ., Feb. 18, 1888.

† Cases of Hoyt, Atlanta Med. and Surg. Journ., Sept. 1875.

Now, cases of this kind are difficult to distinguish from organic disease. We can do so only by laying stress on their strictly periodical nature; by noting that the curious manifestations cease entirely to recur with intensity. This does not happen when the symptoms are not caused by a lurking malarial poison; for idiopathic disorders exhibit the phenomena of structural change or of deranged function at all times,—not merely on certain days or at certain hours. It is true that among the inhabitants of miasmatic districts some complaints, and particularly those of the nervous system, display a well-defined periodicity; but here, too, are found the significant traits of organic or functional disturbance between the decided exacerbations of the symptoms.

Then, again, we must remember that diseases may assume an apparently intermittent character, being worse every second day, and yet not be malarial at all. Even mania, as Schroeder van der Kolk tells us, may take this type. The whole aspect of the symptoms, and a tentative treatment with quinine, will help to inform us as to the true nature of the malady.

The temperature in intermittent fever shows a peculiar record, and one which, in doubtful cases, may be turned to great advantage. Notwithstanding the marked sense of chilliness, the thermometer rises suddenly and rapidly to a high degree; there may be a slight elevation of temperature for an hour before a chill, but the striking rise begins with the chill. Even during the decided chill of the beginning of the paroxysm it indicates  $105^{\circ}$  or more in the axilla. The temperature remains stationary, or continues to rise, though not much, during the hot stage, and during the sweating stage falls at first slowly, then rapidly, until it comes down to about the normal heat. During the chill the peripheral temperature is decidedly lowered; during the hot stage it is increased. But with the ending of the paroxysm it is found that the fall has been equally rapid. In the intermission the thermometer in the axilla marks a natural temperature, or one somewhat lower than in health. It rises again quickly with each paroxysm. No other malady presents these variations.

In some cases of intermittent fever a peculiar intermitting murmur is heard over the spleen. This is ascribed to the movement of the blood in the splenic arteries with the systole, in consequence of the soft, enlarged, and overfilled condition of the spleen.




It is usually detected most distinctly during the febrile period; ceases with the paroxysms,\* and is not heard in chronic malaria.

The diagnosis of an ordinary and regular intermittent is easy. Leaving the other malarial fevers out of consideration, only two morbid states are likely to present recurring rigors and febrile excitement, and are, therefore, apt to be confounded with it: hectic fever, and chills attending upon suppuration in deep-seated parts. Now, *hectic fever* differs in this from an intermittent: it is simply a fever of irritation, the cause of which a careful scrutiny will generally detect. We find it accompanying many chronic diseases in which destruction of tissue occurs, especially phthisis; and the chronic affection has its own signs, which exist at all times, whether the symptomatic fever be present or not. Then its outbreaks are irregular. Several often take place within the twenty-four hours; their intermissions are incomplete; the temperature does not fall as in intermittent fever, for there is not complete defervescence; and al-

though the paroxysms may begin with chilliness, they are not ushered in by a well-defined rigor. Further, they are apt to be morning paroxysms, and are not modified by antiperiodics. Whenever, indeed, we find an intermitting fever not influenced by these agents, it ought to arouse suspicion, and all the internal organs, particularly the lungs, should be carefully explored. Thus only can serious errors in diagnosis be guarded against.

When *pus* forms, and especially when it forms in internal cavi-

FIG. 80.



Temperature-record of a tertian intermittent.

\* Maissurianz, St. Petersburg medicin. Wochenschr., 1882, 12.

ties, it betrays its presence by rigors, followed by more or less fever. But these, unlike the chills of ague, do not repeat themselves at definite periods. Moreover, in the midst of the apparent intermission, febrile signs or other manifestations of a seriously disordered system may be discovered. The chills of ordinary pyæmia are distinguished by the same phenomena; then the rigors, unlike the malarial malady, are often characterized by the profuse sweating which immediately follows them, rather than by an active development of the fever.

But there are other causes which may occasion attacks of fever happening in paroxysms and simulating ague. They may occur in *disease of the heart*, as in ulcerative endocarditis and in valvular affections.\*

*Gall-stones* which form in the radicles of the hepatic duct in the interior of the liver may, as Frerichs shows, give rise to attacks of chills, followed by heat and by sweating, easily mistaken for ague. The fact that these febrile phenomena are preceded in many instances of *intra-hepatic concretion* by dull pain in the hepatic region, and by sudden sharp seizures of pain at the lower part of the thorax on the right side, is very significant. Even gall-stones passing along the gall-duct and the common duct may occasion febrile symptoms like those of an intermittent, and the paroxysms may extend over months, and then the patient recover. Jaundice is apt to be a symptom of this *hepatic fever*, the attacks of which are more likely to be repeated in the afternoon and evening, while the malarial paroxysm more commonly occurs in the morning. We have already pointed out that hepatic fever may occur without the impaction of gall-stones, and have seen how far these different forms can be told apart.

An affection which on account of the chill succeeded by fever might be mistaken for the malarial disorder is the curious so-called *urethral fever* which sometimes arises after the passage of a bougie, and which may even terminate in death.† Our knowledge of the introduction of the instrument, and the non-recurrence at a fixed time of the rigor and febrile phenomena, furnish the points of distinction.

---

\* Ord, St. Thomas's Hospital Reports, 1882; Osler, Practitioner, vol. 1., No. 3, p. 181.

† Roser, quoted in Brit. and For. Med.-Chir. Rev., Oct. 1867.

Yet another affection liable to be mistaken for intermittent fever is *syphilitic fever*. The fever may occur in attacks consisting of a chill, followed by a hot stage and sweating, and be so similar to the malarial disorder as to lead to error.\* The apparent ague-fits happen, however, toward evening, and are succeeded or accompanied by severe headache and pains in the bones,—in fact, by the same symptoms as the more ordinary form of syphilitic fever. In the form in which the febrile symptoms are continuous, these generally precede the eruption for a week or more, and may continue after this appears.

We may also find this syphilitic fever with symptoms like those of malaria in cerebral syphilis.† This, it is well known, may occur, is indeed apt to occur, years after the early manifestations of syphilis, though the brain affection may happen within six months: thus the paroxysmal pyrexias may be met with at very varying times after the infection. The history of cerebral syphilis must often be considered, to understand their meaning. We must bear in mind that disease of the membranes of the brain may exist which may disclose itself with great suddenness or gradually, and which does not unusually appear with apoplectic seizures; that headache is a very marked symptom: that irregular motor palsies and epileptic attacks frequently happen, as well as mental failure and perversion, and symptoms similar to general paralysis, though wanting in the tremulousness. The aphasia which may be met with is said to be very commonly associated with left-sided hemiplegia.

In the puerperal state a malarial outbreak may happen which, as Manson and Fordyce Barker‡ have shown, may be mistaken for puerperal fever. Unlike the latter, however, the *puerperal malarial* fever is attended with pain in the head, back, and limbs, and does not generally appear so soon after parturition,—not, therefore, between the first and fifth days after delivery. Moreover, it has at the beginning a great temperature-rise, and marked

\* See cases of Bassereau, referred to by Bumstead in his *Treatise on Venereal Diseases*; Ord, *loc. cit.*

† Wood, *Transactions of the College of Physicians of Philadelphia*, Feb. 1884; also in *Medical News*, Philadelphia, March, 1881; Janowsky, quoted *ib.*

‡ *Medical Record*, Feb. 1880; *Virginia Med. Monthly*, Nov. 1881.



remissions or intermissions. Puerperal malarial fever may lead, after the twelfth day, to secondary hemorrhage.

**Remittent Fever.**—This is a fever pre-eminently of hot climates and malarial districts. It is the fever of Hungary, of the Pontine Marshes, and particularly of Africa and the southern portion of the North American continent. Occasionally, not often, we meet with it in winter and in early spring; very generally, during the summer and autumn months.


Remittent fever has no well-defined and constant prodromic symptoms, except, perhaps, a singular sense of gastric uneasiness. It is ushered in by a marked chill, soon succeeded by violent fever, which, after a varying period, decreases, and then breaks out again. By this time the symptoms of the disease are very apparent. The patient complains of pain, of fulness and of throbbing in his head. He is restless and distressed; his limbs ache; his tongue has become coated; he suffers from thirst, and rejects the contents of the stomach. After continuing at their height from six to eighteen hours, these symptoms again subside: a sweat breaks out all over the body; the irritability of the stomach lessens; the patient is composed, even cheerful; his headache has nearly ceased, and he falls into a quiet slumber. But this lull is not of long duration, not longer than some hours. Soon the active fever is rekindled: the skin is as hot and dry as before, the pulse as full, frequent, and hard; the spleen is observed to be swollen; and the other symptoms return with increased intensity, again to abate, again to recur, until either the exacerbations are effaced and the fever assumes a continued type and then gradually lessens, or else subsequently the remissions become better and better defined,—more, indeed, like intermissions than remissions. In the progress of the disease at and after its height the pulse is generally quicker and weaker than at first.

The temperature rises markedly with the first chill, and continues to rise during the high fever that follows. With the sweating stage it declines by several degrees, to rise to a greater height than previously with the succeeding febrile phenomena; then again there is a fall in the remission, with another quick rise in the fever, which may attain a very high point, marking from  $105^{\circ}$  to  $108^{\circ}$ . The greatest height is usually reached in the exacerbation of the third day. After this the remissions become



less distinct, and may, indeed, be recognizable only by the thermometer; the whole fever is more like a continuous one. Subsequent to the ninth day usually the remissions are very marked, the difference between the heat in them and the exacerbations being three degrees or more. The exacerbations become less and less high, and soon cease, the temperature falling perhaps previously to below the norm. In cases in which the fever remains for a long time continuous, irregular remissions occur, especially

FIG. 81.



Temperature in a case of remittent fever of moderate severity, ending in recovery on the twelfth day. The chart shows also the pulse and the respiration.

toward the end, though the fever may preserve its continuous type more or less to the end.

The average duration of the fever, unless protracted by complications, is from nine to twelve days. Its most common type is quotidian, or rather, perhaps, double tertian, the exacerbations of alternate days corresponding in severity, in duration, and even in the nature of the symptoms. Sometimes there are two

exacerbations in twenty-four hours,—a duplicated quotidian,—or the paroxysms have a tertian form. The exacerbations may occur any time in the twenty-four hours; in many instances morning exacerbation is noticed, and I have met with more cases in which the paroxysm comes on in the afternoon than in the evening.

The urine in remittent fever presents much the same changes, though in a different degree, as those occurring in intermittent fever. During the active stages of the fever there is an increase of urea, not simply above the standard of health, but even above that in intermittent fever; and this increase of urea is attended with a diminution of uric acid—unlike what happens during the paroxysms of ague—and of the coloring and extractive matter; while, as convalescence sets in, the urea decreases in amount, and the other ingredients mentioned increase.\* A copious deposit of urates, forming with the phosphates as it were a critical discharge, is noticed as the fever subsides, and is analogous to what takes place after the paroxysm in intermittent fever. At no stage does the urine contain albumin, as it often does in typhus, and as it generally does in yellow fever, but, as in intermittent, it may contain sugar.

Remittent fever is readily recognized: the rise and fall of its febrile signs are too striking to escape observation. Its characteristic traits are more closely allied to those of *intermittent fever* than to those of any other disorder. But there are these points of contrast: in intermittent fever each paroxysm begins with a chill, which is not the case in remittent fever; for after the first paroxysm there is rarely a marked chill, and even the chill ushering in the disease is usually not violent. After each febrile exacerbation comes an abatement,—not an intermission, for the thermometer shows that the fever does not wholly leave; the tongue remains coated, and the gastric derangement does not entirely cease; the patient is not well, as after a fit of ague. The symptoms grow and decline; they do not appear and disappear. In both affections we may have herpes labialis at the decline, but it is more common in remittent than in intermittent.

Owing to the jaundice in many cases of bilious remittent fever, the disease is often mistaken for *acute congestion of the liver*.

---

\* Joseph Jones, Observations on Malarial Fever.

Here, again, the exacerbations and remissions in the temperature serve as distinguishing marks; and so, too, in separating the gastric complications of bilious remittent fever from acute gastric inflammation. The severe headache is also a distinctive feature of value; so is the herpes labialis.

Under ordinary circumstances, there is very little likelihood of confounding with each other *typhoid* and remittent fevers. The lines between the two diseases are too strongly drawn: no marked periodicity exists in typhoid fever, and, on the other hand, we find no diarrhoea, no eruption, no thoracic symptoms, no deafness, and no very great prostration, in remittent fever. But instances are met with in which the diagnosis is not easy, because the symptoms of the two maladies are blended. Thus, in a typhoid fever occurring in a malarious region there are often distinct exacerbations and remissions obscuring the real ailment. The malarial influence has set its stamp on the disease, and may for several days completely veil it; but soon its real nature becomes manifest. The great weakness; the low delirium; the tympanitic abdomen; the thin passages, so unlike the dark, hard stools of remittent fever,—all unfold the true character of the disease. Sometimes a certain periodicity is witnessed in typhoid fever as it is approaching a favorable termination; the afternoon or evening rise of temperature is most marked, the morning remission very great. Here a knowledge of the previous history of the case guards against error. We shall presently again refer to the symptoms of periodicity in fevers of low type in examining into *typho-malarial fever*.

Further, not unfrequently, after an attack of remittent fever has lasted for ten or twelve days, these symptoms are noticed: great muscular debility, jerking of the tendons, picking at the bedclothes, dark, dry tongue, and weak pulse, perhaps diarrhoea. The fever becomes of a continued type. It is these cases which have given rise to the opinion that bilious fever often changes into typhoid fever. But in reality it is not so much the specific typhoid fever, with its enteric lesions, as a typhoid condition, that is developed.

During the exacerbations of remittent fever the cerebral symptoms are sometimes almost identical with those of an acute brain-affection. There is severe headache, with violent beating of the arteries of the neck and face, a wild eye, intolerance of light,

and even delirium. Were the patient now seen for the first time, he would be at once pronounced to be laboring under *acute meningitis*. Suddenly the pulse loses its throbbing character, a perspiration covers the surface, and, as if by magic, the cerebral disturbance ceases until the next paroxysm redevelops it. Cases of this kind are readily enough recognized, if we know something of their history. If we are not familiar with it, we have to await the remission for their explanation; and after the sudden withdrawal of the signs of disorder of the brain it is hardly possible to have doubts as to the meaning of the acute nervous symptoms, should they recur. It cannot be a meningitis we are dealing with,—a steady, progressing disease, and one never exhibiting such strange freaks of intermission. But occasionally the symptoms show themselves under circumstances where a malarial poison is not suspected to be at work :

A young gentleman of studious habits, while diligently preparing for a college examination, was seized with violent headache and fever. The sense of fulness in the head was unbearable, the fever was high, there was nausea with great gastric irritability. These symptoms lasted for nearly twenty-four hours, and then subsided in the forenoon, to become aggravated in the evening. Delirium followed by great drowsiness was perceived at an early hour of the third day of the disease. The case now assumed a very alarming aspect. Local blood-letting was resorted to with some relief, and in a few hours the symptoms were, fortunately, favorably modified : the headache was much less, the mind was again quite clear. Although the patient had never suffered from a malarial fever, he had spent part of his summer vacation in the marshy neighborhood of Washington ; but several months had elapsed, and winter was setting in. The time of the year was not in favor of malaria. But the evident remission in the cerebral symptoms, the coated state of the tongue, and the indescribable malarial look of the countenance, that became daily more apparent, decided me upon administering quinine. The evening exacerbation came, but was far less severe. The nature of the case was now evident : the quinine treatment was vigorously pursued, and the patient soon recovered.

The violent headache and delirium were in this case observed to be in connection with well-defined febrile signs. Occasionally



one or both of the symptoms mentioned last during the remission, while the fever abates. I have even met with them occurring in paroxysms without fever being present, as in the following case seen a number of years ago :

A young lady of delicate constitution was attacked, in September, with remittent fever. The disease ran its course without any unusual symptoms ; a violent headache, but little if any wandering of the mind, being observed during the daily exacerbations. After the tenth day the fever lessened, and the disease assumed a continued type ; yet soon afterward, as convalescence seemed to be established, every evening for three days, between five and six o'clock, a boisterous delirium set in, lasting for three or four hours, and once nearly all night. It was followed by a profound sleep, from which she woke up with a clear mind. During these fits the pulse was not accelerated, and there was no fever. The third attack was not so very severe, as the patient was already in part under the influence of decided doses of quinine ; another was prevented by this drug.


In both these cases the symptoms approached those of the congestive type of the disease, and the issue appeared at one time doubtful. Generally speaking, remittent fever, unless it be of the congestive variety, has a favorable prognosis. It is difficult for us, living in a century in which the remarkable effects of bark are so well understood, to believe that the complaint was once so fatal, and that so many deaths should have taken place from a disorder over which we now exercise so undoubted a control. But the long list of distinguished names that have fallen victims to it, among them Cromwell, James I., and the Emperor Charles V.,\* proves the medical skill of former times to have been insufficient for its cure. In our day, the consequences of remittent fever are more to be dreaded than the disease itself. We often find, as its sequelæ, obstinate intermittents, enlargement of the liver and spleen, dropsy, protracted anæmia, headache, and impaired activity of mind.

---

\* From the record of the Emperor's illness, as given by the historian Mignet (*Charles V au Monastère de Yuste*), we may learn, what fortunately now we hardly have an opportunity of observing, the features of remittent fever when left to itself.

It is in this *malarial cachexia* that, on pricking the finger and examining a drop of the blood thus obtained, we detect a large number of those particles and masses of black or dark color and irregular shape to which Frerichs has particularly called attention. Not that the pigment-matter is found merely in the cachexia following remittent fever. We observe it in the blood in the severer forms of any malarial disease; and it is very probable that the spleen is the principal seat of its formation, and that it is chiefly derived from a destruction of the red

FIG. 82.



A drop of blood taken from the finger of a man the subject of malarial cachexia. The granules of pigment, as well as the larger fragments of irregular form, are seen among the blood-globules. The pigment was for the most part black; some of the particles were reddish brown.

globules. The pigment is in great part carried from the spleen to the liver, where it remains; or it passes through this viscus to the lungs, brain, and kidneys. The clogging of the coarser fragments in the capillaries of the liver may, as Frerichs suggests, by interference with the portal circulation, explain the intestinal hemorrhage and diarrhœa which attend some severe cases of remittent fever; while the cerebral phenomena, or albuminuria, hæmaturia, or suppression of urine may also be caused by retention of pigment, in the one case in the capillaries of the brain, in the other in those of the Malpighian bodies. Thus, then, would be solved some of the anomalous symptoms of malarial fevers.

But the abundance of pigment does not occur in all; and whether a peculiar quality or an unusual intensity of the miasm produces

it, is undetermined. In a diagnostic point of view, though from the very evident grayish or ash-colored hue of the skin, and the singular character of the symptoms, we may suspect that we have to deal with the pathological state under discussion, we cannot be sure of it until we have examined the blood microscopically. And here, too, it seems to me that the question of the amount of pigmentary matter present must not be overlooked. For pigment may be found in the blood of those who never, to their knowledge, have had intermittent fever, and who certainly present no signs of malarial poisoning.\*

Parasitic formations have been described by Laveran† as present in the blood of those suffering from malarial fevers; and these minute appearances of the blood, as regards both the free bodies and the masses of protoplasm which are found in the red corpuscles of the blood, are of distinct diagnostic value. In the latter is reddish or black pigment, due to the action on the hæmoglobin. The hæmatozoa of malaria in their varied forms are represented in the accompanying drawings, made from cases mainly under my care at the Pennsylvania Hospital, and drawn by Dr. Joseph Leidy, Jr. Bodies 1, 2, and 3 were found in the blood of a case of malarial paralysis.‡

To the peculiar appearance of the tongue which those under the malarial influence may show, Osborn has directed particular attention.§ There is a distinct lateral boundary of the organ, an appearance of indentation transversely, and the inferior surface appears to have encroached upon the superior and lateral borders.

In children, a fever of remittent type is observed, called *infantile remittent*, which is rarely a miasmatic disorder. It is often a gastro-enteritis connected with verminous irritation or produced by errors in diet; or a typhoid fever,—an affection which now

---

\* J. F. Meigs, Pennsylvania Hospital Reports, vol. i., 1868.


† Bulletin de la Société Médicale de Paris, 1880.

‡ Dr. Leidy called my attention to the effect of heat and of cold on ordinary blood-corpuscles: they are distorted, and changes are produced which might easily be taken for some of the forms of the hæmatozoa of malaria. It is also very important to cleanse the part thoroughly from which the blood is taken, as epithelial cells filled with pigment or free pigment from the skin might otherwise be mistaken for malarial changes in the blood.

§ Transactions of the American Medical Association, vol. xx.

and then occurs even in very young children. What has given rise to this confusion is, that all febrile diseases in children exhibit a much greater periodicity than in adults, and in all cerebral symptoms are apt to be present. To distinguish the two maladies mentioned from true remittent fever, we must study particularly

FIG. 83.



their manner of beginning and their probable origin, and note the peculiarities of the abdominal symptoms. Then we may lay stress on the irregular mode and the unequal duration of the febrile exacerbations. Sometimes, also, by close scrutiny, the characteristic



eruption of a low continued fever may be found in an apparent remittent.

But some of these cases of remittent fever are really of malarial origin ; even in very young children this may be their source. I saw, for instance, some years ago, a little girl, three years of age, who had a distinctly malarial remittent fever, which was checked by antiperiodics. During the violent exacerbations she was very delirious ; her face had a most anxious, frightened look ; her screams could be heard all over the house. In the remissions she was perfectly sensible, but there was gastric irritability, and the bowels were very constipated. I have met with a similar case in an infant of eighteen months.

**Congestive Fever.**—This is a malignant, destructive, malarial fever, which may be either of the intermittent or of the remittent form. The pernicious attacks are of the tertian or of the quotidian type. While they are at their height, there is intense congestion of one or several internal organs, with a dangerous perversion of the function of innervation. From this state the patient may rally, but only to fall a victim to another paroxysm, unless art intervene. The temperature during the chill and subsequent fever ranges from  $104^{\circ}$  to  $108^{\circ}$ . Sugar is apt to be found in the urine much more commonly than in ordinary intermittent fever.

The symptoms of this violent malady vary according to the organ more specially disturbed, and to the extent of the derangement of the nervous system. We have, thus, several distinct varieties, of which I shall describe the most prominent.

The *gastro-enteric* form is common in our Southwestern States. Its distinctive features are nausea and vomiting, purging of thin discharges mixed with blood, intense thirst, and an equally intense desire for air. There is little abdominal pain or tenderness, but a weak, frequent pulse, and very great restlessness. The patient complains of a sense of sinking and of weight, and of burning heat in the stomach. His breathing is deep-drawn ; to each expiration succeed two short inspirations. The face, hands, and feet are pale and cold ; the features shrunk. Sometimes these symptoms continue for several days, and gradually increase in intensity, in spite of nature making efforts at reaction. More frequently reaction does take place ; the temperature is very high,

the pulse feeble, and the stormy symptoms subside or wholly yield, until another outbreak, which is very apt to be deadly, occurs. The usual length of the fatal paroxysm is stated by Parry,\* in his short but interesting sketch of the disease, to be from three to six hours.

The *thoracic* variety of the malady is often combined with the one just described. Its most characteristic trait is violent dyspnœa, caused by overwhelming congestion of the lungs. It is perhaps the most rapidly destructive of all the forms of the disastrous affection.

In the *cerebral* variety there is intense congestion of the brain ; and sometimes effusion of serum into the ventricles takes place, or even rupture of the blood-vessels. The abnormal state of the brain manifests itself either by coma or by delirium. In the former case there is usually preceding stupor with occasional delirium ; the pulse is slow and full ; the face is dull, and either flushed or livid ; indeed, some of the symptoms which are observed in apoplexy show themselves. When, on the other hand, delirium is marked, we have much the same morbid phenomena as in acute meningitis: the patient is wild ; he sings, he cries. He may die in this state without coma supervening ; but a comatose condition generally succeeds rapidly to the fierce excitement. Should recovery take place, the delirium gradually ceases.

Another variety much dwelt upon is the so-called *algid* form. This is not often seen in this country ; Maillot† noticed it in Corsica and Algeria. The disease is more than a mere continuation of the cold stage of a paroxysm : commonly the characteristic symptoms manifest themselves during the period of reaction. The pulse slackens, and finally ceases ; the extremities, face, and trunk become in succession rapidly cold. There is no thirst ; the skin feels like marble ; the breath is cold ; the voice broken. The mind is clear ; the expression of the countenance impassive and like that of a dead man. There may be vomiting and choleraic discharges. These symptoms go on steadily toward death, unless decided reaction be brought about.

In none of these forms of congestive fever is the first paroxysm

---

\* Amer. Journ. Med. Sci., July, 1843.

† Traité des Fièvres intermittentes, Paris, 1836.

apt to be of a pernicious character. In the majority of instances the disease begins as ordinary periodic fever, and it is only in the second or third paroxysm that the alarming symptoms appear. Nor is the first congestive paroxysm likely to prove mortal; generally it is not until the second or third that a fatal issue is to be apprehended. Proper watchfulness will sometimes detect, even at the onset of the attack, by the unusual prolongation of the cold stage, or by the irregularity of the pulse, or by the great sensitiveness in the splenic region and by the pain which pressure there may occasion all over the body, or by an imperfect hot stage, or by the feeling of internal heat while the surface is really cold, the danger that is approaching, and arrest its further steps by the bold use of antiperiodics.

The cause of this desperate disease is evidently a highly active malarial poison; and once in the system, it remains for a long time. Should the patient even weather the first attack completely, he is not wholly out of danger; he may have a second seizure quite as dangerous within the same season. Dock \* has recorded in detail the study of a case of pernicious malarial fever characterized by an enormous development of plasmodia in the blood, with consequent anæmia and melanæmia; parenchymatous degeneration and inflammation in liver, kidneys, and stomach; thrombosis in various organs; hyperplasia of the spleen and lymphatic glands. On micro-chemical examination the pigment in the malarial parasites failed to respond to tests for iron, while deposits in the tissues themselves yielded such reaction.

*Hemorrhagic Malarial Fever.*—Closely connected with congestive fever, indeed in a certain sense a form of it, is that pernicious malady which is attracting in this country more and more attention, and is known as the yellow disease, icterode pernicious fever, malarial hæmaturia, or hemorrhagic malarial fever. It is the same disease as that which some of the French writers have long described as hæmaturic bilious fever, and is found in intensely malarial places, sometimes in epidemics. It usually occurs in those who have already suffered much from malarial fevers, and is almost always ushered in by a marked chill, longer usually and more intense than the patient has had in the pre-

---

\* Amer. Journ. Med. Sci., April, 1894, p. 379.

ceding seizure of intermittent,—for often the dangerous paroxysm is preceded by one of ordinary kind. Soon after the protracted chill, distressing nausea and vomiting are noticed, as well as headache, great restlessness, and rapidly-developed deep jaundice. The fever which follows the chill is not high, the pulse is rarely extremely rapid, the patient is very thirsty. In a few hours after the chill, pain in the right hypochondrium, in the epigastrium, and over the kidneys is encountered, and a dark-colored, bloody urine is voided. Sometimes hemorrhages occur also from the nose and bowels. The type of the fever is either intermittent or remittent; occasionally it is continuous. The bloody urine—for I know the dark-colored urine, judging from the specimens I have examined, to be bloody or to contain large quantities of dissolved hæmoglobin—is at times associated with considerable albumin and with tube-casts.

If the case progress unfavorably, the pulse rises, cold sweats occur, purpuric spots appear on the skin, and the signs of uræmic poisoning are not unusual. In the intermission or remission the symptoms abate considerably, jaundice and bloody urine cease to a great extent, perhaps almost entirely,—at least this is true of the latter symptom,—but they recur in the paroxysms, which may happen every day or every ten or twelve hours.

The disease may prove fatal in three days; but generally it lasts longer. Convalescence is apt to set in slowly, and not until the urine has entirely and permanently cleared. The liver and spleen may remain for a time greatly enlarged.

As regards the diagnosis of the disease, there are but two diseases that closely resemble it. One is *intermittent hæmoglobinuria*. Now, undoubtedly some of the recorded cases of this are cases of the malady under discussion; but in those to which the name can be fairly given the absence of marked malarial elements, of jaundice, of red blood-disks in the urine, and the want generally of fever, supply the distinguishing traits. From *yellow fever*, for which hemorrhagic malarial fever may be mistaken, it differs in the speedy occurrence of marked jaundice, in the bloody urine, in the extreme rarity of black vomit, in the course of the fever with its recurring paroxysms, and in the high degree of malarial poisoning which the history of the case proves.

Then, again, the malarial poison may affect the kidneys, pro-



ducing altered secretion and even transitory albuminuria, and may lead to recognizable organic change.\*

Before proceeding to the discussion of another subject, I shall here devote a few pages to the consideration of some of the irregular forms and modifications of malarial poisoning, and to its share in producing febrile disorders of blurred and uncertain type. Practically, this is of great importance, and specially of importance to American physicians.

In the first place, I shall speak of the *chronic malarial poisoning* so often seen among inhabitants of malarial districts. It manifests itself by lassitude, debility, torpor of the liver, and enlargement of the spleen. The stools are often black, the digestion is impaired, the complexion sallow. Occasionally attacks of jaundice occur, which rather relieve than aggravate the unhealthy state of the system. Sometimes the noxious influence shows itself in another way: the patient is seized with nausea, and with gastric irritability so great that almost everything he takes is instantly rejected. The tongue is coated, the skin dryish; but he has little if any fever. The bowels are confined, the urine is turbid. He is restless, and as weak as if he had typhoid fever; but he has neither an eruption nor diarrhoea. His sleep is disturbed, and he often suffers with hyperæsthesia of the scalp, and neuralgic pain shooting over the forehead and causing twitching of the eyelids. After remaining from six to seven days in this condition, his nails, perhaps at a certain hour every day, are noticed to become bluish; or he feels chilly, and a slight fever immediately afterward sets in. The return of these febrile symptoms is checked by quinine, and the patient enters upon a slow convalescence, remaining for a long time enfeebled. Again, there may be head-

---

\* See papers on malarial changes in the kidneys in Arch. de Phys., 1882, Nos. 1, 2, 3; and on malarial hæmaturia, T. F. Wood, North Carolina Med. Journ., 1884, xiv.; J. Cochrane, Journ. Amer. Med. Assoc., Chicago, 1885, iv.; I. J. Newton, Jr., Transact. Louisiana Med. Soc., 1885, vii.; R. H. Day, Therap. Gaz., 1886, 3d S., ii.; B. H. Riggs, Alabama Med. and Surg. Journ., 1886, i.; W. L. Van Horn, Gaillard's Med. Journ. 1887, xliii.; J. W. McLaughlin, New Orleans Med. and Surg. Journ., 1888-89, N. S., xvi.; I. T. Young, South. Med. Rec., 1889, xix.; Chambless, Med. Brief, St. Louis, May, 1891; McHatton, Times and Register, Phila., June, 1891; Ferreira, Rev. mensuelle des maladies de l'enfance, Paris, March, 1893; Stephanowicz, Wien. Klin. Wochens., No. 8, 1893.

ache, coming on at a certain hour, associated with rise of temperature. We also encounter malarial neuralgias and malarial palsies. In these, as in a case under my care at the Pennsylvania Hospital in 1889, the detection of the malarial corpuscles in the blood led to the diagnosis of the affection.

*Typho-malarial Fever.*—Fever of hybrid character, for the most part of kindred nature to those low states of malaria just described, have long been recognized by practitioners in this country. But it is only since our civil war that, owing to the publications of Woodward, they have been set apart in a separate class. Now, one of the most marked forms of “typho-malarial fever,” to adopt this, from a practical point of view, convenient name, was that curious fever which so many soldiers brought with them from the swamps of the Chickahominy.

The fever generally began with a decided chill, to which febrile excitement soon succeeded. This chill was sometimes, but not always, repeated. Many cases of the disorder showed at first distinct remissions; but if the fever lasted for more than a week it became continued. Diarrhœa was a prominent symptom from the first; sometimes it preceded the disease by several weeks. In the cases that I saw in Philadelphia, nausea, vomiting of bile, and great thirst were often present; the stools were very frequent and offensive; the eye was injected. There was generally mental confusion, and not unusually wild delirium; but no eruption,—certainly no rose-colored spots; the spleen was always much enlarged. Dark purple spots, unchanged by pressure, showed themselves at times all over the body, and often did not appear until long after the fever had left. The diarrhœa was very obstinate, and remained long after the fever. The solitary glands and Peyer’s patches were generally swollen, of dark color, but in a large number of instances not ulcerated.

I have mentioned this fever because it presented on a large scale a striking illustration of the typho-malarial disease. According to Woodward, the fever belonged to the group which was the most frequent form of camp fever during our civil war. It consisted of mixed cases, in which the malarial and typhoid elements were variously combined with each other and with the scorbutic taint, now one, now the other of these elements preponderating. Prominent among the peculiarities of the malady were a decided

tendency to periodicity, hepatic tenderness, with an icteroid hue of the countenance, gastric disturbance, excessive enlargement of the spleen, a very protracted convalescence, and the appearance throughout of the signs of a scorbutic affection. The rose-colored rash and the tympanites of typhoid fever were generally absent. Diarrhœa was ordinarily very marked, and was apt to be persistent.

Now, except the scorbutic symptoms, similar cases are seen by all of us to this day throughout large portions of the United States, and the clinical manifestations are those of a malarial fever with prominent typhoid symptoms. In fact, I have already mentioned these symptoms when describing remittent fever, and I will here only add that they may come on early in the case, as well as develop late. They are cases of malarial fever complicated with a typhoid state, or more generally lapsing into it; and, while they present the symptoms of a typhoid condition, they are lacking in the eruption of enteric fever, and in the abdominal phenomena, if we except diarrhœa and some abdominal swelling, both of which may, however, also be absent. It is these cases, malarial primarily, in which the typhoid condition shows itself, but in which there is not the characteristic lesion of typhoid fever, to which, in my judgment, the term typho-malarial should be restricted. Yet most cases that are now called typho-malarial fever are really typhoid fever associated with malaria. They are true typhoid cases showing unwonted periodicity and greater enlargement of the spleen from a malarial complication, and should not, I think, be called typho-malaria. They are simply cases of typhoid fever with malaria, and, if we are to give them a name, might be distinguished as "malario-typhoids." Thompson\* has reported four cases of concurrent typhoid fever and malarial fever. In three plasmodia and in one malarial pigment were found in the blood.†

**Yellow Fever.**—This formidable malady takes its familiar appellation of yellow fever from the yellow tinge assumed during its course by the skin. It is a distemper met with in hot climates in low and level localities on the sea-coast. Its source is unknown; it is not malaria, nor has a characteristic micro-organism been detected. All we know certainly of the cause is, that the

---

\* Amer. Journ. Med. Sci., Aug. 1894, p. 158.

† See, also, Johnston, Trans. Assoc. Amer. Physic'ans, 1888, vol. iii. p. 8.



malady is due to a specific poison which does not exist without a high temperature, and that frost is its greatest enemy.

Yellow fever is an affection of short duration : it rarely lasts a week ; many die on the third or the fifth day of the disease. It has but one paroxysm, which is never repeated. This paroxysm may be divided into three stages, which are well marked in some epidemics, far less so in others.

The first stage, called that of reaction, is pre-eminently the febrile stage. Its average duration is from thirty-six to forty-eight hours. It usually begins suddenly, and is frequently ushered in by a chill. In rare instances this is protracted, there is great internal congestion, and death ensues before reaction occurs. But much more generally a short chill is followed by marked febrile excitement. The skin is harsh and hot ; the pulse quick and tense, although sometimes it is both easily compressible and not much accelerated ; indeed, as a rule, it declines before the temperature. The face is flushed ; the eye brilliantly injected, yet watery. The patient is conscious, restless, anxious, and complains much of the torturing pains in his forehead, loins, and legs ; and the muscles of the extremities are sore when moved. The breathing is hurried ; the stomach irritable, the epigastrium painful on pressure ; there is great thirst. The bowels are constipated ; the stools very dark-colored. The tongue is more or less coated and moist ; sometimes it is red, while at other times it remains natural throughout the disease. The febrile signs increase toward evening and lessen toward morning, but do not distinctly remit until after from thirty-six to forty-eight hours, when a remission does occur, or when, to speak more correctly, the whole aspect of the case changes.

The disorder now appears in its second stage ; the fever subsides ; the pulse falls and becomes easily compressible ; the headache is relieved ; the breathing is no longer oppressed ; the temperature declines to a little above the norm. But the gastric irritability does not wholly disappear, and a deep yellow or orange hue gradually tinges the eye and the whole surface of the body. The patient is cheerful, and wishes to get out of bed. His sufferings may be, indeed, over, his convalescence may have set in : after a few dark, biliary stools, the yellowness of the skin fades, and he slowly gets well.



But it is not often that the disease relaxes its hold so easily: more generally the deceptive improvement does not last a day, and after a brief lull the struggle for life begins. The patient grows again very uncomfortable and anxious. In truth, the symptoms of the first stage reappear with increased intensity. In addition, new signs, of the gravest import, show themselves; some of which are clearly due to the corruption of the blood that the poison has silently effected. The pulse sinks, and becomes slow and extremely irregular and compressible; the skin is cool, dry, dark, and in some cases of a bronze hue, or livid, and spots may occasionally be seen on its surface. The stomach is as irritable as before, but the act of vomiting is easier; and, without much retching, large quantities of altered blood, or "black vomit," are ejected. Blood oozes from the mouth, from the gums; sometimes from the eyes and nostrils, from the bowels, and from the vagina;\* or hemorrhage takes place into internal cavities, and the blood is retained.†

The phenomena of collapse become now more and more unmistakable: the black vomit often ceases, because the contractile power of the stomach has ceased; a low, muttering delirium sets in, and the patient dies prostrated. Yet the mind may remain clear almost to the last, and the strength be but little impaired. Should reaction take place, recovery is only very gradual.

But yellow fever does not at all times and in all localities present precisely the same degree of intensity or the same group of symptoms. Sometimes it exhibits frank, active febrile phenomena; at other times there is little febrile excitement, but a disposition to internal congestions and to early prostration. This congestive form is far more dangerous than the inflammatory. Yet both are highly destructive. From 10 up to 75 per cent. are the figures representing the mortality of this fearful malady. Omitting the instances of an exceptionally mild type, the average is calculated, in the elaborate work of La Roche,‡ to be 1 in 2.32. The more rapidly the stages succeed one another, the more danger-

---


\* Cases in the epidemic of 1856-57 at Lisbon, reported upon by Lyons, London, 1858; also by Alvarenga, *Fièvre jaune à Lisbonne*, Paris, 1861.

† In a case at the Pennsylvania Hospital the pericardium was filled with blood resembling black vomit.

‡ Yellow Fever, Philadelphia, 1855.

ous the case. The occurrence of black vomit, of great epigastric tenderness, of hiccough, of suppression of urine, of delirium, of early jaundice, of oppression in breathing, of convulsions, of a fiery, glistening eye, and of petechiæ, warrants an unfavorable prognosis. "Walking cases," or those in which the patients walk about until they suddenly eject black vomit, always terminate fatally.

FIG. 84.



Temperature of yellow fever in a case ending in recovery recorded by Bemiss.

As regards the temperature in yellow fever, the maximum elevation is attained upon the first, second, and third days of the disease, ranging from  $102^{\circ}$  to  $110^{\circ}$ ; it then falls in the stage of calm, to rise for the most part again in the stage of collapse, though it never attains the high temperature characteristic of the first stage, and never rises so rapidly. The elevated temperature of the first days may, however, continue with little variation until the sixth day, when the remission becomes marked. A complete remission usually happens on the morning of the third day, but may not occur until the fifth day or the ninth. Whenever it takes place, the speedy defervescence is very characteristic. Even in these longer cases there is a decrease in temperature preceding a fresh rise, which occurs in paroxysms of two days each. Slight rises in temperature are neither uncommon nor grave after the marked fall in the second stage. But when the temperature rises rapidly in this stage of calm it is of most

serious meaning. In the stage of calm the absence of fever may be complete; but generally the defervescence is only partial: a remission, therefore, rather than an intermission.\*

Yellow fever has rarely any complications. It may, however, seize upon those affected with other diseases. It has been specially noticed that it is frequently intercurrent in surgical and obstetrical cases.†

The recognition of yellow fever is, generally speaking, easy. The intense pain in the back, limbs, and forehead; the appearance of the eye; the color of the skin; the short duration of the febrile symptoms; the nausea; the epigastric tenderness; the black vomit; the albuminuria,—constitute a group of symptoms which unmistakably mark the disease.

But let us look at the points of contrast which yellow fever presents to other affections. It differs from *plague* by the absence of buboes and of carbuncles, and by the much more frequent occurrence, on the other hand, of jaundice and black vomit. Then, too, the red, suffused eye and the single paroxysm are not witnessed in *plague*. The febrile malady may run on to a state of collapse as complete as in *Asiatic cholera*; but, unlike this destructive disease, the symptoms of entire prostration are preceded by fever, and not by vomiting or purging of rice-water.

The lines of demarcation between the ordinary forms of continued fever and yellow fever are very broadly drawn. It is distinguished from *relapsing* fever by the different countenance, by the supra-orbital pain, by the soon-occurring remission, and, above all, by the extreme rarity of a relapse and the infinitely greater mortality. To *typhoid fever* it bears so slight a resemblance that it is scarcely possible to confound the two affections: one, a short, severe disease, with its peculiar physiognomy and gastric symptoms; the other, a long-continued malady, of low type, with its characteristic eruption and enteric signs. It is only when yellow

---

\* In addition to the authors quoted, see on the temperature Faget, New Orleans Med. and Surg. Journ., 1873-74; Bemiss, Amer. Journ. Med. Sci., 1880, and article "Yellow Fever" in Syst. of Pract. Med. by American Authors, and the temperature charts of Naegeli, of Rio Janeiro, as given by Jaccoud, Pathologie interne.

† S. M. Bemiss, Clinical Study of Yellow Fever, Amer. Journ. Med. Sci., April, 1880.



fever is protracted beyond the ninth day that the diagnosis is rendered doubtful; and then we have generally the history to guide to a correct understanding of the case. The likeness between yellow fever and *typhus* is much closer. But one is a short fever, with distinct stages; the other is a longer, much more continued fever. One has no marked cerebral symptoms; in the other the cerebral symptoms are the most prominent feature. One has but rarely an eruption, but often hemorrhages; the other has always an eruption, and hardly ever hemorrhages.

The disease most likely to be confounded with yellow fever is *bilious remittent*. In truth, the symptoms are very similar, and many of them differ only in intensity. The diagnosis of the milder forms of yellow fever from remittent fever is, indeed, extremely difficult, unless the epidemic influences prevailing be taken into account. Then, as is well known, the affections may be blended, and yellow fever become obviously periodical in its febrile phenomena. The occurrence of black vomit is not in itself a distinctive sign between the two diseases; for black vomit may be absent in yellow fever, and, on the other hand, it may, although it rarely does, occur in remittent fever, just as it has been known to occur in childbed fever, in the plague, and even in typhus.\* A valuable sign is derived from an examination of the urine. Unlike what happens in bilious fever, a trace of albumin appears in from twelve to fourteen hours after the fever sets in; then the albumin increases, and the urea and the uric acid diminish and gradually disappear, as does the bile-pigment.† The more obvious the suffusion of the countenance in yellow fever, the more marked and early is the albuminuria.‡ In children,

---

\* This statement with reference to typhus fever is made on the authority of Stokes. The occasional occurrence of black vomit in remittent fever is admitted by many authors. Some winters ago, a physician of this city brought to me, for examination, a specimen of black vomit which had the same microscopical characters that I have repeatedly found in the black vomit of yellow fever. The patient undoubtedly had remittent fever, from which he recovered.

† Ballot, Arch. Gén. de Méd., Nov. 1869; see, also, Joseph Jones, New Orleans Med. and Surg. Journ., Jan. 1874; Holland, American Practitioner, Sept. 1879; Sternberg, New Orleans Med. and Surg. Journ., 1880-81, N. S., viii.

‡ Bemiss, *loc. cit.*



albumin may be present only in the evening urine.\* An additional valuable sign is finding the plasmodium of malaria in the blood in cases of remittent fever.

When yellow fever is well marked, it differs in this way from bilious remittent :

## YELLOW FEVER.

Of short duration, ending commonly in from three to seven days.

Period of incubation from five to nine days.

A disease of one paroxysm, terminating in recovery or in collapse.

Very severe nausea and vomiting throughout; early and decided epigastric tenderness; black vomit.

Hemorrhages from gums and various parts of the body.

Tongue clean, or but slightly coated; pulse very variable, frequently becomes slow in last stages.

Highly-injected, humid eyes; often fierce or anxious expression of face.

Supra-orbital pain, and pain in back and in calves of the legs.

Very rarely delirium; mind usually clear.

Urine acid, generally contains albumin, also epithelial and granular casts and blood casts; suppression of urine common; no micro-organism in blood.

Little muscular prostration; often rapid convalescence; no sequelæ.

Almost certain immunity after one attack.

Very high mortality; disease is epidemic.

Treatment unsatisfactory.

## BILIOUS REMITTENT.

Lasts nine days or upward.

Period of incubation very variable; may extend to months.

A disease of several paroxysms, with intervening remissions.

Nausea and vomiting not so severe, and rarely as marked at the onset; neither as early nor as constant and decided epigastric tenderness; vomiting of bile and of the contents of the stomach.

No hemorrhagic tendency.

Tongue heavily coated; pulse varies less, is always quick until convalescence sets in.

Eye not peculiar; different physiognomy.

Headache; sense of fulness in head; often no pain in loins or in legs.

Delirium frequent; mind always dull.

No albumin in urine; suppression of urine rare; plasmodium of malaria in blood.

Much greater muscular prostration; slow convalescence and tedious sequelæ.

One attack seems rather to predispose to others.

Slight mortality; disease more endemic in its nature.

Very amenable to treatment.

---

\* Guitéras, article "Yellow Fever," in Keating's Cyclopædia of the Diseases of Children.

**YELLOW FEVER.**

Autopsy shows inflammation or very great congestion of stomach, and sometimes ulceration or softening. Liver enlarged, of a yellowish color, its secreting cells filled with oil-globules. Kidneys swollen, inflamed. Heart often exhibits disintegration of muscular fibres.

**BILIOUS REMITTENT.**

Autopsy shows congestion of stomach; more rarely inflammation. Liver of an olive or bronze hue, not fatty; accumulation of animal starch in liver of malarial fever, no grape-sugar.\* Kidneys unchanged, or simply congested.

**Eruptive Fevers.**

The eruptive or exanthematous fevers form a group having numerous features in common. They are all characterized by a period of incubation, during which the poison lies dormant in the system; by a fever of more or less intensity preceding the eruption; by an eruption which presents a distinct aspect in each disease, and which pursues a definite, clearly-defined course until it, and with it the febrile malady, disappears. Moreover, they are all very prone to occasion serious sequelæ; are all, in the main, disorders of childhood; rarely attack the same person twice; and are contagious. These remarks apply particularly to the three chief exanthematous fevers: scarlet fever, measles, and smallpox. In great part, too, they hold good in regard to corysipelas, described here in connection with the eruptive fevers.

**Scarlet Fever.**—Scarlatina affects both children and adults, and is marked by great heat of skin, frequent pulse, sore throat, and an early scarlet eruption. These symptoms are preceded by an uncertain, generally a short, period of incubation, but soon exhibit their striking features. The febrile excitement is characteristic; the skin is very hot and generally dry, and the rapidity of the pulse so great that often by this sign alone we may, especially in the midst of an epidemic, predict the coming eruption. Vomiting, too, is a frequent symptom at the beginning of the illness. The temperature, which may reach between  $105^{\circ}$  and  $107^{\circ}$ , does not fall with the appearance of the eruption. The highest temperature occurs on the second or the third day.† The

\* Joseph Jones, Medical and Surgical Memoirs, vol. ii., New Orleans, 1887.

† Hatfield, article "Scarlet Fever," in Amer. Text-Book of Diseases of Children, 1894.

temperature continues high until the eruption is completed and at its height. It slowly declines as this fades, and with the occurrence of desquamation attains the norm; but it may persist, with marked morning remissions and evening exacerbations, when the eruption has gone and during the earlier stages of desquamation.

The *rash* appears on the second day of the disease. It comes out almost simultaneously all over the body, although, on close scrutiny, it may be soonest perceived on the neck and the breast. At first the surface exhibits an almost uniform red blush, which disappears momentarily on pressure, or rather pressure leaves a white stain on the skin, which quickly again reddens from the periphery to the centre. Soon, however, the eruption presents an unequal aspect: it is of more vivid scarlet hue in some parts of the body, as in and around the flexures of the joints, and is not everywhere smooth. Here and there are seen elevated rough points of darker tint, edged by the red integument, and not unfrequently vesicles containing a thin fluid. The skin is very hot and itchy, and tumefied, especially on the hands and feet. The eruption declines on the fourth or the fifth day; by the seventh or eighth, the cuticle begins to come away in large flakes. Sometimes the rash, when at its height, recedes and then appears again. In malignant cases it comes out late, and is either pale and indistinct or dark and livid. In some instances it is wanting. Some years ago, I saw this "*scarlatina sine exanthemate*" in a lady, who, watching over the sick-bed of her daughter, contracted the disease and went regularly through it, even to its sequelæ of disorder of the kidneys and swelling of the salivary glands, but in whom not a trace of an eruption could be detected.

The *sore throat* of scarlatina is almost as constant and as characteristic as the scarlet rash. It shows itself early, sometimes before the eruption, and rarely waits until the third day of the complaint. At first the throat-affection consists in a diffused redness extending over the tonsils, palate, and half-arches, and in a swelling of the tonsils: the patient complains of pain in his throat, augmented by pressure and by swallowing, and of stiffness of the muscles of the neck. After a few days, if the disorder be severe, irritating discharges occur from the inflamed surfaces, and patches of false membrane and superficial ulcerations are

seen in the fauces. The glands of the angle of the jaw become much tumefied, and, by pressing on the cervical vessels, produce a tendency to drowsiness and stupor. These are grave symptoms; their occurrence, indeed, is indicative of one of the main dangers in these "anginose" cases of the disease.

The false membranes which are developed last about five or six days; they form as well as reform in patches, and are very easily removed. Sometimes they extend to the larynx; but this does not often happen, and, even when it does, the symptoms of croup, in the opinion of Barthez and Rilliet, do not arise. The acid discharges and the decomposing membranes often occasion a most fetid breath, and, by being swallowed, a persistent diarrhœa.

The *tongue* has a peculiar look. At first it is thickly coated, and its borders only are red; but soon the fur is cast off, and the whole organ becomes very red and its papillæ prominent. After it has presented this appearance for six or eight days, it returns to its normal condition. In bad cases it is extremely dry and of a brownish hue.

In children the disease frequently sets in with convulsions. In truth, cerebral symptoms of one kind or another are not uncommon at all stages of the malady; yet great differences are observed, in this respect, in different epidemics. In some cases of malignant character, the vomiting, the screams, the grinding of the teeth, the occurrence of delirium and insomnia, make the attack look, at the onset, like one of acute meningitis; but the eruption soon sets all doubt at rest, and, even before it is noticed, the great heat of the skin and the extreme rapidity of the pulse point to the source of the mischief. The nervous symptoms in these dangerous instances of the affection do not, however, cease with the eruption; they may last to the end of the malady. Sometimes they are not noticed until late in the disorder, and after the period of desquamation has fully begun; but the convulsions and stupor—for these are the morbid manifestations then more specially encountered—are owing rather to a diseased state of the kidneys that has been induced, than to the immediate effect of the fever poison.

Occasionally some of the larger joints swell up, and present the appearance of subacute rheumatism. The joints are not, however, very painful on pressure, and generally only two or



three are enlarged. This form of rheumatism is evidently owing to the retention in the blood of some morbid material, and simulates ordinary acute articular rheumatism in presenting endocarditis and pericarditis as complications. Endocarditis and pericarditis occur also irrespective of articular involvement.

Further complications of the disease are dropsies, renal hæmaturia, pleurisy, local gangrene, œdema of the glottis, neuritis, diphtheria,\* and a very low state of the system. These complications are not apt to arise until at or soon after the period of desquamation; sometimes they lead to long-continued disorder, and become thus the most hazardous of the sequelæ. Other consequences of the affection, lasting, it may be, for years after the febrile attack, are a tendency to boils, swelling of the parotid and of the lymphatic glands of the neck, nasal catarrh, diarrhœa, chronic inflammation of the eyelids, and deafness from inflammation extending up the Eustachian tube to the membrane of the tympanum, or from suppurative destruction of portions of the ear. Epilepsy is also a sequel of scarlet fever, more cases being consecutive to it than to all other acute diseases combined.† Optic neuritis may follow scarlet fever, without organic change in the brain.

Of all these morbid states, *dropsy* is the most common. The effusion of fluid may be caused by the altered state of the blood; but much more generally it is owing to the poison producing an acute desquamative nephritis: albumin, tube-casts, epithelial cells, and sometimes blood, are found in the scanty urine; and we meet with severe headache, great restlessness, and œdema of the face and extremities, as the attending symptoms. Still, notwithstanding these grave phenomena, the majority of the cases recover, and the kidneys are rarely permanently injured.

The dropsy is apt to show itself between the tenth and the twentieth day of the malady. The albuminous condition of the urine may precede it by several days; yet dropsy may happen without albuminuria,‡ especially in some epidemics, and albumin in the

---

\* Trousseau, *Clinique Médicale*, tome i.; see, also, article "Scarlet Fever" in Ziemssen's *Cyclopædia* and in *Amer. Syst. of Pract. Med.*

† Gowers, *Diseases of the Nervous System*.

‡ Gee, in *Russell Reynolds's System of Medicine*; also Quinke, *Berlin. Klin. Woch.*, 1882, No. 27; Dyce Duckworth, *St. Barth. Hosp. Rep.*, 1883.

urine is not always associated with dropsy. In most cases of scarlatina it is found at some period of the disease for a short time and in small quantities; but this transitory albuminuria is not, like the albuminuria coexisting with marked anasarca, connected with many tube-casts in the urine and numerous epithelial cells.

The *state of exhaustion* noticeable at the close of the fever and while desquamation is still going on is at times great,—so great that, in young persons especially, the case wears the look of *typhoid fever*. And the resemblance is heightened by the occurrence of diarrhœa associated with a swelling of the solitary and agminated glands. But the signs of desquamation, the sore throat, the enlargement of the cervical glands, and the history of the affection furnish distinctive marks of the utmost value.

The statements that have just been made concerning the diverse complications of the malady are mainly of interest on account of their exhibiting the intricate diagnostic questions which may arise. Of the recognition of the disorder during the febrile stage it is not necessary to say much, as ordinarily it is not difficult. The distinction between it and the other exanthematous fevers may be seen by glancing at the table, to which a place is elsewhere assigned. I shall only here mention, as bearing upon the differences between scarlet fever and *measles*, that cases are occasionally encountered in which the eruption alone is too ill defined to become the sole basis of an opinion, and that then we have to lay the greatest stress on the presence or absence of catarrhal symptoms and sore throat, and on the march of the symptoms. So, too, with reference to *smallpox*. The rash preceding the formation of the pustules may have so strong a resemblance to that of scarlet fever that a scrutiny of all the attending circumstances, and a careful watching of the eruption for at least a day, are requisite for the detection of the true nature of the case.\*

An erythematous rash, appearing in blotches everywhere except on the face, has been noticed in *membranous croup* and in *laryngeal diphtheria* after the operation of tracheotomy.† But it is very irregular, runs a rapid course, and is not followed by desquamation; a point, it may be here mentioned, distinguishing all the forms

---

\* The disorders may also be combined. See the cases of Marson, *Medico-Chirurg. Transact.*, vol. xxx.

† Bericht des k. k. Krankenhauses, Weiden, 1865.

of irregular rashes happening at times—though very rarely—in diphtheria, from the scarlet fever eruption. As the result of gonorrhœa we may have symptoms of a low fever associated with a cutaneous rash like that of scarlet fever. The history and progress of the case chiefly distinguish this *pseudo-scarlatina*.\* The same is true with reference to the so-called surgical scarlet fever. It shows an eruption that may be like that of scarlet fever, though the throat symptoms and the sequelæ are lacking.

Like measles, scarlatina may be mistaken for that curious form of eruptive fever called rubella or rubeola. But this really resembles measles more closely, and in examining it presently the differences between it and scarlet fever will become apparent.

An affection with several features like scarlatina is breakbone fever, or *dengue*. The points of dissimilarity may be learned by referring to the description of the malady given farther on.

Scarlet fever may go on concurrently with other fevers. It has been observed with typhoid fever, and with varicella.† Edington and Shakespeare have described a special bacillus to which they believe the disease to be due.

**Measles.**—The symptoms precursory to the specific eruption of this affection are fever, watery eyes, frequent sneezing, flow from the nose, and cough; in fact, all the manifestations of an acute coryza or catarrh. To these diarrhœa is in many instances added, indicating a simultaneous irritation of the intestinal mucous membrane. On the fourth day after the beginning of the morbid signs, a rash is perceived on the face and neck; thence it continues to extend, until, in the course of two or three days, the whole body is covered. The temperature during the first day of the disease is generally from  $102^{\circ}$  to  $103^{\circ}$ ; if higher, the attack is likely to be severe. On the second or third day—usually on the second, when it may be but  $98.6^{\circ}$  or  $99^{\circ}$ —it is markedly lower, and it rises again on the evening of the third or on the fourth day to decided fever heat. The temperature does not at once decline with the rash. Indeed, it is apt to go on rising for twenty-four to thirty-six hours; the occurrence of the eruption does not alle-

---

\* Ballot, Arch. Gén. de Méd., Sept. 1882. The same author calls attention to a puerperal pseudo-rubeola, a false measles, from blood-infection.

† Church, St. Barthol. Hosp. Rep., 1881; Lond. Med. Record, Nov. 1883.

viate the febrile symptoms ; on the contrary, while it is spreading to the trunk and the lower extremities, the constitutional disturbance lasts, or more generally increases. But as soon as the rash has fully reached its height, the defervescence is rapid ; and from the fifth to the seventh day of the disease the temperature sinks until it is but little above the norm. By the ninth day of the disease both fever and rash have left. Frequently then the cuticle comes away in fine scales, and this desquamation is attended with very annoying itching. The patient, now that he is convalescent, shows his illness : he is pale and somewhat emaciated. Often he still coughs, and his eyes are slightly inflamed. These signs are not unusually the last to disappear. Paralysis, of cerebral, spinal, or peripheral origin, may occur in the sequence of measles.\*

Of all the symptoms mentioned, two are, in a diagnostic sense, of pre-eminent importance : the catarrh and the eruption.

The *catarrh* is nearly constant. It is true that a variety of measles is recognized,—“*rubeola sine catarrho* ;” but this is very rare. Generally speaking, the coryza and catarrh decline with the eruption ; occasionally, however, they remain for some time after the rash has left. The feature which distinguishes these catarrhal symptoms from those of influenza consists in the eruption : before this happens, the diagnosis is uncertain, though we may often suspect measles by the look of the face, the greater intensity of the febrile signs, and the knowledge that the disease is prevailing in the community.

The *eruption* is peculiar : it consists of slightly-raised red spots, which coalesce and form blotches of an irregular, crescentic shape ; between these blotches the skin is of natural color. The eruption disappears first from the face ; in other words, it disappears in the same order in which it appeared. As it fades, which it does on the third or fourth day of its appearance, it becomes brownish, and subsequently of a yellowish tint. In its earliest stages it is similar to the papulæ of smallpox ; and this similarity may be heightened by its being mixed, as it sometimes is, with a few miliary vesicles. But after the first day of the rash there is

---

\* Allyn, Medical News, Nov. 28, 1891, p. 617 ; Carpenter, Medical News Feb. 13, 1892, p. 183.



little room for doubt. In the one case the spots remain as they were ; in the other, they change into pustules.

A question may sometimes arise as to whether the eruption be that of *typhus fever* or of measles. Both are coarse, both often not unlike in color, and both may be developed about the same time. Generally speaking, however, the eruption of typhus fever shows itself several days later than the rash of measles ; and, although coarse, it is not crescentic, and is found on the trunk and extremities rather than upon the face. Moreover, the physiognomy, the excessive prostration of strength, and the marked cerebral symptoms of the low fever are such as to render a differential diagnosis seldom difficult.

Measles is usually met with in children ; but it may be encountered in adults, especially among soldiers, and is in adults a much more severe complaint than in children. In the latter it is not an alarming disease. Only occasionally does it occur in epidemics which present a malignant character. Its greatest danger commonly consists in the eruption disappearing prematurely or appearing but partially, and in the severity of the thoracic complications. These are either acute bronchitis or acute pneumonia.

*Acute bronchitis* may occur at any period of the disorder, and involve the finer tubes. But it does not generally set in with severity until the eruption has reached its height or is beginning to fade. In young children, symptoms of inflammation of the larynx, or of croup, are at the same period apt to manifest themselves. Acute pneumonia, too, either croupous or bronchopneumonia, the latter most often, is met with at this stage of the malady, or sometimes even after convalescence has apparently begun. We may suspect that mischief is going on within the chest, if the breathing be very oppressed and the pulse continue to be rapid ; but to detect early the hazardous and insidious complication we have to depend chiefly on physical exploration.

Occasionally the thoracic affection leaves a chronic bronchial disease, or a persistent cough and night-sweats point to the development of tubercles. It may, in individual cases, be extremely difficult to decide with which of these morbid states we have to deal. Emaciation and a chronic cough are found in both chronic bronchitis and phthisis ; and the physical signs of

tubercular consumption are, in children, notoriously ill defined and untrustworthy. We may be obliged, therefore, to depend upon finding the tubercle-bacillus before coming to a definite conclusion. The pneumonia of measles has been attributed to the bacteria which are detected in the nasal mucus in measles penetrating in large numbers into the lung, and there setting up inflammation.\*

At times we meet with anomalous forms of measles. Such was the "*rubeola notha*" that prevailed extensively in London about thirty years since,† and in which there was a papular eruption like that of ordinary measles, but distinguished from it by its dusky hue, and by the papulæ not being arranged in crescentic clusters, being less obvious, or not appearing at all, on the limbs. A similar anomalous exanthem was common in Philadelphia during the winter of 1865-66, occurring at a time when both measles and scarlatina were frequent, and particularly the former. The eruption, which occurred in patches of dark hue, was principally confined to the face; there was no sore throat.

Perhaps these anomalous forms of measles are rather varieties of rubella than of measles. An affection formerly very common, *miliary fever*, would be also a source of much confusion were it in our day often encountered. But epidemics of miliaria are now extremely rare. Yet we know that it is a disorder with a prodromal stage of two or three days, during which great irritation of the skin, debility, and a feeling of suffocation are usual. The marked disease begins with profuse sweating and with severe fever, and præcordial and epigastric distress. These symptoms last until the appearance of the rash, generally on the third or the fourth day, though sometimes not until much later, and then, as a rule, slowly subside. The rash appears first upon the neck and the breast, and consists of numerous round or irregular spots, in the centre of which vesicles arise that finally burst and form crusts. The disease ends with desquamation and generally in a slow convalescence. The sweating, the oppression and præcordial pain, and the peculiar eruption distinguish chiefly this epidemic disease from measles.

---

\* Cornil et Babès, Arch. de Phys., Aug. 1883.

† Babington, Lancet, May 7, 1864.

**Rubella.**—The most striking resemblance to measles is furnished by rubella. This, called by the Germans *Rötheln*, and often spoken of as “German measles,” is not a hybrid of measles and of scarlet fever, but a special exanthem, which occurs in epidemics. It displays a red eruption, ushered in by a chill, followed by slight fever, which is accompanied by coryza, cough, and sore throat. The fever prior to the eruption lasts for two or three days, but this is far from constant; indeed, it often does not last more than half a day, or it may be of a week’s duration.\* The temperature rarely exceeds 102.5°. The rash may come out all over at once, or spread in a day or two over the body; it generally appears first on the face and neck. It is most distinct on the face, the scalp, the neck, and the trunk, being more scattered on the extremities; it is specially distinct about the mouth. It first resembles measles, but the spots are round or oval, and smaller and paler, and they soon run together in irregular patches, unlike the well-defined crescentic eruption of measles; they show no tendency, however, to become generally confluent. The patches are of variable size, and, unlike the rash of scarlatina, are surrounded by healthy skin; small spots range themselves around the large ones. They are of deepest color in the centre, but not bright-colored as in measles, nor of the dark red of severe scarlatina, are elevated, and very much influenced by pressure. The eruption lasts ordinarily four or five days, but in severe cases eight or ten. It gradually fades, but it may happen that it fades on the face before it has fairly come out on the legs, and desquamation may ensue, though the scales are small, and never in size like those of scarlet fever. During the continuance of the rash, which is attended with much itching, the general symptoms are much aggravated, except the fever, which indeed may be perceptible only at the beginning of the affection; the sore throat and catarrh may be severe, and attended with hoarseness and with inability to swallow; there are congestion of the conjunctivæ and pain in the eyes. Osborn has called attention to enlargement of the small glands at the edge of the hair on the postero-lateral sides of the neck as a pathognomonic sign.† As the rash fades, the other

---

\* Edwards, article “Rubella,” in Keating’s Cycl. of Diseases of Children.

† Weekly Med. Rev., Dec. 24, 1887.

symptoms subside. Swelling and even suppuration of the cervical glands are not uncommon sequelæ.

The disease may be very difficult to distinguish from measles, except when it is epidemic and affects those who have already had measles. The more sudden onset, often almost feverless, the milder course of the complaint, and the peculiarities of the eruption already spoken of, are guides in separating individual cases. But the appearance of the rash may be ill defined and very misleading. The following table exhibits the differences between well-marked cases of rubella, measles, and scarlet fever :

RUBELLA.	MEASLES.	SCARLET FEVER.
Premonitory symptoms often wanting, but frequent sore throat. If attack severe, there are loss of appetite and drowsiness for twenty-four hours before eruption.	Premonitory symptoms common, such as lassitude, loss of appetite, headache, vomiting, watery eyes, catarrh, cough.	Premonitory symptoms; usually feeling of lassitude for a few hours, frequently vomiting. If attack slight, patient feels tired, but complains only of sore throat.
Incubation from nine to twenty-one days; usually eighteen days.	From seven to fourteen days.	From a few hours to seven days; rarely beyond five days.
Eruption is mostly the first noticeable symptom; dots, rosy-red, with well-defined edges, first behind the ears, on scalp and face, around mouth; extends to neck and chest; gradually covers entire body. Dots become larger, coalesce, and form patches.	Appears on fourth day; first behind ears, then on scalp and forehead; gradually spreads all over face, body, and limbs, forming crescentic blotches. Eruption is papular in character and dark red in color; never bright rose-red.	Diffuse, dusky red; papular; just behind ears; presents goose-flesh appearance; appears early about clavicles and on chest, and on covered parts of the body; intensely hot to touch.
Fauces look dry, with a dark mottled red hue; little relation of appearance of fauces to extent of rash. Sore throat may disappear, to recur in last stages of the disease.	Fauces red and swollen throughout activity of the disease.	Fauces vary in appearance from slight to intense dusky redness, with marked swelling, and sometimes with white spots of inspissated secretion; intensity bears direct relation to skin-eruption. Sore throat throughout disease.



## RUBELLA.

Eyes pink-red and suffused.

Lymphatic glands generally enlarged, tender, hard, notably the posterior cervical, the axillary, and the inguinal.

Catarrhal symptoms and cough wanting; there may be a little flaky desquamation; frequently none.

Kidneys rarely affected; may be transient trace of albumin.

Diarrhoea none.

Patient, as a rule, does not feel ill.

Tongue clean or slightly furred.

Pulse normal or slightly accelerated; maintains ratio to temperature.

Temperature varies between 103° and 104°.

Duration short; a few days.

## MEASLES.

Eyes red and watery; photophobia.

Not usually affected; the posterior cervical rarely and slightly; bronchial glands always enlarged.

Catarrhal symptoms and cough constant; a little flaky shedding of the epithelium, varying according to the intensity of the rash.

Kidneys not affected.

Diarrhoea frequent.

Usually feels illness much.

Tongue slightly furred.

Usually accelerated; maintains ratio to temperature.

Temperature usually from 101° to 104°.

Convalescence more protracted; often patient very weak.

## SCARLET FEVER.

Eyes unaffected.

Lymphatic glands of throat and neck scarcely discernible during first few days, but subsequently may be enlarged in proportion to the severity of the faucial affection.

Catarrhal symptoms and cough absent, or slight throat cough, unless a pulmonary complication. Desquamation in proportion to the extent of the eruption; continues for many weeks; begins as a peeling of the tongue, extending to lips, face, and ears; free about hands and feet.

Kidneys often implicated; albuminuria; acute nephritis common.

Diarrhoea not uncommon.

In slight cases light illness; in severe cases grave illness.

Coated with a thick, white fur, peeling off from the tip and edges on the fourth day, leaving the "strawberry" tongue.

Pulse accelerated out of proportion to temperature.

Ranges from 99° to 106°; proportionate to rash, but not to pulse.

Gradually subsides in from four to seven days; desquamation begins as eruption fades, and may continue for seven or eight weeks or more, lasting longest on hands and feet.

RUBELLA.	MEASLES.	SCARLET FEVER.
Infectiveness lasts from ten to fourteen days if disinfection efficient.	Infectiveness does not last for more than from fourteen to twenty days, if disinfection efficient.	At onset only slightly infective; is very infective after first forty-eight hours; infectiveness may continue for six or eight weeks or longer.
Sequelæ few and not frequent; glandular enlargements may follow.	Pneumonia, bronchitis, pleurisy, ophthalmia, otitis, etc.	Nephritis; suppuration of submaxillary and other lymphatic glands; otitis; arthritis; endocarditis.
Usually complete recovery in two weeks.	Usually complete recovery in two weeks; sometimes followed by prolonged period of ill health.	Usually complete recovery; sometimes prolonged convalescence from sequelæ; mortality high in the very young.

*Typhus fever*, at least as regards the eruption, has some similarity to German measles. But the severe fever, the far greater gravity of the constitutional symptoms, the rash not appearing on the face, and the absence of catarrhal symptoms, render it strikingly unlike the latter affection.

Rubella is contagious, and affects especially children; it is extremely uncommon after forty years of age. Second attacks are also very rare. It does not protect from either scarlet fever or measles, nor do they from it.

**Smallpox.**—Smallpox, or variola, attacks both children and adults. It is a highly contagious malady, spreading rapidly among those who are unprotected by vaccination, and among masses of men.

The chief symptoms of the stage of *invasion* are chills, fever, and pain in the back. The fever runs very high, and exacerbates markedly toward evening. The pain in the back is severe, particularly in grave cases; it may be attended by pain in the limbs like those of rheumatism; there are also nausea, vomiting, headache, and great restlessness. All these symptoms subside at the end of the third or on the fourth day, when an eruption shows itself on the lips and forehead, and soon extends to the trunk, and from the trunk to the extremities; and with the appearance and the spread of the eruption there is a very decided fall in temperature.

At first the *eruption* has the appearance of papulæ; but on the second and third days the coarse spots undergo a decided change. At the top of each papule appears a vesicle, which gradually becomes larger, and fills up with a milky, thick fluid; in short, becomes a pustule. By the fifth or sixth day this change has been fully accomplished, and the pustules are spheroidal and lose the umbilicated look which they had while forming. On the eighth day matter begins to ooze from their edges, and a secondary fever sets in, lasting for three or four days,—until, indeed, all the pustules are broken; this secondary fever is sometimes ushered in by a chill; it is of remittent type, and the evening temperature marks between  $103^{\circ}$  and  $105^{\circ}$ . By the time it subsides, crusts form where previously there had been pustules; and as these crusts dry and fall off, the skin beneath is seen to be of a red color, which only very gradually fades, and here and there are noticed those scars and pits which the patient carries during the remainder of his life. Preceding the characteristic eruption in smallpox a red rash is at times noticed in the pubic and the inguinal regions, which is very significant.\*


When the pustules are in great abundance, they run together, constituting *confluent* smallpox. The eruption may be discovered a day earlier than in the discrete form, and the rough, red blotches are often so thickly clustered as to give a uniformly red aspect to the whole surface. When the pustules completely fill up, whole portions of the face or of the trunk seem to be covered by one extensive pustule, which gradually dries into a continuous brownish and most disfiguring crust. While the process of maturation is going on, the features are observed to be greatly swollen; the eyes may be hidden from view; the nose and lips are tumid. The patient complains of the tension of the skin, and not infrequently of sore throat and of a steady flow of saliva from the mouth,—a symptom that may be also met with in measles. The secondary fever is violent, far more so than in discrete variola. It may not appear until a day or two later, but lasts longer, shows a higher temperature, and is the period of danger, since it is at this time that death is most apt to happen. Death is sometimes preceded by extraordinarily high temperature,  $108^{\circ}$  or upwards.

---

\* Fagge, Practice of Medicine, vol. i p. 223.

A fatal issue is often preceded by a dry tongue, by delirium, and by great restlessness; by what, in fact, are called typhoid symptoms. Sometimes death is occasioned by attacks of dysentery or of diarrhoea, by affections of the larynx or the trachea, by

FIG. 85.



Temperature in the severe form of variola; death during the secondary fever.  
(After Wunderlich.)

acute endocarditis, or by plugging of a vessel in the brain; \* by some complication, therefore, which the worn and irritated frame is unable to withstand. A case of variola has been reported complicated, during convalescence, by convulsions, followed by left hemiplegia, in which after death an area of softening was found in

\* Quinquad, Arch. Gén. de Méd., Sept. 1870.



the motor area of the right cerebral hemisphere, due to vascular occlusion.\* Cases of variola have also been observed complicated by purulent peritonitis.† Now and then death takes place from supervening pleurisy or pneumonia or bronchitis; but an unfortunate termination from maladies of the respiratory organs does not occur in the secondary fever only, as these affections are also encountered during the period of eruption. Sometimes the patient sinks at the very onset of the disease. In these *malignant* cases, mostly met with at the beginning of an epidemic, he dies from the virulence of the poison. He is stupid, delirious; the eruption seems, as it were, to struggle to reach the surface, is ill defined and of a livid hue, and may fail to appear until after death. Many of the malignant cases, too, are of the hemorrhagic type, marked by petechial blotches and ecchymoses, and profuse hemorrhages from mucous membranes. Bay‡ has found in the lymph obtained from a case of confluent smallpox, as well as from a large number of vaccine-points, non-motile bacilli with a long diameter of from 0.6 to 1  $\mu$  and a short diameter of from 0.2 to 0.3  $\mu$ , which he believes to be the specific cause of both variola and vaccinia.

Smallpox is occasionally met with during the progress of other disorders, blending its symptoms with those of the complaint to which it becomes superadded. It is thus found as an intercurrent affection in typhoid fever, in typhus, in scarlet fever, and in measles; yet even then there is no difficulty in recognizing its peculiar traits,—its lumbar pain and characteristic eruption. Ordinarily the detection of variola is extremely easy, except at its onset. But the points of similarity it may present, in its early stages, to typhus fever, to erysipelas, and to several other diseases, have been already discussed, and need not be repeated; we have often to wait the course of the eruption before framing a positive diagnosis from the symptoms alone, and without taking into account the epidemic influences prevailing. When the disorder is fully developed, all difficulty in its diagnosis ceases. Let us here look at the marks of distinction between it and the other principal eruptive fevers, premising the statement that in the period of invasion the pain in the loins is the most significant differential sign.

---

\* Davezac and Delmas, *Journal de Médecine de Bordeaux*, 1893, No. 38, p. 421.

† Auché, *Bulletin Médical*, Jan. 25, 1893.

‡ *Medical News*, Jan. 26, 1895, p. 92.

TABLE EXHIBITING THE DIFFERENCES BETWEEN SCARLET FEVER, MEASLES, AND SMALLPOX.

SCARLET FEVER.	MEASLES.	SMALLPOX.
Period of incubation generally a week or less.	Period of incubation generally from seven to fourteen days.	Period of incubation generally about twelve days.
Fever, with very frequent pulse; persists unabated during eruption.	Fever, with moderate frequency of pulse; not relieved, but rather increased, by eruption.	Fever, with bounding pulse, and pain in the loins; great relief from occurrence of eruption.
Eruption on second day, first on neck and chest; spreads rapidly.	Eruption on fourth day, first on face; spreads gradually, in course of about forty-eight hours, to rest of body.	Eruption at end of third or on fourth day; first on lips and forehead; a preceding red eruption on arms, in pubic and inguinal regions.
Eruption uniform or in large patches of scarlet hue, with interspersed raised spots and some vesicles; rash, followed, after the seventh day from its appearance, by complete desquamation.	Eruption in crescentic patches, with intervening portions of healthy skin; lasts about five days; followed by partial and very incomplete desquamation; scales, as a rule, very fine.	Eruption first papular; remains so about a day; then becomes vesicular, then pustular; on the eighth day of eruption, pustules mature.
Sore throat; rarely coryza or bronchitis.	Coryza and bronchitis very constant; rarely sore throat.	Often sore throat and dry cough; bronchitis only as a complication.
Red "strawberry" tongue.	Tongue coated; may be red at edges.	Tongue coated and swollen; may become red at edges.
Cerebral symptoms frequent and grave.	Cerebral symptoms neither frequent nor grave.	Cerebral symptoms, especially convulsions in children, frequent.
Temperature very high; may range from 105° to 110°; no fall soon after eruption, nor decided increase of heat preceding it; high temperature during height of eruption; subsequently gradual decline. In protracted cases, a fall of temperature takes place on the	Temperature during the fever preceding eruption rarely over from 102° to 103°; falls on second day, rises rapidly toward breaking out of the eruption, and remains high during its appearance and spread; then sinks speedily. The defervescence that takes	Temperature during the fever preceding eruption very high, often 106°; then decided defervescence, taking place within thirty-six hours; subsequently thermometer indicating a temperature of about 100°, notwithstanding the progressing development of

TABLE EXHIBITING THE DIFFERENCES BETWEEN SCARLET FEVER, MEASLES, AND SMALLPOX.—*Continued.*

SCARLET FEVER.	MEASLES.	SMALLPOX.
fifth, tenth, and fifteenth days of the disease.* Irregular cases have irregular, though mostly very high, temperatures.	place, generally within from twenty-four to forty-eight hours, is both rapid and complete. A protracted defervescence indicates a severe case; a high temperature after the rash has faded is due to a complication.	the pimples into pustules. Decided rise of temperature during secondary fever, and then gradual and protracted defervescence; a slight rise during desiccation.
No secondary fever.	No secondary fever; although sometimes a slight increase of fever just before eruption leaves.	Always secondary fever.
Pneumonia rare; pleurisy more frequent.	Pneumonia a very frequent complication.	Pneumonia not a very frequent complication.
Sequelæ: Bright's disease; dropsy; conjunctivitis; deafness; phthisis; chronic diarrhœa; glandular enlargements; epilepsy.	Sequelæ: chronic bronchitis; phthisis; conjunctivitis.	Sequelæ: chronic diarrhœa; glandular enlargements; various diseases of the eyeball and eyelids.


The contagion of smallpox does not always manifest itself by an attack of variola. Sometimes it is modified by happening in a person who is partially protected by vaccination. This *varioid* disease is mild and very rarely fatal; it protects against smallpox. It is distinguished from variola by the pustules passing more quickly through all their stages, and, above all, by an absence of secondary fever. Soon after the eruption—within thirty-six hours—the thermometer shows freedom from fever, and, unless serious complications happen, the temperature remains nearly normal. The suppuration is far less deep; and the resulting cicatrices are often scarcely discernible.

*Varicella*.—A specific disorder similar to but not identical with variola or varioid is chicken-pox, or varicella. It differs, as regards its symptoms, from smallpox in the leniency of the intro-

\* Ringer.

ductory fever; in the eruption beginning generally first on the trunk, occurring often on the second day, though it may not show itself until the end of the third, and continuing to appear and disappear in crops, the mass of the eruption, however, having become evident within twenty-four hours; in the vesicles being surrounded by little or no inflammatory redness; in their remaining vesicles and not becoming pustules; in their attaining their height on the third or fourth day of the eruption, and then bursting and shrivelling without presenting depressions at their apices, and in the crust which falls off about five days subsequently being followed by a smooth, shining, round, and irregular pit. Then the eruption is rarely prominent on the face; and the disease does not protect, as mild forms of smallpox do, from a subsequent attack of variola. Sometimes the vesicles may be found, as are the pustules of smallpox, on the roof of the mouth and at the back of the throat. But, although they may be everywhere very

FIG. 86.



Temperature-record in varioloid ending in recovery; the absence of secondary fever is clearly seen. (After Wunderlich.)

plentiful, the disorder is not a grave one. Still, I have known it in one instance to terminate fatally. Spivak\* has described a case of gangrene of the scrotum that followed varicella.

**Dengue.**—This is an arthritic fever with a cutaneous eruption. It has been prevalent in the form of epidemics chiefly in India, and in the West Indies, as well as in Virginia, South Caro-

\* Medical News, March, 1895.



lina, Texas, and other of the Southern States. We owe some of its best descriptions to Dickson.

It usually begins with pain, stiffness, and swelling of some of the smaller joints, or with severe muscular pains, aching in the back, and stiffness of the muscles of the neck. Fever follows, with suffusion of the eyes, violent headache, hurried breathing, and coated tongue; but, as a rule, without nausea and vomiting. The temperature usually attains its height within the first twenty-four hours, and then shows during defervescence marked remissions and exacerbations. On the third day the fever ceases altogether, or subsides markedly, though the muscular and arthritic pains do not pass off entirely. The febrile paroxysm may last somewhat longer, indeed, five to seven days, or only six to twelve hours. In any case it is apt to be succeeded by an interval of two to four days free from absolute suffering, though not from great debility. Then the pain returns, and with it a moderate fever; nausea and vomiting and a thickly-coated tongue, too, are noticed. This new phase of the complaint is generally relieved by the appearance of an eruption, which may be accompanied by a slight rise in temperature. The eruption shows itself on the fifth, sixth, or seventh day of the malady, and, therefore, very much later than the rash of scarlatina, which it resembles in hue and aspect. But not invariably; for it may occur in patches and be papular, or even vesicular or like urticaria. The eruption is attended with a sense of burning and of itching, and disappears after two or three days' duration, with more or less decided desquamation. It is much more pronounced than the slight and inconstant erythematous rash of the period of invasion, which disappears without desquamation with the febrile stage.

With the occurrence of desquamation following the marked rash of the third period of the disease convalescence sets in, marked by considerable muscular weakness and general depression, and frequently with the rheumatic stiffness or soreness persisting for some time. Swellings of the lymphatic glands of the neck, axilla, and groin occur in many instances, and may continue during convalescence, which in any case is apt to be prolonged, and may be interrupted by a relapse.

The cause of this singular malady—the breakbone fever of

parts of our country—is unknown. McLaughlin \* has invariably found in the blood micrococci in great numbers, about one-twentieth to one-thirtieth the diameter of the red corpuscles, of spherical shape, and red or purplish in color.

Dengue is generally a harmless disorder, epidemic, and contagious. Isolated cases are difficult of diagnosis, but when the disease largely prevails its recognition is easy. It differs from *rheumatism* or *gout* by the significant features of the fever and the eruption; from *scarlet fever* by the different character and want of continuousness of the fever, and the arthritic symptoms; from *influenza* by these and the eruption. The remission may cause the disease to be mistaken for a *malarial fever*; but the irregularity of the fever in dengue, the joint and muscle pains, and the absence of hepatic and splenic enlargement are very unlike. Dengue has a closer resemblance to *yellow fever*, and the difficulty of distinction becomes the greater because epidemics of both may be present side by side. But the single paroxysm, the tongue with red edges, the yellow skin, the frequent vomiting, the hemorrhage, the grave nervous symptoms, and the albuminous urine are not met with in dengue.

**Erysipelas.**—This disease, as the physician sees it, is mostly confined to the head and face. It may or may not be found to have been preceded by a scratch or an abrasion. It is an eruptive fever beginning with a chill. Soon a portion of the face is noticed to be red and hot. The redness spreads, a clearly-defined edge marking its onward march; and generally it does not stop until it has occupied the whole of the face and a considerable portion of the scalp. The features are then so tumefied as to be hardly recognizable. The patient is very restless, has high fever, and not unfrequently enlargement of the glands at the angle of the jaw, and sore throat. By the seventh or eighth day the disease is over, and large patches of cuticle fall from the countenance no longer swollen and disfigured. The temperature remains high for a few days, with marked evening exacerbations and morning remissions, and then falls, not to rise again to any considerable height.


This is simple erysipelas; but the affection may extend—as is,

---

\* Journ. Amer. Med. Assoc., June 19, 1886.

in truth, its tendency—from the true skin to the subcutaneous areolar tissue, and give rise there to collections of pus, which reveal their presence by chills and an obscure sense of fluctuation,

FIG. 87.



Temperature-chart in a case of facial erysipelas, seen soon after outbreak of the disease.

and keep up an irritative fever until they are discharged. Irrespective of this, the tumefaction, while the complaint is at its height, is much greater in this phlegmonous variety of the malady, and there is more constitutional disturbance; but, on the other hand, there is not so much local irritation, for the morbid action travels less rapidly, and often remains more circumscribed. In some cases the inflammation extends to the brain, and instead of wandering at night, always a very common symptom, we have violent delirium, soon succeeded by coma and rapid sinking. In other cases, again, and they are by far the most frequent, we may find these active cerebral symptoms and yet not be able to detect, after death, signs of inflammation of the brain or its membranes. Now and then the disorder passes to the throat, reaches the larynx and bronchial tubes, and places life in imminent peril from œdema of the glottis, or from a hazardous form of capillary bronchitis. In some instances a highly-asthenic state becomes developed, and



the patient dies exhausted. The disease often manifests a distinct tendency to recur. The contagious principle of erysipelas is the *streptococcus erysipelatis*, also called after Fehleisen, who has specially described it.

Internal lesions happen not unfrequently in erysipelas. I have found the urine albuminous in the great majority of instances.\* Heart-murmurs are not unusual, and are said to depend upon endocarditis, which is doubtful. Friedreich† speaks of swelling of the spleen being of common occurrence both in erysipelas and in diphtheria.

The diagnosis of erysipelas is not beset with difficulties. *Erythema* resembles it very closely; but there is this manifest difference: in erythema there is scarcely any swelling, not much tendency to spread, and almost no constitutional disturbance. The ordinary *exanthematous fevers* may, at an early stage, be mistaken for erysipelas. But all of them, even *scarlatina*, have a longer period of febrile invasion; in all, too, although the eruption takes its origin at one spot, and generally on the face, it is not limited there. The thickly-clustered blotches of beginning *confluent smallpox* and the swelling attending them give at times to the face the look of erysipelas. But here, also, evidences can be found of a rash about to appear all over the body; and should doubt still exist, it is soon dispelled by the progress of the eruption. Sometimes vesicles and even irregular pustules form in erysipelas, and occasion some misgivings as to whether the malady be not a chronic disease of the skin, such as eczema, pemphigus, or impetigo; but these affections lack the constitutional symptoms and the history of a recent acute disease, and in reality the likeness is not a very striking one, if the inflamed surface be carefully examined. The closest similarity is to herpes zoster of the forehead and face. But the eruption in this does not pass the middle line.‡ The red color of the skin and the attending symptoms distinguish erysipelas from the swelling of *angio-neurotic œdema*.

Erysipelas may break out in one part of the body after another, and the disease be thus kept up for a long period. This

---

\* On the Internal Complications of Acute Erysipelas, Amer. Journ. Med. Sci., Oct. 1877.

† Klinische Vorträge, No. 75, 1874.

‡ Fagge, Practice of Medicine, vol i. p. 271.



*erysipelas migrans* runs its course more rapidly and more completely in one part than in another, and in accordance with a general law which it obeys.\*

Erysipelas may be confounded with *mumps*. This does not seem at first sight likely; but I have known the error to be committed. It was mainly caused by too much stress being laid on the redness which is frequently found beneath one or both ears in parotitis, but which, unlike erysipelas, is attended with much pain on moving the jaw, and with decided glandular tumefaction. The redness, moreover, shows no tendency to spread, and rarely continues for the four or five days during which mumps lasts. In very young children, however, there may be some difficulty in diagnosis. I have seen the glands at the angle of the jaw much swollen for one or two days prior to the slight discoloration over them taking on a deeper blush, and then spreading rapidly as marked erysipelas over the whole face and part of the scalp, reaching the other jaw, where subsequently the glands began to swell. In such cases great weight must be attached to the history of the case, to determine which disorder was primary, and whether the glandular complaint was or was not the complication.

A fever with a distinct pharyngitis as a local manifestation, the so-called *pharyngeal fever*, is probably an epidemic erysipelatous fever of light type. It has been particularly described by Austin Flint, Rochester,† and Harvey E. Brown.‡ The fever lasts from three to six days, and, besides the marked pharyngitis, is ordinarily attended with swelling of the lymphatic glands of the neck, accompanied by pain. The disease shows a certain proportion of cases with erysipelas of the face, and is thought to be a mild form of the fever, known popularly as the "black tongue," which prevailed in this country from 1841 to 1846, and in about one-sixth of the cases of which erysipelas happened.§

---

\* Traced by Pflüger in 70 cases; quoted in Schmidt's Jahrb., No. 7, 1873.

† Buffalo Medical Journal, 1857.

‡ Flint's Principles and Practice of Medicine.

§ Ibid.

## CHAPTER XIII.

### DISEASES OF THE SKIN.

To facilitate the discrimination of diseases of the skin, they have been grouped into classes. These have been arranged by some in accordance with the obvious characters of the eruption, by others in accordance with its presupposed cause and attending structural alteration. An extensively-used system of classification takes for its basis the anatomical seat and arrangement of the cutaneous malady: it is that of Hebra. As developed by him, it is, however, not a purely anatomical, but a mixed system, resting largely on a pathological basis. Similar is the classification of the American Dermatological Association, now much followed. All diseases of the skin are arranged in eight classes: Disorders of the Glands, sweat and sebaceous; Inflammations; Hemorrhages; Hypertrophies, of pigment, epidermal, and papillary layers, and of connective tissue; Atrophies, of pigment, hair, nail, and cutis; New Growths, of connective tissue, vessels, and granulation-tissue; Neuroses; and Parasitic Affections, vegetable and animal. Whatever classification we adopt, when a disease of the skin is presented for examination we generally first endeavor to ascertain the group it belongs to; for instance, is it macular, papular, vesicular, or pustular, or does it present lesions representing more than one group? Having determined this, we next fix which member of the group it is, and then regard its precise seat and its pathological causation. When this has been accomplished, we inquire into the history of the affection and its duration, whether acute or chronic; take into account the presence or absence of fever, and the general condition of the patient; search for the evidences of a cachexia or of some visceral disturbance,—a study the importance of which is as great as that of the recognition of the cutaneous malady; and trace, so far as possible, the cause of the disorder. In many instances microscopical and bacteriologi-

cal examination will be necessary to supplement the clinical evidence and complete the diagnosis. Having done all this, we have a groundwork upon which to institute suitable treatment.

Here is a table in which cutaneous affections, omitting some of the less important ones, are grouped according to their most obvious features, as well as according to their pathological bearings :

## DISEASES OF THE SKIN.

INFLAMMATORY.	{	ERYTHEMATOUS DISEASES...	{	Initial rashes of eruptive fevers.		
				Erythema.		
				Roseola.		
				Urticaria.		
	{	PAPULAR DISEASES.....	{	Papular eczema.		
				Lichen.		
				Prurigo.		
	{	VESICULAR DISEASES.....	{	Eczema.		
				Herpes.		
				Dermatitis herpetiformis.		
	{	BULLOUS DISEASES.....	{	Pemphigus.		
				Hydroa.		
	{	PUSTULAR DISEASES.....	{	Acne.		
				Boils, or Furuncle.		
				Sycosis non-parasitica.		
				Impetigo.		
				Ecthyma.		
				Rupia.		
Glanders.						
{	SQUAMOUS DISEASES.....	{	Psoriasis.			
			Pityriasis.			
			Ichthyosis.			
			Squamous eczema.			
MACULÆ; PIGMENTARY CHANGES.....	{	{	Melasma.			
			Ephelides.			
			Vitiligo.			
			Chloasmata.			
			Nævi.			
			Purpura simplex.			
			Xanthoma, or Xanthe-			
			lasma.			
			{	HYPERTROPHIES OF SPECIAL TEXTURES.....	{	Elephantiasis Arabum.
						Scleroderma.
Keloid.						
Dermatolysis.						
{		{	Warts, Corns, etc.			

DISEASES OF THE SKIN.—*Continued.*

ATROPHIES.....		{ As of the Hair; the Nails. { Senile Atrophy.
		{ Scabies. { Phtheiriasis. { Favus. { Anthrax. { Tuberculosis. { Molluscum epitheliale. { Lepra.
PARASITIC DISEASES.....		{ Mycetoma. { Actinomycosis. { Tinea sycosis, or Men- tagra. { Tinea circinata. { Tinea tonsurans. { Tinea decalvans. { Tinea versicolor, etc.
NEW GROWTHS.....		{ Cancer. { Sarcoma. { Molluscum fibrosum. { Lupus. { Leprosy, etc.
	of Sebaceous Glands.....	{ Seborrhœa. { Comedo. { Sebaceous cyst.
ALTERED GLAND-SECRETION..	of Sweat- Glands.....	{ Hyperidrosis. { Anidrosis. { Chromidrosis. { Bromidrosis. { Miliaria, etc.
		{ Hyperæsthesia. { Anæsthesia. { Pruritus. { Neuroma.
NERVOUS AFFECTIONS.....		{ Dermatitis herpetiformis. { Herpes Zoster. { Peliosis rheumatica. { Plica Polonica. { Alopecia areata.
CONSTITUTIONAL SKIN AFFECTIONS.....		{ Syphilodermata. { Scrofulodermata, etc.

Most diseases of the skin are again subdivided into several varieties, based, for the most part, on their duration, situation, form, feel, and color. Thus, we have constantly recurring the



terms *fugax*, *inveterata*, *capitis*, *facialis*, *palmaris*; *guttata*, when like a drop on the skin; *nummularis*, when like a coin; *larvalis*, like a mask; the qualifying words *læve*, *induratum*; *circinatum*, *annulatum*, *marginatum*, indicating configuration, and the adjectives of color, *nigrum*, *rubrum*, *versicolor*. But these divisions are all of secondary importance; and in this outline not much regard will be paid to them. Premising this statement, let us briefly examine the characteristics of the various cutaneous affections of more common form, beginning with those of inflammatory origin.

**Erythematous Diseases.**—There are only three affections which, strictly speaking, come under this division of cutaneous complaints: *erythema*, *roseola*, and *urticaria*. In all of these the skin is more or less red, and its surface unbroken; the hyperæmia affects chiefly the papillary layer.

*Erythema.*—This is characterized by a uniform and continuous redness of the skin, occurring in irregular patches of some size, attended with some burning, and with but slight swelling, if with any, and disappearing without desquamation or mark or scar. The eruption is chiefly found on the back of the hands, the forearms, the legs, and the face and neck; rarely on the trunk. There is little or no itching. The affection may be due to the action of heat or cold, or of irritants; or it may be connected with some visceral abdominal disorder. It is usually acute. There is only one variety apt to be combined with decided constitutional or febrile symptoms,—the hard, painful, reddish protuberances most commonly seen on the legs, and constituting the so-called “*erythema nodosum*.” This form of the complaint, in which there is a serous effusion into or under the skin, is chiefly observed in those of rheumatic diathesis, and, unlike the simple erythema and the erythema intertrigo, which are looked upon as mere hyperæmias, is classed with the exudations or inflammations. All the exudative forms of erythema may be grouped under the title of *erythema multiforme*, the varieties of which are the papular, bullous, and nodose, the lesions appearing principally on the backs of the hands and feet.

There is a *desquamative form of erythema* resembling scarlet fever, attended with fever of a few days' duration, with epistaxis, and showing an extraordinary tendency to relapse. The erup-

tion is uniform and intensely red, and there is no sore throat, or there is mere redness of the fauces. *Erythema solare*, or superficial dermatitis following exposure to the sun's rays, is usually followed by free desquamation.

A chronic form of erythema results from pressure, or the rubbing together of folds of skin, the *erythema intertrigo*; a slight discharge may coat the rubbed surface. It is liable to acute exacerbations.

*Roseola*.—This term is applied to circumscribed spots of a rose-red color and of a more or less circular form. The spots are smaller than those of erythema simplex. In *erythema congestivum*, or roseola, there is slight fever, and at times redness of the fauces. The affection often exists in connection with a derangement of the stomach, or with rheumatism, is frequent in summer and in autumn, is generally acute, and bears a certain resemblance to scarlatina and to measles; but it is not contagious, its constitutional symptoms are much milder, the rash is rosy, not crescentic, nor present over the whole body, and we find neither the marked sore throat of scarlet fever nor the catarrh of measles. A roserash occurs in the course of typhoid fever, and there is also a syphilitic form.

*Urticaria*.—Nettle-rash gives rise to prominent and perfectly smooth patches, the color of which is either redder or whiter than the surrounding skin; the white wheals may be surrounded by a red border. The wheals are generally small, but they may be of the size of the palm. The eruption is fugitive and capricious, is attended with more itching, burning, and tingling than the other exanthemata, and is much more evanescent, generally disappearing in two days at furthest. It may, however, exist in a chronic form, the wheals coming out in constant succession, especially after scratching or other irritation of the surface. Pigmentation occurs in the variety known as *urticaria pigmentosa*.

The cause of urticaria is irritation of the gastro-intestinal, pulmonary, or urinary mucous membrane. Certain kinds of fish, especially shell-fish, are particularly prone to produce it; so do mushrooms and strawberries. At times it is due to menstrual disorders, or to sudden mental emotion, or to the excessive use of mineral waters, or to antipyrin. It may be secondary to the

itch, or to phtheiriæsis. It occurs in cerebro spinal fever, and is common in dengue, especially in children.\*

Urticaria is thought generally to be an exudative disease of the skin; yet it seems most probable that it is a reflex phenomenon, caused chiefly by reflected irritation to the cutaneous vaso-motor nerves. Urticaria resembles erythema nodosum; but there is no itching in the latter affection, which is chiefly found in the lower limbs, and the swellings change like bruises.

**Papular Diseases.**—A papule, or pimple, is a small elevation of the cuticle with an inflamed base; it does not contain fluid, and usually terminates in desquamation. It results from a small amount of lymph or a newly-formed growth in the derm itself.

*Lichen.*—This furnishes the best-marked example of a papular eruption. It consists of minute conical papulæ, generally of reddish color, and occurring in clusters. It is most frequently encountered in the summer months and in adults, and often in persons who are in good health but who have been exposed to much fatigue or anxiety. Sometimes it is evidently connected with disordered digestion. It is commonly chronic. There is often a mixture of papulæ with an eczematous eruption. Prickly heat, or lichen tropicus, frequently exhibits also sudamina, and is called by some “miliaria papulosa.”

In the *lichen ruber* of Hebra the red papules are of the size of the head of a pin; they spread by peripheral growth, are flat, irregular, and have a glazed look and very slight scales; there is considerable itching. The disease, which is an inflammatory one, is chronic; its common site is on the forearm. It resembles psoriasis, but at the edge of the patch are the characteristic papules. Poor nutrition and nervous exhaustion are its main causes.

In the *lichen scrofulosorum* the eruption consists of little pale papules, which are chiefly found on the trunk. There is no itching; but we find marked signs of scrofula.

The lesions of *lichen planus* are small, hard, red papules, that may be umbilicated and coalesce into patches. In the latter case scaling occurs, and more or less itching is a frequent accompaniment. The smallness of the recent lesions, which are

---

\* J. C. Wilson, Treatise on the Continued Fevers, 1881.

at first of the same color as the surrounding skin, the flat glazed tops of the older papules, the pure white color of the silvery scales, which are not heaped up, and the unsymmetrical character and distribution of the patches, none of which are circular, will serve to distinguish this from *psoriasis*. *Papular eczema* has its lesions in groups upon an inflamed base, and vesiculation and desquamation are apt to occur.

*Prurigo*.—This is characterized by a papular affection of the skin attended with excessive itching. It is a very rare disease in this country.\* The pimples are generally torn by the fingernails, and are surmounted by black scabs. They are not red, as those of lichen usually are, and are, as a rule, larger, and accompanied by much more pruritus and by thickening of the skin. The affection, which is uncommon, may or may not be attended with constitutional symptoms. It is very obstinate, especially when happening in old persons. It generally affects the legs, the arms, and the trunk, rarely the face and the neck, never the palms and the soles. The skin of the anterior and outer part of the leg is most changed; that over the flexors in the forearm is always healthy. The distressing disorder may be purely local, occurring around the anus, or on the scrotum and the root of the penis, or on the pudenda. Some of these cases, however, though called prurigo, present no papulæ, and the disorder is due to perverted sensibility of the cutaneous nerves alone, and is really a pruritus. Prurigo is often attended with eczema, and may follow urticaria, especially of the chronic variety.

A good many supposed instances of the malady are not really prurigo, but phtheiriasis, due to the irritation of body-lice, that produce papules, whose apices are scratched off and show little points of dried blood. True prurigo is frequently found to be connected with deterioration of the health, and is chiefly met with among the poor and the neglected. It may last a lifetime, beginning in childhood. Its local forms are associated with irritation of the bladder, the rectum, or the uterus.

Papules and tubercles, or large papules, occur in the later stages of syphilis; they are often preceded by the pigmented erythematous syphiloderm. Gumma is a tertiary manifestation,

---

\* Only 34 cases in 123,746 of skin-disease: Van Harlingen on Skin-Diseases.



mostly appearing in the subcutaneous or submucous connective tissue without inflammation, irritation, or itching, gradually involving the tissues above and below, the lesions ultimately attaining a considerable size. At first the color of the skin is not changed, but ultimately it becomes deeply congested and glazed, and as the contents of the lesion soften, the overlying skin is thinned, and finally it breaks down, discharging the purulent material.

**Vesicular Diseases.**—These are characterized by an effusion of a clear or a sero-purulent fluid beneath the epidermis, which is generally raised in small elevations. To the class of vesicular diseases belong especially eczema and herpes.

*Eczema.*—The malady consists of minute vesicles collected together in irregular patches. The vesicles are often confluent, and it then appears as if the whole surface were secreting fluid. This may harden, from exposure to the air, in scabs of various thickness and color. The skin itself is often of a vividly red hue; indeed, it is inflamed, and a new cell-growth takes place both in the rete mucosum and in the papillary layer of the derm. It is there that the effusion of serum begins. In chronic cases the inflammatory infiltration extends deeper into the skin.

Eczema is the most common of all the cutaneous maladies; but it is not contagious. It may affect the whole body, yet is ordinarily limited to some portion of it. It is acute or chronic. The former is generally seen as the effect of local irritants, and may be met with in young and healthy persons. Chronic eczema is more usual, is often the consequence of constitutional disturbance, and is frequently found to be associated with some disorder of the digestive system. It has as a frequent seat the flexor surfaces of the limbs. Dentition and unhealthy milk are common sources of the affection in very young children. In them the disease is extremely apt to attack the scalp and face, forming the complaint often described as “crusta lactea;” or, if the secretion be partly purulent, or early become so, and dry into large, dark scabs, the malady is designated as *eczema impetiginodes*. This is most often met with in scrofulous subjects. There is less heat and itching than in other forms of eczema.

In some of the forms of eczema, especially in its chronic varieties, the vesicles supposed to characterize the disorder can often

not be found. This and other reasons have caused several dermatologists, especially Hebra and Anderson, to deny that eczema need be vesicular at all. Infiltration of the skin, exudation on its surface, the formation of crusts, and itching, are held to be its distinctive signs while the eruption is at its height; but the eruption may consist of clusters of papules, vesicles, or pustules, or there may not be a vestige of any of these, the skin being thickened, red, and smooth and secreting a sticky discharge, or covered with green or gummy crusts, or fissured with deep cracks; yet there are no ulcerations. Not unfrequently the disorder begins as an erythema. A scaly form of eczema, *eczema squamosum*, is apt to be confined to the hands and feet. In all the forms of eczema, as Hebra insists upon, there is severe itching. This itching is especially violent in the form with the deep-red and weeping surface, named *eczema rubrum*. It is in this variety that we find the signs of local inflammation very marked, and we often see it in gouty or in dyspeptic subjects. It has a predilection for the flexures of the joints.

Eczema, especially when it affects the scalp and face, must not be confounded with the morbid secretion from the sebaceous follicles that gives rise to soft crusts. *Seborrhœa* by preference attacks the parts mentioned; but its crusts, as Hardy has shown, are unlike those of eczema in the readiness with which they are detached, and are susceptible of being moulded between the fingers. The surface beneath the crusts, too, is dissimilar. It has an oily, glistening look; there is no discharge. Unna\* has distinguished a seborrhœic form of eczema, which, beginning usually on the scalp, spreads to other portions of the cutaneous surface; but he attributes the source of the fatty scales and crusts to disorder of the sudoriparous, rather than to the sebaceous glands. Patches of seborrhœic eczema are also found on the sternal region, which, after the scalp, is the most frequent locality to be affected. The scales are yellowish in color, the patches spreading by small papules at the border, leaving the centre less scaly and even smooth, while the margin is a red, scale-covered, bow-formed wall.

Eczema may be confounded with *pityriasis rubra*. But this very rare disease speedily involves the whole surface of the body,

---

\* Journal of Cutaneous and Genito-Urinary Diseases, 1887.

is very chronic, and is not accompanied by discharge ; and there are large, thin epidermic scales.

*Herpes*.—Like eczema, herpes is classed as a vesicular affection, although it differs from the obviously vesicular form of eczema by the larger size of the vesicles. These are generally of a globular form, and are symmetrically arranged in clusters upon an inflamed patch of skin. Each vesicle is distinct, and remains so throughout its course. It lasts about eight to twelve days, and often terminates by the formation of a thin incrustation. The eruption is attended with burning, and in the acute variety with some fever.

Herpes has seldom a longer duration than three weeks ; though it may be a chronic disease. It happens usually in persons of delicate skin ; is generally very local, having its seat on the lips, eyelids, prepuce, or pudenda ; and is very often associated with an internal disease, especially with irritation of some portion of the gastro-pulmonary mucous membrane. *Herpes labialis* mostly appears at the decline or termination of fevers ; sometimes at the height of acute maladies, as in pneumonia. *Herpes gestationis*, or the herpes of pregnancy, is identical with dermatitis herpetiformis of Duhring. The most distressing form of herpes is that usually extending around one-half of the trunk,—*herpes zoster*, an acute disorder, which may show itself over the course of any of the superficial nerves, and is attended by nerve-pain. Indeed, herpetic or bullous eruptions often happen over the course of the nerves, and any nerve-lesion the result of disease or of an injury will produce them ; the vesicles are seated upon a highly-inflamed base. In herpes zoster around the chest, the severe pain preceding the eruption is often mistaken for pleurisy, but inspection of the surface and palpation will reveal local spots of tenderness along the course of the affected intercostal nerve.

Herpes and eczema may both be confounded with *scabies*, which, like them, occasions a vesicular eruption that is apt to be found on the inner surface of the limbs and flexures of the joints. The distinction consists in the more severe itching, especially at night ; in the small conical vesicles, torn, as they usually are, by scratching ; and in the presence of the acarus, which may be removed from its burrow with the point of a needle or of any sharp instrument.

**Bullous Diseases.**—Bullæ differ from vesicles only in their size. The typical bullous disease is *pemphigus*. This affection is not often met with ; it is more common in children than in adults. It appears in very large vesicles or bullæ surrounded by a slight zone of erythematous redness. The blebs occur in crops, and look like small blisters filled with serum. They are not met with on the scalp. Where there are few bullæ, we generally find them on the ankle or on the hand. The disorder may be acute or chronic. It is ordinarily chronic, and happens in persons of enfeebled constitution. Relapses are frequent, and a fatal result is common. Pemphigus may be produced by the administration of iodide of potassium,\* or by syphilis. *Syphilitic pemphigus* is mainly met with on the soles of the feet and the palms of the hands of newly-born syphilitic children. There is a form of extensive pemphigus with flaky incrustations like eczema,—*pemphigus foliaceus* ; but we can still find bullæ, and there is great attending prostration. Neurotic vesicular erythema occurs after injury to a nerve, and sometimes causes blebs which may be mistaken for *herpes* or *pemphigus*, as in a case reported by Shields,† in which recurrent attacks of vesicular (or bullous) erythema were observed in the forearm, following the crush of a finger. The symptoms were entirely obviated by amputation of the stump of the finger, after the affection had existed for a period of three years.

*Hydroa*.—This is a disease like herpes, only occurring in a more diffused manner and presenting larger vesicles, arranged for the most part in the form of crescentic rings. It is a chronic condition, lasting usually from five to eight months, and there are in this period many acute or subacute outbreaks, in which the large vesicles form and then dry away. These attacks are non-febrile, and are attended with marked itching. The base of the vesicle is red, and it forms out of a red papule. The disorder happens chiefly in persons of depressed nervous system or gouty taint. It has been confounded with the eruption of bullæ from iodide of potassium ; but these are much larger, are more persistent, and leave a marked scar. Van Harlingen considers cases of hydroa to be examples either of erythema iris or of dermatitis herpetiformis.

---

\* Bumstead, Amer. Journ. Med. Sci., July, 1872.

† The Cincinnati Lancet-Clinic, May 25, 1895.



**Pustular Diseases.**—These are marked by circumscribed elevations of the cuticle which contain pus. Acne, impetigo, and cethyma belong to this group. Rupia, too, although often classed among the bullous disorders, appertains more strictly to the pustular or to the syphilides.

*Acne.*—This is an eruption of hard, isolated, red elevations, due to chronic inflammation of the sebaceous follicles and the areolar tissue around them; plugs of sebum are retained in the ducts. At the apices of many of these elevations pus forms, which is discharged, leaving a hardened base, that only gradually disappears. Acne is generally seen on the face and shoulders. Men of sedentary occupations and drunkards are very liable to it. In women it is frequently associated with uterine disturbances; in men, with some digestive or genito-urinary disorder. An acne eruption also follows the use of the bromides and the iodides internally, and the local use of tar. In *acne rosacea*, lymph is generally effused into the papillary layer of the skin, and some acne pustules are seen, surrounded by the reddened, altered skin. It is a disease of years' duration, but no ulcerations happen, although scarring is a not infrequent result from the small abscesses.

*Impetigo.*—This is a malady often happening in persons of good general health, and mostly soon ending in recovery. It presents small pustules occurring in successive crops and arranged in clusters. The pustules are isolated, are little raised above the surface, break, and a thick yellowish or greenish crust is developed; no scar follows. When the disorder attacks the scalp and face, especially in infants and children, it gives rise to very extensive incrustations, and constitutes, particularly if conjoined with eczema, the affection formerly designated as "porrigo larvalis." There is a contagious form of it, described by Tilbury Fox, which occurs acutely, is epidemic, preceded by fever, and unattended with pain or itching. Another form of impetigo, first mentioned by Hebra, consists in a multiform eruption of vesicles, vesico-pustules, and pustules.

*Impetigo contagiosa* is characterized by vesico-pustules or blebs drying into flat, straw-colored crusts. It is contagious, and is especially encountered in children. The lesions occur especially on the face and hands, and vary in size from a pin-head to a quarter of a dollar; their contents, at first serous, become sero-

purulent in the process of drying. *Dermatitis herpetiformis*, or *Duhring's disease*, differs in being not contagious, and in its happening in older persons who are frequently of an hysterical type. Leredde and Parin, at the Hôpital St.-Louis, found in the skin, at the site of the lesions, numbers of eosinophile granules and cells. These cells, unusually rare in the skin, play a considerable part in this affection, and by their extraordinary frequency serve to distinguish this from other cutaneous diseases. Suppuration did not occur, and no micro-organism was found in the vesicles. There is a close connection between this disease and herpes gestationis, in which the same cellular elements have been found.\* There is deficiency of urea in the urine of Duhring's impetigo, and M. Bar found that the toxicity of the urine was increased at the time that the eruption occurred. Stephen Mackenzie † considers Duhring's disease a cutaneous neurosis, and Leredde suggests that the exciting cause may be deficient elimination by the kidneys, as nephritis has been found by Gaston ‡ in two autopsies. The eruption upon the skin is due to nervous influence owing to an alteration of the blood and a functional disorder of the kidneys. §

*Ecthyma*.—This differs from impetigo by the larger size and greater prominence of the pustules and their inflamed base. When the crust that forms on each pustule falls, a highly-congested surface or a superficial ulceration is seen, which leaves a cicatrix. This disorder is painful, generally chronic, and connected with a cachectic state of the system; irritation of the skin may excite it. It bears a certain resemblance to sycosis; but the limitation to the hairy portions of the face, the yellow color of the pustules, their conical form and smaller size, and the brown crusts they occasion, distinguish this malady.

*Rupia*.—This affection produces at first bullæ, but soon very large pustules, which desiccate into thick, brownish crusts, often of conical shape or resembling the shell of an oyster, which, when thrown off, expose ulcerations of various depth that are

---

\* Anatomie pathologique de la Dermatose de Duhring : Annales de Dermatologie et de Syphillographie, No. 4, April, 1895.

† British Journal of Dermatology, Jan. 1893.

‡ Annales de Derm. et de Syph., Paris, April, 1895.

§ Loc. cit.

slow to heal, and on which fresh crusts arise. The disease runs a chronic course. It occurs especially on the lower extremities, is due to syphilis, and coexists with a deteriorated constitution. It is very like ecthyma, and can be distinguished only by the history of the case, the evidences of syphilitic taint, the persistent ulcerations, and the prominent, peculiarly-shaped crusts.

**Squamous Diseases.**—The predominant characteristic of these is the formation of small, whitish patches of unhealthy cuticle covering red papular elevations or a deep-red, dry, somewhat thickened surface; the scales are generally very freely cast off. Psoriasis is the main disorder belonging to the group. *Pityriasis* is included by many, while others regard it as merely a variety of chronic erythema, or of eczema. It differs from lepra and psoriasis by the production of minute scales, which are constantly thrown off and reformed, and which are seated on a reddened integument: hence its chief variety is designated *pityriasis rubra*. It begins at a special point, and, unlike psoriasis, spreads over the whole body. The skin is very red, and not thickened except in instances of long standing; there is no discharge, as in eczema, nor itching or burning; the scales are loosely adherent to the surface, and at times come off in large flakes. The disease is most apparent on the body and the limbs; in chronic cases the general health deteriorates, and a fatal result is the rule. *Pityriasis rubra* is to be distinguished from exfoliative dermatitis, which is an acute affection and more amenable to treatment. In this disease the scales are thicker, larger, and more abundant than in *pityriasis rubra*; there may be some spots of moist eczema, the lesions being papular at first and then vesicular, ending in profuse exfoliation, large casts coming away from the fingers and toes. Alopecia and shedding of the nails are common. Savill\* reports an epidemic of this disorder in an infirmary affecting twenty per cent. of all the patients.

*Pityriasis rosea*, or *pityriasis maculata et circinata* of Duhring, is recognized by the presence of maculæ, or very slightly elevated patches, varying from a pin-point to a half-dollar in size, the color being rosy, or pink, with a yellowish tint. The surface of the lesions is dry and slightly scaly; the appearance is circinate.

---

\* British Medical Journal, Jan. 2, 1893.

The eruption usually appears on the trunk, is moderately acute, and may last two or three months or longer. It is not contagious, and apparently is not parasitic in origin, though this is a matter of doubt. The general health is not impaired, and the patches give no annoyance except by the itching, which is not excessive, and by their appearance, which may lead to their being taken for lesions of syphilis, or for ringworm, lichen ruber, psoriasis, or one of the eruptive fevers, as in cases reported by Duhring and Stelwagon.

*Psoriasis*.—Here we find patches of a red hue raised above the surrounding integument and covered by scales of dried epidermis. The patches are infiltrated and thickened, and they often have a circular or circumscribed shape, with large pearly white scales. But generally the scales which completely cover the morbid portion of skin are small, though thick; the patches are large or consist of small ones which have coalesced into a single large one, are not of an annular form, or completely separated by healthy skin; they are very symmetrical. Psoriasis generally first appears on the extensor surfaces of the elbow- and knee-joints, and finally on the face, where the scales are usually very small. As Beverley Robinson has proved, the morbid change begins in the cells of the epidermis. There is no watery discharge, and scarcely any itching, attending the affection.

Psoriasis is often hereditary; in old persons it is frequently of gouty origin. It is a chronic affection, and extremely obstinate. It is liable to be mistaken for *lichen*, especially the isolated circular form of it, the so-called lepra. It is, however, distinguished by the distinct, dry, and silvery scales, and by the smooth, red, perhaps bleeding skin which is at once perceived when the scales are detached. Psoriasis has a predilection for the vicinity of the joints, especially the elbow- and knee-joints. Sometimes it appears exclusively on the palm of the hand; and in this form especially we are apt to find deep cracks. Palmar psoriasis is rare; but a condition resembling it in the production of scales and fissures occurs in constitutional syphilis, the so-called *syphilitic psoriasis*, which is common and is usually attended by other symptoms of syphilis. Psoriasis differs from *eczema squamosum* by the preceding vesicles, severe itching, and the want of uniformity of lesion of the latter. In scaly syphilitic eruption the scales are



comparatively few and fine; when they are removed, the dense skin underneath does not bleed; and the eruption is not likely to be met with on the elbows and the knees.

*Ichthyosis*.—Fish-skin is also a squamous disease; but it differs from the others of this class in being much more general, involving as it does often the whole integument, and in the absence of reddening or any signs of inflammation of the harsh, dry surface; it is, indeed, an hypertrophy of the cuticle. The skin is dry, dirty, and rough, and covered with thickened and exfoliating cuticle and with sebum; there may also be fissures and cracks. *Ichthyosis* is almost always of congenital origin and begins in childhood; it affects the whole body, though the face but very slightly.

Among the inflammatory diseases of the skin, those resulting from medicines may be here mentioned. This *dermatitis medicamentosa* is brought about by a variety of drugs, and differs according to the special drug. Among the principal ones producing morbid appearances of the skin are arsenic, quinine, belladonna, opium, chloral, salicylic acid, antipyrin, arnica, copaiba, the bromides, and the iodides. The acneiform eruption due to the bromides, with the dusky-red color of parts of the skin, or the ulcers they may occasion; the papular or bullous eruption caused by the iodides, especially by iodide of potassium, and the scarlet rash of belladonna,—are well known.

**Maculæ**.—These include stains on the skin which are due to chemical substances, such as nitrate of silver, or blood-spots, as in purpura, or spots in consequence of parasitic formations, as in *tinea versicolor*. But their chief cause is increased pigmentation; and it is this cause that we shall look at more particularly.

First, *lentigo* may be mentioned. This consists of the little yellow or yellowish-brown spots which are so often met with on the face and on the arms in children under eight years of age, and which, if they have persisted, disappear in middle life. Similar spots are *ephelides*, or freckles; these, though aggravated by exposure to the sun, may exist all the year round. *Melasma* is a very dark pigmentation, which, although it has been met with in an epidemic form, is commonly seen in connection with Addison's disease.

*Chloasma* consists of a brownish or yellowish-brown pigmen-

tation, giving rise to the so called liver spots. They are smooth and well-defined maculæ without scales, and may result from any local irritation or from exposure to the sun or heat. They may also happen in cases of faulty digestion with torpor of the liver, in uterine disorders, and in the pregnant state. *Tinea versicolor* is constantly confounded with these so-called liver spots. But it is almost entirely a disease of the trunk, is much more itchy, is slightly raised, and in the scales we scrape off is found the characteristic fungus.

**New Growths.**—These are hard, indolent, and often permanent tumors of the skin, which in their main forms consist of granulation tissue. Lupus, fibroma molluscum, and elephantiasis of the Greeks mainly illustrate this group.

*Lupus.*—In lupus the new growth mostly takes place in the form of isolated tubercles. These may or may not ulcerate. They are of a dull-red color, elevated above the surface, with a well-defined outline, spread outward into normal textures, and, if they ulcerate, destroy the tissues in which they are situated. The ulcers also spread, and occasion much devastation. When they heal, they leave a strongly-marked whitish cicatrix and unhealthy-looking skin. The disorder occurs in syphilitic or in scrofulous persons,—generally in the latter,—appears often in childhood, is attended with some pain and itching, and pursues a very slow course. The nose and cheek are the favorite sites. There is a form of lupus occurring only in strumous subjects, and characterized by warty formations. This *lupus verrucosus* is without pain or itching, but cicatrices form, though there has been no previous ulceration.\* In *lupus erythematodes* the disease is superficial, and the sebaceous glands particularly are distended. The surface is somewhat raised, the centre of the diseased patch is pale and sinks in. The tubercles form late, if at all, and there is no ulceration. The most common site of the disease is under the eye. It does not generally appear until after puberty, and is preceded by erythema of the affected parts. The diagnosis of *lupus vulgaris* depends principally upon the small nodules deeply embedded in the corium of the skin, that have a tendency to undergo ulceration, leaving upon healing a peculiar cicatricial tissue

---

\* McCall Anderson, *Journal of Cutaneous Medicine*, vol. i.

of uneven thickness. Tubercle-bacilli and giant cells have been found in these lesions, but Virchow has shown that the giant cells may be found in normal granulations and elsewhere. Baum \* has determined the bacilli to be comparatively rare ; they are only demonstrable in those nodules which have undergone retrograde metamorphosis.

*Lepra*.—Leprosy is a chronic constitutional disorder, and the symptoms of general depression may precede the characteristic local features. The true leprosy, the elephantiasis of the Greeks, is distinguished by tubercles, from the size of a pea to that of a walnut, of reddish or whitish or bronze-like hue, which slowly ulcerate, and which are preceded by erythematous patches ; ulceration is apt to take place about the fingers and toes. Like lupus, the tubercles have the structure of granulation tissue. Often, too, there are symptoms of defective innervation, especially deficient sensation of the surface, anæsthesia of the fingers being an early symptom. The nerve-trunks are invaded, cutaneous eruptions in their course result, and the blood is seriously affected. Muscular weakness and wasting may be also present. The face is most frequently the seat of the malady, and becomes very much thickened and disfigured ; similar changes may also be seen in the limbs. Pemphigus like blebs are among the earliest signs. When marked nodules form, the skin is decidedly discolored, often copper-colored, and the face is distorted and has a fierce expression. Sometimes anæsthesia is the main symptom ; the uneven thickening may occur in circular patches like psoriasis, but without tubercles, and be markedly anæsthetic. The disease is often hereditary.

Two forms of the disease are generally recognized,—the tubercular and the anæsthetic ; but there is no absolute distinction between them. The disease is common in tropical regions, especially the East, and in Africa and Brazil ; it is also found in Norway and in the Hawaiian Islands, and has become naturalized in the United States.†

**Hypertrophies.**—There are many forms of these, according

---

\* International Clinic, Fourth Series, vol. ii. p. 349.

† See Transact. Amer. Dermatol. Assoc., 1879, and Henry Dickson Bruns, Archives of Medicine, New York, Dec. 1881.

to whether the connective tissue, the epidermis, the arteries and veins, or the lymphatic vessels are affected. I shall notice particularly two; and first, elephantiasis Arabum.

*Elephantiasis of the Arabs.*—This, the Barbadoes leg, is an enormous increase in size of the limb, usually dependent upon an indurated swelling of the subcutaneous tissues, with some alteration of the skin proper, and lymphangitis. The tumefaction may be in swellings separated by deep furrows, giving somewhat of a tuberculated look to the part, or it may be uniform; it chiefly attacks males, and gives rise to great deformities. It is a disease of the tropics. The cases, especially of elephantiasis of the scrotum, have been frequently traced to filariæ.

There is a form of enlargement of the leg to which we may here briefly refer,—one in which the overgrowth of the affected limb is associated with *disease in the lymphatic system*. Vesicles form, which are connected by ridge-like elevations, and which from time to time discharge a chylous fluid.\* The subcutaneous lymphatics near the groin are usually found to be distended.

*Scleroderma.*—Scleroderma, or sclerema, is an induration of the skin and areolar texture, which may be partial or general, affecting nearly the whole body.† The skin is dense and hard, and in the true skin and the subcutaneous tissue the fibrous elements are much increased. The true skin shrinks and binds down and is bound to the parts beneath. If the malady seize upon the fingers, it renders them rigid and immovable. The disease is generally symmetrical, and is much more common in women than in men. In cases reported by Goldzieher,‡ the disease appeared in the extremities after unusual exposure to cold, and resulting frost-bite. It frequently coexists with feeble health; and in time the internal organs become affected, or these may be from the first deeply implicated.§ The general health may, however, remain good.

\* W. H. Day, Transact. Clin. Soc. Lond., vol. ii., 1869.

† See the cases collected by Van Harlingen, Amer. Journ. of Syphilography and Dermatology, 1873.

‡ Beiträge der Berlin. Dermatolog. Gessellsch., March, 1893.

§ Harley, Med.-Chir. Transact., 1877.



I had some years since a marked case of this strange affection under my charge at the Pennsylvania Hospital, in a woman, forty-two years of age, who, admitted with œdema of the feet, was at the same time noticed to have a swelling of both wrists and fore-arms as well as of the cheeks. The swelling was firm and resistant, and did not pit on pressure. The skin covering it was very smooth, and of redder hue than at other portions of the body; there was well-preserved sensibility. The œdema disappeared from the feet, but the signs of the indurated cellular tissue did not leave the affected parts. On the contrary, the condition of these parts became worse, though the general health was excellent, all the internal viscera being in a normal state. Gradually the hands, particularly the fingers, were found to be more and more resisting and immovable, and she could scarcely bend them; occasionally they were the seat of pain. The skin lost all suppleness, and could not be raised up. At no time while under observation was albumen present in the urine. She left the hospital unimproved by the sulphur baths, the bichloride of mercury, and the various other alteratives she took; and I afterward learned that she died of an acute pleurisy succeeding an attack of acute meningitis. Prior to her death, so great was the pressure exerted by the dense and contracting cellular tissue that dry gangrene of a finger ensued, as well as of a toe, the disease having been also noticed in the lower extremities. She died about one year from the beginning of the complaint. Examined after death, the skin over the diseased parts was found to be firmly united to the muscles beneath by the dense and augmented areolar textures.

Scleroderma is very similar in many of its features to *myxœdema*. But the marked anæmia of this, the decided nervous symptoms, and the fact that we do not find the stiff, hard skin compressing the parts beneath and bound to them causing in time marked atrophies, distinguish the two maladies. The recent successful treatment of *myxœdema* by thyroid extract suggests a means of diagnosis between the two affections. Goldzieher\* considers scleroderma a progressive chronic dermatitis accompanied by permanent œdema. Repeated attacks of *erysipelas*

---

\* Beiträge der Berlin. Dermatolog. Gesellsch., March 12, 1893.

thicken the skin, but we do not find atrophies from compression.

Scleroderma is closely related to *morphœa*. This occurs over the course of nerve tracts, the thickening being in circumscribed patches and lacking the peculiar hardness of sclerema; then changes in the structure of the skin, hyperæmic appearances at first, pigmentation and cicatrization afterward, occur in *morphœa*, with pain and tingling in the affected parts. The color of the patch of *morphœa* is characteristic. The central part is usually of a yellowish-white or ivory color, which is surrounded by a zone of lilac, due to enlarged capillaries. By some, *morphœa* is regarded as a local expression and an early stage of scleroderma, each being a form of tropho-neurosis.

**Parasitic Diseases.**—These may be caused by the presence either of parasitic animals or of parasitic plants. To affections of the former origin, or the epizoa, belongs especially scabies; though the various forms of lice producing the ailment presenting a pruriginous eruption with little hemorrhagic marks—*phtheiiriasis*—must be mentioned. Another animal parasite, the entozoon or *demodex folliculorum*, inhabits the sebaceous and hair follicles, but does not, so far as is known, cause disease.

The complaints associated with the vegetable parasites, the *epiphytes*, or, as those on the skin are called, the *dermatophytes*, also known by the generic name of *tinea*, are chiefly favus, mentagra, pityriasis versicolor, and some of the forms of ringworm, *tinea circinata*, and *tinea tonsurans*. *Pellagra*, also supposed to be due to a vegetable parasitic growth, is not an affection met with in this country. Nor does the presumed parasitic fungus lodge in the skin. It is said to be found in diseased Indian corn or maize, which, when eaten, causes the general cachexia or cutaneous eruption that characterizes the malady, of which the eruption, moreover, is determined by exposure to the sun. Belmondo found lesions of the spinal cord in a number of cases of this grave nutritive disorder.

As stated by Eichhoff, many cases of *eczema* are of parasitic origin, and indeed it is asserted by others that all *eczemas* are of this character, and that the non-parasitic forms are simply cases of dermatitis.

*Scabies.*—Scabies, or the itch, is owing to the *acarus scabiei*. This burrows into the skin, particularly between the fingers and between the toes, about the wrists, and on the buttocks and abdomen and the upper part of the penis. The channels produced are generally somewhat curved, and may be traced as whitish or more generally black streaks several lines in length, in the situations just indicated. The disease is attended with excessive itching, which is increased at night, and with the eruption of conical vesicles, or even of a marked eczema and of papules and pustules; most of the rash is due to the irritation of scratching.

At the close of our civil war we had a form of itch very prevalent in this country, which was spread far and wide, as is presumed, by contact with the troops,—the so-called *army itch*. It was a chronic and distressing affection, and no age or social state was exempt from it. Indeed, so prevalent was it that it almost appeared as an epidemic. The itching was intense; the eruption, as by far most frequently met with, was like prurigo, but vesicles, or even an eczematous condition of the skin, or pus-

FIG. 88.



A female *acarus*, taken from a photograph from nature; magnified 220 diameters. The ventral surface is shown.

tules, attended the intolerable itching; and in cases of very long duration the appearance of the skin was altered, and all trace of a distinctive eruption was gone. The eruption was seen on the arms, forearms, chest, abdomen, and lower extremities, particularly on the ulnar side of the forearm and the inner aspect of the thigh.

It was sometimes found on the scalp, but very seldom in the groins, in the axillæ, on the hands, or between the fingers. It was benefited by sulphur; for almost all the preparations recommended for it contained sulphur. Whether it was due to the same *acarus* as ordinary scabies, or to a different species, I am unable to say.

*Tinea Favosa*.—*Tinea favosa*, or *favus*, is a chronic disease which gives rise to bright-yellow umbilicated crusts, of circular shape and smooth surface, which often form yellow rings around the hair follicles and are not much elevated above the skin. There is no discharge. The disease rarely affects any other part of the body than the scalp, and produces baldness; when the nails are attacked, they become brittle and yellow. In cases of doubt, the microscope furnishes us with a certain means of diagnosis, by exhibiting the cryptogamic plants.

*Tinea Sycosis*.—*Tinea sycosis*, or barber's itch, is to be distinguished from a non-parasitic form. The distinctive marks of the disease consist in the development of yellowish pustules, having a bright-red base, around the roots of the hair of the beard; the hairy portion of the neck may also be affected. The crusts may run together, and more or less inflammatory thickening of the skin exists. This is especially seen in the parasitic form of the disease, in which, however, less suppuration happens, and less pain or itching, but in which the hairs become brittle and lose their healthy look. The upper lip is rarely implicated in *tinea sycosis*. Non-parasitic *sycosis* consists chiefly in an inflammation around the follicles, which always starts in these parts.\*

*Tinea Circinata and Tinea Tonsurans*.—The *trichophyton tonsurans* is the parasite met with in *tinea circinata*, the ringworm of the body, and in *tinea tonsurans*, the ringworm of the scalp. This is common in children, and spreads by contagion. It exists in circular scaly patches, on which are dry broken hairs. In ringworm of the body the patches are also circular and scaly; but they are red and very itchy, and much paler in the centre than at the edge. Examining the scurf, we find the fungous growth. *Tinea kerion* is a suppurative form of *tinea tonsurans*.

*Tinea Versicolor*.—This parasitic affection, also known as pityri-

---

\* Robinson, New York Medical Journal, Aug. and Sept. 1877.



asis versicolor, occasions those yellow or yellowish-brown discolorations which may be not infrequently seen on various parts of the body. The affection is to be distinguished from chloasma, which is common in women, especially in pregnant women. The microsporon furfur of Eichstädt is the parasite present in this disorder; and it is found abundantly in the scales which can be scraped from the raised, itching patches. In pityriasis affecting the scalp we may also find parasitic growths of vegetable nature; they are often the cause of baldness, as in *tinea decalvans*. Pityriasis capitis, or dandruff, is readily distinguished from ordinary seborrhœa, in which the oily element predominates in the scales, that are aggregated in masses. This is in marked contrast to the fine pearly scales of pityriasis, which are due to epithelial exfoliation. This condition often leads also to baldness. Some forms of *alopecia* are of parasitic nature.

The diagnosis of *actinomyces* of the skin depends upon the history, the mode of invasion, and the distribution of the tumors in necklace-like series, either in lines or in circles, or in groups. The disease pursues a rapid course, with fever, sometimes septicæmic in character. Majocchi \* recognizes two forms, the anthracoid and the ulcero-fungous. In the former the lesions are flat-topped, with a multitude of small openings from which thick yellow pus exudes; in the latter ulceration occurs early, with large granulations. In both forms the lymphatic glands in the neighborhood of the lesions are not involved. The lesions are situated chiefly around the buccal cavity. Microscopical examination of the yellow granules reveals the characteristic actinomyces.

**Altered Gland-secretions.**—One of these, *seborrhœa*, or increased secretion from the sebaceous glands mixed with epidermic scales, has been already mentioned. It is especially found on the scalp, nose, and genitals, and is often seen among those who have menstrual disorders. It is unattended by itching; the crusts are readily removed by strong alkaline soaps, and the skin beneath is healthy, or pale and glistening or slightly reddened.

Where the sebum is retained in the follicle, giving rise to little prominences, apt to be discolored by dirt, and without, as happens in acne, decided inflammation around the gland and its duct, the

---

\* Annales de Derm. et de Syph., Paris, 1892, p. 310.

disorder is called *comedo*. The plug of sebum can be easily squeezed out. The disorder is most common on the face and shoulders of young persons of lymphatic temperament.

The *sweat-glands* are often altered in their activity, and excessive perspiration results. This may be general, or confined to particular localities, as to the hands and feet. This local sweating is often offensive, and makes the parts very tender. The disease formerly known as *lichen tropicus* is now regarded as due to congestion or inflammation of the sweat-glands, and is therefore termed more correctly *miliaria papulosa*. The strophulus or "red gum" of infants is *miliaria vesiculosa*. At times there is sweating of blood from the skin, as in the case recorded by Hart.\* This condition, known as *hæmidrosis*, is due to some alteration in innervation, and may be preceded or accompanied by a localized erythema or eczema; or the bleeding may come from the follicles of the skin; it is not a secretion of the sweat glands, but is a hemorrhage, or an exudation.

*Molluscum epitheliale* presents numerous globular or flattish nodules, sometimes seated on a broad base or attached to a pedicle. They are due to the presence of a psorosperm in the deep layers of the skin and the sebaceous glands. The lesions occur chiefly in groups on the face and neck, or on the trunk; they have often a doughy feel, vary in size from that of a pea to that of a pigeon's egg, show no tendency to inflame or ulcerate, and are not attended with increased sensibility of surface. They are of the color of the skin or of brownish hue. They may last during life and increase slowly without affecting the general health. There is a variety met with especially in children, which has at the top or the side of each tubercle a small orifice, from which a creamy, fatty fluid can be pressed. This variety is regarded as contagious; though there are many who doubt the contagious nature of "*molluscum contagiosum*." The little tumors are distinguished from so-called *molluscum fibroma* by the central aperture, and by the substance resembling sebum that can be squeezed out of them.

Although *plica polonica* is of rare occurrence in this country, yet among immigrants it is occasionally met with, and may be

---

\* Louisville Medical Journal, Jan. 1875.

recognized by the mass of matted hair, which is felted and defies efforts to untangle the mass. Associated with it is some inflammation of the scalp, from which serous oozing occurs. The mass of hair affords refuge for vermin, and the secretions from the scalp produce a peculiar odor, which has led to the supposition that the disease is caused solely by dirt. Dumesnil considers it a neurosis and the dirt only incidental.\* Under the microscope the hairs show decided change, affecting mainly the medulla, which in the most diseased parts is entirely lost. Jarochevski has pronounced the disease a disturbance of nutrition of neurotic origin. It is related on the one hand to gray hair and on the other to alopecia.

**Nervous Affections.**—These are of many varieties. Several of these, such as herpes zoster, have already been considered. The large group of itching affections in which no obvious local affection exists, find here their place. Such are, for instance, the various forms of *pruritus*, either local or general, which are specially apt to affect elderly persons. Sometimes the itching is very violent and obstinate, and we cannot even trace it to reflected irritation, though this is often its cause. Again, diabetes, gout, lithæmia, or jaundice may lie at the root of the pruritus. At times we can find no cause for it. Season influences it much, as seen in the winter itching, the *pruritus hiemalis*, described by Dühring. It happens particularly about the thighs and legs, and there may be prominence of the hair follicles. Among the other manifestations of nervous skin affections are *dermatalgia*, *hyperæsthesia*, *anæsthesia*; then there are undoubtedly cutaneous diseases which are being more and more recognized as of nervous origin.

The disorders of the skin which we have been considering do not always occur isolated; they may be combined. Again, they are altered by the existence of a special taint, as by the syphilitic. Now, without making any attempt to describe *syphilitic* diseases of the skin, it may briefly be stated that they differ chiefly by their copper-colored tint, by the stained aspect they leave, and by the absence of pain and of itching. In syphilitic erythema the eruption runs a very chronic course, and is very distinct generally on the trunk. It belongs to early syphilis. Syphilitic lichen has

---

\* International Clinics, Fourth Series, vol. ii., 1894.

better-defined, more obvious papules than simple lichen. The ulcerations in the pustular affections are deeper; while in the squamous disorders the scabs are smaller and the papules larger than in the non-syphilitic eruptions. A furunculoid eruption is occasionally met with in hereditary syphilis. Syphilitic affections of the skin are very apt to be mixed, and light is thrown on them by this fact, as well as by the history of the case, the sore throat, the falling of the hair, and the nerve- and bone-pains.



## CHAPTER XIV.

### POISONS AND PARASITES.

TOXIC symptoms from causes arising within the body, either from fermentative or putrefactive changes of the food within the intestinal tract, or from micro-organisms causing infectious diseases, septicæmia, sapræmia, and the like, have been referred to in other chapters, especially those on Diseases of the Blood, the Acute Infectious Diseases, and Gastro-Intestinal Affections and Fevers. In this section will be considered only those disorders due to poisons or parasites, the morbid phenomena of which are clearly occasioned by causes introduced into the system from without. Thus they agree in being affections of external origin; and as regards both the diagnosis and the treatment, our chief aim is to ascertain precisely to what foreign substance the symptoms are due.

### POISONS.

Cases of poisoning may arise from accident, attempt at suicide, or criminal intent. It is not necessary here to enter at any length into the subject of toxicology, but merely to set forth the main signs by which the most common poisons may be recognized and distinguished. For this purpose it will be convenient to consider the cases as divided into acute and chronic, subdividing these classes according to the character and effects of the different substances.

#### Acute Poisoning.

The attack comes on suddenly, the patient, previously in perfect health, having taken some food, drink, or medicine which has been followed by the severe symptoms. It is always, in a case of suspected poisoning, of the utmost importance to be able to make out these points.

**Irritant Poisons.**—The chief articles which give rise to acute poisoning belong to the class of irritant poisons. The symptoms are generally those of acute gastritis, attended often with more or less inflammation of the mouth, the fauces, and the œsophagus. Sometimes the air-passages may be involved, either directly or by sympathy, and we find hoarseness and cough. Convulsions are occasionally observed, and collapse is apt to occur sooner or later.

The acute pain, the tenderness, and the vomiting come on shortly after a meal, or at least after something has been swallowed. This distinguishes the acute gastritis caused by poisons from *idiopathic acute gastritis* or from *acute gastric catarrh*. Sometimes several persons are similarly affected,—a circumstance always strongly in favor of the idea of poisoning. From *perforation of the stomach or intestines*, irritant poisoning is discriminated by noting that the acute signs in the former case follow upon the manifestations of some gastric or intestinal disorder; and the attending phenomena of collapse are not, as in poisoning, associated with cramps or convulsions. *Cholera morbus* is separated by the history of the case, by the absence of epigastric tenderness, and by the purging and vomiting often coming on simultaneously. *Cholera* resembles poisoning in the suddenness and the violence of the attack, but is distinguished by the rice-water discharges and by its epidemic character. Bacteriological examination of the stools also affords a means of diagnosis. In *strangulated hernia*, the comparatively gradual onset, the pain, the tumor, and the constipation will be significant. As regards the separation of those cases of poisoning in which blood is ejected, from ordinary *hemorrhage from the stomach*, we find that pain and purging are both absent in the latter, while in irritant poisoning they are well-marked symptoms.

Let us now examine some special poisons. Strong acids are sometimes used in self-destruction. *Nitric acid* stains the lips and mouth orange-yellow wherever it touches them. *Sulphuric acid* stains the skin or mucous membrane white or even dark gray; the pain is excessive, and nervous symptoms are not infrequent. If the vomited matter be mixed with a solution of barium nitrate, a dense white precipitate of barium sulphate is thrown down. *Hydrochloric acid* is less irritant and corrosive than sul-

phuric acid ; in the ejected matter silver nitrate produces a copious white precipitate insoluble in nitric acid. *Oxalic acid*, when concentrated, is rapidly fatal. The irritant effects are those of the mineral acids ; but we also meet with dyspnœa and with nervous phenomena, such as anæsthesia, paræsthesia, palsies, and convulsions.

The strong *alkalies*, when taken into the stomach, cause inflammation of the organ and of the fauces and the œsophagus. Should the case end in recovery from the poisonous influence, thickening of the œsophagus is apt to occur. *Ammonia* may also induce severe nervous symptoms, similar to those of tetanus ; its vapor sometimes acts powerfully on the air-passages, producing harassing cough. *Potassium* and *sodium hydroxides*—commonly known as caustic potash and caustic soda—give rise to violent local inflammation in the mouth, œsophagus, and stomach. The vomited matter has an alkaline reaction. *Potassium nitrate* is a strong cardiac sedative.

*Potassium iodide*, *iodine*, *bromine*, and *chlorine* are all capable of destroying life by their intensely irritant effect.

*Phosphorus*, which is not infrequently taken as a poison, imparts to the breath, to the fæces, and even to the urine an alliaceous smell, and may make them luminous in the dark. It acts as an irritant, causing obstinate vomiting and purging, pain at the epigastrium, rapid, weak pulse, jaundice, and unquenchable thirst. The local pain and inflammation are usually extreme, and collapse, with or without convulsions, comes on early. In some cases painful cramps in the limbs occur, and various disturbances of sensibility, and, later, violent delirium and convulsions, eventuating in coma and in death. In other cases hemorrhage is a striking feature, the blood is very fluid, and issues from all the passages, and petechiæ form beneath the skin. The temperature remains normal until near death. The pulse becomes feeble and small ; the first sound of the heart almost disappears ; peptonuria is observed.\* Jaundice is a constant symptom ; it seldom, however, comes on before the third day, and is rarely intense ; it may be associated with urticaria. The spleen increases in size

---

\* L'Abeille Médicale, July, 1882, quoted in Medical News, Phila., vol. ii., 1882 ; also Jaksch, Wien. Med. Presse, Oct. 1882.

simultaneously with the liver. The urine becomes very scanty. Albumin, blood, and casts are occasionally present in the secretion, and the biliary coloring-matter is usually; urea is very defective. In cases of phosphorus poisoning, acute and extreme fatty degeneration of the tissues happens. It occurs with astonishing rapidity. It has been seen, in the bodies of persons poisoned by phosphorus, within so short a period as forty-eight hours, and has been found to affect the heart, the smaller blood-vessels and capillaries, the liver, the kidneys, the glands of the stomach, and the voluntary muscles.\* The liver is always principally implicated.

Various compounds of *potassium*, *copper*, *zinc*, *silver*, *lead*, and *iron* occasionally cause death. They act, for the most part, as irritants merely; but some of them are powerfully astringent, and even caustic, as, for instance, zinc chloride or silver nitrate. If the toxical phenomena are due to the nitrate of silver, the staining of the lips may afford a clue to the nature of the case. There are no really distinctive symptoms produced by large doses of *arsenic*, of *antimony*, of *mercury*, or of their compounds, which are among the best known of irritant poisons: the peculiar effects of each of these substances, when insidiously introduced into the economy, will be presently mentioned. In *acute arsenical poisoning*, besides the pain and the gastro-enteric symptoms, convulsions, delirium, palsies, and bloody or albuminous urine have been specially noticed. Arsenical poisoning is a very common form of self-destruction. It is also observed among those who accidentally take Scheele's green, or among children who swallow arsenical paints. There is in the internal organs a fatty degeneration similar to that in phosphorus poisoning. In the recognition of the cause of the symptoms, Reinsch's test, applied to the vomited matter, is very convenient and satisfactory. In poisoning by *corrosive sublimate*, epigastric pain, vomiting, diarrhoea, bloody stools, and finally collapse, are met with.

Among animal substances, *cantharides* has sometimes been productive of poisonous effects; strangury, bloody urine, albuminuria, more permanent than that produced by turpentine, priapism, and spasm of the glottis, are the most marked symptoms; while

---

\* Tardieu, *Etude médico-légale sur l'Empoisonnement*, 1867, p. 445.



the shining, green particles of the drug, if taken in substance, have been detected in the vomited matters.

*Sausage, milk, cheese, eggs*, especially in articles of confectionery, such as *cream puffs*, frequently produce violent symptoms suggesting some of the more powerful irritants, although chemical examination fails to reveal any mineral poison. The researches of Vaughan have shown the main cause of these actions. Under the influence of certain micro-organisms, the albuminous matters undergo rapid decomposition, producing a nitrogenous body which has been identified as *diazobenzene*. Vaughan originally called this body tyrotoxicon, cheese poison. It is highly poisonous, but also very unstable. It is produced early in the decay of the albuminous articles, and is decomposed subsequently. We can therefore understand why articles of food may be less irritating when decidedly decomposed than when decomposition has just set in. Besides the signs of gastro-intestinal irritation, vertigo, headache, marked anxiety, and muscular weakness have been noticed among the effects of these ptomaines.

The vegetable irritants are mainly articles commonly used as purgatives. Thus, *elaterium*, *aloes*, *colocynth*, and *colchicum* have all proved fatal when taken too freely. The symptoms do not differ materially from those caused by other poisons of this class. *Tobacco* and *lobelia* are powerful local excitants, occasioning emesis and purging, with a speedy collapse of the system. The former, when the nicotine produces acute symptoms of poisoning, gives rise also to salivation, cold sweats, slow pulse, colicky pains, and at times convulsions. *Savin* not only produces inflammation of the alimentary canal, but is apt also to give rise to strangury; it is most frequently resorted to with the view of bringing on abortion. *Ergot* is also used for the same purpose; the most striking symptoms of acute ergot poisoning are colic, vomiting, diarrhoea, increased salivation, retardation and weakening of pulse, muscular weakness, and, in severe instances, stupor. The poisoning rarely ends fatally.

*Poisonous fungi*, such as the fly fungus, which are eaten by mistake for mushrooms, produce violent symptoms of irritant poisoning attended with other phenomena. The poisonous agent in the fly fungus is *muscarine*, and it gives rise to vomiting, violent colic, and diarrhoea, besides slowing of the pulse and the breathing, and

violent excitement followed by stupor and somnolency. The case generally lasts two or three days, and may then end in recovery or in collapse; but it may terminate fatally in six or seven hours, hæmoglobinuria being among the symptoms. Finding the fungi in the vomited matter or in the stools greatly facilitates the diagnosis. Other poisonous fungi produce much the same symptoms; and even the usually-eaten and innocuous kinds of mushrooms may, if at all spoiled, or in certain individuals, or when eaten raw, occasion similar symptoms.

**Narcotic Poisoning.**—The symptoms of narcotic poisoning vary more, according to the special article taken, than those caused by irritants. Narcotic poisons affect chiefly the nervous system and the circulation. Many of them produce phenomena like apoplexy and intoxication, from which they need to be carefully distinguished. Narcotic poisoning is, for the most part, of the acute form.

*Opium* is by far the most important of narcotic poisons. It induces giddiness, stupor, and lethargic sleep, from which, however, the patient can at first be roused, if sharply spoken to. Subsequently this sleep deepens into coma and cannot be broken; the skin is relaxed and perspiring; the face is usually pale; the pupils are contracted and insensible to light; erections of the penis are common. A more or less evident odor of opium may often be perceived about the person or on the breath. No distinction can be drawn between the effects of different forms of this poison: the stronger the preparation, however, the more marked and the more rapid will be the progress of the case. Morphine, codeine, narcotine, and the other alkaloids give rise to similar symptoms, but the smell of opium is absent; convulsions are most likely to occur from narcotine, papaverine, and thebaine.

The diagnosis of opium poisoning from *apoplexy* and from the coma of *uræmia* has been discussed in a former chapter. We may merely recall that the contracted pupil caused by opium is of very great significance, and does not, with the exceptions there referred to, exist in the other states. Moreover, the coma of apoplexy is at once developed; while in narcotic poisoning it is not sudden, but is preceded by drowsiness or stupor, which gradually passes into coma. These phenomena occur also in the same sequence in *uræmia*; but they are even slower in their progress, and are

frequently associated with convulsions and with markedly albuminous urine and dropsy.

From *acute alcoholism* we discriminate opium poisoning chiefly by the absence of the alcoholic odor, the slow respiration, and the presence of morphine in the urine. The characteristic smell of *chloroform*, the great pallor of the countenance, the complete and speedy collapse, and the absence of contracted pupils distinguish chloroform poisoning from opium poisoning. It is the same with *ether*. Poisoning by chloroform and by ether is mostly encountered during surgical operations.

*Chloral*, in excessive doses, produces heavy sleep, with contracted pupils; but they dilate on awaking.\* There is some reduction of temperature, with rapid pulse, giddiness, inability to walk straight, double vision, and headache, in cases in which consciousness, sensibility, and muscular power have not been entirely suspended by the drug. Weak action of the heart is another of the dangers of chloral poisoning, and I have known a dilated heart almost paralyzed even by small doses. In some instances a stage of excitement like alcoholic intoxication precedes the narcotism. The urine may or may not contain sugar.† Chloral itself simulates sugar in the copper and bismuth tests. It is occasionally used for drugging liquor for purposes of robbery or rape.

*Benzene*, when taken internally, occasions noises in the head, muscular tremor and twitchings, and deep sleep; but the narcotic depression ends in recovery.

*Alcohol*, if taken in large quantities and not much diluted, gives rise to symptoms like those caused by opium. The eye is injected and the seat of ecchymosis; the pupils are, as a rule, dilated and very sluggish; the breathing is irregular and stertorous; the temperature lowered; the insensibility may alternate with convulsions; the breath has a strong smell of alcohol or may be quite free from spirituous odor. This absence of odor of the breath, although not usual, may give rise to a confusion between alcoholic poisoning and apoplexy, and the discrimination of these condi-

---

\* Taylor, On Poisons, 3d edit., 1875.

† See a case of mine recorded in a Clinical Lecture on Chloral Poisoning, Phila. Med. Times, March, 1883.

tions must then depend in some measure upon evidence furnished by the history of the occurrence of the insensibility, and by the presence or absence of palsy.

Alcohol may readily be detected in the urine. Woodbury's \* modification of Ainstie's test is very convenient. Into a tube containing a gramme of sulphuric acid, which should be colorless or nearly so, twice as much of the urine to be tested is poured. A small crystal of bichromate of potassium is then dropped in, and the liquid slowly mixed by rotating the test-tube. If alcohol be present in proportion as large as two or three parts per thousand, a permanent green discoloration will result; if there be less than this, the liquid will remain of ruby color. Chloral in the urine does not produce the peculiar reaction.

*Belladonna*, or its active principle, *atropine*, and *hyoscyamus* produce more marked excitement of the brain than opium does, causing delirium of active kind, perhaps with convulsions. The pupils are greatly dilated, and vision is singularly deranged; there is intense thirst, with great dryness, redness, spasm, and burning in the throat; the breathing is rapid, thus differing from apoplectic conditions. The temperature is always lowered; the pulse becomes quick and compressible; a scarlet efflorescence may happen. The surest test of poisoning by atropine is to take some of the urine passed, and with it to dilate the pupil in the eye of a cat.

*Conium* occasions stupor, paralyzes the muscular system, and dilates the pupils; there is dyspnoea, while the heart, though rendered slower, is not much affected. Convulsions may come on. These help to distinguish conium poisoning from curare poisoning, which it much resembles. In the latter, however, the palsy is greater.

*Carbolic acid*, if taken in poisonous doses, produces rapidly dangerous symptoms, which in bad cases terminate in death in a few hours. Vomiting, slow pulse, noisy breathing, loss of consciousness, deepening into profound coma, abolition of reflex movements, cool skin, suppression of urine, are the main symptoms. When the urine is obtained, it is of dark-green or black color; this and the odor of carbolic acid about the patient are

---

\* Phila. Med. Times, March, 1883.



very significant features. The discolored urine is apt to contain blood-corpuscles, casts, epithelium, and tube-casts.

*Aniline* poisoning is met with among the workers in factories in which the aniline colors are made. It is the breathing of the aniline vapor, especially, which occasions the toxic effect. Vertigo, headache, a sense of suffocation, vomiting, anæsthesia, pain in the extremities, somnolency, and a dark cyanotic discoloration of the ears, the nails, and the mucous membrane of the nose, have been especially noticed.

*Hydrocyanic* or *prussic acid* usually leads to convulsive contractions of the muscles of the limbs and trunk, and destroys life by stopping the circulation and the respiration. Sometimes the odor of the acid, resembling that of bitter almonds, is perceptible in the breath; but too much reliance must not be placed upon this point. Unfortunately, the diagnosis of this poison has generally to be made after death, for medico-legal purposes.

The gases arising from burning *coal* and *charcoal* may cause death by asphyxia; and a knowledge of this fact has, particularly in France, led to many suicides. In those cases in which the asphyxia has not a fatal termination, yet has been decided, disorders in the peripheral nerves may manifest themselves, either by the signs of neuritis, or by pain and swelling simulating a phlegmon, or by vesicular eruptions in the course of an affected nerve. The peripheral disturbances may appear at once or not until after some days. The signs of disorder of the vaso-motor nerves do not last long; those of the motor or sensitive nerves have a longer duration; the complaint induced may be incurable, extending from the centre to the periphery, or in the reverse direction; or, lastly, the affection may cause an acute ascending paralysis.\*

The poisonous action in these cases is due largely to carbon monoxide, carbonic oxide, a gas which has a strong affinity for hæmoglobin, and suspends the oxygen-absorbing function of the blood, thus establishing a chemical asphyxia. The gas, being non-irritating, may be inhaled without exciting immediate suspicion. The so-called water-gas contains large amounts of carbon monoxide. Experiment has shown that such gas is much more dan-

---

\* Leudet, Arch. Gén. de Méd, May, 1865.

gerous when inhaled than the ordinary illuminating gas, which consists almost entirely of compounds of carbon and hydrogen.

*Antipyrine* given in large doses may produce extreme lowering of the temperature and collapse. Cyanosis, frequency of respiration and of pulse, dyspnoea, a feeling of extreme heat over the body, and an erythematous, urticarial, or measly eruption, have also been noticed. In one instance reported, the use of the drug led to the formation of membranes in the mouth and to symptoms of laryngeal spasm, which was not the case when phenacetin, antifebrin, or exalgin was substituted.\*

*Petroleum* taken in excessive quantities produces giddiness, faintness, and palpitation, with tonic and clonic convulsions, contracted pupils, hot skin, and slow pulse; it does not occasion either stupor or vomiting; the urine has an aromatic odor. Recovery is the rule.

*Nitroglycerine* occasions a throbbing headache increased by motion, mental confusion, flushing of the face, pulsations all over the body, arterial relaxation, and collapse.

Following these poisons, which are in the main narcotic poisons or belong to the group of poisonous carbon compounds, we shall examine some forms of acute poisoning produced by certain powerful vegetable poisons.

*Aconite* has a strongly sedative influence upon the action of the heart, brain, and spinal cord, as well as an irritant action upon the alimentary canal; slow pulse, giddiness, delirium, numbness, and tingling of the skin, loss of power in the legs, with formication, tingling of the tongue, vomiting, and purging, are followed by syncope and death.

*Digitalis* causes dilatation of the pupil, generally with vomiting, often with purging and with headache, giddiness, and suppression of urine; its chief effect, however, is upon the pulse, which is strikingly lessened both in frequency and in force, and becomes irregular; the action of the heart, too, becomes weak, and blood-pressure is diminished. The skin is cold, pale, and covered with sweat; the mind is generally clear, though there is great lassitude, with muscular debility, a tendency to sleep, and at times convulsions. The action of the poison generally extends over

---

\* Salinger, Amer. Journ. Med. Sci., May, 1890.

days. *Veratrum viride* resembles *digitalis* in its action. It markedly reduces the pulse, and gives rise to vomiting, to great prostration, and to irregular breathing. The temperature is much lowered.

*Calabar bean* acts as a direct sedative to the spinal marrow, particularly to the medulla, and produces great muscular debility or relaxation, or even paralysis, extending to the heart and respiratory muscles. The mental faculties remain unaffected, and in this its action differs from that of the cerebral sedatives. It is, however, irritant to the alimentary canal, causing vomiting or purging, a peculiar epigastric sensation is generally experienced, and increased salivation is met with. Calabar bean contracts the pupil and also the ciliary muscle, thus making the eye myopic. The condition of the eye is the main diagnostic sign that distinguishes poisoning by calabar bean from poisoning by curare or by conium.

*Strychnine* and *brucine*, the active principles of *nux vomica* and of several allied plants, give rise to phenomena strongly resembling those of *tetanus*. A very short time, however,—from a few minutes to an hour or two,—will determine the issue of a case of poisoning; while *tetanus* may run a course of several weeks. The first symptoms of strychnine poisoning are apt to be a sense of suffocation and dyspnoea, followed by spasms of the respiratory muscles, by starting and twitching and rigidity of the arms and legs, especially of the extensor muscles, but not by lock-jaw; *tetanus*, on the other hand, comes on with setting or locking of the jaws, and the limbs are not at first affected with spasms; indeed, the arms remain throughout nearly free from them, and the paroxysms of spasm do not follow one another so rapidly as in strychnine poisoning, and are of shorter duration. Again, idiopathic *tetanus* is extremely rare; almost always there has been some wound or injury as a proximate cause of the malady. But we need not pursue these points of diagnosis further: they have been already mentioned in connection with *tetanus*. From *epilepsy* strychnine poisoning differs by the unimpaired consciousness; from *hydrophobia*, by the absence of spasm of the œsophagus and of the terrible dysphagia.

*Pierotoxin* also produces convulsions which may be mistaken for those caused by strychnine. But they are not of a reflex nature,

and reflex spasms are not induced. The breathing is rapid; the contraction of the heart is retarded; there are often somnolence and muscular debility. A scarlatinal eruption has been noticed.

### Chronic Poisoning.

When the patient has been subjected to the continuous action of a noxious substance, the case is said to be one of chronic or slow poisoning. Any of the irritant poisons, given in small and repeated doses, will keep up a morbid condition of the stomach and bowels much like ordinary chronic inflammation.

The narcotics, taken in the same manner, act upon the vasomotor nerves and the cerebro-spinal system, and through this upon the alimentary canal, so deranging digestion and nutrition as even indirectly to cause death. *Opium* is the most important of the articles thus used; it is often administered to infants for the purpose of quieting their cries, and the frequent repetition of the dose induces a series of phenomena closely allied to those observed in the adult. With the effects, on the mind, of opium taken persistently for the sake of intoxication, the reading world is familiar through the published experiences of De Quincey and of Coleridge.

The habit is here and in Europe generally acquired only by persons who have begun the practice for the relief of some painful affection; in the East, opium is used much more commonly, and, in many Oriental countries, to smoke it is a favorite amusement. Those who employ it constantly are pale, or have a sallow, haggard countenance and a dull eye. They lose their power of will and their energy, and are troubled by loss of appetite, giddiness, anomalous neuralgic pains, sleeplessness, and low spirits, which they remove by resorting to the opiate. Though, in spite of the pernicious custom, the general health may remain for many years good, yet sooner or later it gives way, and the opium-eater dies worn out; or death may be the consequence of disease of the liver, of palsy, or of inveterate diarrhoea, produced by long addiction to the vice. Persons who consume large quantities of opium are apt to have, from time to time, attacks of extreme nervous prostration, attended, perhaps, with violent headache, and requiring free stimulation for their relief. The employment of morphine hypodermically has become an alarmingly frequent form of the



opium habit, especially among members of the medical profession. Besides the general symptoms of chronic opium poisoning, we may have extensive ulcers and other local signs of skin irritation to deal with.

*Ether* and *chloroform*, habitually made use of, also cause serious disturbance of the nervous system; and so does *alcohol*. The abuse of spirituous liquors gives rise to a disorder of the mental, motor, and sensory functions, producing sleeplessness, headache, giddiness, hallucinations, imbecility, anæsthesia, disordered vision, and palsies. There results a fine irregular tremor, affecting particularly the hands, lips, and tongue, and occurring only on attempted movement. Multiple neuritis is also a common sequel. *Chronic alcoholism* also occasions a sensation of choking, a diminished vitality, a persistent catarrh of the gastro-intestinal membrane, a tendency to fatty degeneration, especially of the liver and kidneys; in short, the symptoms met with in drunkards and constituting the state described as chronic alcoholism. Chronic alcoholism in the parent may produce epilepsy in the child.

*Chloral* has proved, like opium and like chloroform, a very fascinating drug to many. The chief symptoms of chronic chloral poisoning are digestive disorders, irregular breathing, impairment of intelligence and of memory, persistent drowsiness, almost stupor, striking enfeeblement of will, want of power in the legs amounting at times to paralysis, and occasional tremor. Defective co-ordination with marked ataxic symptoms, similar to those of locomotor ataxia, and loss of knee-jerk, occur from the habit of taking chloral.\* I have known delirium tremens to follow its use, when large quantities of it had been taken and the medicine stopped. Feeble, irregular action of the heart, and sweating, I have also found among the symptoms of chloral poisoning. An erythematous inflammation of the skin of the fingers, with desquamation and ulceration around the borders of the nails, has been pointed out as a result;† and various forms of eruption, such as urticaria, lichen, and purpurous spots, as well as bed-sores, have been observed after its prolonged use.

*Paraldehyde* is abused like chloral and morphine. It gives

---

\* J. C. Wilson, article "Opium Habit and Kindred Affections," *System of Practical Medicine by American Authors*, vol. v.

† Smith, *Lancet*, vol. ii., 1871.

rise, when taken habitually, to gastric disorder, diarrhœa, sleeplessness, feeble circulation, sweating, and delirium tremens.

*Tobacco* used in excess gives rise to tremors, to giddiness, to emaciation, to impaired digestion, and to intermittence in the pulse, with irregular cardiac action and palpitations, which may become very annoying and originate the belief of an organic disease of the heart. Like the persistent abuse of alcoholic drinks, tobacco may occasion amaurosis; and it is also affirmed that an insidious, obstinate form of otitis is developed in inveterate smokers, and is attended with very minute granulations of the pharynx, nasal fossæ, tubes, and middle ear.\* When taken in large quantities by those previously unaccustomed to it, tobacco produces colic, diarrhœa, weakness, sleeplessness, dull hearing, vomiting, difficulty in breathing, cold sweats, feeble action of the heart, and will even cause collapse and death. The peculiar odor of tobacco may assist us in the diagnosis of tobacco poisoning; but it must be remembered that this may attend other morbid states in those who use tobacco largely.

*Ergot* long continued, particularly when taken contained in impure flour, gives rise to the well-characterized disease, chronic ergotism. This appears mainly in two forms: the first is marked by convulsions with disturbance of sensation; the second by gangrene; both are apt to show themselves in epidemics. In the convulsive form there is at first formication, which lasts, whether attended with anæsthesia or not, throughout the whole illness. Soon muscular twitchings and cramps followed by painful contractions happen, and the convulsions may become very general. These spasms especially affect the flexors of the arm, and, unlike those of strychnine, they are not reflex spasms. There is no fever; the circulation is slow and feeble; the appetite is insatiable; we find nausea, vomiting, and diarrhœa. The disease generally lasts one or two months. In severe cases delirium occurs as a precursor to death. In gangrenous ergotism the same symptoms happen; but in addition we meet with gangrene without fever or signs of inflammation. The gangrene may be in the extremities or in the face. Where ergot is being taken in diabetes, the gangrene results from the malady, not from the drug.

---

\* Triquet, Le Briert.

Let us now examine some of the features of slow poisoning by the metals.

*Mercury*, in any of its preparations, may lead to chronic poisoning. The mouth is inflamed, the gums are sore and swollen, the salivary glands act inordinately, and the breath is very offensive. Colicky pains, and sometimes diarrhœa, occur. Tremors of the limbs when any motion is attempted are particularly frequent in cases where the poison has been inhaled in the form of vapor; they come on by degrees, and are associated with loss of power of locomotion and with digestive disturbances. The tremors may be incessant and the movements involuntary, like those of chorea, and so rapid as to prevent the patient from obtaining rest at night.\* In some cases an eczematous affection is observed.

Poisoning by mercury is generally the result of the exposure to its action incidental to certain occupations, such as glass-plating, gilding, and working in quicksilver-mines.

*Lead poisoning* is by no means uncommon among painters, plumbers, type-setters, and other workers in lead. Sometimes it may be caused by accidental circumstances, as when the patient has drunk water passed through leaden pipes, or taken snuff which has been impregnated with lead for the purpose of coloring it. Poisonous properties are also acquired by snuff wrapped in lead-foil; and lead poisoning has been observed after the use of cosmetics; and among those engaged in the manufacture of lucifer matches, of brushes, of lace, or working in glass enamel or glass powder;† and in consequence of food adulteration, especially of the use of lead chromate to color cakes.‡

In such cases, the physician may have to depend entirely upon a correct appreciation of the symptoms for the diagnosis. Pain and uneasiness in the course of the colon, constipation, loss of appetite, anæmia, weakness, mental depression, and emaciation are the earlier signs. A metallic taste is perceived; the breath is

---

\* As in a case reported by Taylor, in which the patient died from the effects of the poison, without, however, having presented salivation or mercurial fetor of the breath, or a blue line on the gums. Guy's Hospital Reports, 3d Series, vol. x.

† Lacharrière, Arch. Gén. de Méd., Dec. 1859.

‡ Stewart, Clinical Analysis of Sixty-Four Cases of Poisoning by Lead Chromate, Medical News, Dec. 31, 1887, and *ib.*, Jan. 26, 1889.

fetid, the tongue pale and furred; the gums are edged with a narrow blue line of sulphide of lead, deposited mainly outside loops of blood-vessels. Colicky pains are felt from time to time, and a severe and long-continued attack of colic may form the culmination of the disease. The muscles atrophy; electro-muscular contractility to the faradaic current is greatly diminished, to the galvanic current it is frequently unaltered or increased; the sensibility of the skin is but little affected. Occasionally wrist-drop or paralysis of the extensor muscles of the forearms, the well-known phenomenon of lead poisoning, occurs among the first symptoms; but it is more generally preceded by one or more attacks of colic. The right arm mostly suffers first. We also find at times lesions of the tendons in saturnine palsy.\* Yet as regards this palsy we must bear in mind that a paralysis of the extensors occurs which is not due to lead.†

Another manifestation of lead poisoning is found in the severe pains in the joints and the neighboring muscles. These pains have violent exacerbations, and may be associated with cramps of the painful muscles. They are most common in the lower extremity, especially over and near the knee-joints. There are no signs of inflammation of the affected joints and muscles; pressure tends to relieve the pains.

Sometimes, in cases of saturnine poisoning, there is evidence of grave cerebral disorder: epileptiform convulsions, attacks resembling apoplexy, or general tremors and extended paralysis of the muscles, with acute delirium, inequality of the pupil, optic neuritis, retinal hemorrhages, loss of sight, and other signs of nervous disturbance, are noticed. Of course the diagnosis, under these circumstances, will be materially assisted by an accurate knowledge of the previous history of the patient as regards exposure to the action of the poison. The tremors are, like those caused by mercury, peculiar in ceasing when the limbs are supported or at rest; they are increased by movement. There may be tremor in the muscles of the face, which, however, are not affected by paralysis. Another result of lead poisoning is that it leads to granular degeneration of the kidneys. This is apt, again, to

---

\* Medical Times and Gazette, May, 1868.

† St. George's Hospital Reports, 1868, p. 86.



coexist with a gouty condition, which, as Garrod has shown, is one of the results of the absorption of lead. But the kidney affection may be found whether or not the joints are markedly affected. Lancereaux \* has attributed most of the cerebral symptoms and the dyspnœa that may be met with to the diseased condition of the kidneys, which may, however, exist without albuminous urine.

In instances in which the symptoms of lead poisoning are obscure or conflicting, we may search for lead in the urine. The detection of small amounts of lead is simple in principle, but practically cannot be undertaken except by a professional chemist. A considerable proportion of the lead is eliminated by the bowels.

*Copper* poisoning gives rise to dyspeptic symptoms, to diuresis, to loss of flesh, to lassitude and giddiness, to a peculiar greenish-blue perspiration, and to a green line on the gums and teeth. It is said that workmen in copper are singularly insusceptible to cholera or choleraic diarrhœa,† and that wounds in them heal with extraordinary rapidity. Copper appears to be somewhat less liable than mercury, lead, arsenic, or antimony to cause serious chronic poisoning, possibly because it is less cumulative. Small amounts of copper are frequently present in the liver and brain of man and some of the lower animals, also in some articles of food. Dr. Leffmann informs me that, in the examination of viscera from the cases of lead poisoning which occurred in Philadelphia, copper in minute amounts was frequently encountered, and in one case, that of a child four years of age, an appreciable quantity was obtained from a portion of the liver.

*Arsenic*, administered in small doses for a lengthened period, produces a state of chronic inflammation of the alimentary canal. Conjunctivitis, œdema of the face and the limbs, in some instances associated with albuminous urine, irritability of the stomach, diarrhœa, sleeplessness, increasing weakness, numbness, formication, alterations of sensation, and even paralysis, mark the progress of these cases; the hair and the nails occasionally fall out, and there is much frontal headache. Similar effects are noticed to follow the pernicious habit of arsenic-eating, and will be also encountered

---

\* Arch. Gén. de Méd., Dec. 1881.

† Clapton, Clinical Society's Transactions, vol. iii.

among persons employed in making artificial flowers and toys, in dyeing cloths, in manufacturing and hanging green wall-papers, or in the sublimation of arsenical ores; those, too, who live in rooms hung with papers containing much arsenic have exhibited the influences of the poison.\* Besides the phenomena of internal poisoning, cutaneous eruptions occur from arsenic. The extensors of the hands and feet are especially affected. In some instances, said to be not uncommon in Russia,† paralysis of the extremities, with muscular atrophy, happens. Arsenical paralysis may have mainly the symptoms of poliomyelitis, as I have had occasion to observe.‡ In other cases there are severe darting pains in the arms and legs, defective cutaneous sensibility, loss of knee-jerk, and the appearances of locomotor ataxia.§ The palsies of arsenical poisoning are now generally thought to be due to peripheral neuritis.

The inhalation of the fumes of *zinc* gives rise to a peculiar form of poisoning, characterized by a sense of weariness, by a feeling of tightness in the chest, and by attacks of shivering, followed by heat of skin and a profuse sweating-stage. This irregular form of ague is common among brass-founders.||

*Carbon disulphide* produces toxical effects of a singular character, conspicuous among which are gastric disturbances, inordinate appetite, loss of muscular strength, a cachectic condition, a feeling of icy coldness in the lower limbs, severe cramps in the calves of the legs, impotence, and, in severe cases, amaurosis, impaired hearing, hallucinations, loss of memory, and complete perversion of the intellect.¶ These phenomena are met with among workers in india-rubber.

*Phosphorus* is often seen, particularly among those who work in lucifer-match factories, to give rise to serious lesions. When the poisoning is caused by inhaling the vapor, it may occasion, as acute phosphorus poisoning does, alteration of the composition

\* James Putnam, Analysis of Twenty-Six Cases, Bost. Med. and Surg. Journ., March, 1889.

† Sculosuboff, Arch. de Phys., Sept. 1875.

‡ Phila. Med. Times, March and July, 1881.

§ Dana, Brain, vol. ix.

|| Greenhow, Med.-Chir. Transact., 1862.

¶ Delpech, Mémoires de l'Académie de Médecine, 1856; and Heurtaux, Recueil de la Société Médicale d'Observation, 1860.

of the blood, a hemorrhagic diathesis, a fatty degeneration of several organs, as well as of the voluntary muscles,\* and peptonuria. It also produces chronic bronchial catarrh, but especially necrosis of the jaw, for which the whole lower jaw has been removed.† The disease begins in carious teeth, and may extend to the cranial bones. Osteophytes form freely in the affected bones. Phosphorus taken internally in doses that gradually exert a poisonous effect leads to chronic inflammation and thickening of the stomach, colicky pains, diarrhœa, hectic fever, general emaciation, falling out of the hair, and to palsies, which are generally the precursors of a fatal termination.

*Animal poisons.*—These may give rise to chronic as well as to acute poisoning. We find, for instance, syphilis, gonorrhœa, hydrophobia, dissecting wounds, snake-bites, acute glanders, and farcy,—all disorders exhibiting the effect of an animal virus. But we have already discussed some of these as far as is admissible in a work of this kind; and of the others it need only be said that the antecedent circumstances generally place the diagnosis beyond a doubt,

Yet there are a few illustrations of animal poisons and their effects which must here, however briefly, be mentioned.

One of these is the *malignant pustule*, or *anthrax*, a terrible malady, which is the cause of many deaths on the Continent of Europe, and which is identical with the *charbon* of animals. The disorder is also prevalent in New Mexico.‡ It is communicated to man by direct inoculation; or by means of the skin or hair of the diseased beast, or by eating its flesh; or by insects which, sucking the poison from the sick animal, implant it on the skin of man. The poison produces a red speck, which develops into a vesicle, under and around which an extremely hard spot forms that becomes gangrenous. The surrounding skin inflames, new vesicles or pustules spring up, and the gangrene spreads rapidly, the patient speedily sinking; or the death of the parts is arrested, and separation takes place between the living and the gangrenous textures. In some cases it is attended with extended œdematous

---

\* Lancereaux, L'Union Médicale, 1863.

† Cases of Hunt and Boker, Amer. Journ. Med. Sci., April, 1865; Wells, New York Med. Journ., Jan. 1866; Wegener, Virchow's Archiv, Bd. xl.

‡ A. H. Smith, Amer. Journ. Med. Sci., April, 1867.

swelling and infiltration of the areolar tissue spreading from the anthrax pimple. It is remarkable how little local pain attends the grave constitutional disturbance, and the signs of low, irritative fever. The disease is found on the exposed portions of the body, as on the neck and hands. It has been traced by Davaine to the presence of filiform bacteria, bacillus anthracis. The blood swarms with these bacilli; and, as Koch has proved, they propagate themselves by spores, which finally grow into bacteria. The researches of Pasteur and of Koch fully confirm the parasitic view of the origin of the disease.

Closely connected with malignant pustule is the so-called "*wool-sorter's disease*." The wool from sheep is not nearly so dangerous as the hair from the goat, the alpaca, and the camel. The mohair from the Lake Van district, Asia Minor, is the most dangerous. The symptoms may be those of malignant pustule with secondary splenic fever, or there often is an utter absence of either external or internal pustule.\* The manifestations of the disease are frequently a low fever with secondary abscesses, pyæmic symptoms, and pleuro-pneumonia. The complaint is a dangerous one, and often fatal; when ending in recovery, convalescence is slow.

Another disease transmitted from infected animals, and due, it is thought, to fungi, is the so-called *actinomycosis hominis*, described chiefly by Israel† and by Ponfick.‡ The disease first appears in the lower part of the face, in the shape of little abscesses containing yellowish granules, which consist of ray fungi. These vegetations are readily detected by the microscope. The disease spreads to the ribs and vertebræ, and produces great destruction of tissue; it is also found in the liver and the lungs, in the brain, in the intestines, and in the skin; there are the symptoms of chronic pyæmia. Various forms of it, as of the liver and of the lungs, have been already described in connection with those organs.

The *foot and mouth disease* is an affection from which especially children suffer who have drunk the milk from infected cows. The

---

\* Bell, Lancet, June 12, 1880.

† Virchow's Archiv, Bde. lxxxv., lxxviii.

‡ Die Actinomykose des Menschen, Berlin, 1882.



poison produces an aphthous stomatitis with digestive disorder, and frequently also a vesicular eruption on the face and on the fingers and hands, which gradually dries into brownish scales, and at times a similar eruption between the toes. The disorder is not a serious one. It is due to a micro-organism, the streptocytus of Schottelius.

There is another form of animal poisoning which may be in this connection briefly considered,—namely, *milk-sickness*. Now, its phenomena are so variously described by writers that its characteristic signs are difficult to define. It prevails in the southern and southwestern portions of North America, and is brought on by drinking the milk or eating the flesh of cattle which have been exposed to certain influences the nature of which is as yet unknown. Gastritis and enteritis seem to be more or less blended in the early stage of the disorder, which at a later period is said strongly to resemble typhus fever. The symptoms dwelt upon more especially are lassitude, nausea and vomiting, with a sense of burning at the epigastrium, great oppression, intense thirst, hot, dry skin, obstinate constipation, and obvious abdominal pulsation. If at all, recovery takes place very tardily, the tone of the stomach being often left impaired for life.

Other forms of animal poisons originate in *alkaloids* generated during decay. The poisoning by these *ptomaines* from milk and eggs and other substances has already been mentioned. Frequently the ptomaine poisoning resembles that of the vegetable alkaloids, such as of morphine, codeine, and veratrine.

Besides these forms of animal poisoning, which are produced by the direct contact with the virus, or at all events by its introduction into the system through the stomach, we find morbid states occasioned by animal poisons which arise from decomposing bodies or excretions, or from the crowding of many together, particularly of those of uncleanly habits, or of the wounded. These poisons reach the blood for the most part by the lungs, in the shape of *poisonous exhalations*. They are very depressing in their action, may lead to low fevers, or to septicæmia, and in the case of the wounded to pyæmia and to hospital gangrene. Persistent nausea, too, and a lowering of all vital energy are not uncommonly observed in those who breathe continuously the foul air under the circumstances alluded to,—as in hospitals and in prisons

in which thorough cleanliness is not enforced and due regard is not paid to ventilation.

In some persons deleterious emanations from the human body give rise to a form of toxæmia, one of the chief features of which is the marked anorexia which attends the great debility.\*

The exposure to animal effluvia may also excite violent diarrhœa, or even symptoms like those of cholera, certainly like those of severe attacks of cholera morbus. Of the occurrence of the former we have an illustration in the *dissecting-room diarrhœa*, which is usually attended with very fetid discharges, and may be accompanied by colicky pains, by nausea and vomiting, and by headache. The same kind of diarrhœa also happens in those who clean privies, or who are exposed to the emanations arising from sewers; or dysentery or choleraic attacks may follow the exposure. Nay, as in instances recorded by Becquerel, the instant disengagement of large quantities of putrid gases, arising from bodies far advanced in decomposition, where coffins have been opened, has caused sudden deaths, or has resulted in so serious a state of poisoning as to give rise to very grave illnesses, having mostly a fatal termination.† In individuals who, in consequence of their vocation, are habitually brought in contact with animal effluvia and are liable to inhale noxious gases, besides the attacks of diarrhœa referred to, chronic disturbances of the stomach and liver, with marked impairment of the general health, may happen. Cases occur of self-infection from ptomaines resulting from decomposition of fæcal matter lodged in the cæcum, or by perforations taking place from the intestine into abscesses near by, into which the contents of the bowel find their way.

### PARASITES.

Parasites are organisms which become secondarily implanted within or upon the body. Some parasites give rise to no symptoms at all; many occasion phenomena closely resembling those of other irritations. In any case, the only absolutely convincing evidence of the presence of a parasite is obtained by seeing it.

---

\* See Dr. Hunt's case, described by himself, in *Pennsylvania Hospital Reports*, vol. i.

† *Traité d'Hygiène*, 3d edit., p. 218.

**Vegetable Parasites.**—The chief vegetable parasites have been mentioned in connection with diseases of the skin; the *oïdium albicans*, present in thrush, and the *sarcinæ ventriculi*, have also been described. All these vegetable growths can be detected only by the microscope; and, particularly in those involving the skin or the hair, it is of the utmost use to employ the liquor potassæ, under the action of which the structures become transparent.

The fungus that penetrates the internal tissues, the *chionyphe Carteri*, gives rise to that terrible disease known as *podelcoma*, or the *fungus foot* of India,—a complaint found among the natives of India who go about with naked feet. The fungus, introduced either through a scratch or passing through the pores of the skin, soon spreads, eating its way into the bones of the tarsus and metatarsus, and into the lower end of the tibia and fibula, producing a species of caries, or rather a breaking up and absorption of the osseous tissues. The fungous particles or masses are generally of deep-black color, firm and globular, varying in size from that of a pea to that of a pistol-bullet; or the fungus presents the appearance of sloughing tissue, and exhibits chiefly white granules; or it consists of particles of pinkish color. The foot is enlarged about the ankle and over the instep; and on each side of the ankle-joint, and on the dorsum as well as on the sole of the foot, are small, soft swellings, having pouting openings that lead to fistulous canals communicating with the bones, which they perforate in every direction. The fungous mass is for the most part situated in the cavities in the bones, and from the canals passing to them transudes a discolored, glairy, or purulent and fetid fluid. The toes are distorted, and the muscles of the leg atrophied; but the fungus does not spread up the leg. The tendency of the disease is to cause death by exhaustion; the only remedy is amputation.\* The affection has also been observed in this country.†

A similar disease, leading to local destruction, is the *perforating ulcer of the foot*. It is very uncommon in this country, although I have known of cases; in France it is not uncommon. It is sup-

---

\* See Carter, in *Transact. Bombay Med. and Phys. Soc.*; and on *Mycetoma*, or the Fungus Disease of India, London, 1874; Aitken, *Practice of Medicine*; Lewis and Cunningham, *Arch. of Dermatol.*, Oct. 1880.

† Kemper, *American Practitioner*, Sept. 1876.

posed to be due to defective vitality of the parts from altered nerve-supply. Local anæsthesia, lowered temperature, and a tendency to profuse perspiration exist. The ulcer leads down to diseased bone. It is generally situated on the first or the last toe, over the articulation of the metatarsal bone with the phalanx.\*

The toes sometimes drop off from a disease which constricts them and enlarges them beyond the point of constriction. The affection is not unusual in Brazil, and seems to be peculiar to the negro. It is known as *ainhum*.†

**Animal Parasites.**—When speaking of the affections of particular structures, some of these intruders have been mentioned,—those found in the skin or in the liver, for instance. It remains to consider chiefly such of the more important ones as inhabit the hollow viscera, certain solid organs, and the muscles.‡

*Intestinal worms* are the most common of all parasites. The general symptoms induced by them are those of intestinal irritation with disordered digestion. The appetite is capricious; the bowels are irregular, sometimes constipated, sometimes relaxed; the abdomen is frequently swollen and hard, and the seat of distressing uneasiness or of colicky pains; the tongue is furred; the breath is fetid; and there is constant itching about the nostrils and anus. The patient, furthermore, grits his teeth during sleep, and is often annoyed by nightmare. Phenomena indicative of a greater or less degree of nervous disturbance are also met with; they may range from mere fretfulness up to delirium, convulsions, chorea, epilepsy, or insanity. Strabismus and amaurosis may be also due to worms.§

There are many kinds of worms known to infest the alimentary canal of man, and they belong to the order of *nematoda*, or round worms, or to that of *cestoidea*, or tape-worms.

The round worms are parasites of an attenuated or cylindrical form, and present these varieties:

\* Savory and Butlin, Med.-Chir. Transact., 1879.

† Da Silva Lima, Arch. of Dermatol., Oct. 1880; Duhring, Amer. Journ. Med. Sci., Jan. 1884.

‡ For full description, see the admirable works of Joseph Leidy, A Flora and Fauna within Living Animals, Smithsonian. Pub., vol. v.; Davaine, Traité des Entozoaires et des Maladies vermineuses; Cobbold, Entozoa; Leuckart, Die Menschlichen Parasiten, Leipsic; Küchenmeister, Manual of Parasites.

§ Hogg, Brit. Med. Journ., July, 1888.



1. The *ascaris lumbricoides*, or *round worm*, bears a considerable resemblance to the common earth-worm, from which it is, however, anatomically different. It inhabits the small intestine, sometimes finding its way into the stomach, or even into the œsophagus, or being discharged through the abdominal parietes.\* When it ascends to the stomach and œsophagus, it causes, before it is expelled by the mouth, sudden attacks of fever and gastric derangement, with nausea and vomiting; and even, at times, marked delirium.† The worms have been known to be so numerous as to obstruct the intestine.

2. The *oxyuris vermicularis*, *thread-worm* or *seat-worm*, is very small, the male being about two lines, the female about five lines in length. The parasite is white, slender, and extremely active; it is found in the anus, and causes intense itching of this part. The annoyance is sometimes such as to excite a suspicion of the existence of piles. It may creep into the vagina, giving rise there to profuse discharges; or into the urethra. It affects children frequently, but is not uncommon in adults.

3. The *ascaris mystax*, a parasite which inhabits the cat, may also infest the human body. It is a moderate-sized nematode, from two to three inches long, though the female may reach about four inches. Its head end is spear-shaped.

4. The *trichocephalus dispar*, or *long thread-worm*, is detected in very large numbers in the ileum near its termination, or in the colon, particularly at its head. It has been found in persons laboring under typhus or typhoid fever, or dying from cholera or diarrhœa. It is from an inch and a half to two inches in length, and is characterized by the hair-like appearance of the head, which is generally buried in the mucous membrane of the intestine. It is not a common parasite, and it is doubtful whether its presence gives rise to any marked derangement.

The *tape-worms*, or *cestoidea*, are jointed entozoa, of a ribbon-like form. They embrace the true tape-worms, or *tæniadæ*, and the *bothriocephali*. Of the former there are eight varieties, all of which have been found in man, though only two—the *solium* and the *mediocanellata*—are at all common; the *tænia saginata*,

---

\* Garnier, L'Union Médicale, Oct. 1861.

† Schmidt's Jahrbücher, No. 10, 1868.

however, has spread over parts of Western Europe.\* The *bothriocephalus latus* is the usual species of *bothriocephalus* met with in the human intestine; it, too, is increasing greatly in Europe, and, it is said, in Texas, particularly in the western portions.†

The *tænia solium*, or *pork tape-worm*, consists of an immense number of joints in connection with a single head. It may attain an enormous length, and inhabits chiefly the small intestines. The researches of Küchenmeister,‡ Von Siebold,§ and others have shown that its eggs become developed into the *cysticercus cellulosæ* discerned in the muscles of the pig, rabbit, and other animals whose flesh is used as food. Cysticerci have also been detected in the muscles, the cellular tissue, the brain, the spinal cord, the heart, and the liver of man, and are most commonly met with in middle age and in the destitute; they are the most frequent parasite in the eye.|| They cannot, as a rule, be diagnosed, except they be in positions in which they can be seen or felt, or the little tumors they occasion in the subcutaneous tissues are extirpated and examined. In the brain their chief symptom is violent and rapidly-increasing epilepsy. Being once introduced into the alimentary canal, they find there a nidus in which to undergo development into the tape-worm.

The tape-worm is nourished from its head, the newly-created segments pushing those already formed before them, so that the caudal extremity is the oldest portion of the animal. Each segment is flat and rectangular, and contains both a male and a female organ, the orifices of which are joined at the apex of a lateral papilla. In the *tænia solium*, the papillæ are arranged alternately at one side and the other. The size of the segments increases gradually toward the caudal extremity, the largest being three or four lines in breadth. There may be upwards of eight hundred segments, and the worm may measure above

---

\* Von Zehender, Parasitical Diseases of the Eye, Bowman Lectures, Deutsch. Med. Wochenschr., No. 50, 1887.

† Colman, quoted in Sajous' Annual, vol. i., 1890.

‡ See Manual of Animal and Vegetable Parasites, Syd. Soc. transl., 1857.

§ Origin of Intestinal Worms, *ib.*, 1857.

|| Béranger-Féraud, Leçons de Clinique sur les Taenias de l'Homme, Paris, 1888.

ten feet; it has been stated even to be above thirty. Upon the head, which is about as large as that of a pin, is a double circle


FIG. 89.



Segments of *tænia solium*. Drawn from a specimen.

of hooks contained in sacs, and around this circle are arranged four sucking-cups or mouths. The slender neck exhibits no segmentation. The sucking-disks in the *tænia mediocanellata* are larger than those in the *tænia*

FIG. 90.



Heads of *tæniæ*, magnified, except the small central figure, which represents the head and neck of *tænia solium*, natural size. The figure to the left is the *tænia solium*, that to the right the *mediocanellata*.

*solium*, but the head, which is of blackish appearance, and obtuse, has no hooks.

The form of tape-worm most frequently seen in this country is the *tænia mediocanellata*, which is usually found in beef. Leidy states, as the result of a large experience, that he has rarely encountered the pork tape-worm, *tænia solium*, as a parasite in the human intestines. The habit of eating partially-cooked beef is the cause of much of the infection with tape-worm.

*Tænia* occasions disordered digestion, colic, cramps, a feeling of

uneasiness in the abdomen, irritation of the mouth, nose, and anus, anæmia, headache, dizziness, disturbed sleep, mental depression, emaciation, cough, fainting-fits, cutaneous eruptions, and various cerebro-spinal affections, such as convulsions and epilepsy; yet there are no absolute data for the diagnosis of this parasite, except its appearance in the discharges. In order that relief be permanent, the head must be expelled.

The *bothriocephalus latus*, *tænia lata*, or *broad tape-worm*, differs from the common tape-worm in having no lateral papillæ alternately arranged, but a single one at the centre of each segment; the segments themselves are much broader, and with the breadth greatly preponderating over the length; the head is of elongated form, has no hooks upon it, and only a pair of fissures instead of the four mouths of the *tænia solium*, and we find no traces of joints until about three inches from the head. The parasite is of yellow or grayish-white color.

*Echinococci* belong also to the family of the *tæniadæ*. They may take up their abode in the substance of almost any organ in the body, and are the immature brood of a species of *tænia*. They consist of a vesicle having at one portion of its wall a head, upon which are six hooklets circularly arranged. The whole animal is surrounded by an investing membrane, which may burst and allow it to escape; the term *hydatid* designates the enveloping cyst. It forms when the *tænia* embryo has bored its way to its resting-place in the liver, or has been carried with the circulation to other organs. The *echinococcus*, unlike other larval *tæniæ*, retains a more or less globular figure, in place of exhibiting a head, neck, and body. When the *echinococci* are arrested in their normal development and are barren, not attaining to the production of scolices, they give rise to cysts with walls consisting of distinctly-developed, concentric layers, and having a peculiar gelatinous trembling,—the so-called *acephalocysts*; and the same may be said of abortive cysticeri, embryonic forms of *tænia*, which, some suppose, may also occasion the *hydatid* cysts; though others maintain that the *hydatids* proceed from only one form of *tænia*,—the *tænia echinococcus*.

The family of the *distomata*, belonging to the order of fluke-like parasites, is not at all uncommon in man.

A species of *distoma*, measuring from eight to fourteen lines in



length, called the *distoma hepaticum*, usual in the liver and gall-bladder of the sheep, has been seen in the human liver and gall-duct, and also, it is said, in abscesses of the scalp. Other species of distoma have been found in the portal vein, ureters, kidneys, and bladder, and upon the intestinal mucous membrane; yet in the portal vein and its larger branches—a common seat of the distoma—the parasite produces little or no appreciable derangement; but when in the intestine it may give rise to congestion of the membrane, extravasation of blood, and the symptoms of dysentery. This has been specially noticed of the distoma hæmatobium, or Bilharzia hæmatobia, a worm which is common in Egypt, and which has been found to be the cause of the hæmaturia prevalent at the Cape of Good Hope and at the Mauritius. The entrance into the body is mainly through the urethra in persons bathing.

*Filariae* have been met with in the urine. Lewis\* regards the hæmatozoon he has described as a filaria. The *filaria sanguinis hominis* is supposed to get into the system chiefly through the bites of mosquitoes, or by entering the skin of bathers. It gives rise to considerable pain in the loins, and leads to both bloody and chylous urine, and, according to Manson, to the elephantiasis of the tropics. Mastin† proves that the filaria in the United States may be the cause of chylocele of the tunica vaginalis testis. A worm called the *strongylus gigas* has been observed in the kidneys. It produces hæmaturia, continuous pain, and an abdominal tumor,‡ and may lead to dropsy and death.§

The *dochmius duodenalis* is a worm producing a peculiar anæmia by sucking blood from the walls of the duodenum. It has been found especially among brickmakers, miners, and men working in tunnels, and the disorder has been identified by Leichenstern|| with the so-called Egyptian chlorosis, tropical chlorosis, and brickmaker's anæmia. It has spread largely through Italian

\* Lancet, vol. ii., 1873; see also Manson, Medical Times and Gazette, 1881; Mackenzie, On the Periodicity of Filarial Migration, Lancet, 1881.

† Medical Record, Sept. 1888.

‡ Magner, Journ. de Méd. de Bordeaux, Feb. 1888.

§ George, Med. and Surg. Reporter, Aug. 1888.

|| Schmidt's Jahrbücher, Sept. 1888; also, Internationale Klinische Rundschau, Oct. 1888.

and Polish laborers employed in building tunnels, in mining, and in brickmaking. *Anchylostomiasis*, as the disease caused by the parasite is called, is characterized by marked anæmia, by digestive disorder, abdominal pains, and bleeding from the bowels. There is a greater tendency to retinal hemorrhage than in simple anæmia.\* Sandwith speaks of the marked sleepiness and dense stupidity.†

*Fly parasites* may be found in the dejections from the bowel and in the urine, producing local irritation of the intestine or the bladder.

The parasites which chiefly occupy the areolar tissues or the muscles remain to be described. Of these there are two of special importance.

One is the *filaria medinensis*, *dracunculus*, or *Guinea-worm*. This is a very slender, flat, finely-ringed worm, which introduces itself into the subcutaneous cellular tissue: here it grows rapidly, and gives rise to swelling, with more or less inflammation; and sometimes to severe constitutional disturbance. After a time the swelling points, and breaks, and the worm may be laid hold of and carefully twisted around a little piece of stick or a quill until it is extracted entire; if broken off, the eggs with which it is filled, getting into the wound, will become the agents of fresh mischief. Many of these worms may be found in the same patient, occasioning great annoyance and distress, even fatal exhaustion; but it is stated that there is often only one present. The number may vary between this and fifty. Some worms are twelve, others forty inches long, or even more. According to Busk, the parasite grows in the human areolar tissue at the rate of about an inch a week. Though it is most frequently found in the lower extremities, it has been observed to appear in the socket of the eye, in the mouth, the cheeks, the ears, and under the tongue and the scalp. It migrates rapidly from one part of the body to another. Where it exists, a pricking or an itching heat is felt; a vesicle forms when the worm is about coming to the surface, and this vesicle opens, leaving an angry-looking ulcer, in the centre of which the parasite shows itself. Phlegmonous

---

\* Discussion at the Brit. Gynæcol. Soc., Brit. Med. Journ., June, 1888.

† Trans. Eleventh Internat. Medical Congress, 1894.

spots may appear all over the body in which specimens of dracunculus are found.\* The period of incubation is from eight to twelve months: a year often elapses before the Guinea-worm makes itself manifest in the human body.† The disorder, common in Asia and in Africa, is, fortunately, one with which we are unacquainted.

*Trichina spiralis*.—This parasite was discovered by Owen in 1835 in human muscles taken from the dissecting-room; it was

FIG. 91.



*Trichina* in recent human muscle, taken the thirteenth day of illness. (After Dalton.)

subsequently found by Leidy in the animal which it most infests, the pig; but it was not looked upon as other than harmless until

---

\* Woskresensky, quoted in Sajous' Annual, vol. i., 1889.

† Aitken's Practice of Medicine, vol. i.

in 1860 Zenker proved that trichinæ may exist free in the muscles of man, that they are encapsuled only after some time, and that they are the cause of a very serious disease.


The parasite is always introduced into the body by eating ham, pork, or sausages made from the flesh of pigs containing trichinæ. It is very probable that the hogs themselves obtain them from rats, in which they are common. It has also been stated that trichinæ may exist in beef; but this is not generally admitted.

The trichina spiralis is the juvenile condition of a small nematode worm. It becomes fruitful only when introduced into the intestine. After being swallowed, the female trichina begins to throw off minute embryos, which migrate to the muscular structures.\* They pass to them through the intestinal walls, the mesentery, and the blood-vessels. When they reach the muscles they grow there, but do not generate others.

When the young trichina arrives in the muscles, it begins at once to destroy the muscular texture. It penetrates and irritates the sarcolemma, leading to its gradual thickening and to an exudation that fixes the worm to a particular spot. Thus is formed the cyst which encapsules the parasite, and which plays so important a part in its subsequent destruction. It takes a month or months for the cyst to form completely. Several trichinæ may wander in the same track, and ultimately be enclosed in the same mass of exuded matter. Two are not infrequently seen intimately coiled up, and the number may rise to five.†

After the perfect formation of the cyst, further changes take place in it; particles of calcium and magnesium carbonate are deposited. The calcareous mass extends, and gradually covers the whole parasite, while around the prolongations of the cyst fat-cells are deposited. The whole process is very destructive to the flesh-worm, and it is thus that the dis-

FIG. 92.



Trichina capsule with shell-like calcareous deposits. (After Leuckart.)

\* Leuckart, Untersuchungen über Trichina Spiralis, Leipsic, 1866.


† Thudichum, Blue Book, Seventh Report of the Medical Officer of the Privy Council, p. 367.



order is cured. But it is apt to be months before this result is accomplished. Nay, as we know from two cases recorded by Virchow, neither the encapsuling nor the calcareous transformation kills the worms of necessity at all speedily; for in the one case they had remained alive for eight, in the other for thirteen and a half years after the infection,\* and in one instance mentioned by Turner† they were alive and active after twenty-six years.

The number of trichinæ in the muscles may be from several hundreds to as many millions. Now, in accordance with their number in the muscles, with the character of the changes which there take place, and with the quantity in the intestines, will vary the extent of constitutional derangement and the signs of local irritation. Thus the symptoms and the dangers of *trichiniasis* are not always the same: we find, indeed, all the degrees of the malady. When merely a few thousand trichinæ occupy the muscles, there are chiefly muscular pains with stiffness and general debility; signs which gradually cease as the worms become fully encapsuled and cretaceous alterations occur. When the muscles are occupied by many millions of the flesh-worms, the local phenomena are much more severe; there may be almost complete immobility of the whole body, the muscles of respiration and of deglutition are implicated, irritative fever and the general cachexia are marked, and the patient is apt to perish by gradual exhaustion, or in consequence of the disordered respiratory function, or of some pulmonary complication. The presence of large numbers of trichinæ in the intestine produces diarrhœa, vomiting, abdominal pain and tenderness; or the worms may shortly after being swallowed give rise to a kind of cholera mor-

FIG. 93.



Encapsuled chalky concretions in muscle, due to dead trichinæ. Magnified about thirty times. (After Leuckart.)

\* Virchow, *op. cit.*, p. 40.

† Lancet, London, May, 1889.

bus. Should the signs of the affection not appear until from twenty-one to twenty-five days after the use of the infected meat, and take the form similar to acute rheumatism of the joints, there are not as many trichinæ present as in the choleroïd or the typhoid variety of the malady, each of which Rupprecht\* has told us shows from five to ten millions.

Speaking generally, we may recognize in trichiniasis three stages: the first, lasting about a week, during which the trichinæ are being generated in the intestines and in which we find only signs of gastro-intestinal irritation; the second, the passage of the brood into the muscular textures, and the disturbances it there occasions; the third, the retrogressive formation, which fairly sets in about three or four weeks after the beginning of the second. Now, it is this stage which yields the most striking manifestations of the malady:—loss of appetite; pasty taste in the mouth; nausea or vomiting; dry, somewhat coated tongue; diarrhœa; abdominal pain and meteorism; prostration; fever, with a quick pulse and copious sweating; œdematous swelling of the face, followed in grave cases by almost general anasarca; sensitiveness of the skin and the muscles to the touch, or painfulness when the latter are moved, and their contraction and difficult motion; dyspnœa; apathy; sleepless nights; nocturnal attacks of abdominal neuralgia; and emaciation.

Let us examine some of these phenomena more in detail:

The *fever* is a marked symptom. It sets in early, owing to the intestinal irritation, though it is not until the end of, or after, the first week, after therefore the migration of the young trichinæ has fairly begun, that it is strikingly developed. It is then, except in those cases in which fresh importations of trichinæ from the intestine in considerable numbers produce exacerbations, a continuous fever, with a pulse ranging from 100 to 130, with scanty urine and profuse perspirations having a very unpleasant odor and which may continue in certain parts of the body after the general sweating has entirely ceased. The temperature is about 101°, though it may pass to 104° and 105°; yet it does not, as a rule, reach the high heat which is observable in other continuous fevers. But it is especially in the

---

\* Vierteljahrsschrift für Ges. Med., Oct. 1880.

profuse perspirations, the absence of enlargement of the spleen and of an eruption, the swelling of the face, the muscular symptoms, and in a very red color of the visible mucous membranes, that the points of difference lie between the febrile excitement of

FIG. 94.



*Trichina spiralis*. Magnified 300 times. (After Virchow.)

trichiniasis and *typhoid fever*,—a malady which, on account of the continuous fever, the prostration, the diarrhoea, and the sudamina, it resembles. In light cases of trichiniasis there may be no fever, or there may be a fever more of intermittent or remittent character. The appearance of the face may be like that of *typhus fever*; here, however, the muscular pains are wanting.\*

The *œdema* marks the beginning of the second stage of the affection. It manifests itself first in the eyelids, about the seventh day of the disease, and is attended with a catarrhal state of the conjunctiva, with dilated pupils, great susceptibility to light, diminished power of accommodation, and pain in moving the

---

\* See Clinical Lecture on Acute Trichiniasis, reported in Medical News and Abstract, March, 1881.

eye. The swelling may extend over the whole face, and is sometimes associated with flushing. It is uninfluenced either by the sweats or by the diarrhœa, but lessens generally very much, or even disappears, after lasting eight or nine days, though it may vanish in a few days; at the same time, too, the diarrhœa is apt to diminish, or even gradually to cease. But instead of the œdema subsiding, it may extend to the chin, to the arms and legs, and to the back; or it may show itself in the extremities subsequently to the disappearance from the face, and shortly afterward become perceptible over the trunk. In some cases an anasarca condition, beginning at the ankles and extending upward, occurs during convalescence, and is of long duration. It is then probably connected with the state of the blood; whereas the œdema happening earlier in the malady is thought to be due to the pressure upon the arteries, exerted by the parasites and the exudation of plastic material they produce, or, in accordance with the observations of Thudichum, to their presence within the lymphatic spaces, vessels and glands, and blood-currents.\* The dropsical swelling of trichiniasis is not associated with albumin in the urine, for, except an increased quantity of uric acid, the urinary secretion contains no abnormal ingredient. Still, we find occasionally some albumin in the urine, as well as polyuria, though generally the quantity of urine is diminished. The trichinæ may at times be detected in the passages from the bowels. Boils, acne, and ecthyma are often noticed after the œdema has passed away.†

The *muscular symptoms* begin in the second stage, at about the tenth day, with pain and stiffness in the limbs. Soon at all parts of the body the muscles give the impression of being swollen; they are extremely painful when touched or moved; and the patient lies in consequence as quietly as possible, or, in very severe instances of the affection, like a paralyzed person. The immobility is also due partially to the retracted state of the muscles which occurs in bad cases, and which produces a condition similar to a true spasm, manifest for instance in the semi-flexed position of the extremities, and in the rigid, trismus-like setting of the jaws.

---

\* Thudichum, *loc. cit.*, pp. 362 and 386.

† Meissner, Schmidt's *Jahrbücher*, No. 4, 1868.



The disturbance of function of certain muscles becomes particularly evident. The disorder of the muscles of the eye has already been spoken of; we encounter, besides, impaired hearing, difficulty of deglutition, and loss of voice, from the muscles of the ear, of the pharynx, and of the larynx being filled with trichinæ. The respiratory muscles are commonly much affected, and we find hurried and shallow breathing, and at times considerable distress in respiration. The muscles of the heart usually, and the unstriped muscles of organic life constantly, escape infection; and, as the trichinæ wander to the front of the body rather than to the back, the muscles anteriorly are more infested than those posteriorly. A flabby condition of the muscles, with a certain want of power and painful sensation on motion, has been noticed as an early symptom and preceding their marked implication.\*

The marked muscular pain, the stiffness, the fever, the profuse sweats, the acid urine, simulate the signs of *acute rheumatism*; but we find in trichiniasis diarrhœa, no articular swelling, and no heart-complications. Error is more apt to happen with reference to acute muscular rheumatism. But the signs of prostration and of gastro-intestinal irritation are here wholly wanting.

The condition of the respiratory muscles gives rise, as already stated, to the embarrassed respiration, but it is not the only cause of the *pulmonary symptoms*. Yet, whether it alone leads to congestion of the lung and to bronchitis or pleuritis, or other causes concur in producing them, it is certain that these states are usual. They are not uncommonly combined with pneumonia, which appears suddenly, and selects the lower portion of the left lung by preference, occurs about the twenty-sixth day of the disease, and is apt to prove fatal. The sputa consist of dark unmixed blood; and the pneumonia is thought to be due to a trichinous embolism, the clots being derived from thrombi, which, forming in the venous system, are sent through the heart into the lungs.† Limited catarrhal pneumonia may be also met with.

If the patient escape a serious pulmonary complication, if he have strength enough to withstand the weeks of irritative fever and exhaustion, he enters at the end of a month or of five or six

---

\* Kratz, Die Trichinen-Krankheit im Hadersleben, Leipsic, 1867.

† Rupprecht, Trichinen-Krankheit, 1864.

weeks of suffering upon a gradual convalescence. The fever declines; the respiration is less accelerated; the perspirations are far less copious; the urine increases in quantity; the pains decrease; and by about the sixth week of the malady the patient is sufficiently free from pain to lie on his side, and is thus able to sleep. The pallor of his countenance gives way to a healthier hue; his appetite becomes insatiable; and he moves his limbs with more and more freedom. But it is a long time before he regains his full muscular power. Indeed, this may be always somewhat impaired; though we have the authority of Rupperecht for the statement that it may entirely return, and perfect health be recovered. In some cases convalescence does not set in for four months; in others it is greatly retarded by boils, by inflammation of the lymphatic glands, and by dropsy. The change in the power of accommodation of the eye may also alter but slowly. Children convalesce more quickly than adults. They suffer, in truth, less from the disease, and are not very subject to it.

The diagnosis of the malady has been made evident while discussing the symptoms. At first the signs of gastro-intestinal catarrh, the vomiting, the slight fever, the perspiration, the muscular feebleness, are the most significant, and these early manifestations might be mistaken for *irritant poisoning*; we can tell their meaning prior to the marked development of the phenomena in the muscles only by the detection of trichinæ in the stools. The same may be said of *cholera morbus*. Again, it must be borne in mind that in some cases of trichiniasis the first symptoms of the complaint do not happen for two or three weeks after the infected meat has been eaten; and that in others it runs a chronic course and the whole disease is very protracted. The so-called "*sausage poisoning*," not dependent on trichinæ, differs from trichiniasis in its rapid course and in the quick appearance of the symptoms after the spoiled sausages have been partaken of. In *periarteritis nodosa* the severe muscular pains are associated with thickening of the vessels, and an examination of the muscles will explain the cause of the muscular affection. Indeed, in any instance, no matter what be the complaint trichiniasis may simulate, there is but one means of determining the presence of the flesh-worms positively,—to examine a piece of muscle. This may be effected by cutting down upon a muscle and removing sufficient

of its structure for a microscopical examination, or by using Middeldorpf's harpoon or Duchenne's or Hart's trocar.

Owing to the œdema, and particularly the œdema of the eyelids and face, the malady may be confounded with *Bright's disease*. But the utter absence of albumin in the urine distinguishes it. The physical signs separate the dyspnœa it occasions from that of *cardiac disease*; and the sweats and the muscular symptoms of trichiniasis tell us what we are dealing with.

The chief epidemics of trichiniasis have occurred in Germany; but we have not escaped in this country. Nor can we claim that our hogs are not infested. On the contrary, the report of the Chicago Academy shows that about one in fifty contains trichinæ in the muscles.\* Our comparative immunity from the affection is due to the pork being much more generally cooked thoroughly before it is eaten; for the only prophylactic is thorough cooking, prolonged exposure to high temperature killing the trichinæ. Pickling has little if any effect. Salting and smoking are preventive means of some value, but do not insure safety.

---

\* Chicago Medical Examiner, May, 1866; quoted in Medical and Surgical Reporter, June 2, 1866; see also Billings, New York Med. Journ., 1883, xxxviii.; Mary T. Davis, Nashville Journ. Med. and Surg., 1884, N. S., xxxiii.; J. A. Close, Transact. Internat. Med. Cong., ix., Wash., 1887; E. L. Mark, Rep. Board of Health Massachusetts, 1887-88, Boston, 1889.





# INDEX.

## A.

- |                                       |               |                                      |          |
|---------------------------------------|---------------|--------------------------------------|----------|
| Abdomen, abscess in walls of,         | 616, 628      | Acarus . . . . .                     | 1009     |
| auscultation of . . . . .             | 550           | Acephalocysts . . . . .              | 1054     |
| diseases of . . . . .                 | 541           | Acetone . . . . .                    | 766, 824 |
| attended with pain . . . . .          | 596           | Acetonuria . . . . .                 | 203      |
| simulated by hysteria . . . . .       | 619           | Achillodynia . . . . .               | 897      |
| dropsy of . . . . .                   | 702, 710      | Acidity of stomach as a symptom      | 556      |
| enlargement of, general . . . . .     | 542, 710      | Acne . . . . .                       | 1011     |
| partial . . . . .                     | 718           | rosacea . . . . .                    | 1011     |
| inflammation of muscles of . . . . .  | 616           | Acromegaly . . . . .                 | 251      |
| inspection of . . . . .               | 542           | Acroparesthesia . . . . .            | 80       |
| movements of . . . . .                | 543           | Actinomycosis hominis . . . . .      | 1046     |
| palpation of . . . . .                | 543           | laryngeal . . . . .                  | 285      |
| percussion of . . . . .               | 544           | of the liver . . . . .               | 684      |
| pulsation in . . . . .                | 731           | of the skin . . . . .                | 1023     |
| retraction of parietes of . . . . .   | 542           | pulmonary . . . . .                  | 381      |
| rheumatism of walls of . . . . .      | 618           | Addison's disease . . . . .          | 869      |
| tumors of . . . . .                   | 606, 718      | confounded with acanthosis           |          |
| Abscess, hepatic . . . . .            | 682           | nigricans . . . . .                  | 871      |
| lumbar . . . . .                      | 734           | with disorders of liver . . . . .    | 871      |
| of abdominal walls . . . . .          | 616           | with discoloration of lac-           |          |
| of brain distinguished from           |               | tation and pregnancy . . . . .       | 871      |
| softening . . . . .                   | 238           | with fever-hues . . . . .            | 871      |
| distinguished from tumor . . . . .    | 241           | with hereditary hue . . . . .        | 871      |
| of kidney . . . . .                   | 816           | with diseases of suprarenal          |          |
| distinguished from cystitis . . . . . | 818           | capsules . . . . .                   | 872      |
| from pyonephrosis . . . . .           | 822           | with pernicious anæmia . . . . .     | 872      |
| of larynx . . . . .                   | 284           | with phthisis . . . . .              | 871      |
| of liver . . . . .                    | 667, 675, 682 | with pityriasis versicolor . . . . . | 871      |
| of mediastinum . . . . .              | 508           | with sun-bronzing . . . . .          | 870      |
| of thoracic walls confounded          |               | with syphilis . . . . .              | 871      |
| with chronic pleurisy . . . . .       | 422           | with vagrants' disease . . . . .     | 871      |
| perinephritic . . . . .               | 818           | Adenoid vegetations . . . . .        | 857      |
| peritoneal . . . . .                  | 730           | Adhesions, pericardial . . . . .     | 475, 517 |
| peritonsillar . . . . .               | 285           | Ægophony . . . . .                   | 325      |
| perityphlitic . . . . .               | 626           | Æsthesiometer . . . . .              | 77, 79   |
| post-cæcal . . . . .                  | 624           | Agaphia . . . . .                    | 61, 207  |
| psoas, confounded with aneu-          |               | Ague, dumb . . . . .                 | 949      |
| rism . . . . .                        | 734           | quintan . . . . .                    | 949      |
| confounded with cæcal abscess         | 628           | Ainhum . . . . .                     | 1050     |
| pulmonary . . . . .                   | 378, 379      | Akataphasia . . . . .                | 208      |
| pyæmic . . . . .                      | 687           | Albumin in the urine . . . . .       | 767      |
| retrolaryngeal . . . . .              | 284           | tests for . . . . .                  | 768-771  |
| retropharyngeal . . . . .             | 283, 537      | Albuminometer . . . . .              | 769      |
| tonsillar . . . . .                   | 285           | Albuminose . . . . .                 | 769      |
| tropical . . . . .                    | 667           | Albuminuria, cyclic . . . . .        | 795      |
| Acanthosis nigricans . . . . .        | 871           | from excessive uric acid for-        |          |
|                                       |               | mation . . . . .                     | 795      |
|                                       |               | in laryngeal diphtheria . . . . .    | 532      |
|                                       |               | of uric acid and oxaluria . . . . .  | 796      |

- Albuminuria, simple . . . . 94, 794  
 Albuminuric ulceration of bowel 646  
 Alcoholism, acute, distinguished  
     from apoplexy . . . . 201  
     from opium poisoning . . 1033  
     chronic . . . . . 128, 1039  
 Alexia . . . . . 207  
 Algesimeter . . . . . 551  
 Allochiria . . . . . 79  
 Alopecia . . . . . 1013, 1023  
 Alvine discharges . . . . . 593  
 Amaurosis . . . . . 139, 1040  
     from gastric hemorrhage . . 565  
     uremic . . . . . 801  
 Amblyopia . . . . . 61, 96  
 Amenorrhœa . . . . . 579, 856  
 Ametropia . . . . . 88  
 Amœba coli . . . . 341, 595, 650, 687  
     dysenteria . . . . . 650  
 Amphoric voice . . . . . 326  
     sound . . . . . 306  
 Amygdalitis, follicular . . . . 276  
 Anæmia . . . . . 94, 95, 855  
     as a cause of dropsy . . . . 831  
     cerebral . . . . . 188  
     in Bright's disease . . . . 800  
     essential . . . . . 857  
     from parasites . . . . . 855  
     idiopathic . . . . . 857  
     of amœbic dysentery . . . . 650  
     pernicious . . . . . 95, 294, 857  
     retinal . . . . . 95  
     spinal . . . . . 131  
     state of blood in . . . . . 851  
 Anæsthesia . . . . 74, 103, 107, 125  
     crossed . . . . . 76  
     dolorosa . . . . . 77  
     extended . . . . . 75  
     from poisoning . . . . . 74  
     from reflex action . . . . . 77  
     hysterical . . . . . 75  
     in affections of nervous centres 75  
     localized . . . . . 75, 76  
     muscular . . . . . 79  
     of spinal origin . . . . . 75  
     one-sided . . . . . 76  
     tests for . . . . . 77  
     trigeminal . . . . . 77  
 Anæsthetics in feigned aphonia . 289  
 Analgesia . . . . . 78  
 Anasarca . . . . . 830  
 Anchylostomiasis . . . . . 856, 1056  
 Anchylostomum duodenale . . . 856  
 Aneurism, abdominal . 606, 699, 732  
     physical signs and symptoms  
         of . . . . . 732  
     intracranial . . . . . 245  
     multiple, of renal artery . . 823  
     of abdominal aorta confounded  
         with aortic pulsation . . 733  
 Aneurism of abdominal aorta  
     confounded with colic, 606, 733  
     with disease of the spine . . 733  
     with lumbar and psoas ab-  
         scess . . . . . 734  
     with neuralgia . . . . . 733  
     with non-aneurismal pul-  
         sating tumors . . . . . 734  
     with rheumatism . . . . . 733  
 of ascending aorta . . . . . 506  
 of descending aorta . . . . . 516  
 of heart . . . . . 517  
 of hepatic artery . . . . . 699  
 of innominate artery . . . . . 517  
 of pulmonary artery . . . . . 518  
 phantom . . . . . 518  
 thoracic . . . . . 506  
     confounded with abscess of  
         the mediastinum . . . . 508  
     with chronic laryngitis,  
         287, 507, 514  
     with intra-thoracic morbid  
         growths . . . . . 507  
     with malformation of chest 513  
     with pulsation of pulmo-  
         nary artery . . . . . 512  
     eructation as a symptom of . 558  
     tracheal tugging a sign of . 509  
 Angina pectoris . . . . . 448  
     distinguished from cardiac epi-  
         lepsy . . . . . 451  
     from intercostal neuralgia . . 451  
     from irritability of heart . . 451  
     Ludovici . . . . . 534  
     rheumatic . . . . . 276  
     simple acute . . . . . 522  
     ulcero-membranous, dis-  
         tinguished from diphtheria 530  
 Animal parasites . . . . . 1050  
 Ankle clonus . . . . . 102, 132, 140  
 Ano, fistula in . . . . . 362  
 Anorexia . . . . . 171, 555, 582  
 Anoxæmia . . . . . 857  
 Anthracosis . . . . . 702  
 Anthrax . . . . . 1045  
 Aorta, aneurism of abdominal.  
     See *Aneurism, abdominal*.  
     aneurism of thoracic. See  
         *Aneurism, thoracic*.  
     coarctation of . . . . . 511  
     constriction of . . . . . 511  
     inflammation of . . . . . 466, 508  
     malposition of . . . . . 513  
     pulsation of . . . . . 466, 731  
     valves of . . . . . 497  
         insufficiency of . . . . 510  
 Aortitis . . . . . 466  
 Apepsia, hysterical . . . . . 556  
 Aphasia . . . . . 61, 121, 200, 206  
     amnesic . . . . . 207

- Aphasia, auditory** . . . . . 208  
     **motor** . . . . . 119, 207  
     **sensory** . . . . . 207  
     **visual** . . . . . 208  
**Aphemia** . . . . . 61, 207  
**Aphonia, feigned** . . . . . 289  
     **nervous** . . . . . 287  
     **of hysteria** . . . . . 287  
**Aphthæ** . . . . . 520, 530  
     distinguished from diphtheria 530  
**Apoplexy** . . . . . 71, 75, 194  
     attended with paralysis . . . 195  
     cerebellar . . . . . 197  
     confounded with acute soften-  
         ing of brain . . . . . 204  
     with asphyxia . . . . . 203  
     with catalepsy . . . . . 212  
     with cerebral hysteria . . . 205  
     with diabetic coma . . . . 203  
     with epilepsy . . . . . 199, 218  
     with insensibility from drink 201  
     with insensibility from nar-  
         cotics . . . . . 201, 1032  
     with meningitis . . . . . 199  
     with obstruction of the cere-  
         bral arteries . . . . . 200  
     with protracted sleep . . . 204  
     with sudden extensive paral-  
         ysis . . . . . 204  
     with sun-stroke . . . . . 210  
     with syncope . . . . . 203  
     with tumors . . . . . 199  
     with uræmic coma . . . . . 203  
     hemorrhage the cause of . . 197  
         cerebral . . . . . 198  
         seat of . . . . . 197  
     of the cord . . . . . 125  
     pulmonary . . . . . 394  
         mistaken for acute pneu-  
             monia . . . . . 394  
     serous . . . . . 197  
     spinal . . . . . 125  
     temperature in . . . . . 195  
**Appendicitis** . . . . . 622  
     bacterium coli commune in . . 622  
     chronic . . . . . 624  
     confounded with abscess of  
         liver . . . . . 629  
         with colic . . . . . 626  
         with distention of cæcum . 629  
         with dropsy of the gall-  
             bladder . . . . . 628  
         with extra-uterine preg-  
             nancy . . . . . 628  
         with invagination of bowels 627  
         with obstruction of bowels . 627  
         with pelvic hæmatocele . . 628  
         with pneumonia . . . . . 630  
         with typhoid fever . . . . 626  
         with ulceration of ileum . . 627
- Appendix cæci, diseases of** . . . 622  
     perforation of . . . . . 624  
**Appetite, loss of, as a symptom** . 555  
     perverted . . . . . 592  
**Apyrexia** . . . . . 949  
**Arcus senilis** . . . . . 90, 485  
**Areolar tissue, irritation of** . . . 833  
**Argyll-Robertson pupil** . . . . 93, 160  
**Army itch** . . . . . 1021  
**Arteries, atheromatous changes**  
     in . . . . . 838  
     cerebral, obstructions of, con-  
         founded with apoplexy . . 200  
     coagulation in . . . . . 880  
     diseases of . . . . . 836  
         aphasia in . . . . . 209  
     inflammation of coats of . 506, 836  
     pulmonary, aneurism of . . . 518  
     pulsation of . . . . . 512  
**Arterio-sclerosis** . . . . . 837  
**Arteritis** . . . . . 836, 838, 893  
**Arthritis, rheumatic** . . . . . 900  
     associated with hæmophilia . . 888  
     distinguished from locomotor  
         ataxia . . . . . 902  
         from paralysis agitans . . . 902  
     spurious . . . . . 902  
**Ascaris lumbricoides** . . . . . 1051  
     mystax . . . . . 1051  
**Ascites** . . . . . 691, 703, 710  
     confounded with cancer of peri-  
         toneum . . . . . 715  
     with chronic peritonitis . . . 714  
     with chronic tympanites . . . 716  
     with distention of the blad-  
         der . . . . . 716  
     with gravid uterus . . . . . 716  
     with ovarian dropsy . . . . 711  
     with tubercular peritonitis . 714  
**Asiatic cholera** . . . . . 32, 656, 973  
**Asphyxia distinguished from apo-  
     plexy** . . . . . 203  
     from coal and charcoal gases . 1035  
     local . . . . . 840  
**Astasia-abasia** . . . . . 87  
**Asthenia, cardiac** . . . . . 459  
**Asthma** . . . . . 265, 331  
     cardiac . . . . . 333  
     causes of . . . . . 332  
     diagnosed from croup . . . 333  
         from dyspnoea of disease of  
             the heart . . . . . 333  
         from enlarged glands . . . 333  
         from goitre . . . . . 333  
         from œdema and spasm of  
             the glottis . . . . . 333  
         from paralysis of vocal ap-  
             paratus . . . . . 333  
         from pressure of aneurismal  
             tumor . . . . . 333

- Asthma, dyspnœa in . . . . . 331  
   hay . . . . . 354  
   in Bright's disease . . . . . 334, 802  
   spasmodic . . . . . 332  
   thymic . . . . . 334  
 Astigmatism . . . . . 81, 88, 89  
 Ataxia . . . . . 159  
   Friedreich's . . . . . 163, 174  
   hereditary . . . . . 163  
   locomotor . . . . . 129, 135, 159  
     See *Locomotor Ataxia*.  
   progressive . . . . . 163  
     distinguished from dis-  
       eases of the spinal cord . 163  
 Atheromatous changes in vessels . 838  
 Athetosis . . . . . 157, 223  
 Atrophy from overuse of muscles . 152  
   idiopathic . . . . . 154  
   in joint-inflammation . . . . . 152  
   of brain . . . . . 239  
   of liver, acute yellow . . . . . 676  
   chronic . . . . . 709  
   of optic nerve . . . . . 95  
   progressive muscular . . . . . 149, 156  
   unilateral progressive, of the  
     face . . . . . 152  
 Auscultation . . . . . 310  
   immediate . . . . . 311  
   intra-thoracic . . . . . 312  
   mediate . . . . . 310  
   of abdominal viscera . . . . . 550  
   of children . . . . . 328  
   of the voice . . . . . 325
- B.**
- Bacillus coli communis . . . . . 595  
   of anthrax . . . . . 1046  
   of cholera . . . . . 657  
   of cholera morbus . . . . . 655  
   of diphtheria . . . . . 524  
   of smallpox . . . . . 991  
   of tuberculosis . . . . . 342  
   of typhoid fever . . . . . 917  
   of typhus fever . . . . . 930  
 Bacteria a cause of disease of the  
   kidneys . . . . . 793, 820  
 Barbadoes leg . . . . . 1018  
 Bedsores . . . . . 174  
 Bell's palsy . . . . . 121, 144  
 Beriberi . . . . . 153, 833  
 Bile, inspissated . . . . . 601, 676  
   in the stools . . . . . 594, 655, 677  
   in the urine . . . . . 760  
 Bile-duct, obstruction of . . . . . 676  
 Bilharzia hæmatobia . . . . . 777, 1055  
 Biliary abscesses . . . . . 685  
   acids . . . . . 664, 761  
     Oliver's test for . . . . . 762  
     Pettenkofer's test for . . . . 761  
 Biliary calculi . . . . . 601, 663, 675  
   passages, inflammation of . . . 673  
     confounded with acute  
       hepatitis . . . . . 673  
 Bilious attack . . . . . 573  
   pneumonia . . . . . 403  
   remittent fever . . . . . 974  
 Black tongue . . . . . 999  
 Bladder, distended, confounded  
   with ascites . . . . . 716  
   with peritonitis . . . . . 615  
   fistula into . . . . . 624  
   hemorrhage from . . . . . 773  
   inflammation of . . . . . 815  
     confounded with peritonitis . 615  
   neuralgia of, distinguished  
     from acute inflammation . 815  
   paralysis of . . . . . 133  
   spasm of, confounded with colic . 603  
 Blindness . . . . . 246  
 Blisters . . . . . 174  
 Blood, air in . . . . . 884  
   coagulation of . . . . . 201  
     in arteries . . . . . 880  
     in heart . . . . . 465  
   diseases of . . . . . 842  
   Filaria sanguinis hominis in . . 781  
   impoverished . . . . . 73  
   microscopic examination of, 842, 899  
   pigment in . . . . . 883, 960, 969  
   plasmodia in . . . . . 965, 969  
   spirilla of relapsing fever in . . 947  
   sweating of . . . . . 1024  
   vomiting of . . . . . 563, 579, 1028  
 Blood-casts . . . . . 774, 792  
 Blood-corpuscles . . . . . 851  
 Blood-globule-counting . . . . . 843  
 Blood-vessels, diseases of . . . . . 836  
 Bloody stools in typhoid . . . . . 917  
   sweating . . . . . 175, 1024  
 Body, position of, as a symptom . . 30  
 Bothriocephalus latus, 858, 1052, 1054  
 Bowels, hemorrhage from . . . . . 651  
   inflammation of . . . . . 606  
   intussusception of . . . . . 635  
   invagination of . . . . . 627, 635  
   lithæmic pain in . . . . . 900  
   morbid discharges from . . . . . 643  
   obstruction of . . . . . 627, 630  
     from internal strangulation . 636  
   paralysis of . . . . . 634  
   strictures in . . . . . 544  
   ulceration of . . . . . 646  
 Brachycardia . . . . . 453  
 Brain, abscess of . . . . . 238  
   in cardiac malformation . . . 491  
   metastatic . . . . . 356  
   anæmia of . . . . . 237  
   and spinal cord, table of dis-  
     orders of . . . . . 177



- Brain, atrophy of . . . . . 239  
 concussion of, causing jaundice . . . . . 665  
 congestion of . . . . . 236  
 cortical centres in . . . . . 60  
 cysts in . . . . . 245  
 diseases of . . . . . 57, 177  
   headache as a symptom of . . . . . 81  
 dropsy of . . . . . 187  
 emboli in . . . . . 881  
 geographical centre in . . . . . 61  
 hemorrhage into . . . . . 236  
 hypertrophy of . . . . . 250  
   distinguished from enlarge-  
   ment of the head . . . . . 250  
 inflammation of . . . . . 237  
   confounded with pericarditis . . . . . 473  
 lesions of gray central ganglia . . . . . 118  
 meningitis of base of . . . . . 180  
 sensory centres in . . . . . 63  
 softening of . . . . . 75, 235, 241  
   acute . . . . . 181, 204  
   chronic . . . . . 235  
 syphilis of . . . . . 142, 953  
 thrombosis of sinuses of . . . . . 242, 878  
 tumor of . . . . . 240  
   distinguished from abscess . . . . . 241  
   from chronic meningitis . . . . . 242  
   from softening . . . . . 241  
   gliomatous . . . . . 245  
   intracranial . . . . . 182  
   seat of . . . . . 243  
 Brain-power, exhaustion of . . . . . 239  
 Breathing. See *Respiration*.  
 Breath-sound, metamorphosing . . . . . 320  
 Bright's disease, acute . . . . . 34, 241, 791  
   distinguished from acute ne-  
   phritis . . . . . 794  
   from coma . . . . . 796  
   from dropsy . . . . . 796  
   from epileptiform convul-  
   sions . . . . . 796  
   from hæmaturia . . . . . 794  
   from pericarditis . . . . . 796  
   from pleurisy . . . . . 796  
   from pulmonary œdema . . . . . 796  
   from purulent urine . . . . . 794  
   from simple albuminuria . . . . . 794  
   from suppurative nephritis . . . . . 794  
 chronic . . . . . 798  
   confounded with anæmia . . . . . 800  
   with asthma . . . . . 802  
   with cancer of kidney . . . . . 803  
   with cardiac dropsy . . . . . 802  
   with chronic bronchitis . . . . . 802  
   with chronic consecutive  
   nephritis . . . . . 805  
   with chronic rheumatism . . . . . 802  
   with cysts of kidney . . . . . 805  
   with gastro-intestinal dis-  
   orders . . . . . 803  
 Bright's disease, chronic, con-  
   founded with neuralgia . . . . . 801  
   with renal inadequacy . . . . . 806  
   with tubercle . . . . . 804  
 contracting form of . . . . . 808  
 nervous symptoms in . . . . . 801  
 prealbuminuric stage of . . . . . 772  
 retinitis in . . . . . 95, 801  
 table of clinical differences in . . . . . 812  
 Bronchial dilatation . . . . . 379, 381  
 glands, tuberculization of . . . . . 337  
 phthisis, distinguished from  
   hooping-cough . . . . . 337  
 Bronchiolitis exudativa . . . . . 355  
 Bronchitis, acute . . . . . 348, 396, 983  
   associated with measles . . . . . 983  
   diagnosed from capillary  
   bronchitis . . . . . 351  
   from hooping-cough . . . . . 337  
   from pneumonia . . . . . 350  
   from tuberculosis . . . . . 350, 385  
   of large and middle-sized  
   tubes . . . . . 348  
   physical signs of . . . . . 349  
   sputa in . . . . . 349  
 capillary . . . . . 285, 348, 350, 359  
   confounded with acute lobar  
   pneumonia . . . . . 351  
   with acute miliary tuber-  
   culosis . . . . . 385  
   with broncho-pneumonia . . . . . 351  
   with catarrhal pneumonia . . . . . 351  
   with phthisis . . . . . 351  
 chronic . . . . . 352  
   confounded with Bright's dis-  
   ease . . . . . 802  
   with nasal catarrh . . . . . 353  
   with phthisis . . . . . 369  
 idiopathic, distinguished from  
   typhoid fever . . . . . 928  
 of the finer tubes . . . . . 348, 385  
 plastic . . . . . 341, 354  
   sputa in . . . . . 341  
   putrid . . . . . 339, 355, 381  
 Bronchophony . . . . . 325  
 Broncho-pneumonia . . . . . 351, 386  
   distinguished from tuberculosis . . . . . 386  
   mistaken for collapse . . . . . 360  
   tuberculous inspiration . . . . . 398  
 Bronchorrhœa . . . . . 353  
 Bruit de moulins . . . . . 476  
 Bulbar paralysis . . . . . 147, 152  
 Bulimia . . . . . 457, 556  
 Bulk of body . . . . . 32  
 Bullous diseases . . . . . 174, 1010
- C.**
- Cachexia strumipriva . . . . . 863  
 Cæcum, appendix of, diseases of . . . . . 625

- Cæcum, cancer of . . . . . 629  
   distention of . . . . . 629  
   inflammation of . . . . . 625  
   solitary ulcer of . . . . . 646  
 Calcium oxalate . . . . . 757, 786  
 Calculi, biliary . . . . . 601, 663, 675  
   renal . . . . . 56, 590, 602, 786  
     irritation of . . . . . 821  
   of the pancreas . . . . . 722  
 Cancer of brain . . . . . 245  
   of cæcum . . . . . 629  
     melanotic . . . . . 804  
   of cardiac orifice . . . . . 586  
   of gall-bladder . . . . . 695  
   of intestine . . . . . 731  
   of kidney confounded with  
     Bright's disease . . . . . 803  
   of larynx . . . . . 294  
   of liver . . . . . 690  
     confounded with acute con-  
       gestion . . . . . 693  
     with acute hepatitis . . . . . 693  
     with cancer of omentum . . . . . 696  
     with cancer of stomach . . . . . 696  
     with catarrhal jaundice . . . . . 693  
     with chronic congestion . . . . . 692  
     with disease of gall-blad-  
       der . . . . . 694  
     with enlarged kidney . . . . . 697  
     with fatty liver . . . . . 692  
     with syphilitic liver . . . . . 694  
     with waxy liver . . . . . 692  
   of lungs . . . . . 375, 424  
     confounded with chronic  
       pleurisy . . . . . 424  
     with phthisis . . . . . 375  
   of lymphatic glands . . . . . 867  
   of lymphatic glands lying by  
     side of vertebræ . . . . . 727  
   of omentum . . . . . 590, 696, 715  
     confounded with cancer of  
       liver . . . . . 696  
   of pancreas . . . . . 721  
   of peritoneum . . . . . 715, 730  
   of retro-peritoneal glands . . . . . 715  
   of pleura . . . . . 420  
   of stomach . . . . . 582, 696  
     confounded with cancer of  
       liver . . . . . 696  
       with chronic gastritis, 582, 585  
       with cirrhosis of liver . . . . . 708  
       with gastric ulcer . . . . . 582, 585  
       detection of tumor in . . . . . 586  
       situation of . . . . . 586  
   of the pylorus . . . . . 586  
   of the tongue . . . . . 521  
   of the tonsils . . . . . 523, 536  
 Cancrum oris . . . . . 520  
 Capillaries, diseases of . . . . . 840  
 Capillary pulsation . . . . . 43
- Carcinoma, gastric, test for . . . . . 584  
 Cardiac asthenia . . . . . 459  
   epilepsy . . . . . 451  
   nerve storms . . . . . 453  
 Cardioscope . . . . . 433  
 Cartilage, aortic . . . . . 436  
 Catalepsy accompanying hys-  
   teria . . . . . 212  
   associated with hypnotism . . . . . 213  
   confounded with apoplexy . . . . . 212  
     with ecstasy . . . . . 212  
   daymare form of . . . . . 213  
   feigned . . . . . 213  
 Cataract . . . . . 90, 824  
 Catarrh, gastric . . . . . 573, 575, 1028  
   in measles . . . . . 982  
   intestinal . . . . . 606  
   nasal . . . . . 262, 353  
   suffocative . . . . . 348  
   vesical . . . . . 816  
 Catarrhal fever . . . . . 909  
   distinguished from dengue . . . . . 996  
   from hay-fever . . . . . 912  
   lung complications in . . . . . 910  
   sequelæ of . . . . . 910  
 Cavernous voice . . . . . 325  
 Cavity in lungs . . . . . 367, 377, 379  
 Cell, granular, of ovarian fluid . . . . . 713  
 Cellulitis, pelvic . . . . . 728  
 Cerebellum, diseases of . . . . . 165, 243  
   gait in . . . . . 165, 681  
   tumor of . . . . . 681  
 Cerebral affections, table of . . . . . 177  
   pain in, distinguished from  
     hemisrania . . . . . 256  
   localization . . . . . 57  
   symptoms in congestive fever . . . . . 964  
   thermometry . . . . . 123  
   tumors . . . . . 243  
 Cerebritis . . . . . 237  
   confounded with meningitis . . . . . 181  
 Cerebro-spinal disorders, 86, 88,  
   163, 189  
   fever . . . . . 937  
     causes of . . . . . 940  
     confounded with congestive  
       fever . . . . . 941  
       with inflammation of cord . . . . . 941  
       with malignant measles . . . . . 942  
       with pneumonia . . . . . 942  
       with rheumatism of cer-  
         vical muscles . . . . . 943  
       with scarlatina . . . . . 942  
       with sporadic cerebro-  
         spinal meningitis . . . . . 942  
       with tetanus . . . . . 941  
       with tubercular meningitis . . . . . 942  
       with typhoid fever . . . . . 940  
       with typhus fever . . . . . 943  
       with uræmia . . . . . 943

- Cerebro-spinal fever, pulse in . . . 938  
state of blood in . . . 940  
swelling of whole body fol-  
lowing . . . 944  
urticaria in . . . 1005  
meningitis . . . 189, 937, 942  
typhus . . . 937  
Cestoidea . . . 1051  
Charbon . . . 1045  
Charcot's disease . . . 167  
Charcot-Leyden crystals . . . 332, 340  
Chest, alterations of form, size,  
etc., of, in disease . . . 356  
contusions of, followed by  
pneumonia . . . 401  
dilatation of, diseases present-  
ing . . . 410  
inspection of, in diagnosis . . . 299  
malformation of . . . 513  
mapping out of, for physical  
diagnosis . . . 298  
mensuration of . . . 300  
motions of, in diseases of . . . 299  
retraction of, diseases attended  
with . . . 423  
tumor in . . . 419  
Cheyne-Stokes respiration . . . 335, 449  
Chicken-pox . . . 993  
See *Varicella*.  
Chilblains . . . 841  
Childbed fever . . . 612  
Chloasma . . . 1015, 1023  
Chloral poisoning . . . 1033, 1039  
ataxic symptoms following . . . 1039  
Chlorides in the urine . . . 755  
Chlorosis . . . 34, 43, 856  
confounded with pernicious  
anæmia . . . 863  
Choked disk . . . 95, 241, 244  
Cholera . . . 656, 1028  
Asiatic, distinguished from yel-  
low fever . . . 973  
associated with uræmia . . . 660  
comma-bacillus of . . . 657  
infantum . . . 654  
morbus . . . 655, 660  
distinguished from irritant  
poisoning . . . 656, 660, 1028  
nostras . . . 655  
reaction . . . 657  
subnormal temperature in . . . 659  
with typhoid symptoms . . . 660  
Cholérine . . . 659  
Chorditis tuberosa . . . 291  
Chorea . . . 219  
attended with salaam convul-  
sions . . . 224  
caused by eye-strain . . . 88, 221  
distinguished from athetosis . . . 223  
from cerebro-spinal sclerosis . . . 222  
Chorea distinguished from con-  
vulsive tremor . . . 222  
from epilepsy . . . 222  
from facial spasm . . . 223  
from hysteria . . . 224  
from paralysis agitans . . . 222  
from spasms of acute cerebral  
disease . . . 222  
from tetanus . . . 222  
from writer's cramp . . . 223  
electrical . . . 221  
following rheumatic fever . . . 220  
habit . . . 221  
Huntington's . . . 221  
hysterical . . . 221  
paralytic . . . 221  
post-hemiplegic . . . 157, 223  
post-paralytic . . . 120  
relations of, to rheumatism . . . 220  
Choroid coat, inflammation of . . . 96  
tubercles of . . . 96  
Chromocytometer . . . 852  
Chylous urine . . . 781  
Circulation, derangements of, in  
cardiac disease . . . 446, 460  
paralysis from interference  
with . . . 105  
portal, disturbance of . . . 661  
Cirrhosis of liver . . . 702  
confounded with cancer of  
stomach . . . 708  
with chronic peritonitis . . . 708  
with inflammation of por-  
tal vein . . . 708  
distinguished from cancer of  
liver . . . 707  
from hydatids . . . 706  
from malarial infection . . . 706  
from red atrophy . . . 707  
from simple induration . . . 707  
hypertrophic . . . 702, 705  
of children, syphilitic . . . 705  
of lung confounded with  
chronic pleurisy . . . 425  
Clergyman's sore throat . . . 535  
Clots, fibrinous, in the heart . . . 464  
Coffee-ground vomit . . . 564, 583  
Colic as a symptom . . . 596  
bilious . . . 597, 626  
confounded with abdominal  
aneurism . . . 606  
with abdominal neuralgia . . . 605  
with abdominal tumors . . . 606  
with appendicitis . . . 626  
with enteritis . . . 606  
with gall-stones . . . 601  
with gastralgia . . . 600, 605  
with hepatic neuralgia . . . 601  
with hernia . . . 600  
with nephralgia . . . 602

- Colic confounded with neuralgia  
of dorsal and lumbar  
nerves . . . . . 605  
with perforation of the intes-  
tine . . . . . 600  
with peritonitis . . . . . 606,  
620  
with spasm of the bladder . . 603  
with spinal disease . . . . . 606  
with uterine colic . . . . . 604  
copper . . . . . 598  
flatulent . . . . . 597  
from disease of the bowel . . 599  
lead . . . . . 598  
malarial . . . . . 598  
metallic . . . . . 598  
nervous . . . . . 598  
renal . . . . . 626, 786, 822  
simple . . . . . 596  
spasmodic . . . . . 596  
uterine . . . . . 604  
Colitis, croupous . . . . . 390  
ulcerative . . . . . 646  
Collapse, delirium in . . . . . 69  
in acute poisoning . . . . . 1028  
in appendicitis . . . . . 625  
in cholera . . . . . 656, 659  
in relapsing fever . . . . . 946  
in yellow fever . . . . . 971  
of the lung . . . . . 359  
confounded with chronic  
pleurisy . . . . . 426  
Colon, dilatation of . . . . . 549, 638, 730  
disease of, associated with  
heart disease . . . . . 647  
inflammation of . . . . . 608  
percussion of . . . . . 549  
solitary ulcer of . . . . . 646  
Color-blindness . . . . . 76, 96  
Coma . . . . . 67, 70, 704  
diabetic . . . . . 71, 203  
from narcotic poisoning . . 71, 202  
in typhoid . . . . . 920  
occurring in Bright's disease, 71, 796  
uræmic . . . . . 71, 203, 797  
Coma-vigil . . . . . 931  
Comedo . . . . . 1024  
Comma-bacillus of Koch . . . . 657  
Concretion, intra-hepatic . . . . 952  
Congestion, pulmonary . . . . . 394  
of brain discriminated from  
softening . . . . . 236  
of features, as a symptom . . 33  
passive . . . . . 564  
Congestive fever . . . . . 963  
algid . . . . . 964  
cerebral . . . . . 964  
confounded with cerebro-spinal  
fever . . . . . 941  
gastro-enteric . . . . . 963  
thoracic . . . . . 964  
Conjunctiva, tuberculosis of . . . 90  
Conjunctivitis, from lithæmia . . 900  
Consciousness, diseases marked  
by sudden loss of . . . . . 194  
Constipation as a symptom . . . . 630  
from mechanical changes . . . 642  
habitual . . . . . 641  
ulcers from . . . . . 643  
Consumption. See *Phthisis*.  
galloping . . . . . 384, 388  
Contractility, electro muscular,  
109, 168  
Contracture . . . . . 174  
Convulsions . . . . . 171  
See also *Spasms*.  
diseases marked by . . . . . 214  
distinguished from epilepsy . . 218  
epileptic . . . . . 141, 151, 255  
from irritant poisoning . . . . 1028  
from purulent otitis . . . . . 217  
hysterical . . . . . 138, 151  
in Bright's disease . . . . . 796  
in scarlet fever . . . . . 978  
in typhoid . . . . . 920  
salaam . . . . . 224  
uræmic . . . . . 798  
Convulsive tic . . . . . 223  
Cord. See *Spinal Cord*.  
Coryza . . . . . 261  
Cough . . . . . 335  
from nasal affections . . . . . 353  
hooping . . . . . 285, 336  
in laryngeal affections . . . . 267, 336  
Countenance, expression of, as a  
symptom . . . . . 32  
Crackling in tubercle of lungs . . 323  
Cramp caused by various occupa-  
tions . . . . . 223  
of stomach . . . . . 566  
writer's . . . . . 223  
Cranial reflexes . . . . . 100  
Craniotabes . . . . . 905  
Crepitation . . . . . 322, 396  
Croup . . . . . 275, 278  
catarrhal . . . . . 278  
diphtheritic . . . . . 278  
diseases confounded with . . 283, 333  
false . . . . . 278, 282  
membranous, or true . . . . . 280  
distinguished from abscess of  
larynx . . . . . 284  
from acute laryngitis . . . . . 283  
from diphtheria . . . . . 285, 531  
from false croup . . . . . 282  
from œdema of the larynx . . 283  
from retrolaryngeal ab-  
scesses . . . . . 284  
from retropharyngeal ab-  
scesses . . . . . 283  
from scarlet fever . . . . . 980



- Croup, membranous, or true, distinguished from secondary laryngitis of the exanthemata . . . . . 283  
 non-diphtheritic membranous . . . . . 285  
 spasm of glottis in . . . . . 279  
*Crus cerebri*, lesions of . . . . . 117  
*Crusta lactea* . . . . . 1007  
 Crystals, Charcot-Leyden . . . . . 332, 340  
 Cyanosis . . . . . 34, 392  
 Cyrtometer . . . . . 300  
*Cysticercus cellulosæ* . . . . . 1052  
 Cystine . . . . . 786  
 Cystitis, acute . . . . . 815  
     confounded with abscess of  
     kidney . . . . . 818  
     with acute nephritis . . . . . 815  
     with metritis . . . . . 816  
     with neuralgia of bladder . . . . . 815  
     with peritonitis . . . . . 615  
     chronic . . . . . 816  
 Cysts of kidneys . . . . . 700, 805, 823  
     confounded with hydronephrosis . . . . . 823  
     of nose . . . . . 264  
     of vocal cords . . . . . 294  
     ovarian . . . . . 711
- D.**
- Day-blindness . . . . . 96  
 Daymare . . . . . 213  
 Dead fingers . . . . . 840  
 Deafness . . . . . 76, 98, 528  
 Debility confounded with typhoid fever . . . . . 924  
 Deep reflexes . . . . . 100  
 Delirium . . . . . 67, 179  
     active . . . . . 68  
     confounded with delirium tremens . . . . . 193  
     fierce . . . . . 68  
     hysterical . . . . . 70  
     in chorea . . . . . 220  
     in typhoid . . . . . 920  
     mistaken for insanity . . . . . 69  
     of cerebral rheumatism . . . . . 893  
     of inanition . . . . . 69  
     of pneumonia . . . . . 68, 388  
     prominent as a symptom, acute affections with . . . . . 179  
     simulated . . . . . 70  
     tremens . . . . . 72, 190, 388  
     confounded with acute mania . . . . . 193  
     with acute meningitis . . . . . 191  
     uræmic . . . . . 68, 798  
 Dengue . . . . . 994  
     distinguished from influenza . . . . . 996  
     from malarial fever . . . . . 996  
     from rheumatism or gout . . . . . 996  
 Dengue distinguished from scarlet fever . . . . . 996  
     from yellow fever . . . . . 996  
 Dermatalgia . . . . . 1025  
 Dermatitis exfoliative . . . . . 1013  
     herpetiformis . . . . . 1009, 1012  
     medicamentosa . . . . . 1015  
 Dermatophytes. See *Tinea*.  
 Diabetes . . . . . 94, 823  
     distinguished from chronic polyuria . . . . . 827  
     fatty diarrhœa in . . . . . 653  
 Diabetes insipidus . . . . . 826  
     intermitting . . . . . 826  
     phosphatic . . . . . 753  
     with coexisting albuminuria . . . . . 826  
 Diacetic acid . . . . . 766  
 Diagnosis by exclusion . . . . . 23  
     differential . . . . . 22  
     methods of arriving at . . . . . 21  
     physical . . . . . 297  
     sources of error in . . . . . 23  
 Diaphragm . . . . . 334  
     fatty degeneration of . . . . . 334  
     hernia of . . . . . 414  
     inflammation of . . . . . 334  
     paralysis of . . . . . 334  
     rheumatism of . . . . . 334  
 Diarrhœa . . . . . 643  
     acute . . . . . 644  
     bilious . . . . . 644  
     choleraic . . . . . 657  
     chronic . . . . . 82, 644  
     fatty . . . . . 653  
     in pulmonary consumption . . . . . 362  
     intermittent . . . . . 647  
     in typhoid fever . . . . . 915  
     membranous . . . . . 647  
     of dissecting-room . . . . . 1048  
     of soldiers . . . . . 645  
     strumous, of children . . . . . 647  
     tubercular . . . . . 646  
 Diazo-reaction . . . . . 771  
 Digestion, disorders of . . . . . 661  
 Dilatation, bronchial, confounded with phthisis . . . . . 377  
     with pulmonary abscess . . . . . 378  
     with pulmonary gangrene . . . . . 380  
     of heart . . . . . 482  
     confounded with fatty degeneration . . . . . 484  
     with pericardial effusion . . . . . 488  
 Diphtheria . . . . . 524  
     catarrhal . . . . . 529  
     confounded with aphthæ . . . . . 530  
     with croup . . . . . 531  
     with erysipelas of the fauces . . . . . 530  
     with gangrene of the mouth . . . . . 529  
     with pharyngitis and tonsillitis . . . . . 529

- Diphtheria confounded with scar-  
   latina . . . . . 532  
   with thrush . . . . . 530  
   with ulcerative stomatitis . . 529  
   with ulcero-membranous an-  
   gina . . . . . 530  
 croupous . . . . . 526  
 faucial . . . . . 262  
 intercurrent . . . . . 533  
 laryngeal . . . . . 285, 532  
   confounded with scarlet  
   fever . . . . . 980  
   nasal . . . . . 262, 533  
   paralysis in . . . . . 141, 164, 528  
   sequelæ of . . . . . 528  
 Diphtheria-bacilli . . . . . 524  
 Diplococcus exanthematicus . . 930  
   in mumps . . . . . 534  
   pneumoniæ . . . . . 398, 940  
 Discharges, alvine, as a symptom 593  
 Displacements of heart . . . 410, 505  
 Distoma hæmatobium . . . . . 1055  
   hepaticum . . . . . 1055  
 Dittrich's plugs . . . . . 855  
 Diuresis, chronic . . . . . 826  
   in hysterical women . . . . . 827  
 Doehmius duodenalis . . . . . 1055  
 Dracunculus . . . . . 1056  
 Drink, insensibility from . . . 201  
 Dropsy . . . . . 830  
   abdominal . . . . . 702, 710  
   acute or active . . . . . 835  
   cardiac . . . . . 446, 802, 832  
   causes of . . . . . 832  
   chronic . . . . . 835  
   dependent upon a tumor . . . 831  
   diseases marked by . . . . . 791  
   from anæmia . . . . . 831  
   from malarial poisoning . . . 831  
   from scarlet fever . . . . . 979  
   general . . . . . 832  
   from irritation of areolar tis-  
   sue . . . . . 833  
   from peripheral multiple  
   neuritis . . . . . 833  
   hepatic . . . . . 833  
   in beriberi . . . . . 833  
   in Bright's disease . . . . . 796  
   internal . . . . . 831  
   of brain . . . . . 187, 249  
   ovarian . . . . . 711  
   pericardial, confounded with  
   cardiac dilatation . . . . . 488  
   renal . . . . . 832  
 Dühring's disease . . . . . 1012  
 Duodenum, catarrh of . . . . . 608  
   ulcer of . . . . . 581  
 Dysentery . . . . . 648  
   acute . . . . . 648  
   amœbic . . . . . 650  
 Dysentery, catarrhal . . . . . 650  
   chronic . . . . . 651  
   confounded with piles . . . . 649  
   with proctitis . . . . . 649  
   diphtheritic . . . . . 651  
   distinguished from diarrhœa . . 650  
   epidemic . . . . . 649  
   tropical . . . . . 595  
 Dyspepsia as a symptom, 570, 575, 641  
   atonic . . . . . 570  
   nervous . . . . . 570  
 Dysphagia . . . . . 284, 540  
 Dyspnoea . . . . . 30, 284, 288, 330, 515  
   caused by aneurismal tumor,  
   . . . . . 333, 515  
   by goitre . . . . . 333  
   diseases presenting . . . . . 410  
   from disease of the dia-  
   phragm . . . . . 334  
   from enlarged glands of neck . 333  
   in asthma . . . . . 331  
   in plastic bronchitis . . . . . 354
- E.**
- Ear, disease of . . . . . 217  
   causing abscess of brain . . . 238  
 Ecchymoses . . . . . 665  
 Echinococci . . . . . 697, 700, 805, 1054  
 Ecstasy . . . . . 212  
   distinguished from catalepsy . . 212  
 Ecthyma . . . . . 1012  
 Eczema . . . . . 175, 1007, 1020  
   distinguished from pityriasis  
   rubra . . . . . 1008  
   from scabies . . . . . 1009  
   from seborrhœa . . . . . 1008  
   impetiginodes . . . . . 1007  
   papular . . . . . 1006  
   rubrum . . . . . 1008  
   squamosum . . . . . 1008, 1014  
 Effusions, pericardial . . . 422, 469, 488  
   diagnostic sign of . . . . . 470  
   peritoneal . . . . . 610  
   pleural . . . . . 374, 405, 411, 700  
 Electricity in paralysis . . . . 108  
   faradaic . . . . . 110  
   galvanic . . . . . 110  
   static . . . . . 111  
 Electro-muscular sensibility, 112, 115  
 Elephantiasis of the Arabs . . . 1018  
   of the Greeks . . . . . 1017  
 Emaciation as a symptom . . . . 32  
 Embarras gastrique . . . . . 573  
 Embolism . . . . . 465, 877, 880  
   cerebral . . . . . 201  
   fat . . . . . 883  
   from accumulations of pigment  
   in the blood . . . . . 883  
   from acute endarteritis . . . . 884

- Embolism of arteries of the ex-  
 tremities . . . . . 882  
 of cerebral arteries . . . . . 881  
 of pulmonary artery . . . . . 880  
 of renal artery . . . . . 881  
 of vessels of liver . . . . . 881  
 splenic . . . . . 881  
 trichinous . . . . . 1063  
 Embryo-cardia . . . . . 453  
 Emphysema . . . . . 332, 356, 411, 435  
 chest sounds in . . . . . 413  
 coexisting with tubercle . . . . . 369  
 compensatory . . . . . 358  
 confounded with aneurismal  
 tumor . . . . . 357  
 with chronic pleurisy . . . . . 419  
 diagnosed from pleuritic  
 effusions . . . . . 358  
 from pneumothorax . . . . . 358  
 interlobular . . . . . 358  
 Emprosthotonos . . . . . 229  
 Empyema, pulsating, confounded  
 with aneurism . . . . . 511  
 Encephalitis . . . . . 235  
 acute hemorrhagic . . . . . 238  
 Endarteritis . . . . . 836, 884  
 gouty . . . . . 899  
 Endocardial murmurs . . . . . 439, 461  
 Endocarditis, acute . . . . . 461  
 confounded with pericarditis . . . . . 471  
 complications of . . . . . 464  
 diabetic . . . . . 824  
 puerperal . . . . . 466  
 rheumatic . . . . . 462  
 ulcerative . . . . . 466  
 associated with pneumonia . . . . . 399  
 with pyemia . . . . . 875  
 head-symptoms of, con-  
 founded with acute men-  
 ingitis . . . . . 185  
 confounded with typhoid  
 fever . . . . . 467, 927  
 Engorgements, pulmonary, in  
 fevers, mistaken for acute  
 pneumonia . . . . . 393  
 Enteralgia . . . . . 600  
 Enteritis confounded with colic . . . . . 606  
 with peritonitis . . . . . 614  
 with typhoid fever . . . . . 926  
 croupous or diphtheritic . . . . . 609  
 membranous . . . . . 647  
 muco- . . . . . 606, 608, 644  
 Enteroptosis . . . . . 592, 856  
 Ephelides . . . . . 1015  
 Epidemic meningitis . . . . . 937  
 Epigastrium, tumors of . . . . . 721  
 Epiglottis, disease of . . . . . 279, 290  
 Epilepsy . . . . . 67, 214  
 abortive . . . . . 215, 218  
 associated with vertigo . . . . . 215  
 Epilepsy, aura preceding . . . . . 214  
 cardiac . . . . . 451  
 consecutive to scarlet fever . . . . . 979  
 distinguished from apoplexy . . . . . 218  
 from chorea . . . . . 222  
 from convulsions . . . . . 217  
 from hysteria . . . . . 218  
 from strychnine poisoning . . . . . 1037  
 eccentric . . . . . 216  
 feigned . . . . . 218  
 idiopathic . . . . . 217  
 Jacksonian . . . . . 217  
 masked . . . . . 217  
 nocturnal . . . . . 215  
 post-hemiplegic . . . . . 215  
 sequelæ of . . . . . 215  
 syphilitic . . . . . 217  
 tested by administration of  
 ether . . . . . 219  
 Epiphytes . . . . . 1020  
 Episcleritis, from lithæmia . . . . . 900  
 Epistaxis . . . . . 345, 457, 533, 565, 921  
 Epithelioma . . . . . 539  
 Ergotism . . . . . 1040  
 Eructation as a symptom . . . . . 558  
 Eruption, herpetic . . . . . 132  
 in measles . . . . . 982  
 in typhoid . . . . . 921  
 Erysipelas . . . . . 90, 262, 530, 996  
 associated with pharyngeal  
 fever . . . . . 999  
 distinguished from angio-neu-  
 rotic œdema . . . . . 998  
 from confluent smallpox . . . . . 998  
 from erythema . . . . . 998  
 from mumps . . . . . 999  
 from scarlatina . . . . . 998  
 from scleroderma . . . . . 1019  
 in aortitis . . . . . 466  
 migrans . . . . . 999  
 of the fauces confounded with  
 diphtheria . . . . . 530  
 Erythema . . . . . 175, 1003  
 congestivum . . . . . 1004  
 desquamative . . . . . 1003  
 distinguished from erysipelas . . . . . 998  
 intertrigo . . . . . 1004  
 marginatum . . . . . 464  
 multiforme . . . . . 1003  
 neurotic vesicular . . . . . 1010  
 nodosum . . . . . 1003  
 solare . . . . . 1004  
 Erythromelalgia . . . . . 841  
 Examination of patients, methods  
 of . . . . . 27  
 analytical . . . . . 27  
 by anamnesis . . . . . 27  
 synthetical . . . . . 27  
 Exanthematous fevers . . . . . 976, 998  
 Excitation of muscles . . . . . 109

- Excitation of muscles, direct and indirect . . . . . 109
- Exhalations, poisonous . . . . . 1047
- Exophthalmic goitre . . . 90, 456, 513
- Eye, abnormal changes in fundus of . . . . . 94
- abnormalities of pupils of, 89, 92, 219
- appearance of, in disease . . . . . 88
- conjugate lateral deviation of . . . 91
- spasm of . . . . . 116
- derangements of mechanism of, 88, 89
- disorders of, from lithæmia . . . 900
- embolism of . . . . . 95
- hyperæmia of . . . . . 94
- paralysis of accommodation of, 93, 96
- paresis of . . . . . 96
- ptosis of . . . . . 92
- reflex neuroses of . . . . . 96
- refraction, errors of . . . . . 900
- sixth nerve of, affections of . . . 92
- strain of . . . . . 88, 96, 900
- subjective visual derangements of . . . . . 96
- third nerve of, affections of . . . 92
- Eyeball, protrusion of . . . . . 456
- Eyelids, drooping of . . . . . 162
- Eye-strain . . . . . 88, 96, 900
- as a cause of chorea . . . 88, 89, 220
- of epilepsy . . . . . 89
- of gastric derangements . . . 89
- of hysteria . . . . . 89
- of melancholia . . . . . 89
- F.**
- Face, moon-shaped . . . . . 251
- spasm of . . . . . 223, 255
- Facial hemiatrophy . . . . . 152
- palsy . . . . . 117, 200
- Fæcal discharges . . . . . 593
- vomiting . . . . . 562
- Fæces, accumulation of, 638, 720, 730
- impacted, simulating gall-stones . . . . . 602
- Faradaic excitability . . . . . 110
- Farcy, acute, confounded with pyæmia . . . . . 874
- Fat in intestinal discharges . . . 653
- in urine . . . . . 653, 780
- Fatty degeneration of heart . . . 484
- confounded with chills . . . . . 486
- with dilatation . . . . . 485
- of pancreas . . . . . 721
- of tissues, in poisoning . . . 1030
- Fauces, diseases of . . . . . 522
- erysipelas of . . . . . 530
- inflammation of . . . . . 522
- ulcers of, syphilitic . . . . . 536
- Favus . . . . . 1022
- Feigned aphonia . . . . . 289
- delirium . . . . . 70
- epilepsy . . . . . 218
- hysteria . . . . . 226
- rheumatism . . . . . 897
- sciatica . . . . . 259
- Fever, bilious remittent . . . . . 974
- bilious typhoid . . . . . 946
- breakbone, or dengue . . . . . 995
- catarrhal . . . . . 909
- cerebro-spinal . . . . . 937
- Chickahominy . . . . . 968
- congestive . . . . . 963
- enteric . . . . . 926
- erysipelatos . . . . . 999
- gastro-enteric . . . . . 963
- hectic . . . . . 951
- hemorrhagic malarial . . . . . 965
- hepatic . . . . . 675, 952
- icterode pernicious . . . . . 965
- infantile remittent . . . . . 961
- intermittent . . . . . 948
- malarial-typhoid . . . . . 969
- miasmatic . . . . . 948
- miliary . . . . . 984
- nervous . . . . . 919
- of erysipelas . . . . . 996
- pharyngeal . . . . . 999
- puerperal malarial . . . . . 953
- relapsing . . . . . 944
- remittent . . . . . 954
- scarlet . . . . . 976
- simple continued . . . . . 908
- spotted . . . . . 937
- syphilitic . . . . . 953
- typhoid . . . . . 913
- typho-malarial . . . . . 957, 968
- typhus . . . . . 929
- urethral . . . . . 952
- yellow . . . . . 969
- Fevers . . . . . 906
- classification of . . . . . 907
- continued . . . . . 907
- head-symptoms of, con-founded with meningitis . . 182
- eruptive . . . . . 976
- exanthematous . . . . . 976, 948
- periodical . . . . . 908, 948
- type of . . . . . 907
- Fibrin, clots of, in the heart . . . 464
- in the urine . . . . . 781
- Fifth nerve, painful anæsthesia of . . . . . 255
- Filaria medinensis . . . . . 1056
- sanguinis hominis . . . 214, 777, 781, 1055
- Fissure of Rolando . . . . . 58, 59, 63
- of Sylvius . . . . . 58, 63, 881
- Fistula, gastro-pulmonary . . . . 577
- rectal . . . . . 624



- Flagellate protozoa in influenza . 876  
 Flatulency as a symptom . . . . . 557  
 Fly parasites . . . . . 1056  
 Follicular tonsillitis . . . . . 524  
 Foot and mouth disease . . . . . 1046  
 Foot clonus . . . . . 168  
   fungus, of India . . . . . 1049  
   perforating ulcer of . . . . . 1049  
 Frenum linguae, ulceration of . . 337  
 Fremitus, bronchial . . . . . 326  
   cavernous . . . . . 326  
   friction . . . . . 303  
   pleural . . . . . 326  
   rhonchal . . . . . 303  
   vocal . . . . . 303, 326  
 Friction, pericardial . . . . . 444  
   pleural . . . . . 323  
 Friedreich's ataxia . . . . . 163  
 Frontal sinus, diseases of . . . . 260  
   neuralgia in . . . . . 261  
 Fungi . . . . . 339, 538, 561, 595  
   poisonous . . . . . 1031  
 Fungus foot of India . . . . . 1049
- G.**
- Gait as a symptom . . . . . 31  
   in diseases of cerebellum . . . . 165  
   in general paralysis . . . . . 247  
   in hysterical hemiplegia . . . . 139  
   in locomotor ataxia . . . . . 31, 160  
   waddling . . . . . 154  
 Gall-bladder, cancer of . . . . . 694, 695  
   diseases of . . . . . 673, 694  
   confounded with cancer of  
     liver . . . . . 694  
   distention of . . . . . 674, 694  
   distinguished from hydatids of  
     liver . . . . . 699  
   dropsy of . . . . . 628  
   inflammation of . . . . . 673  
   rupture of . . . . . 628  
 Gall-duets, inflammation of, 602, 673  
 Gall-stones . . . . . 590, 601, 695  
   impacted . . . . . 676  
   passage of, confounded with  
     cancer of the liver . . . . . 696  
     with colic . . . . . 601  
     with faecal accumulations . . 602  
     with intermittent fever . . . . 952  
 Galvanic excitability . . . . . 110  
 Gangrene of ergot poisoning . . 1040  
   of the mouth . . . . . 520  
   confounded with diphtheria . . 529  
   pulmonary, confounded with  
     phthisis . . . . . 380  
   symmetrical . . . . . 840  
 Ganglia, central gray, lesions of . 118  
 Gastralgia . . . . . 566, 580  
   confounded with colic . . . . . 600, 605
- Gastric cancer . . . . . 582  
   crises . . . . . 162, 580, 725  
   juice, examination of . . . . . 553, 557  
   super-secretion of . . . . . 592  
   motormeter . . . . . 552  
   tubules, atrophy of . . . . . 576, 583  
   ulcer . . . . . 576, 580  
 Gastritis, acute idiopathic . . . . 571  
   distinguished from gastritis  
     by poisoning . . . . . 571, 1028  
   chronic . . . . . 575  
   confounded with gastric can-  
     cer . . . . . 582, 585  
   with gastric ulcer . . . . . 576, 585  
   with hepatic congestion . . . . 680  
   with peritonitis . . . . . 613  
   of young children . . . . . 574  
 Gastrodiaphane . . . . . 551  
 Gastrodynia . . . . . 566  
 Gastrograph . . . . . 552  
 Gastro-intestinal disorders asso-  
   ciated with congestive  
   fever . . . . . 963  
   confounded with Bright's  
   disease . . . . . 803  
 Gastromalacia . . . . . 574  
 Gastroptosis . . . . . 592, 856  
 Gastroxynsis . . . . . 557  
 German measles . . . . . 985  
 Girdle pain . . . . . 133, 168  
   sense . . . . . 129  
 Gland of axilla, enlarged . . . . . 508  
   of neck, enlarged . . . . . 333, 508  
   retroperitoneal, cancer of . . . . 715  
   retropharyngeal . . . . . 537  
   scrofulous . . . . . 647, 868  
   thymus . . . . . 334  
   thyroid . . . . . 251, 456, 513  
   tuberculous . . . . . 868  
 Glanders . . . . . 263  
   acute, confounded with pyæ-  
     mia . . . . . 264, 874  
   distinguished from rhinitis . . . . 264  
   from typhoid fever . . . . . 264  
 Gland-secretions, altered . . . . 1023  
 Glaucoma . . . . . 90  
 Glénard's disease . . . . . 592  
 Glossoplegia . . . . . 122  
 Glottis, œdema of . . . . . 333, 997  
   spasm of . . . . . 279, 333  
 Glycosuria . . . . . 265, 826  
 Goitre . . . . . 90, 333, 513  
   exophthalmic . . . . . 90, 456, 513  
 Gonorrhœal rheumatism . . . . . 891  
 Gout . . . . . 73, 466, 889, 898  
   blood examination in . . . . . 899  
   distinguished from rheumatism . 898  
   rheumatic . . . . . 900  
 Gram's tests for micro-organisms . 345  
 Gravel . . . . . 603, 750

- Graves's disease . . . . . 456, 457  
Guinea-worm . . . . . 1056  
Gummata . . . . . 153  
Gums, bleeding from . . . 706, 855  
    red line of . . . . . 362  
Gyromele . . . . . 551
- H.**
- Habit-chorea . . . . . 221  
Habit-spasm . . . . . 221  
Hæmacytometer, forms of . 842, 845  
Hæmatemesis . . . . . 346, 563, 917  
    alcoholic . . . . . 704  
Hæmatinuria, intermittent . . 776  
Hæmatocele, retro-uterine . . 728  
Hæmatokrit . . . . . 843, 848, 849  
Hæmatoma . . . . . 199  
Hæmatoscope . . . . . 852  
Hæmatozoa of malaria . . . 961  
Hæmaturia . . . . . 772  
    confounded with acute Bright's  
    disease . . . . . 794  
    intermittent . . . . . 776  
    malarial . . . . . 776, 966  
    parasitic . . . . . 777  
    renal tubal . . . . . 774, 777  
    diagnostic sign of . . . 775  
    vesical . . . . . 778  
Hæmidrosis . . . . . 1024  
Hæmoglobin . . . . . 777, 849, 856  
    apparatuses for estimating . 852  
Hæmoglobinometer . . . . . 852  
Hæmoglobinuria . . . . . 776, 1032  
    intermittent . . . . . 966  
Hæmometer . . . . . 849, 852, 853  
Hæmophilia distinguished from  
    leukæmia . . . . . 888  
    from purpura . . . . . 887  
Hæmoptysis . . . . . 345, 355, 425  
    in typhoid . . . . . 917  
Handwriting, alteration of . . 247  
Hay asthma . . . . . 262  
Hay fever . . . . . 90, 262, 354  
    distinguished from catarrhal  
    fever . . . . . 912  
Head, enlargement of, diseases  
    characterized by . . . 249  
    gouty inflammation in . . 899  
    shapes of, in disease . . 249, 250  
Headache . . . . . 81  
    congestive . . . . . 82  
    from astigmatism . . . 81  
    from Bright's disease . . 802  
    from eye-strain . . . . 88  
    from lithæmia . . . . . 900  
    from poisoning . . . . 83  
    in Bright's disease . . . 83  
    in diseases of the brain . 81  
    nervous and neuralgic . . 82
- Headache, sick . . . . . 82  
    sympathetic . . . . . 83  
Hearing, derangement of . . 98  
Heart, anatomy and physiology  
    of . . . . . 428  
    aneurism of . . . . . 478  
    atrophy of . . . . . 460, 487  
    auricle of, dilated . . . . 512  
    auscultation of . . . . . 435  
    cavities of, accumulation of  
    blood in . . . . . 488  
    chronic diseases of, with in-  
    creased percussion dullness 478  
    clots of fibrin in . . . . . 464  
    dilatation of . . . . . 425, 482  
    diseases of . . . . . 428  
        associated with asthma . 333  
        with diseases of colon . 647  
        confounded with intermittent  
        fever . . . . . 952  
        with pernicious anæmia . 862  
        symptoms of . . . . . 445  
    displacements of, diseases pre-  
    sented . . . . . 410, 505  
    dropsy caused by disease of,  
        . . . . . 446, 802  
    enlargement of, symptoms of . 489  
        mistaken for aneurism . . 512  
    examination of . . . . . 431  
    fatty accumulation on . . 335, 487  
    fatty degeneration of . . . 484  
    functional disorders of . . 453, 491  
    gouty . . . . . 481, 899  
        associated with contracting  
        kidney . . . . . 481  
    hemisystole of . . . . . 455  
    hypertrophy of . 356, 425, 478, 802  
    impulse of . . . . . 431, 453  
    inflammation of . . . . . 476  
    inspection of . . . . . 431  
    irregularity of action of . . 455  
    irritable . . . . . 451, 467  
    malformations of . . . . . 490  
        causing abscess of brain . 491  
    mitral disease of . . . . . 489, 497  
    murmurs . . . . . 489, 489  
    organic diseases of . . . . 460  
    overaction of . . . . . 441, 459  
    pain in region of . . . 448, 451, 460  
    palpation of . . . . . 432  
    palpitation of . . . . . 452, 454  
    paralysis of, in relapsing fever 945  
    percussion of . . . . . 433  
    percussion dullness of, increased 478  
    rhythm of . . . . . 454  
    rupture of . . . . . 487  
    sounds of . . . . . 438, 460  
    starvation . . . . . 487  
    strain . . . . . 459  
    valvular affections of, 456, 489, 496

- Heart, valvular affections of,  
table of . . . . . 499
- Heart-burn . . . . . 557
- Heat exhaustion . . . . . 211
- Hectic fever distinguished from  
intermittent fever . . . . . 951
- Heller's test . . . . . 768
- Hemeralopia . . . . . 96
- Hemianopsia . . . . . 61, 93, 97, 122  
bilateral . . . . . 61, 118
- Hemianæsthesia . . . . . 76, 139
- Hemicrania . . . . . 82, 255, 801  
distinguished from pain of or-  
ganic cerebral affections . . . . . 256  
from periostitis . . . . . 256  
from rheumatism of the  
scalp . . . . . 256
- Hemiparaplegia . . . . . 115
- Hemiplegia . . . . . 113  
alternating . . . . . 114  
anatomical diagnosis of . . . . . 115  
appearance of muscles in . . . . . 119  
cerebral . . . . . 115, 150, 200  
corpus striatum in . . . . . 116  
cortical . . . . . 118  
electricity as a test of . . . . . 115, 120  
feigned . . . . . 120  
following epilepsy . . . . . 215  
lesions of capsule . . . . . 118  
of crus cerebri . . . . . 117  
of gray central ganglia . . . . . 118  
of optic tract . . . . . 118  
of pons Varolii . . . . . 114, 116  
of præfrontal lobes . . . . . 119  
hysterical . . . . . 139  
ovarian tenderness in . . . . . 139  
in diphtheria . . . . . 528  
in the course of typhoid fever . . . . . 921  
nature of lesions in . . . . . 119  
optic thalamus in . . . . . 116  
pain in . . . . . 120  
pathological diagnosis of . . . . . 119  
rigidity in . . . . . 119  
seat of lesion in . . . . . 114  
spinal . . . . . 115
- Hemorrhage a cause of apoplexy . . . . . 194  
between brain membranes . . . . . 199  
cerebellar . . . . . 198  
cerebral . . . . . 199, 200  
cortical . . . . . 198  
from aneurism . . . . . 346  
from the bladder . . . . . 773  
from the intestines . . . . . 581, 917  
from the kidneys . . . . . 773  
from the larynx, trachea, etc. . . . . 346  
from the lungs . . . . . 346, 372  
from the œsophagus . . . . . 346  
from the oral cavity . . . . . 345  
from the prostate gland . . . . . 778  
from the stomach, 346, 563, 579, 609
- Hemorrhage from the stomach  
distinguished from irritant  
poisoning . . . . . 1028  
from the urethra . . . . . 778  
from the uterus in myxœdema . . . . . 834  
from the ventricles of the brain . . . . . 197  
in apoplexy, seat of . . . . . 197  
in fevers . . . . . 845  
in yellow fever . . . . . 971  
into cerebrum ovale . . . . . 198  
into lung texture . . . . . 394  
into the corpora quadrigemina . . . . . 198  
into the internal capsule . . . . . 197  
into the pons . . . . . 198  
into the pons Varolii . . . . . 202  
into the subarachnoid spaces . . . . . 198  
into the thalamus . . . . . 198  
limited to the arachnoid . . . . . 198  
to one crus cerebri . . . . . 198  
nasal . . . . . 263  
of the bowels . . . . . 651  
punctiform . . . . . 238  
relations of, to brain-softening . . . . . 236  
renal, clots in . . . . . 774  
spinal . . . . . 125, 133  
vicarious . . . . . 345, 564, 652
- Hemorrhagic diathesis . . . . . 1045  
malarial fever . . . . . 965  
confounded with intermit-  
tent hæmoglobinuria . . . . . 966  
with yellow fever . . . . . 966
- Hemorrhoids . . . . . 641, 649, 652
- Hepatic abscess . . . . . 342  
diseases, chronic and acute,  
confounded . . . . . 670  
dropsy . . . . . 833  
fever . . . . . 675  
confounded with intermit-  
tent fever . . . . . 952  
neuralgia . . . . . 601
- Hepatitis, acute . . . . . 667, 675  
confounded with acute in-  
fectious jaundice . . . . . 671  
with acute non-hepatic  
diseases with jaundice . . . . . 670  
with acute yellow atrophy,  
678, 678  
with cancer of liver . . . . . 693  
with chronic hepatic dis-  
ease with acute symp-  
toms . . . . . 670  
with diaphragmatic pleu-  
risy . . . . . 670  
with inflammation of the  
biliary passages . . . . . 673  
with inflammation of the  
portal veins . . . . . 669  
with perihepatitis . . . . . 668  
with pigment liver . . . . . 669  
chronic . . . . . 681

- Hepatitis, interstitial . . . 702, 705  
   subacute infectious . . . . . 705  
   suppurative . . . . . 670  
 Hernia, diaphragmatic, con-  
   founded with pneumo-  
   thorax . . . . . 414  
   irreducible . . . . . 632  
   omental . . . . . 590  
   strangulated, confounded with  
     colic . . . . . 600  
     with intestinal obstructions . 632  
     with irritant poisoning . 1028  
   through the recti muscles . 687  
 Herpes . . . . . 388, 1009  
   gestationis . . . . . 1009, 1012  
   labialis . . . . . 1009  
   zoster . . . . . 174, 1009  
     distinguished from erysipelas 998  
     ophthalmicus . . . . . 90  
     pain in, mistaken for pleurisy 1009  
 Hiccup in diaphragmatic pleu-  
   risy . . . . . 671  
   in diphtheritic dysentery . . 651  
 Hip-joint affections . . . 258, 629  
   confounded with sciatica . . 258  
 Hodgkin's disease . . . . . 866  
   confounded with lymphatic  
   cancer . . . . . 867  
   with a malarial affection . . 868  
 Hooping-cough . . . . 285, 336, 359  
   diagnosed from acute bron-  
   chitis . . . . . 337  
   from bronchial phthisis . . 337  
 Hydatid cysts . . . . . 421  
 Hydatids of the liver . . 683, 697, 706  
   distinctive character of fluid in 700  
   multilocular . . . . . 701  
 Hydroa . . . . . 1010  
 Hydrocephaloid disease . . . 187  
 Hydrocephalus, acute . . . 185, 187  
   chronic . . . . . 187, 249  
 Hydrochloric acid in gastric juice,  
   . . . . . 557, 567, 575  
 Hydronephrosis . . . . 700, 822, 828  
   confounded with hydatid tumor  
   of kidney . . . . . 822  
   with renal cysts . . . . . 823  
   intermitting . . . . . 725  
 Hydrophobia . . . . . 232, 539  
   distinguished from strychnine  
   poisoning . . . . . 1037  
   from tetanus . . . . . 232  
 Hydrorrhœa, nasal . . . . . 262  
 Hydrothorax . . . . . 331, 422  
   confounded with chronic pleu-  
   risy . . . . . 422  
 Hyperæmia of the disk . . . . 94  
 Hyperæsthesia . . . . . 72, 107  
   general . . . . . 72  
   hysteria as a cause of . . . . 73  
 Hyperæsthesia, one-sided . . . 74  
 Hyperalgesia . . . . . 73  
 Hyperopia . . . . . 81, 88  
 Hypertrophy of brain . . . . 250  
   of heart . . . . . 478  
   of mucous membrane . . . . 264  
   of skin . . . . . 1017  
 Hypochondriasis . . . . . 167, 569  
 Hypochondrium, tumors of . . . 718  
 Hypogastric region, tumors of . . 729  
 Hypo-leukæmia . . . . . 866  
   false . . . . . 866  
 Hysteria . . . . . 34, 73, 151, 224  
   abdominal, confounded with  
   peritonitis . . . . . 619  
   with tubercular meningitis . 189  
   after railway accidents . . . 227  
   associated with catalepsy . . 212  
   with membranous diarrhœa . 648  
   with muscular atrophy . . . 151  
   cerebral, distinguished from  
     apoplexy . . . . . 205  
     from chorea . . . . . 224  
     from epilepsy . . . . . 225  
   feigned . . . . . 228  
   resembling locomotor ataxia . 165  
   spasms in . . . . . 171  
   toxic . . . . . 227  
   traumatic . . . . . 140  
 Hysterical anorexia . . . . . 556  
   complaints, local . . . . . 226  
   contractures . . . . . 231  
   delirium . . . . . 70  
   headache . . . . . 227  
   hydrophobia . . . . . 233  
   locomotor ataxia . . . . . 165  
   paralysis . . . . . 138  
   pseudo-maladies . . . . . 227  
   tetanus . . . . . 229  
   urine in . . . . . 228  
 Hystero-epilepsy . . . . . 226  
   aura in . . . . . 226  
   ovarian . . . . . 226

## I.

- Ichthyosis . . . . . 175, 1015  
 Icterus . . . . . 665  
   catarrhalis . . . . . 673  
   distinguished from abscess of  
   liver . . . . . 675  
   from acute atrophy of liver . 675  
   from biliary calculi . . . 675  
   from cancer of liver . . . 675  
   from cirrhosis . . . . . 675  
   from congestion of liver . 674  
   from hepatic inflammation . 675  
   neonatorum . . . . . 665  
 Ileum, catarrh of . . . . . 608  
   ulceration of . . . . . 627



- Iliac fossa, disease in . . . . . 622  
     region, tumor of . . . . . 727  
 Illusions . . . . . 87  
 Impetigo . . . . . 1011  
     contagiosa . . . . . 1011  
 India-rubber poisoning . . . . . 1044  
 Indigestion, functional . . . . . 570  
 Infantile paralysis . . . . . 155  
     scurvy . . . . . 885  
 Infaret, hemorrhagic . . . . . 395, 466  
 Influenza . . . . . 262, 876, 909  
     See also *Catarrhal fever*.  
 Innominate, aneurism of . . . . . 517  
 Inosite . . . . . 766  
 Inosuria . . . . . 766  
 Insanity . . . . . 67  
     chronic . . . . . 181  
     confounded with delirium . . . 69  
     following acute rheumatism . . 893  
 Insensibility from drink distin-  
     guished from apoplexy . . . 201  
     from narcotics distinguished  
     from apoplexy . . . . . 201  
 Insolatio. See *Sun-stroke*.  
 Insomnia . . . . . 67, 72  
     with delirium . . . . . 72  
 Inspiration, jerking . . . . . 316  
 Insufficiency of aortic valves  
     confounded with aneurism . . 510  
 Intellection deranged . . . . . 67  
 Intermittent fever . . . . . 948  
     distinguished from chills of  
     pus formations . . . . . 951  
     from diseases of the heart . . 952  
     from hectic fever . . . . . 951  
     from hepatic fever . . . . . 952  
     from passage of gall-  
     stones . . . . . 952  
     from puerperal malarial  
     fever . . . . . 953  
     from remittent fever . . . . . 956  
     from syphilitic fever . . . . . 953  
     from urethral fever . . . . . 952  
     types of . . . . . 949  
 Intestinal worms . . . . . 1050  
 Intestines, cancer of . . . . . 731  
     contraction of . . . . . 634, 638  
     dilatation of . . . . . 592, 640  
     confounded with dilatation of  
     stomach . . . . . 592  
     diseases of . . . . . 593  
     hemorrhage of . . . . . 651, 706, 917  
     inflammation of . . . . . 606  
     internal strangulation of . . . 636  
     intussusception of . . . . . 635  
     invagination of . . . . . 627, 635  
     obstruction of . . . . . 630, 637  
     causes of . . . . . 634  
     confounded with peritonitis . . 631  
     with strangulated hernia . . 632  
 Intestines, obstruction of, fre-  
     quency of . . . . . 639  
     from passage of gall-stones . . 637  
     from stricture . . . . . 638  
     from volvulus . . . . . 637  
     location of lesion in . . . . . 639  
     percussion of . . . . . 547, 549  
     perforation of . . . . . 612  
     distinguished from colic . . . 600  
     from irritant poisoning . . . 1028  
     sloughing of . . . . . 636  
     stricture of . . . . . 638  
     tubercular disease of . . . . . 646  
     worms in . . . . . 1050  
 Intoxication . . . . . 71  
     uræmic . . . . . 796  
 Intra-hepatic concretion . . . . . 952  
 Iris-contraction . . . . . 100  
 Iritis . . . . . 900  
 Irritant poisoning . . . . . 1028  
 Itch . . . . . 1005, 1021  
     army . . . . . 1021
- J.**
- Jaundice . . . . . 601, 662  
     acute infectious . . . . . 671  
     catarrhal . . . . . 673  
     diagnosis of . . . . . 662  
     fatal forms of . . . . . 665  
     from blood-poison . . . . . 664  
     from mental emotion . . . . . 665  
     in acute non-hepatic diseases . . 670  
     in phosphorus poisoning . . . 1029  
     in pneumonia . . . . . 390  
     obstructive . . . . . 663  
     of the new-born . . . . . 665  
 Jaw-jerk . . . . . 100  
 Joint affection, hip . . . . . 258, 629  
     pyæmic . . . . . 892  
 Joint-inflammations . . . . . 152
- K.**
- Kakke . . . . . 153  
 Keratitis . . . . . 90  
 Kidney, abscess around . . . . . 627, 818  
     abscess of . . . . . 627, 816  
     distinguished from cystitis . . 818  
     affections of, with swelling . . 590  
     calculus in, symptoms of . . . 790  
     cancer of . . . . . 700  
     distinguished from enlarged  
     spleen . . . . . 720  
     contracted . . . . . 481, 646, 808, 810  
     associated with albuminuric  
     ulceration of bowel . . . . . 646  
     with gout . . . . . 899  
     confounded with myxædema . . 834  
     with pernicious anæmia . . . 862

- Kidney, contracted, diuresis in . . . 827  
 displacement of . . . . . 591  
 cysts of . . . . . 805  
 enlarged, chronically inflamed . . . 807  
   confounded with cancer of  
     liver . . . . . 697  
     with hydatids of liver . . . 700  
     with ovarian tumor . . . 728  
 fatty, enlarged . . . . . 807  
 floating . . . . . 627  
 hemorrhage from . . . . . 773  
 hydatids of . . . . . 822  
   confounded with hydrone-  
     phrosis . . . . . 822  
 inflammation of . . . . . 784  
   of pelvis of . . . . . 819  
 movable or displaced . . . 724, 725  
   simulated by malignant dis-  
     ease of colon . . . . . 726  
 neuralgia of . . . . . 786  
 pain in . . . . . 785  
   confounded with colic . . . 602  
   paroxysmal . . . . . 786  
   persistent . . . . . 788  
 percussion of . . . . . 547  
 sarcoma of . . . . . 804  
 suppurative inflammation of . . 818  
 surgical . . . . . 806  
 syphilomata of . . . . . 804  
 tubercle of, confounded with  
   Bright's disease . . . . . 804  
   tumors of . . . . . 627, 805  
   waxy or amyloid, enlarged . . 808  
 Klebs-Loeffler bacillus . . . 262, 524  
 Knee-jerk . . . . . 101, 182  
 Kreatin and kreatinin . . . . 757
- L.**
- Lactic acid . . . . . 584, 589  
 Landry's paralysis . . . . . 126  
 Laryngeal affections, acute . . . 274  
   cough . . . . . 267  
   crises . . . . . 162  
   diphtheria . . . . . 285  
   paralysis . . . . . 288  
   phthisis . . . . . 292  
   rheumatism . . . . . 276  
   spasm . . . . . 280  
     from use of antipyrine . . . 1036  
   stenosis . . . . . 293  
   stridor . . . . . 266, 514  
   vertigo . . . . . 86, 280  
 Laryngismus stridulus . . . . 279  
 Laryngitis, acute . . . . . 267, 274  
   confounded with croup . . . 275  
   distinguished from acute pul-  
     monary affections . . . 275  
   from pharyngitis . . . . . 275  
   from tonsillitis . . . . . 275
- Laryngitis, chronic . . . . . 286  
   aneurism of aorta confounded  
     with . . . . . 287, 514  
   combined with syphilis . . . 286  
   with tuberculosis . . . . . 286  
   confounded with altered  
     voice . . . . . 287  
   with nervous aphonia . . . 287  
   of epiglottis . . . . . 290  
   diffuse cellular . . . . . 275  
   diseases confounded with . . . 287  
   erysipelatosus . . . . . 275  
   feigned . . . . . 289  
   hemorrhagic . . . . . 276  
   hiemalis . . . . . 275  
   membranous . . . . . 285  
   oedematous . . . . . 277  
   secondary, of the exanthemata . 283  
   sicca . . . . . 275  
   spasmodic . . . . . 280  
   syphilitic or tubercular . . . 292  
 Laryngoscopy . . . . . 267, 271  
 Larynx, abscess of . . . . . 284  
   actinomycosis of . . . . . 285  
   acute diseases of . . . . . 274  
   affections of nerves of . . . 288  
   cancer of . . . . . 294  
   cartilages and perichondrium  
     of, diseases of . . . . . 291  
   changes in breathing in dis-  
     eases of . . . . . 266  
     in voice in diseases of . . . 266  
   chronic diseases of . . . . . 286, 514  
   cough in diseases of . . . . 267  
   extirpation of . . . . . 291  
   growths in . . . . . 293  
   hyperæsthesia of . . . . . 289  
   inflammation of . . . . . 274  
   myxomata of . . . . . 274  
   oedema of . . . . . 277  
   organic diseases of . . . . . 273  
   pachydermia of . . . . . 292  
   pain in diseases of . . . . . 267  
   polypi in . . . . . 87  
   rupture of . . . . . 358  
   stenosis of . . . . . 293  
   table of diseases of . . . . . 273  
   tubercle of . . . . . 292  
   tumors of . . . . . 293  
   ulcers in . . . . . 292, 293, 539  
   venous congestion of . . . . 290  
   ventricular bands of, hyper-  
     trophy of . . . . . 290  
 Lead poisoning . . . . . 598, 609, 810,  
   1030, 1041  
   paralysis from, 105, 141, 151, 1042  
 Lentigo . . . . . 1015  
 Lepra, or leprosy . . . . . 1017  
 Leptothrix . . . . . 339, 537  
 Lethargy, African . . . . . 214

- Leucine . . . . . 679, 745, 759  
 Leucocytosis . . . . . 625  
 Leukæmia . . . . . 95, 863  
   distinguished from hæmophilia  
     and purpura . . . . . 888  
   from pernicious anæmia . . . . . 863  
   false hypo- . . . . . 866  
   lymphatic . . . . . 864, 865  
   medullary . . . . . 866  
   myelogenous . . . . . 864  
   of liver . . . . . 690  
   pseudo-leukæmia distinguished  
     from . . . . . 867  
   splenic . . . . . 864, 865  
 Lichen . . . . . 1005, 1014  
   planus . . . . . 1005  
   ruber . . . . . 1005  
   scrofulosorum . . . . . 1005  
   syphilitic . . . . . 1026  
   tropicus . . . . . 1005, 1024  
 Lipoma of intestine . . . . . 636  
 Lithæmia . . . . . 73, 85, 900  
   associated with disorders of  
     vision . . . . . 900  
   with pain in stomach and  
     bowels . . . . . 900  
 Liver, abscess of . . . . . 667, 675, 682  
   distinguished from hydatids . . . . . 699  
   actinomycosis of . . . . . 684  
   acute affections of . . . . . 667, 676  
     confounded with pyæmia . . . . . 874  
     congestion of . . . . . 667, 674  
     confounded with cancer of  
       liver . . . . . 693  
     with remittent fever . . . . . 956  
   inflammation of . . . . . 667, 675, 693  
     distinguished from catarrhal  
       icterus . . . . . 675, 693  
   yellow atrophy of . . . . . 672, 675, 676  
     confounded with hepatitis . . . . . 678  
     with typhoid fever . . . . . 678  
     with yellow fever . . . . . 678  
   from phosphorus-poisoning . . . . . 679  
   nervous symptoms in . . . . . 677  
   cancer of . . . . . 675, 690, 699, 707  
   chronic affections of . . . . . 679  
   atrophy of . . . . . 709  
   congestion of . . . . . 674, 679, 689  
     confounded with cancer of  
       liver . . . . . 692  
     with catarrhal icterus . . . . . 674  
     with chronic gastritis . . . . . 680  
     with hypertrophy of  
       liver . . . . . 680  
     with torpor of liver . . . . . 680  
     nervous symptoms in . . . . . 681  
   cirrhosis of . . . . . 621, 675, 702  
   hypertrophic . . . . . 705, 709  
   decrease in size of . . . . . 676, 702  
   diseases of . . . . . 661, 666  
 Liver, diseases of, dropsy in . . . . . 702  
   from inflammation of portal  
     vein . . . . . 708  
   jaundice in . . . . . 661, 676  
   malarial infection in . . . . . 676  
   pain in . . . . . 661, 702  
   with absence of jaundice, 679, 701  
   displacement of, diseases pre-  
     sented . . . . . 410  
   from tight lacing . . . . . 680  
   elastic tumors of . . . . . 698  
   enlargement of . . . . . 589, 679, 689  
     confounded with chronic  
       pleurisy . . . . . 421  
   fatty . . . . . 672, 688  
     confounded with cancer . . . . . 692  
   fibro-fatty . . . . . 706  
   hydatids of . . . . . 683, 697, 706  
   hypertrophy of . . . . . 680  
   infarcts in . . . . . 467  
   inflammation of . . . . . 668, 681  
   leukæmic . . . . . 690  
   movable . . . . . 726  
   percussion of . . . . . 545  
   pigment, confounded with  
     acute hepatitis . . . . . 669  
   pyæmic abscess of . . . . . 678  
   red atrophy of . . . . . 707  
   simple induration of . . . . . 707  
   syphilitic . . . . . 689, 694  
     confounded with cancer of  
       liver . . . . . 694  
   table of diseases of . . . . . 666  
   torpor of . . . . . 680  
   tropical abscess of . . . . . 667, 687  
   waxy . . . . . 688, 692  
     confounded with cancer . . . . . 692  
     diseases confounded with . . . . . 689  
 Lock-jaw. See *Tetanus*.  
 Locomotor ataxia . . . . . 129, 135, 159  
   arthropathies of, distinguished  
     from rheumatic arthritis . . . . . 902  
   crises in . . . . . 162  
   diminution or loss of muscular  
     sense in . . . . . 164  
   distinguished from diseases of  
     the spinal cord . . . . . 163  
   from general paralysis of the  
     insane . . . . . 162  
   eruptions in . . . . . 162  
   gait in . . . . . 31, 160, 161  
   of syphilitic origin . . . . . 160, 165  
   resembling hysteria . . . . . 165  
   station in . . . . . 32  
 Lumbago . . . . . 895  
 Lumbar region, tumors of . . . . . 727  
 Lungs, actinomycosis of . . . . . 381  
   acute affections of . . . . . 384  
   confounded with tubercular  
     meningitis . . . . . 188

- Lungs, acute affections of, in  
  typhoid fever . . . . . 927  
  cancer of . . . . . 375  
  cirrhosis of . . . . . 425, 426  
  collapse of . . . . . 359, 426  
  diseases of . . . . . 299, 329  
  fistulous opening into . . . . . 427  
  gangrene of . . . . . 339, 355, 380  
  gray hepatization of . . . . . 392  
  hydatids of . . . . . 421  
  hypostatic congestion of . . . . . 654  
  inflammation of . . . . . 388  
  oedema of . . . . . 393, 796  
  scrofulous disease of . . . . . 383  
  syphilitic disease of . . . . . 375  
  tuberculosis of . . . . . 344, 385  
  See also *Phthisis*.
- Lupus . . . . . 1016  
  erythematodes . . . . . 1016  
  verrucosus . . . . . 1016  
  vulgaris . . . . . 1016
- Lymphadenoma . . 137, 420, 508, 866  
  distinguished from lymphatic  
  cancer . . . . . 867
- Lymphatic glands, cancer of, 727, 867  
  sarcoma of . . . . . 867  
  system, disease of . . . . . 1018
- Lymphomas, local gland . . . . . 868  
  distinguished from Hodgkin's  
  disease . . . . . 868  
  of mediastinum . . . . . 508
- M.**
- Maculæ . . . . . 1015
- Malaria, appearance of tongue in . 961  
  chronic . . . . . 95, 967  
  examination of blood in . . . . . 960  
  poisoning by . . . . . 787
- Malarial cachexia . . . . . 960  
  changes in kidneys . . . . . 967  
  cirrhosis . . . . . 706  
  corpuscles in blood . . . . . 968  
  fever, distinguished from den-  
  gue . . . . . 996  
  hæmaturia . . . . . 776  
  neuralgia . . . . . 968  
  palsy . . . . . 968  
  poisoning, chronic . . . . . 95, 967  
  septicæmia . . . . . 876  
  symptoms, in pneumonia . . . . . 402
- Malformations of heart con-  
  founded with valvular af-  
  fections . . . . . 490
- Malignant pustule . . . . . 1045
- Mania, acute . . . . . 192  
  alcoholic . . . . . 194  
  associated with mumps . . . . . 534  
  with typhoid fever . . . . . 919
- Mania, confounded with acute  
  meningitis . . . . . 193  
  with delirium tremens . . . . . 193  
  uræmic . . . . . 798
- Marasmus . . . . . 243
- Measles . . . . . 95, 398, 981  
  anomalous forms of . . . . . 984  
  associated with acute bronchitis . 983  
  with pneumonia . . . . . 984  
  catarrh in . . . . . 262, 982  
  complications of . . . . . 398, 983  
  distinguished from miliary  
  fever . . . . . 984  
  from rubella . . . . . 986  
  from scarlet fever . . . . . 980  
  from smallpox . . 980, 982, 992  
  from typhus fever . . . . . 983
- eruption in . . . . . 982
- German . . . . . 985
- malignant, confounded with  
  cerbro-spinal fever . . . . . 942
- Mediastinum, abscess of . . . . . 508  
  fluid in . . . . . 420  
  inflammatory thickening of . . 509  
  tumor of . . . . . 420
- Megrim. See *Migraine*.
- Melæna . . . . . 651
- Melancholia, acute . . . . . 194
- Melasma . . . . . 1015
- Memory, disordered . . . . . 67
- Ménière's disease . . . 84, 85, 98, 681
- Meningitis, acute . . . . . 179, 378  
  confounded with acute mania . 193  
  with acute softening of  
  brain . . . . . 181  
  with apoplexy . . . . . 199  
  with cerebritis . . . . . 181  
  with delirium tremens . . . . . 193  
  with head-symptoms of  
  acute rheumatism . . . . . 184  
  of acute ulcerative endo-  
  carditis . . . . . 185  
  of continued fevers . . . . . 182  
  of pericarditis . . . . . 185  
  of pneumonia . . . . . 185  
  of remittent fever . . . . . 958  
  of typhus fever . . . . . 932  
  with typhoid fever . . . . . 927
- cerebro-spinal . . . . . 189, 937  
  diseases confounded with . . . 940  
  distinguished from purpura . . 887  
  sporadic . . . . . 942
- chronic, distinguished from tu-  
  mor . . . . . 242
- epidemic . . . . . 937
- idiopathic . . . . . 942
- of the base of the brain . . . . . 180
- of the convexity of the brain . 180
- ordinary . . . . . 186
- purulent . . . . . 705



Meningitis, spinal . . . . .	131
sporadic cerebro-spinal . . . . .	942
tubercular . . . . .	185, 942
distinguished from acute hydrocephalus . . . . .	187
from cerebro-spinal fever . . . . .	942
from chronic hydrocephalus . . . . .	187
from hysteria . . . . .	189
from inflammatory affections of the lungs . . . . .	188
from typhoid fever . . . . .	188
rose-spots in . . . . .	188
Meningo-encephalitis . . . . .	246
Mensuration of chest . . . . .	300
Mental faculties, diseases characterized by gradual impairment of . . . . .	235
Mercurial tremor . . . . .	1041
Mercyism . . . . .	565
Metritis confounded with acute cystitis . . . . .	816
with peritonitis . . . . .	615
Migraine . . . . .	82, 256, 260, 802
Miliaria papulosa . . . . .	1005, 1024
vesiculosa . . . . .	1024
Miliary fever . . . . .	984
Milk-leg. See <i>Phlegmasia alba dolens</i> .	
Milk-sickness . . . . .	1047
Mind-blindness . . . . .	98, 209
Mollities ossium . . . . .	904
Molluscum contagiosum . . . . .	1024
epitheliale . . . . .	1024
fibroma . . . . .	1024
Monoplegia . . . . .	120
brachio-facial . . . . .	120, 121
crural . . . . .	122
facial . . . . .	120
facio-lingual . . . . .	122
oculo-motor . . . . .	122
Morphine habit . . . . .	1038
Morphœa . . . . .	1020
Morton's disease . . . . .	897
Morvan's disease . . . . .	153, 175
Motion, deranged . . . . .	103
voluntary diseases marked by sudden loss of . . . . .	194
Motor path of Gowers . . . . .	64
Mouth, diseases of . . . . .	519
gangrene of . . . . .	520
inflammation of . . . . .	519
morbid appearances of . . . . .	519
Mucus, vomiting of . . . . .	561
Multiple neuritis, 128, 164, 528, 833, 1044	
See also <i>Neuritis</i> .	
Mumps . . . . .	523, 534
See also <i>Parotitis</i> .	
confounded with erysipelas . . . . .	999

Mumps, diplococcus in . . . . .	534
Murmur, respiratory . . . . .	314
vesicular . . . . .	313
absence of . . . . .	316
changes in . . . . .	314
Murmurs, cardiac . . . . .	439
endocardial . . . . .	439, 461
from lung changes . . . . .	492
from misdirection of current . . . . .	492
functional valvular . . . . .	491
in the course of fevers . . . . .	463
musical . . . . .	493
over thyroid gland . . . . .	457
pericardial . . . . .	444
without valvular lesion . . . . .	492
Muscæ volitantes . . . . .	87
Muscle, rectus, contraction of . . . . .	722
Muscles, appearance of, in paralysis . . . . .	108
morbid states of, paralysis from overuse of . . . . .	105
overuse of . . . . .	152
Muscular contraction, paradoxical . . . . .	103
movements, irregular forms of . . . . .	164
rigidity . . . . .	131
sense . . . . .	64, 79
diminution or loss of . . . . .	164
Myalgia . . . . .	253, 409, 896
Myasthenia gravis pseudo-paralytica . . . . .	105
Myelitis . . . . .	126, 132
acute, contrasted with acute ascending paralysis . . . . .	129
contrasted with multiple neuritis . . . . .	129
central . . . . .	134
disseminated . . . . .	134
from compression . . . . .	134
hemorrhagic . . . . .	134
transverse . . . . .	134
Myocarditis . . . . .	476
acute . . . . .	477
associated with gonorrhœa . . . . .	477
chronic . . . . .	478
rheumatic . . . . .	892
Myoclonus multiplex . . . . .	222
Myopia . . . . .	88, 96
Myotone . . . . .	234
Myxedema . . . . .	251, 834, 1019
distinguished from acute nephritis . . . . .	834
from contracted kidney . . . . .	834
from scleroderma . . . . .	1019
uterine hemorrhage in . . . . .	834
Myxomata of larynx . . . . .	294

N.

Nails, appearance of, in disease, 363, 417, 427, 1039	
---	--

- Nails in typhoid fever relapses . . . 923  
   shedding of . . . . . 1013  
 Narcolepsy distinguished from  
   trance . . . . . 205, 214  
 Narcotics, insensibility from . . . 202  
   poisoning by . . . . . 1032  
 Nasal catarrh . . . . . 262  
   hydrorrhœa . . . . . 262  
   polypi . . . . . 265  
 Nausea as a symptom . . . . . 558  
 Neck, cellulitis of . . . . . 534  
 Nematoda . . . . . 1050  
 Nephralgia . . . . . 785  
   confounded with colic . . . 602, 786  
   from passage of calculi . . . 785  
 Nephritis . . . . . 784  
   acute, parenchymatous . . . 791  
   confounded with acute  
     Bright's disease . . . . 794  
     with acute cystitis . . . 815  
     with myxœdema . . . . . 834  
   bacillosa interstitialis primaria 793  
   chronic consecutive . . . . 805  
     distinguished from Bright's  
       disease . . . . . 805  
   infectious . . . . . 705, 793  
   interstitial . . . . . 810  
   suppurative . . . . . 794  
 Nerve-storm . . . . . 256  
   cardiac . . . . . 453  
 Nerves, diseases of . . . . . 57  
   paralysis from affections of . . 104  
   wounds of . . . . . 174  
 Nervous affections, classification  
   of . . . . . 177  
   deranged nutrition and se-  
     cretion in . . . . . 174  
   centres, diseases of, anæsthesia  
     a symptom of . . . . . 75  
     paralysis from . . . . . 104  
   system, diseases of . . . . . 67  
 Nettle-rash . . . . . 1004  
 Neuralgia . . . . . 252  
   abdominal . . . . . 605, 733  
     due to aneurism . . . . . 733  
   as a cause of headache . . . . 82  
   cerebral . . . . . 256  
   confounded with aneurism . . . 733  
     with pain of rheumatism, 253, 894  
   dorso-intercostal . . . . . 174, 605  
   epileptiform . . . . . 255  
   facial . . . . . 254  
     distinguished from painful  
       anæsthesia of fifth nerve 255  
     from spasm of face . . . . 255  
   hepatic . . . . . 601  
   in Bright's disease . . . . . 801  
   intercostal . . . . . 410, 451  
     confounded with acute pleu-  
       risy . . . . . 410  
 Neuralgia, lumbo-abdominal . . . 605  
   of frontal sinus disease . . . . 261  
   of kidney . . . . . 786  
   of spinal nerves confounded  
     with colic . . . . . 605  
   of stomach . . . . . 566  
   ovarian . . . . . 604  
   reflex . . . . . 253  
   supra-orbital . . . . . 260  
 Neurasthenia . . . . . 240  
   associated with neuralgia . . . 254  
 Neuritis . . . . . 252, 257, 921  
   acute progressive . . . . . 127  
   ascending . . . . . 105, 137  
   multiple . . . 128, 164, 528, 833, 1044  
     distinguished from acute  
       ascending paralysis . . . 129  
       from acute myelitis . . . 129  
       from locomotor ataxia . . 129  
       from rheumatism . . . . 129  
   optic . . . . . 95, 141, 180, 241, 244  
 Neuroses, reflex . . . . . 96  
 Neurosis, cardiac . . . . . 450  
   cutaneous . . . . . 1012  
 Night-blindness . . . . . 96  
 Night palsy . . . . . 840  
   terrors . . . . . 69  
 Nodules, connected with rheuma-  
   tism . . . . . 893  
   in leprosy . . . . . 1017  
   subcutaneous fibrous . . . . . 462  
   syphilitic, in paralysis . . . . 143  
     of lung . . . . . 375  
     of tongue . . . . . 521  
 Nose, diphtheria of . . . . . 262  
   diseases of . . . . . 260  
   foreign body in . . . . . 261  
   gonorrhœal infection of . . . . 264  
   hemorrhage from . . . . . 263, 706  
 Numbness of extremities . . . . 80, 164  
 Nutrition, deranged . . . . . 174  
 Nystagmus . . . . . 119, 173, 222
- O.**
- Occlusion, vascular . . . . . 201, 237  
 Occupation neurosis . . . . . 223  
 Oedema, angio-neurotic . . . . . 176  
   distinguished from erysip-  
     elas . . . . . 998  
   myxoid . . . . . 276  
   of forehead and eyelids . . . . 243  
   of the ankles . . . . . 584, 830  
   of the larynx . . . . . 277  
     diagnosed from croup . . . 278  
   pulmonary . . . . . 393, 796  
     mistaken for pneumonia . . 893  
     occurring in Bright's disease 796  
 Œsophagus, auscultation of . . . 540  
   cancerous narrowing of . . . 539

Esophagus, cicatrices of . . . . .	539
compression of . . . . .	539
dilatation of . . . . .	539
diseases of . . . . .	519, 537
inflammation of . . . . .	538
rupture of . . . . .	539
stricture of . . . . .	514, 538, 586
spasmodic . . . . .	538
thickening of, from alkaline poisoning . . . . .	1029
Omentum, cancer of . . . . .	590, 696
tumor of . . . . .	589
Ophthalmia . . . . .	90
Ophthalmoplegia . . . . .	94
Ophthalmoscope in diseases of the nervous system . . . . .	89, 94
Opisthotonus . . . . .	229, 232
Opium poisoning . . . . .	1032, 1038
Optic nerve, atrophy of . . . . .	95
neuritis . . . . .	95, 141, 180, 241, 244
tract, diseases of the . . . . .	118
Orthopnea . . . . .	30, 331
Orthotonos . . . . .	229
Osteitis deformans . . . . .	251
Otitis from inveterate smoking . . . . .	1040
Ovarian cysts . . . . .	700, 711
dropsy confounded with ascites . . . . .	711
fluid, chemical character of . . . . .	713
inflammation . . . . .	604, 627
neuralgia . . . . .	605
tumors simulating renal growths . . . . .	728
Oxalate of lime in the urine, . . . . .	575,
. . . . .	757, 786
Oxaluria . . . . .	758, 795
Oxyuris vermicularis . . . . .	1051

**P.**

Pachymeningitis, hemorrhagic, . . . . .	181, 199
spinalis interna . . . . .	132
Paget's disease . . . . .	251
Pain as a symptom . . . . .	55
abdominal, in enteritis . . . . .	607
in typhoid . . . . .	918
cardiac . . . . .	448
from organic structural disease . . . . .	894
gastric, as a symptom . . . . .	565
in appendicitis . . . . .	623, 625
in diseases of the liver . . . . .	661, 702
in laryngeal affections . . . . .	267
paroxysmal, diseases characterized by . . . . .	252
Palate, paralysis of . . . . .	528
Pallor of malignant disease . . . . .	862
Palpation of the chest . . . . .	303
Palpitation, cardiac, diseases attended with . . . . .	452

Palsy. See <i>Paralysis</i> .	
associated with typhoid fever . . . . .	921
Bell's . . . . .	121, 144
by compression . . . . .	146
cerebral . . . . .	156
facial . . . . .	140, 144
double . . . . .	146, 148
functional . . . . .	76, 126
hysterical . . . . .	139
lead . . . . .	141
limited . . . . .	121, 138
local . . . . .	144
motor . . . . .	121
shaking . . . . .	167, 248
wasting . . . . .	149
Pancreas, abscess of . . . . .	615
calculous disease of . . . . .	722
diseases of . . . . .	721
fatty diarrhœa in . . . . .	653
ulcerating . . . . .	565
uniform simple hardening of . . . . .	721
Pancreatic fat necrosis . . . . .	721
Pancreatitis, acute . . . . .	614
confounded with peritonitis . . . . .	614
chronic . . . . .	721
hemorrhagic . . . . .	614, 641
suppurative . . . . .	721
Papillitis . . . . .	95, 244
Papilloma . . . . .	264, 272, 294
Papular diseases . . . . .	1005
Paracentesis . . . . .	423
Paræsthesia . . . . .	80
Paralysis . . . . .	64, 65, 66, 75, 103
See also <i>Palsy</i> .	
ascending . . . . .	126
contrasted with acute myelitis . . . . .	129
with multiple neuritis . . . . .	129
acute atrophic . . . . .	155
agitans . . . . .	167, 248, 902
distinguished from chorea . . . . .	222
from general paralysis . . . . .	248
from rheumatism . . . . .	902
bulbar . . . . .	147, 152
from compression . . . . .	146
clinical investigation of . . . . .	107
convulsions in . . . . .	117
crossed . . . . .	114, 116
diphtheritic . . . . .	141, 164, 528
from multiple neuritis . . . . .	142
electro-muscular contractility . . . . .	
in . . . . .	109
sensitivity in . . . . .	112
essential . . . . .	155
eye symptoms in . . . . .	91, 116
facial . . . . .	144
following measles . . . . .	982
from affection of nerves at their extremities . . . . .	104
from apoplexy . . . . .	194
from chronic softening . . . . .	235

- Paralysis from interference with  
the circulation . . . . . 105  
from lead poisoning . 105, 141, 151  
from lesion in the course of a  
nerve . . . . . 104  
from lesion of crus cerebri . . 117  
from lesion of nervous centres . 103  
from lesion of spinal cord . . 124  
from morbid state of the mus-  
cles . . . . . 105  
from poisoning . . . . . 105  
from progressive muscular  
atrophy . . . . . 149  
from reflex action . . . . . 104  
functional . . . . . 104, 106  
general . . . . . 103, 246  
distinguished from other pal-  
sies . . . . . 247  
of insane . . . . . 162, 247  
glosso-labio-laryngeal . . . . 148  
hereditary . . . . . 104  
hyperpyrexia in . . . . . 116  
hysterical . . . . . 104, 138  
infantile . . . . . 155  
intermitting . . . . . 106  
local . . . . . 144, 149, 151  
malarial . . . . . 106  
motor, from exposure to cold . 104  
of individual muscles . . . . 288  
of nerves of the arm . . . . 146  
of leg only . . . . . 122  
of arms . . . . . 121, 134  
of vocal apparatus . . . . 289, 333  
partial . . . . . 103  
periodic . . . . . 106  
peripheral . . . . . 104  
pseudo-hypertrophic muscular,  
31, 154, 478  
reflex . . . . . 104  
rheumatic . . . . . 140  
spastic spinal . . . . . 135  
spinal, general . . . . . 125, 151  
sudden, distinguished from  
apoplexy . . . . . 125, 204  
syphilitic . . . . . 137, 142  
from inherited taint . . . . 144  
tabular view of . . . . . 167  
tremor in . . . . . 167  
vaso-motor . . . . . 115  
with muscular wasting . . . 149  
without coma . . . . . 204  
Paramyoclonus multiplex . . . 173  
Paramyotone . . . . . 234  
Paraplegia . . . . . 124, 133  
ataxic . . . . . 163  
cervical . . . . . 134  
following accidents . . . . 140  
from hypnotic suggestion . . 140  
from spinal hemorrhage . . . 125  
gradual . . . . . 130  
Paraplegia, hysterical . . . . . 75  
reflex, from intestinal worms . 137  
seat of lesion in . . . . . 124  
spastic . . . . . 163  
sudden . . . . . 125  
Parasites . . . . . 1027, 1048  
animal . . . . . 595, 1050  
diseases caused by . . . . . 1020  
fly . . . . . 1056  
in hæmaturia . . . . . 777  
in sputum . . . . . 341  
in stomatitis . . . . . 520  
intestinal . . . . . 855  
vegetable . . . . . 1049  
Paresis . . . . . 103  
spastic . . . . . 153  
spinal . . . . . 126  
Parotitis . . . . . 534  
associated with pneumonia . . 390  
secondary . . . . . 523, 534  
See also *Mumps*.  
Patellar tendon reflex . . . . 101  
Pectoriloquy, whispering . . . 325  
Pellagra . . . . . 1020  
Pelvic cellulitis . . . . . 728  
hæmatocele . . . . . 628  
Pemphigus . . . . . 74, 1010  
foliaceus . . . . . 1010  
syphilitic . . . . . 1010  
Peptonuria in phosphorus poi-  
soning . . . . . 1029, 1045  
Percussion . . . . . 303  
auscultatory . . . . . 307  
clearness of, as a diagnostic  
sign . . . . . 348  
dulness of, diseases accom-  
panied by . . . . . 361  
in pneumonia . . . . . 396  
mediate . . . . . 303  
of abdominal viscera . . . . 544  
of healthy chest . . . . . 309  
respiratory . . . . . 308  
sounds elicited by . . . . . 304  
Percussion hammer . . . . . 306  
Perforation, intestinal, confound-  
ed with colic . . . . . 600  
Periarteritis . . . . . 836  
nodosa . . . . . 836, 1064  
Pericardial effusion . . . . 422, 469, 488  
diagnostic signs of . . . . . 470  
mistaken for dilatation of  
heart . . . . . 488  
murmurs . . . . . 444  
Pericarditis, acute . . . . . 464, 468  
caused by scurvy . . . . . 885  
diagnosed from endocar-  
ditis . . . . . 471  
from gastric irritation . . . 473  
from inflammation of brain . 473  
from pleuritis . . . . . 471



- Pericarditis, cancerous . . . . . 474  
friction sounds of . . . . . 464  
head-symptoms of, confounded  
with meningitis . . . . . 185  
hemorrhagic . . . . . 474  
in Bright's disease . . . . . 796  
indurated mediastino- . . . . . 475  
tubercular . . . . . 474  
Pericardium, adhesions of . . . . . 475  
dropsy of . . . . . 473  
effusion of, confounded with  
chronic pleurisy . . . . . 422  
ulcerative perforation of . . . . . 476  
Perihepatitis . . . . . 668  
Perinephritis . . . . . 818  
distinguished from inflamma-  
tion of psoas muscle . . . . . 819  
Periosteum, rheumatism of . . . . . 897  
Periostitis . . . . . 256  
Peripheral irritation . . . . . 105  
Peritoneum, abscess of . . . . . 730  
carcinoma of . . . . . 715, 729  
colloid cancer of . . . . . 730  
diseases of . . . . . 593, 714, 730  
fatty tumor of . . . . . 730  
hydatid disease of . . . . . 730  
perforation of . . . . . 600  
sarcoma of . . . . . 730  
Peritonitis, acute . . . . . 606, 609  
associated with acute pancre-  
atitis . . . . . 614  
confounded with abdominal  
hysteria . . . . . 619  
with acute enteritis . . . . . 614  
with acute gastritis . . . . . 613  
with colic . . . . . 606, 620  
with cystitis . . . . . 615  
with distention of bladder . . . . . 615  
with inflammation and ab-  
scess of abdominal mus-  
cles . . . . . 616  
with intestinal obstruction . . . . . 631  
with metritis . . . . . 615  
with rheumatism of ab-  
dominal walls . . . . . 618  
with typhoid fever . . . . . 926  
chronic . . . . . 621, 708, 714  
attended with ascites . . . . . 622  
cancerous deposits in . . . . . 622  
distinguished from dropsical  
effusion . . . . . 714  
from collections of pus in the  
cavity . . . . . 617  
from extravasation into the sac . . . . . 611  
local . . . . . 612  
perforative . . . . . 611  
puerperal . . . . . 612  
tubercular . . . . . 589, 622, 714  
Perityphlitis . . . . . 626  
Pernicious anæmia . . . . . 857  
Pernicious anæmia associated with  
alterations in spinal cord . . . . . 861  
confounded with Addison's dis-  
ease . . . . . 872  
with contracted kidney . . . . . 862  
with disease of heart . . . . . 862  
with organic disease of  
stomach . . . . . 862  
distinguished from ordinary  
anæmia . . . . . 863  
from chlorosis . . . . . 863  
from leukæmia . . . . . 863  
from pseudo-leukæmia . . . . . 863  
state of blood in . . . . . 859  
Petit mal . . . . . 215, 218  
Pettenkofer's test . . . . . 761  
Phantom tumors . . . . . 723  
Pharyngeal fever . . . . . 999  
tonsil . . . . . 263  
Pharyngitis . . . . . 275  
confounded with diphtheria . . . . . 529  
Pharyngo-mycosis . . . . . 537  
Pharynx and œsophagus, diseases  
of . . . . . 519, 537  
adenoid vegetations in . . . . . 857  
Phlebitis . . . . . 477, 624, 839, 878  
Phlegmasia alba dolens, 831, 839,  
856, 857, 878  
associated with chlorosis . . . . . 891  
confounded with rheumatism . . . . . 890  
Phosphates in the urine . . . . . 752  
Phosphatic diathesis . . . . . 753  
Photophobia . . . . . 337, 900  
Phthetiriasis . . . . . 1005, 1006, 1020  
Phthisis . . . . . 361  
See also *Tuberculosis of Lungs*.  
acute . . . . . 384, 396, 928  
distinguished from menin-  
gitis . . . . . 387  
from typhoid fever . . . . . 387, 928  
acute pneumonic . . . . . 387  
bronchial . . . . . 337  
cavity from . . . . . 379  
distinguished from pulmo-  
nary abscess . . . . . 380  
chronic . . . . . 613  
confounded with actinomy-  
cosis . . . . . 381  
with bronchial dilatation . . . . . 377  
with chronic bronchitis . . . . . 369  
with chronic pleurisy . . . . . 374, 418  
with chronic pneumonic con-  
solidation . . . . . 370, 372  
with emphysema . . . . . 369  
with phthisis of old people . . . . . 370  
with pulmonary abscess . . . . . 378  
with pulmonary cancer . . . . . 375  
with pulmonary gangrene . . . . . 380  
with syphilitic disease of the  
lungs . . . . . 375

- Phthisis, cough in . . . . . 361  
   fibroid . . . . . 425  
   laryngeal . . . . . 292  
   of old people . . . . . 370  
   pneumonic . . . . . 373, 387  
   pulmonary . . . . . 338, 418  
   rapid progress of . . . . . 388  
   retrogression of . . . . . 382  
   symptoms of . . . . . 368  
   temperature in . . . . . 363  
 Physical diagnosis . . . . . 297, 299  
   signs . . . . . 297  
 Picric acid test . . . . . 770  
 Pigment in the blood . . . . . 883, 960  
   liver . . . . . 669  
 Pityriasis capitis . . . . . 1023  
   maculata et circinata . . . . . 1013  
   rosea . . . . . 1013  
   rubra . . . . . 1008, 1013  
   versicolor . . . . . 871, 1022  
 Plague distinguished from ty-  
   phus fever . . . . . 936  
   from yellow fever . . . . . 973  
 Plasmodium malarie . . . . . 876  
 Pleura, cancer of . . . . . 420  
   effusion into . . . . . 422  
   fistula of . . . . . 427  
   friction sound in . . . . . 472  
   liquid in . . . . . 472  
 Pleurisy, acute . . . . . 360, 396, 404  
   confounded with acute  
     Bright's disease . . . . . 796  
     with acute pneumonia . . . . . 407  
     with intercostal neural-  
       gia . . . . . 410  
     with pericarditis . . . . . 422, 472  
     with pleurodynia . . . . . 409  
   bilious . . . . . 403  
   chronic . . . . . 374, 415, 423  
     confounded with abscess in  
       thoracic walls . . . . . 422  
     with cancer . . . . . 420, 424  
     with chronic interstitial  
       pneumonia . . . . . 424  
     with cirrhosis of lung . . . . . 425  
     with collapse of lung, 359, 426  
     with emphysema . . . . . 419  
     with enlargement of liver . . . . . 421  
     with enlargement of  
       spleen . . . . . 421  
     with fistula of pleura . . . . . 427  
     with hydatid cysts . . . . . 421  
     with hydrothorax . . . . . 422  
     with intra-thoracic tumor . . . . . 419  
     with pericardial effusion . . . . . 422  
     with phthisis . . . . . 418, 425  
     with pneumothorax . . . . . 415  
     with tubercle . . . . . 416, 424  
   diseases confounded with . . . . . 419  
   circumscribed . . . . . 420  
 Pleurisy, diaphragmatic, con-  
   founded with acute hepa-  
   titis . . . . . 670  
   double . . . . . 374  
   dry . . . . . 405, 426,  
     associated with typhoid fever . . . . . 928  
   different forms of . . . . . 418  
   fluid of, microscopical and  
     bacteriological examina-  
     tion of . . . . . 418  
 Pleuritic effusion . . . . . 358, 374, 405, 685  
 Pleurodynia . . . . . 316, 409  
   confounded with acute pleurisy . . . . . 409  
 Pleurothotonos . . . . . 229  
 Plica polonica . . . . . 1024  
 Pneumatometry . . . . . 303  
 Pneumococcus of Fraenkel . . . . . 344, 398  
   of Friedlaender . . . . . 400  
 Pneumo-hydro-pericardium . . . . . 476  
 Pneumonia acute . . . . . 388  
   confounded with acute bron-  
   chitis . . . . . 396  
   with acute phthisis . . . . . 396  
   with acute pleurisy . . . . . 407  
   with appendicitis . . . . . 630  
   with bilious pneumonia . . . . . 403  
   with cerebro-spinal fever . . . . . 942  
   with hypostatic congestion . . . . . 394  
   with pulmonary apoplexy . . . . . 394  
   with pulmonary engorge-  
     ment in fevers . . . . . 393  
   with pulmonary edema . . . . . 393  
   with typhoid pneumonia . . . . . 401  
   head-symptoms of, confound-  
   ed with meningitis . . . . . 185  
   apex . . . . . 399  
   associated with measles . . . . . 983  
   with typhoid fever . . . . . 928  
   with typhus fever . . . . . 934  
   with ulcerative endocarditis . . . . . 399  
   auscultation in . . . . . 390  
   bilious . . . . . 401  
   broncho . . . . . 351, 360, 396, 934  
     following hemorrhage from  
       cavities . . . . . 398  
   caseous . . . . . 338  
   catarrhal . . . . . 351, 396  
   chronic, confounded with  
     phthisis . . . . . 370  
     catarrhal . . . . . 373  
   croupous . . . . . 388, 398  
   deglutition . . . . . 398  
   delirium in . . . . . 68  
   dissecting . . . . . 380  
   double . . . . . 399  
   gangrenous . . . . . 294  
   hypostatic . . . . . 394  
   interstitial . . . . . 424  
   latent . . . . . 400  
   lobar . . . . . 360

- Pneumonia, malarial . . . . . 403  
   massive . . . . . 408  
   migratory . . . . . 400  
   physical signs of . . . . . 390  
   respiration in . . . . . 330  
   tuberculous inspiration bron-  
     cho- . . . . . 398  
   typhoid . . . . . 401  
     articular symptoms of . . . . . 402  
 Pneumothorax . . 411, 476, 580, 686  
   chest sounds in . . . . . 413  
   diagnosed from chronic  
     pleurisy . . . . . 419  
     from diaphragmatic hernia . . 414  
     from emphysema . . . . . 358  
     from pneumo-pericardium . . 476  
     without perforation . . . . . 415  
 Pneumotypus . . . . . 934  
 Podelcoma . . . . . 1049  
 Poisoning, aconite . . . . . 1036  
   acute . . . . . 1027  
   alcohol . . . . . 1033, 1039  
     urinary test for . . . . . 1034  
   alkaline . . . . . 1029  
   aloes . . . . . 1031  
   ammonium . . . . . 538, 1029  
   aniline . . . . . 1035  
   antimony . . . . . 664, 1030  
   antipyrine . . . . . 1036  
   arsenic . . . 128, 609, 1030, 1043  
   atropine . . . . . 1034  
   belladonna . . . . . 1034  
   benzene . . . . . 1033  
   bromine . . . . . 1029  
   brucine . . . . . 1037  
   by poisonous exhalations . . . 1047  
   by ptomaines . . . . . 1047  
   Calabar bean . . . . . 1037  
   cantharides . . . . . 1030  
   carbolic acid . . . . . 1034  
   carbon disulphide . . . . . 1044  
     monoxide or carbonic oxide,  
       762, 1035  
   charcoal fumes . . . . . 1035  
   cheese, egg, milk . . . . . 1031  
   chloral . . . . . 1033, 1039  
   chlorine . . . . . 1029  
   chloroform . . 202, 664, 1033, 1039  
   chronic . . . . . 1038  
   coal gas . . . . . 1035  
   colchicum . . . . . 1031  
   colocynth . . . . . 1031  
   conium . . . . . 1034  
   copper . . . . . 598, 664, 1030, 1048  
   corrosive sublimate . . . . . 1030  
   cream puff . . . . . 1031  
   diazobenzene . . . . . 1031  
   digitalis . . . . . 1036  
   elaterium . . . . . 1031  
   ergot . . . . . 1031, 1040  
 Poisoning, ether . . 664, 1033, 1039  
   from alkaloids . . . . . 1047  
   from animal effluvia . . . . . 1048  
   fungi . . . . . 1031  
   hydrochloric acid . . . . . 1028  
   hydrocyanic acid . . . 203, 1035  
   hyoscyamus . . . . . 1034  
   iodine . . . . . 1029  
   iron . . . . . 1030  
   irritant . . . . . 656, 660, 1028  
   lead, 287, 598, 609, 810, 1030, 1041  
   lobelia . . . . . 1031  
   malarial . . . . . 831  
   mercurial . . . . . 609, 1030, 1041  
   muscarine . . . . . 1031  
   mushroom . . . . . 1032  
   narcotic . . . . . 287, 797, 1032  
     insensibility from, distin-  
       guished from alcohol-  
       ism . . . . . 1033  
       from apoplexy . . . 201, 1032  
       from uræmia . . . 797, 1032  
   nitric acid . . . . . 538, 1028  
   nitrobenzole . . . . . 202  
   nitro-glycerine . . . . . 1036  
   opium . . . . . 1032, 1038  
   oxalic acid . . . . . 1029  
   paraldehyde . . . . . 1039  
   petroleum . . . . . 1036  
   phosphorus, 664, 679, 780,  
     1029, 1044  
   picrotoxin . . . . . 1037  
   potassium hydroxide . . . . . 1029  
     iodide and nitrate . . . . . 1029  
   producing anæsthesia . . . . . 74  
     coma . . . . . 71  
     headache . . . . . 83  
     paralysis . . . . . 105  
   prussic acid . . . . . 1035  
   sausage . . . . . 1031  
   savin . . . . . 1031  
   sewer-gas . . . . . 791  
   silver . . . . . 1030  
   slow, by metals . . . . . 1041  
   sodium hydroxide . . . . . 1029  
   strychnine . . . . . 1037  
     confounded with epilepsy . . 1037  
     with hydrophobia . . . . . 1037  
     with tetanus . . . . . 232, 1037  
   sulphuric acid . . . . . 538, 1028  
   tobacco . . . . . 1031, 1040  
   tyrotoxicon . . . . . 1031  
   veratrum viride . . . . . 1037  
   zinc . . . . . 1030, 1044  
 Poisons . . . . . 1027  
   animal, diseases caused by . . 1045  
   irritant . . . . . 1028  
 Poliomyelitis . . . . . 134, 174  
   acute anterior . . . . . 156  
 Polyæsthesia . . . . . 79

- Polyarthritis . . . . . 276  
 Polypi, nasal . . . . . 264, 265, 332  
   of larynx . . . . . 294  
 Polyuria . . . . . 826  
   chronic, distinguished from  
     true diabetes . . . . . 827  
 Porencephalus . . . . . 250  
 Porrigo larvalis . . . . . 1011  
 Portal veins, inflammation of,  
   confounded with acute  
     hepatitis . . . . . 669  
   with cirrhosis of liver . . . . . 708  
   inflammation of, with coagula . . . . . 708  
   thrombosis of . . . . . 709  
 Position as a symptom . . . . . 30  
 Posterior sclerosis . . . . . 135, 160  
 Potassium ferrocyanide test . . . . . 770  
 Pregnancy, extra-uterine, mis-  
   taken for appendicitis . . . . . 628  
 Presbyopia . . . . . 88, 89  
 Pressure-points . . . . . 233  
 Progressive muscular atrophy,  
   149, 156  
   distinguished from bulbar pa-  
     ralysis . . . . . 152  
   from cerebral hemiplegia . . . . . 150  
   from general spinal paraly-  
     sis . . . . . 151  
   from idiopathic atrophy . . . . . 154  
   from infantile paralysis . . . . . 156  
   from local paralysis . . . . . 151  
   from pseudo-hypertrophic  
     muscular paralysis . . . . . 154  
   from syringo-myelitis . . . . . 152  
   from unilateral atrophy of  
     the face . . . . . 152  
 Prostate gland, hemorrhage from . . . . . 778  
 Prurigo . . . . . 1006  
 Pruritus . . . . . 1006, 1025  
   hiemalis . . . . . 1025  
 Pseudo-disseminated sclerosis . . . . . 169  
 Pseudo-leukæmia . . . . . 863  
 Pseudo-scarlatina . . . . . 981  
 Pseudo tabes mesenterica . . . . . 556, 724  
 Psoas muscle, inflammation of . . . . . 819  
 Psoriasis . . . . . 1006, 1013, 1014  
   distinguished from eczema  
     squamosum . . . . . 1014  
   from lichen planus . . . . . 1006, 1014  
   syphilitic . . . . . 1014  
 Ptomaines . . . . . 83, 641, 676  
 Ptoxis . . . . . 92  
 Puerperal malarial fever . . . . . 953  
 Pulmonary affections, con-  
   founded with typhoid  
     fever . . . . . 927  
   disease, physical signs of . . . . . 299  
   engorgement in fevers . . . . . 393  
 Pulsation, abdominal . . . . . 731  
   aortic . . . . . 731  
   Pulsation, aortic, confounded  
     with aneurism of abdomi-  
       nal aorta . . . . . 733  
 Pulse, condition of, in disease . . . . . 35  
   dicrotic . . . . . 38, 922  
   frequency and rhythm of . . . . . 35, 36  
   irregular . . . . . 36  
   resistance of . . . . . 37  
   respiration-ratio, perverted . . . . . 388  
   strength and volume of . . . . . 36  
 Pulsus alternans . . . . . 36  
   paradoxus . . . . . 475  
 Purging, diseases attended by,  
   643, 653  
 Purpura . . . . . 886  
   acute, distinguished from cere-  
     bro-spinal meningitis . . . . . 887  
   from hæmophilia . . . . . 887  
   from scurvy . . . . . 896  
   hæmorrhagica and rheumat-  
     ica . . . . . 886  
 Purulent urine . . . . . 794  
   diseases associated with . . . . . 814  
 Pus in internal cavities . . . . . 951  
   in stools . . . . . 593  
   in urine . . . . . 779  
   in vomit . . . . . 562  
   presence of, in appendicitis . . . . . 624  
 Pustular diseases . . . . . 1011  
 Pustule, malignant . . . . . 1045  
 Pyæmia . . . . . 872  
   arterial . . . . . 875  
   associated with myocarditis . . . . . 477  
   chronic or relapsing . . . . . 875  
   confounded with acute affec-  
     tions of liver . . . . . 874  
   with acute glands or acute  
     farcy . . . . . 874  
   with intermittent fever . . . . . 952  
   with rheumatic fever . . . . . 874  
   with typhoid fever . . . . . 873  
   idiopathic . . . . . 875  
   joint-affection of . . . . . 892  
   metastatic or embolic abscesses  
     of . . . . . 874  
   spontaneous septico- . . . . . 875  
 Pyelitis . . . . . 780, 794, 819  
   catarrhal or rheumatic . . . . . 821  
   from irritation of calculi . . . . . 821  
   tuberculous . . . . . 821  
 Pylorus, cancer of . . . . . 586  
   fibroid thickening of . . . . . 587  
 Pyonephrosis . . . . . 821  
   confounded with abscess of the  
     kidney . . . . . 822  
   with suppurative nephri-  
     tis . . . . . 822  
 Pyopneumothorax, subphrenic,  
   580, 686  
 Pyrosis . . . . . 561



## Q.

Quinsy, distinguished from secondary parotitis . . . . .	523
from tonsillitis . . . . .	522

## R.

Rachitis . . . . .	249
Radial nerve, paralysis of . . . . .	146
Rales . . . . .	321, 390
varieties of . . . . .	322
Rash, mulberry, of typhus . . . . .	930
of scarlet fever . . . . .	977
rose . . . . .	467
Raynaud's disease . . . . .	776, 840
mistaken for chilblains . . . . .	841
Records of cases, plans for . . . . .	29
Red gum of infants . . . . .	1024
Reflex, abdominal . . . . .	99
arc . . . . .	99
cremaster . . . . .	99
epigastric . . . . .	99
excitability . . . . .	112, 133
irritability . . . . .	134
paraplegia . . . . .	137
tendo Achillis . . . . .	100, 247
Reflexes, aural . . . . .	100
cranial . . . . .	100
crossed . . . . .	101
deep . . . . .	100, 132
derangements of . . . . .	99, 150
laryngeal and pharyngeal, . . . . .	100, 272
nasal . . . . .	100
reinforcement of . . . . .	102
superficial . . . . .	99
tendon . . . . .	99
Regurgitation of fluid or food . . . . .	565
Relapsing fever . . . . .	944
bilious typhoid form of . . . . .	946
distinguished from typhoid and typhus fever . . . . .	947
from Weil's disease . . . . .	672
from yellow fever . . . . .	947, 973
renal disease in . . . . .	946
spirilla of . . . . .	947
Remittent fever . . . . .	954
bilious . . . . .	974
distinguished from acute congestion of the liver . . . . .	956
from acute meningitis . . . . .	958
from intermittent fever . . . . .	956
from typhoid fever . . . . .	957
from yellow fever . . . . .	974
infantile . . . . .	961
sequelæ of . . . . .	959
Renal abscess . . . . .	794
artery, multiple aneurisms of . . . . .	823
asthma . . . . .	334
colic . . . . .	786, 822
concretions, forms of . . . . .	786

Renal concretions, passage of, 602, 789	
cysts . . . . .	823
disease in relapsing fever . . . . .	946
enlargements . . . . .	700
growths simulated by ovarian tumors . . . . .	728
hæmaturia . . . . .	774
inadequacy . . . . .	806
vein, thrombosis of . . . . .	828
Respiration, amphoric . . . . .	320, 476
bronchial . . . . .	313, 318
broncho-cavernous . . . . .	320
cavernous . . . . .	319
Cheyne-Stokes . . . . .	195, 335, 449
feeble . . . . .	314
harsh . . . . .	317
in children, peculiarities of . . . . .	328
jerkings . . . . .	316
metallic . . . . .	320
metamorphosing breath-sound . . . . .	320
prolonged . . . . .	316
puerile . . . . .	314
sounds of, in health . . . . .	313
supplementary . . . . .	314
vesiculo-bronchial . . . . .	317
vesiculo-cavernous . . . . .	320
Respiratory movements . . . . .	299
percussion . . . . .	308
Retina, embolism of central artery of . . . . .	95
Retinal hemorrhage . . . . .	94, 95
Retinitis, albuminuric . . . . .	95, 801
anæmic . . . . .	95
diabetic . . . . .	95, 824
leukæmic . . . . .	95
pigmentosa . . . . .	94
Retropharyngeal abscesses . . . . .	283, 537
Rheumatic arthritis . . . . .	900
fever distinguished from pyæmia . . . . .	874
gout . . . . .	900
paralysis . . . . .	140
Rheumatism . . . . .	129, 253, 889
acute articular . . . . .	461, 889
confounded with acute synovitis . . . . .	890
with cerebro-spinal fever . . . . .	943
with milk-leg . . . . .	890
with rickets . . . . .	905
head-symptoms of, confounded with meningitis . . . . .	184, 892
heart-symptoms in . . . . .	892
cerebral . . . . .	893
chronic . . . . .	894
confounded with abdominal aneurism . . . . .	733
with neuralgia . . . . .	253, 894
with pain of organic structural disease . . . . .	894

- Rheumatism, chronic, con-  
     founded with paralysis  
         agitans . . . . . 902  
         with sciatica . . . . . 258  
     in Bright's disease . . . . . 802  
     distinguished from dengue . . . . . 996  
     from gout . . . . . 898  
     feigned . . . . . 897  
     gonorrhœal . . . . . 477, 891  
     hyperpyrexia in . . . . . 893  
     muscular . . . . . 894  
         distinguished from achillo-  
         dynia . . . . . 897  
         from Morton's disease . . . . . 897  
         from myalgia . . . . . 896  
         from tetanus . . . . . 230  
         from trichiniasis . . . . . 897, 1063  
     of abdominal walls . . . . . 618  
     of cervical muscles . . . . . 943  
     of scalp . . . . . 256  
     periosteal . . . . . 897  
     relations of, to chorea . . . . . 220  
     subacute . . . . . 893  
 Rheumatoid arthritis . . . . . 900  
 Rhinitis . . . . . 262, 332  
     caseosa and fibrinosa . . . . . 262  
     membranous . . . . . 534  
     œdematosa . . . . . 262  
     scrofulous . . . . . 264  
 Rhinoliths . . . . . 261  
 Rhinoscleroma . . . . . 265  
 Rhinoscopy . . . . . 273  
 Rhythm of respiration, changes  
     in . . . . . 316  
 Rickets . . . . . 250, 513, 902  
     combined with scurvy . . . . . 905  
     confounded with craniotabes . . . . . 905  
     with hereditary syphilis . . . . . 904  
     with mollities ossium . . . . . 904  
     with rheumatism . . . . . 905  
 Rigidity, local, confounded with  
     tetanus . . . . . 230  
 Risus sardonicus . . . . . 228  
 Romberg symptom . . . . . 161  
 Rose-cold . . . . . 262, 354  
 Roseola . . . . . 1004  
 Rötheln . . . . . 985  
 Rubella . . . . . 985  
     distinguished from measles . . . . . 986  
     from scarlet fever . . . . . 986  
     from typhus fever . . . . . 988  
 Rubeola notha . . . . . 984  
     sine catarrho . . . . . 982  
 Rumination . . . . . 565  
 Rupia . . . . . 1012
- S.**
- Salaam convulsions . . . . . 224  
 Salivation . . . . . 519
- Sapræmia . . . . . 877  
 Sarcinæ ventriculi . . . . . 339, 561, 591  
 Sarcoma, mediastinal . . . . . 420, 508  
     of kidney . . . . . 804  
     of larynx . . . . . 294  
 Scabies . . . . . 1009, 1021  
 Scalp, œdema of . . . . . 201  
     rheumatism of, confounded  
         with hemicrania . . . . . 256  
 Scarletina . . . . . 90, 976  
     anginose . . . . . 978  
     associated with rheumatism . . . . . 978  
     complications and sequelæ of . . . . . 979  
     distinguished from cerebro-spi-  
         nal meningitis . . . . . 942  
     from dengue . . . . . 981, 996  
     from laryngeal diphtheria,  
         . . . . . 532, 980  
     from erysipelas . . . . . 998  
     from measles . . . . . 980  
     from membranous croup . . . . . 980  
     from rubella . . . . . 986  
     from smallpox . . . . . 980, 992  
     from typhoid fever . . . . . 980  
     exhaustion in . . . . . 980  
     followed by dropsy . . . . . 979  
     nervous symptoms in . . . . . 978  
     pseudo- . . . . . 981  
     rash of . . . . . 977  
     sine exanthemate . . . . . 977  
     sore throat of . . . . . 977  
     surgical . . . . . 981  
     tongue in . . . . . 978  
     urine of . . . . . 979  
 Scarlet fever. See *Scarlatina*.  
 Sciatica . . . . . 77, 256  
     distinguished from hip-joint  
         affections . . . . . 258  
         from irritation of the kidney . . . . . 258  
         from rheumatism . . . . . 258  
     feigned . . . . . 259  
     pressure of fluid on nerve in . . . . . 258  
     rheumatic . . . . . 257  
 Scirrhus . . . . . 582  
 Scleroderma or sclerema . . . . . 1018  
     distinguished from erysipelas . . . . . 1019  
         from morphea . . . . . 1020  
         from myxœdema . . . . . 1019  
 Sclerosis, cerebro-spinal . . . . . 167, 248  
     diffused . . . . . 239  
     disseminated . . . . . 135, 167  
     lateral amyotrophic . . . . . 135  
     multiple . . . . . 167  
         eye-symptoms in . . . . . 168  
     posterior . . . . . 135, 160  
 Scrofula . . . . . 34, 1005  
     associated with disease of in-  
         testines . . . . . 647  
         with disease of lungs . . . . . 383  
     psoas abscess from . . . . . 628

- Scrofulous glands distinguished  
from lymphadenoma . . . . . 868
- Scurvy . . . . . 346, 646, 884  
combined with rickets . . . . . 905  
confounded with purpura . . . . . 886  
hemorrhage in . . . . . 564  
infantile . . . . . 885  
sore mouth of . . . . . 519
- Seborrhæa . . . . . 1008, 1023
- Secretion, deranged . . . . . 174, 204
- Senile dementia . . . . . 248
- Sensation, deranged . . . . . 72  
gnawing, in vertebræ . . . . . 516
- Sensations of patients . . . . . 55  
tests of . . . . . 77
- Senses, special, derangement of . . . . . 87
- Sensibility, electrical . . . . . 74  
impairment of . . . . . 64  
perverted . . . . . 79, 466
- Sensory centres . . . . . 63
- Septicæmia . . . . . 876  
from absorption of toxines . . . . . 877  
malarial . . . . . 876  
puerperal . . . . . 878  
typhoid . . . . . 876
- Shock . . . . . 33
- Sight, dimness of . . . . . 61
- Sinus, frontal, diseases of . . . . . 260
- Skin, atrophies of . . . . . 1019  
condition of, as a symptom . . . . . 34  
discoloration of, following fe-  
vers . . . . . 871  
from lactation and preg-  
nancy . . . . . 871  
hereditary . . . . . 871  
in Addison's disease . . . . . 870  
in psoriasis . . . . . 872  
of hands and feet in typhoid . . . . . 919
- diseases . . . . . 1000  
bullous . . . . . 1010  
constitutional . . . . . 1002  
erythematous . . . . . 1003  
from altered gland-secretion . . . . . 1023  
nervous . . . . . 1025  
papular . . . . . 1005  
parasitic . . . . . 1020  
pustular . . . . . 1011  
squamous . . . . . 1013  
syphilitic . . . . . 1006, 1025  
vesicular . . . . . 1007  
glazing of . . . . . 74  
hypertrophies of . . . . . 1017  
maculæ of . . . . . 1015  
new growths of . . . . . 1016  
pigmentation of . . . . . 902
- Sleep, protracted, distinguished  
from apoplexy . . . . . 204
- Sleeping sickness . . . . . 214
- Smallpox . . . . . 988  
bacillus of . . . . . 991
- Smallpox, confluent . . . . . 989  
distinguished from erysipelas . . . . . 998  
from measles . . . . . 992  
from scarlet fever, 980, 982, 992  
from varioloid . . . . . 993  
eruption of . . . . . 989  
invasion of . . . . . 988  
malignant . . . . . 991
- Softening of the brain . . . . . 235  
acute, confounded with acute  
meningitis . . . . . 181  
with apoplexy . . . . . 204  
chronic . . . . . 235  
discriminated from abscess . . . . . 238  
from atrophy . . . . . 239  
from congestion . . . . . 236  
from exhaustion of brain-  
power . . . . . 239  
from neurasthenia . . . . . 240  
from tumor . . . . . 240  
paralysis from . . . . . 236  
red . . . . . 235  
relations of, to hemorrhage . . . . . 236  
white . . . . . 235  
of spinal cord . . . . . 135
- Sore throat . . . . . 522  
chronic . . . . . 535  
chronic rheumatic . . . . . 536  
clergyman's . . . . . 535  
follicular . . . . . 535  
in scarlet fever . . . . . 977  
syphilitic . . . . . 536
- Sound, bronchial . . . . . 313, 317  
elicited by percussion . . . . . 304, 412  
Hippocratic, or succussion . . . . . 303  
in chest, adventitious . . . . . 320  
œsophageal . . . . . 540  
splashing, in stomach . . . . . 591  
tympanitic . . . . . 305
- Spasm, asthmatic or bronchial . . . . . 331  
facial, distinguished from  
chorea . . . . . 223  
masticatory, of the face . . . . . 230  
of arterioles . . . . . 840  
of bladder confounded with colic . . . . . 603  
of glottis in croup . . . . . 279  
of vocal cord tensors . . . . . 288
- Spasmodic dorsal tabes . . . . . 135
- Spasms . . . . . 171  
See also *Convulsions*.  
clonic and tonic . . . . . 171  
diseases marked by . . . . . 214  
functional . . . . . 233  
mobile . . . . . 157, 223  
of acute cerebral disease . . . . . 222  
saltatory . . . . . 173
- Spectroscope . . . . . 773
- Sphygmo-chronograph . . . . . 43
- Sphygmogram . . . . . 41  
in gouty heart . . . . . 482

- Sphygmogram of aortic insufficiency . . . . . 498  
   of contracted kidney . . . . . 811  
   of mitral regurgitation . . . . . 498  
   of thoracic aneurism . . . . . 509  
 Sphygmograph . . . . . 39, 433, 482  
   of Dudgeon . . . . . 40  
   of Marey . . . . . 39  
 Spinal anæmia . . . . . 131  
   cord, congestion of . . . . . 131  
   degeneration of columns of . . . . . 861  
   diseases of . . . . . 57  
   gout of . . . . . 900  
   hemorrhage into . . . . . 125  
   inflammation of . . . . . 131  
     distinguished from epidemic cerebro-spinal meningitis . . . . . 941  
   morbid conditions of, as a cause of paraplegia . . . . . 125  
   sclerosis of . . . . . 79, 135  
   softening of . . . . . 135  
   syphilis of . . . . . 143  
     from inherited taint . . . . . 144  
   table of diseases of . . . . . 177  
   tumors of . . . . . 136  
   curvature . . . . . 251  
   irritation . . . . . 131  
   localization . . . . . 65  
   meningitis . . . . . 126, 131  
     acute . . . . . 132  
   paresis . . . . . 126  
   sclerosis . . . . . 135  
     disseminated . . . . . 135  
     lateral amyotrophic . . . . . 136  
     primary . . . . . 135  
 Spine, concussion of . . . . . 134  
   deviation of . . . . . 153  
   in chronic pleurisy . . . . . 416  
   disease of, confounded with  
     aneurism . . . . . 733  
     with colic . . . . . 606  
   irritable . . . . . 131  
 Spirals . . . . . 332, 340, 355  
 Spirilla of relapsing fever . . . . . 946  
 Spirometer . . . . . 302  
 Splanchnoptosis . . . . . 592  
 Spleen, affections of . . . . . 719  
   displacement of . . . . . 725  
   enlargement of . . . . . 421, 689, 998  
     chronic . . . . . 719  
     confounded with chronic pleurisy . . . . . 421  
     distinguished from cancer of kidney . . . . . 720  
   gastric hemorrhage in . . . . . 564  
   infarct in . . . . . 467  
   inflammation of . . . . . 719  
   in typhoid fever . . . . . 918  
   percussion of . . . . . 546  
 Spotted fever . . . . . 937  
 Sputa . . . . . 338  
   albuminous . . . . . 423  
   constituents of . . . . . 339  
   crystals in . . . . . 341, 355  
   elastic fibres in . . . . . 340  
   fibrinous coagula in . . . . . 340  
   microscopical examination of, . . . . . 339, 342  
   nummular . . . . . 361, 378  
   of acute pneumonia . . . . . 388  
   of bronchitis . . . . . 353  
   of phthisis . . . . . 362  
   parasites in . . . . . 341  
   resembling currant-jelly . . . . . 508  
   spirals in . . . . . 332, 340, 355  
 Squamous diseases . . . . . 1013  
 Staphylococcus, 398, 467, 529, 816, 873  
 Starvation from stricture of  
   œsophagus . . . . . 538  
 Station . . . . . 31  
 Stenosis, bronchial . . . . . 296  
   laryngeal . . . . . 293  
   of pylorus . . . . . 589  
 Stethometer . . . . . 301  
 Stethoscope . . . . . 311  
 Stomach, acidity of . . . . . 556, 584  
   acute diseases of . . . . . 571  
   cancer of . . . . . 582, 696  
     contrasted with chronic gastritis . . . . . 585  
     with cirrhosis of liver . . . . . 708  
     with gastric ulcer . . . . . 582  
   catarrh of . . . . . 573  
   chronic affections of . . . . . 574  
   cirrhosis of . . . . . 621  
   cramp of . . . . . 566  
   dilatation of . . . . . 590  
     associated with nervous symptoms . . . . . 591  
     confounded with dilatation of large intestine . . . . . 592  
     connected with tetany . . . . . 231, 591  
   dislocation of . . . . . 590, 592  
   distention of . . . . . 503, 591  
   examination of contents of, . . . . . 552, 575, 584, 591  
   fibroid thickening of . . . . . 589  
   gout in . . . . . 899  
   hemorrhage from, 563, 577, 609, 1028  
   inflammation of . . . . . 571  
   inspection of interior of . . . . . 551  
   irritation of, confounded with pericarditis . . . . . 473  
   lithæmic pain in . . . . . 900  
   membrane of, absorptive activity of . . . . . 555, 589  
   secondary inflammation of . . . . . 574



- Stomach, motor activity of . . . 555  
   neuralgia of . . . 566  
   organic disease of, confounded  
     with pernicious anæmia . . . 862  
   pain in, as a symptom . . . 565, 571  
   palpation of . . . 543  
   percussion of . . . 547  
   perforation of . . . 612  
     distinguished from irritant  
       poisoning . . . 1028  
   peristaltic disturbance of . . . 600  
   physical examination of, in-  
     struments for . . . 551  
   rupture of . . . 172  
   softening of . . . 574  
   suppurative inflammation of . . . 574  
   tests in diseases of, 551, 576,  
     583, 589, 592  
   ulcer of . . . 346, 576  
 Stomatitis, aphthous . . . 520, 1047  
   gangrenous . . . 520  
   mercurial . . . 519  
   ulcerative, confounded with  
     diphtheria . . . 529  
 Stools as symptoms . . . 593  
   examinations of . . . 595  
   fat in . . . 653  
 Strabismus . . . 91, 206, 654  
 Streptococcus . . . 467, 529  
   erysipelatis . . . 998  
   pyogenes . . . 398, 873  
 Streptocytus of Schottelius . . . 1047  
 Stricture of the œsophagus . . . 514, 538  
 Stridor, laryngeal . . . 266, 514  
 Strongylus gigas . . . 1055  
 Strychnine poisoning . . . 1037  
   confounded with tetanus,  
     232, 1037  
 Stupor . . . 67, 70  
   as result of poisoning . . . 70  
   in uræmia . . . 797  
 St. Vitus's dance. See *Chorea*.  
 Sugar in the urine . . . 762  
   tests for . . . 763  
   by fermentation . . . 766  
   by phenylhydrazin . . . 765  
 Sugar of milk . . . 766  
 Sulphates in urine, pathology of . . . 756  
 Sun-bronzing confounded with  
   Addison's disease . . . 870  
 Sun-stroke . . . 210  
   distinguished from apoplexy . . . 210  
 Supra-renal capsules, disease of . . . 872  
 Surface thermometry . . . 45  
 Sweat-glands . . . 1024  
 Sweating, bloody . . . 175, 1024  
   excessive . . . 176, 507  
 Sycosis . . . 1012, 1022  
 Symptoms, feigned . . . 23  
   pathognomonic . . . 21  
 Symptoms, similarity of, in dis-  
   eases . . . 25  
 Syncope . . . 71  
   distinguished from apoplexy . . . 203  
 Synovitis, acute, confounded  
   with acute rheumatism . . . 890  
 Syphilis combined with laryngitis . . . 292  
   constitutional . . . 1014  
   hereditary . . . 34, 904  
   distinguished from rhinoscle-  
     roma . . . 266  
 Syphilitic disease of the bowels . . . 638  
   of the brain . . . 142, 245, 827, 953  
   of the kidney . . . 804  
   of the liver . . . 689, 694  
   of the lungs . . . 362, 375  
   of the skin, 871, 1006, 1010,  
     1014, 1025  
   of the spinal cord . . . 137  
   of the throat . . . 536  
   fever confounded with inter-  
     mittent fever . . . 953  
   stenosis . . . 539  
   ulcers of fauces . . . 536  
 Syphiloderm . . . 1006  
 Syringo-myelitis . . . 152

## T.

- Tabes dorsalis . . . 160, 164  
   See *Locomotor Ataxia*.  
   distinguished from diphtheritic  
     paralysis . . . 164  
   mesenterica . . . 724  
   pseudo-mesenterica . . . 556, 724  
   spasmodic dorsal . . . 135  
 Tachycardia . . . 453  
 Tactile sense, impairment of . . . 78  
 Tænia lata . . . 1054  
   mediocanellata . . . 1053  
   solium . . . 1052  
 Tape-worms . . . 1051  
   of pork . . . 1052  
 Temperature of body as a symp-  
   tom . . . 43  
   cerebral . . . 45, 123  
   extraordinary range of . . . 51  
   in acute encephalitis . . . 238  
   in apoplexy . . . 195  
   in cerebro-spinal fever . . . 939  
   in children . . . 49  
   in cholera . . . 659  
   in hydrophobia . . . 233  
   in intermittent fever . . . 950  
   in measles . . . 50, 981  
   in phthisis . . . 363  
   in pneumonia . . . 50, 389  
   in pyæmia . . . 873  
   in relapsing fever . . . 945  
   in remittent fever . . . 954

- Temperature in rheumatism . . . 893  
   in scarlatina . . . . . 50, 956  
   in smallpox . . . . . 989  
   in spinal injury . . . . . 51  
   in tetanus . . . . . 51, 229  
   in trichiniasis . . . . . 1060  
   in typhoid fever . . . . . 50, 914  
   in typhus fever . . . . . 931  
   in yellow fever . . . . . 50, 972  
   of surface . . . . . 45  
     epigastric . . . . . 579  
     over gall bladder . . . . . 602  
   of tetanus . . . . . 51  
 Tenderness as a symptom . . . . . 56  
 Tenesmus . . . . . 635, 639  
 Tetanus . . . . . 126, 172, 228  
   confounded with hydrophobia . . . . . 232  
   with local rigidity . . . . . 230  
   with muscular rheumatism . . . . . 230  
   with spasms in scarlet fever . . . . . 230  
   with strychnine poisoning, . . . . . 232, 1037  
   distinguished from cerebro-  
     spinal fever . . . . . 230, 941  
   from chorea . . . . . 222  
   hysterical . . . . . 229  
   idiopathic . . . . . 228  
   symptomatic . . . . . 230  
   traumatic . . . . . 228  
 Tetany . . . . . 206, 231, 591  
   associated with laryngismus . . . . . 279  
 Thermal impressions, paths of . . . . . 64  
   sense . . . . . 79, 157  
 Thermometer, clinical use of . . . . . 43  
   See also *Temperature*.  
 Thermometry, cerebral . . . . . 45, 123  
   general . . . . . 46  
   surface . . . . . 45  
 Thirst as a symptom . . . . . 556  
 Thomsen's disease . . . . . 234  
   gait in . . . . . 31  
 Thoracic aneurism . . . . . 506  
   confounded with abscess of the  
     mediastinum . . . . . 508  
   with chronic laryngitis . . . . . 514  
   with dilated auricle . . . . . 512  
   with insufficient aortic valves . . . . . 510  
   with intra-thoracic morbid  
     growth . . . . . 507  
   with malformation of the  
     chest . . . . . 513  
   with malposition of the aorta . . . . . 513  
   with pulsating empyema . . . . . 511  
   with pulsation of pulmonary  
     artery . . . . . 512  
   eructations in . . . . . 558  
 Throat, inflammation of . . . . . 522, 535  
   soreness of, in scarlet fever . . . . . 977  
   rheumatic . . . . . 536  
   syphilitic ulcers of . . . . . 536  
 Thrombosis . . . . . 877  
   cerebral . . . . . 201  
   from chlorosis . . . . . 879  
   from exhausting diseases . . . . . 879  
   of brain sinuses . . . . . 242, 878  
   of cerebral arteries . . . . . 200  
   of renal vein . . . . . 828  
 Thrombus, changes in . . . . . 884  
 Thrush . . . . . 520, 530  
 Thyroid gland . . . . . 251, 456  
   swelling of . . . . . 456  
 Tic douloureux . . . . . 73, 255  
 Tinea . . . . . 1020  
   circinata . . . . . 1020, 1022  
   decalvans . . . . . 1023  
   favosa . . . . . 1022  
   kerion . . . . . 1022  
   sycosis . . . . . 1022  
   tonsurans . . . . . 1020, 1022  
   versicolor . . . . . 1016, 1022  
 Tinnitus aurium . . . . . 98  
 Tobacco amblyopia . . . . . 96  
 Tongue, cancer of . . . . . 521  
   coating of . . . . . 54  
   condition of, in disease . . . . . 52, 521  
   dryness of . . . . . 53  
   inflammation of . . . . . 521  
   movements of . . . . . 52  
   syphilis of . . . . . 521  
 Tonsil, abscess of . . . . . 285  
   cancer of . . . . . 523, 536  
   enlargement of . . . . . 523  
   herpes of . . . . . 530  
   pharyngeal . . . . . 263  
 Tonsillitis . . . . . 275, 285, 522  
   acute follicular . . . . . 524  
   confounded with diphtheria . . . . . 529  
 Torticollis . . . . . 896  
 Toxines, absorption of . . . . . 877  
 Trachea, affections of . . . . . 266, 278  
   foreign body in . . . . . 285  
   morbid growths in . . . . . 295  
   narrowing of . . . . . 295  
   symptoms of diseases of . . . . . 267  
   ulcers in . . . . . 295  
 Tracheal tugging . . . . . 509  
 Trance distinguished from nar-  
   colepsy . . . . . 213  
 Tremor . . . . . 166  
   alcoholic . . . . . 169  
   arsenical . . . . . 170  
   asthenic . . . . . 170  
   convulsive . . . . . 222  
   essential . . . . . 170  
   functional . . . . . 169  
   hereditary . . . . . 171  
   hysterical . . . . . 170  
   in exophthalmic goitre . . . . . 170, 456  
   in spasmodic tabes . . . . . 169  
   lead . . . . . 170, 1042

- Tremor, mercurial . . . . 170, 1041  
 post-hemiplegic . . . . 169  
 senile . . . . 169  
 tobacco . . . . 170  
 Trial meal . . . . 552, 575, 583  
 Trichina spiralis . . . . 1057  
 Trichiniasis . . . . 897, 1059  
 distinguished from Bright's dis-  
 ease . . . . 1065  
 from cardiac disease . . . . 1065  
 from cholera morbus . . . . 1064  
 from irritant poisoning . . . . 1064  
 from periarteritis nodosa . . . . 1064  
 from rheumatism . . . . 897, 1063  
 from sausage poisoning . . . . 1064  
 from typhoid fever . . . . 1061  
 from typhus fever . . . . 1061  
 fever of . . . . 1060  
 muscular symptoms of . . . . 1062  
 cedema in . . . . 1061, 1065  
 pulmonary symptoms in . . . . 1063  
 Trichloracetic acid test . . . . 770  
 Trichocephalus dispar . . . . 1051  
 Tricuspid regurgitation . . . . 502, 602  
 Trismus . . . . 229, 941  
 Tube-casts in the urine . . . . 809, 946  
 Tubercle . . . . 343, 364, 372  
 associated with scrofula . . . . 383  
 bacilli . . . . 342  
 tests for . . . . 342  
 calcareous transformation of . . . . 382  
 in brain . . . . 245  
 in intestines . . . . 387, 646  
 Tubercular meningitis . . . . 185, 942  
 pleurisy . . . . 374  
 Tuberculosis of kidney . . . . 804  
 distinctive signs of . . . . 804  
 Tuberculosis of lungs . . . . 344, 385  
 See also *Phthisis*.  
 acute miliary . . . . 385, 928  
 bacillus of . . . . 342, 368  
 combined with laryngitis . . . . 292  
 Tuberculous inspiration broncho-  
 pneumonia . . . . 398  
 Tumors, abdominal . . . . 718  
 confounded with colic . . . . 606  
 in hypochondria . . . . 718  
 aneurismal . . . . 333, 357  
 dyspnœa in . . . . 333  
 cerebral . . . . 240  
 distinguished from abscess . . . . 241  
 from apoplexy . . . . 199  
 from chronic meningitis . . . . 242  
 from softening . . . . 241  
 from thrombosis of sinuses . . . . 242  
 nature of . . . . 245  
 seat of . . . . 243  
 syphilitic . . . . 245  
 unilateral symptoms of . . . . 244  
 in hypogastric region . . . . 729  
 Tumors in iliac and lumbar re-  
 gions . . . . 727  
 in umbilical region . . . . 724  
 intra-thoracic, confounded  
 with chronic pleurisy . . . . 419  
 mediastinal . . . . 420, 508  
 non-aneurismal, confounded  
 with abdominal aneurism . . . . 734  
 with thoracic aneurism . . . . 509  
 of epigastrium . . . . 721  
 of larynx . . . . 293  
 of liver . . . . 697  
 of spinal cord . . . . 136  
 of spleen . . . . 719  
 ovarian . . . . 728  
 phantom . . . . 723  
 retro-peritoneal . . . . 715  
 tuberculous . . . . 241  
 Tympanites . . . . 636, 717  
 chronic, confounded with  
 ascites . . . . 716, 717  
 Typhlitis . . . . 625  
 Typhoid conditions confounded  
 with typhoid fever . . . . 925  
 Typhoid fever . . . . 913  
 abortive . . . . 672, 928  
 bacillus of . . . . 877, 917  
 colic in . . . . 600  
 confounded with acute atrophy  
 of liver . . . . 678  
 with appendicitis . . . . 626  
 with cerebro-spinal fever . . . . 940  
 with enteritis . . . . 925  
 with general debility . . . . 924  
 with meningitis . . . . 188, 927  
 with peritonitis . . . . 925  
 with pulmonary affections . . . . 927  
 with pyæmia . . . . 873  
 with relapsing fever . . . . 947  
 with remittent fever . . . . 957  
 with scarlet fever . . . . 980  
 with trichiniasis . . . . 1061  
 with typhoid conditions . . . . 925  
 with typhus fever . . . . 934  
 with ulcerative endocarditis,  
 467, 927  
 convulsions in . . . . 920  
 delirium in . . . . 920  
 diarrhœa in . . . . 915  
 distinguished from yellow fever . . . . 973  
 enlargement of the spleen in . . . . 918  
 epistaxis in . . . . 921  
 eruption in . . . . 921  
 febrile symptoms of . . . . 914  
 mild form of . . . . 928  
 nervous symptoms in . . . . 919  
 pain a symptom in . . . . 918  
 palsy in . . . . 921  
 relapses in . . . . 922, 929  
 septicæmic . . . . 877

- Typhoid fever, sequelæ of . . . 923  
   spinal symptoms in . . . 921  
   temperature in . . . 914  
   ulcers of . . . 647  
   urine of . . . 915  
   walking . . . 924  
 Typhoid septicæmia . . . 876  
 Typho-malarial fever . . . 957, 968  
 Typhus fever . . . 929, 988  
   acute tubercular deposits in . . . 934  
   cerebral symptoms in . . . 931  
   cerebro-spinal . . . 937  
   coma-vigil of . . . 931  
   compared with typhoid fever . . . 934  
   complications in . . . 934  
   confounded with measles . . . 983  
   with yellow fever . . . 974  
   distinguished from acute men-  
     ingitis . . . 932  
   from cerebro-spinal fever . . . 943  
   from plague . . . 936  
   from relapsing fever . . . 947  
   from rubella . . . 988  
   eruption in . . . 930  
   exanthematic . . . 930  
   gastric hemorrhage in . . . 564  
   maculated or spotted . . . 930  
   physiognomy of . . . 930  
   pulse in . . . 933  
   temperature in . . . 931  
   urine in . . . 933  
 Tyrosine . . . 679, 745, 759
- U.**
- Ulcer of duodenum . . . 581  
   gastric . . . 346, 576, 590  
   confounded with chronic  
     gastritis . . . 576, 585  
   with gastric cancer . . . 582, 585  
   with ulcer of duodenum . . . 581  
   laryngeal . . . 293  
   of the bowel, follicular . . . 646  
   syphilitic . . . 638  
   of the stomach . . . 577, 588  
   of typhoid, unhealed . . . 647  
   peptic . . . 581  
   perforating, of the foot, 162,  
     175, 1049  
   stercoral . . . 643  
 Umbilical region, tumors of . . . 724  
 Uræmia . . . 796, 1032  
   associated with cholera . . . 660  
   convulsions in . . . 798  
   delirium in . . . 68, 798  
   distinguished from cerebo-  
     spinal fever . . . 943  
   mania in, acute . . . 798  
 Uræmic coma distinguished from  
   apoplexy . . . 203  
 Uræmic coma distinguished from  
   narcotic poisoning . . . 797, 1032  
 Urates, pathology of . . . 750  
   tests for . . . 750  
 Urea, pathology of . . . 744  
   table for estimation of . . . 748  
   tests for . . . 745  
 Ureometer . . . 746  
 Ureter, inflammation and ulcera-  
   tion of . . . 786  
 Urethra, hemorrhage from . . . 778  
 Urethral fever confounded with  
   intermittent fever . . . 952  
 Uric acid in gout . . . 898  
   detection of . . . 898  
   in lithæmia . . . 864  
   murexide test for . . . 749  
   pathology of . . . 747  
 Urinary organs, diseases of . . . 736, 784  
 Urine . . . 736  
   abnormal substances in . . . 757  
   acetone in . . . 766, 824  
   acidity of . . . 742  
   albumin and other proteids in . . . 767  
   principal tests for . . . 768  
   albuminous condition of, dis-  
     eases marked by . . . 791, 811  
   alkalinity of . . . 743, 816  
   analysis of . . . 736  
   bile in . . . 760  
   biliary acids in . . . 761  
   blood in . . . 672, 772  
   guaiacum test for . . . 772  
   blood-corpuscles in . . . 774  
   blood-extractives in . . . 767  
   calcium oxalate in, 575, 757, 776, 786  
   casts, mucous, in . . . 809  
   chlorides in . . . 390, 590, 755  
   chylous . . . 781  
   color of, changes in . . . 740  
   constituents of, changes in  
     quantity of . . . 744  
   cystine in . . . 786  
   diacetic acid in . . . 766  
   diazo-reaction of . . . 771  
   estimate of solids in . . . 741  
   fat in . . . 780  
   fibrin in . . . 781  
   hæmoglobin in . . . 664, 772, 777  
   in Bright's disease . . . 791  
   in carbolic acid poisoning . . . 1034  
   increased discharge of . . . 823  
   in diabetes . . . 823  
   indican in . . . 623, 639  
   in gastric disease . . . 590  
   ingredients of . . . 738  
   in hooping-cough . . . 337  
   in jaundice . . . 664, 760  
   inosite in . . . 766  
   kreatin and kreatinin in . . . 757



Urine, kyestein pellicle on . . .	781
lactic acid in . . . . .	904
leucine in . . . . .	679, 745, 759
mucin or nucleo-albumin in, . . . . .	767, 771
nitrogen in . . . . .	746
normal . . . . .	737, 767
of the insane . . . . .	826
oxalate of lime in . . . . .	575, 757, 786
paraglobulin in . . . . .	770
peptones in . . . . .	767, 771
pigment in . . . . .	740, 933
phosphates in . . . . .	575, 752, 786, 816
alkaline and earthy . . . . .	752, 754
purulent, confounded with . . . . .	
acute Bright's disease . . . . .	794
diseases associated with . . . . .	814
pus in . . . . .	779, 814, 816
quantitative examination of . . . . .	745
reaction of . . . . .	742
retention of . . . . .	829
sediments in . . . . .	782
specific gravity of . . . . .	740
sugar in . . . . .	704, 762, 823
sulphates in . . . . .	756
suppression of . . . . .	822, 828
table showing action of tests . . . . .	
upon . . . . .	782
tyrosine in . . . . .	679, 745, 759
urates in . . . . .	750, 786, 956
urea in . . . . .	744, 776, 933
uric acid in . . . . .	747, 786
xanthine in . . . . .	786
Urinometer . . . . .	741
Urobilin . . . . .	705, 740, 857
pathological . . . . .	740
Urochrome . . . . .	740
Uroerythrine . . . . .	740
Urticaria, pigmentosa . . . . .	1004
Uterus, colic of . . . . .	604
gravid, confounded with ascites . . . . .	716
hemorrhage from, in myxo-	
dema . . . . .	834
tumor of . . . . .	628

V.

Vagrants' disease . . . . .	871
Valve, aortic, disease of . . . . .	497
insufficiency of . . . . .	510, 602
mitral, disease of . . . . .	503
sigmoid, rupture of . . . . .	465
tricuspid, affections of . . . . .	496, 602
Valvular affections of the heart . . . . .	489
confounded with functional car-	
diac disorder . . . . .	491
with malformations of heart . . . . .	490
with misdirection of current . . . . .	492
diagnosis of, before develop-	
ment of murmur . . . . .	504

Valvular affections of the heart . . . . .	
from rupture of a valvule . . . . .	
of a papillary muscle . . . . .	504
table of . . . . .	499
Varicella . . . . .	993
followed by gangrene . . . . .	994
Varicocele . . . . .	804
Variola. See <i>Smallpox</i> . . . . .	988
Varioloid . . . . .	993
Veins, diseases of . . . . .	839
enlargement of . . . . .	704
portal, inflammation of . . . . .	669, 708
thrombosis of . . . . .	709, 878
renal, thrombosis of . . . . .	828
Vena cava, occlusion of . . . . .	517
Venous enlargement . . . . .	517
hum . . . . .	338
pulsation . . . . .	43
Vertigo . . . . .	83, 223
aural . . . . .	85
cerebral . . . . .	84, 86, 241
essential . . . . .	86
from overwork of brain . . . . .	86
laryngeal . . . . .	86, 280
lithæmic . . . . .	85
of malassimilation . . . . .	85
precursor of epilepsy . . . . .	86
stomachal . . . . .	84
syphilitic . . . . .	142
Vesicular murmur, absence of . . . . .	316
changes in . . . . .	314
Vessels, amyloid degeneration of . . . . .	590
Vibrio proteus . . . . .	655
Viscera, abdominal percussion . . . . .	
and auscultation of . . . . .	544, 550
Vision, derangement of . . . . .	87, 900
double . . . . .	91
Vocal cord, cysts of . . . . .	294
diseases of . . . . .	290, 291
extirpation of . . . . .	291
fremitus, absence of . . . . .	326
paralysis of . . . . .	333
resonance . . . . .	325
Voice, altered . . . . .	266, 291
amphoric . . . . .	326
auscultation of . . . . .	325
cavernous . . . . .	325
changes in, in laryngeal dis-	
eases . . . . .	266
loss of . . . . .	289
metallic . . . . .	290
strain of . . . . .	287, 289
whispering . . . . .	290
without vocal cords . . . . .	291
Vomit, black . . . . .	563, 612, 971
coffee-ground . . . . .	564, 583
different forms of . . . . .	560
sarcinæ and fungi in . . . . .	561
Vomiting as a symptom . . . . .	558
disease accompanied by . . . . .	571, 653

Vomiting, faecal . . . . .	562, 631
gastric . . . . .	580
in brain diseases . . . . .	580
nervous . . . . .	559
of bile . . . . .	562, 655
of blood . . . . .	563, 579
in irritant poisoning . . . . .	1028
of food or liquid . . . . .	560

**W.**

Water-brash . . . . .	561
Weil's disease . . . . .	671
bacillus of . . . . .	672
bloody urine in . . . . .	672
Whitlows, painless . . . . .	153
Windpipe, foreign bodies in . . . . .	285
Wool-sorters' disease . . . . .	1046
Word-blindness . . . . .	61, 208
Word-deafness . . . . .	61, 208
Worms, intestinal . . . . .	137, 855, 1050
Wrist clonus . . . . .	103
Writer's cramp . . . . .	223
Wry-neck . . . . .	896

**X.**

Xanthelasma . . . . .	698, 1001
Xanthine . . . . .	786

**Y.**

Yellow fever . . . . .	947, 969
confounded with Asiatic chol-	
era . . . . .	973
with atrophy of liver . . . . .	678
with bilious remittent fever . . . . .	974
with dengue . . . . .	996
with hemorrhagic malarial	
fever . . . . .	966
with plague . . . . .	973
with relapsing fever . . . . .	947, 973
with typhoid fever . . . . .	973
with typhus fever . . . . .	974
examination of urine in . . . . .	974
walking . . . . .	972

**Z.**

Zinc poisoning . . . . .	1030, 1044
--------------------------	------------

THE END.















NATIONAL LIBRARY OF MEDICINE



NLM 00105585 0